


주요 *in vivo* 정보와 이를 활용한 인간 약동학 예측 이론

So Jin Lee, PharmD
Researcher/Team leader

Q-fitter

Dept. of Clinical Pharmacology & Therapeutics /
The Catholic Univ. of Korea Seoul St.Mary's Hospital




Contents

1. Dallphin 에서 사용하는 in vivo 정보
2. In vivo 정보를 통한 PK parameter 예측
 - Basic info – “fraction unbound”
 - Distribution – “allometry”
 - Elimination – “human CL”
3. Allometry
 - General allometric approaches (V, CL)
 - Dallphin allometric approach (V)
4. Examples
 - Various allometries (V, CL)
 - Dallphin CL vs. allometry CL predictions



What is Dallphin-AtoM

- Drugs with
- **ALL**ometry and
- **PH**ysiology
- **IN**side
- **Animal to**
- HuMan



PBPK software for human
PK parameters prediction

using physicochemical
properties, in vitro and
animal PK data



Dallphin: Inputs & Outputs

Input

In vitro data

LogP
Cb/Cp
fu
pKa
Papp
CLmet

Animal in vivo PK

CLh
Vc
Vp
CLr
Q

$$CL_{total} = CLh + CLr$$



Output

Human PK parameter

Clearance (CL)
Central volume (Vc)
Peripheral volume (Vp)
Intercompartment
clearance (Q)
Absorption rate constant
(Ka)
Bioavailability (F)

Vss




Major PK parameters Dallphin predicts

- [Absorption] K_a
- [Distribution] V_d (V_{ss} , V_c , V_p), and Q
- [Elimination] CL
- [Absorption] F (if drug follows first-order kinetics)

In vivo data provides...

- ✓ Allometric methods to predict V_d , Q
- ✓ One of the methods to predict CL_h & F_h (ex vivo data)
- ✓ Two methods to predict CL_r




Dalphin methods: Clearance & F

$$CL_{hepatic} = \frac{Q_H \cdot fu \cdot CL_{int}}{Q_H + fu \cdot CL_{int}}$$

CL total = CL hepatic + CL renal

Microsome

1. MPPGL
2. Rat IVIV ratio
3. PS
4. PS(logD)

Hepatocyte

1. Human hepatocyte

1. CL renal_rat
2. CL renal_dog
3. CL renal_user input

$$CL_{r_{Human}} = CL_{r_{Species}} \times \frac{fu_{Human}}{fu_{Species}} \times \frac{KBF_{Human}}{KBF_{Species}}$$

$$Fh = 1 - \frac{CL_{hepatic}}{Q_H}$$

F total = Fh x Fa

Microsome

1. MPPGL (CL_{int})
2. Rat IVIV ratio
3. PS
4. PS(logD)

Hepatocyte

1. Human hepatocyte

1. Fa_Caco
2. Fa_PSA

CL_{int}: intrinsic metabolic or hepatic clearance
 Fh: hepatic availability
 Fa: fraction absorbed from intestinal tract
 PS: permeability surface area coefficient
 Q_H: hepatic blood flow
 PSA: dynamic polar molecular surface area



Dallphin methods: Vd & Q


Allometric approaches

1. Vss: 1-compartment model
2. Vss, Vc, Vp, Q: 2-compartment model


PBPK approach^{12,13}

1. Vss: 1-compartment model (Vc:well-perfused organs)

PBPK approach




$$V_{ss} = V_p + \sum (V_t \times K_{pt})$$



1. Dallphin 에서 사용하는 in vivo 정보

- [Basic info] Plasma protein binding (ex-vivo data)
 - Rat fraction unbound \Rightarrow human hepatic CL(2), hepatic availability(2), human renal CL
 - Dog fraction unbound \Rightarrow human renal CL
 - Human fraction unbound \Rightarrow human hepatic CL (1)–(4), hepatic availability (1)–(4)
 - Blood to plasma ratio
- [Distribution] Allometric approaches: Animal IV PK parameters
 - Weight, Q, Volume of distributions (V_{ss} , V_c , V_p)
- [Elimination] Microsome:
 - Rat $CL_{int, initial}$, rat hepatic CL (observed) \Rightarrow human hepatic CL(2), hepatic availability(2)



2. Input parameter for human PK prediction^{1,2}

- [Basic info] Plasma protein binding
- fu_Rat → CL int, rat(in vivo observed) → CL int_rat_iviv_ratio
 ⇒ human hepatic CL $CL_{h_fromRativivc}$
 ⇒ human bioavailability $F(CL_{h_fromRativivc})$
 ⇒ human renal CL CL_r (rat to human prediction)

$$CL_{h_fromRativivc} = \frac{Q_{H,B} \cdot fu_B \cdot CL_{u_{int,H}}}{Q_{H,B} + fu_B \cdot CL_{u_{int,H}}} \times \frac{CL_{int_rat, IVIV\ ratio}}{CL_{int_rat, IVIV\ ratio}}$$

- fu_Dog ⇒ human renal CL CL_r (dog to human prediction)

$$CL_{r_{Human}} = CL_{r_{Species}} \times \frac{fu_{Human}}{fu_{Species}} \times \frac{KBF_{Human}}{KBF_{Species}}$$



[Basic info]

- Plasma protein binding
 - fu_Rat
 - fu_Dog

Dallphin-AtoM(ver 0.8.5)

Basic Info | Absorption | Distribution | Elimination | Final Parameters | Concentration Prediction | About

Physicochemical Properties

Neutral

Log P

pKa

Log D7.4

☒ PSA Å²

Plasma Protein Binding

☒ fu_Rat

☒ fu_Dog

fu_Hum

Cb/Cp

CL renal²

- Three animal scaling techniques for the prediction of human CL renal:

1. **Direct correlations** between CLr in man and each of the two main preclinical species (rat and dog)

$$CLr_{Human} = CLr_{Species} \times \frac{fu_{Human}}{fu_{Species}} \times \frac{KBF_{Human}}{KBF_{Species}}$$

2. Simple allometry $CLr = a(W)^b$

$$CLr_{Predicted\ total} = CLr_{Predicted\ unbound} \times fu_{Human}$$

3. Mahmood's renal clearance scaling method

$$SSF = \frac{(\text{Glomerular filtration} \times \text{kidney blood flow})}{(\text{body weight} \times \text{kidney weight})}$$

$$\text{Correction factor} = \frac{SSF_{Species}}{SSF_{Human}}$$



CL renal²




FIG. 2. Measured human renal clearance versus measured renal clearance from rat (a), dog (b), rat corrected (c) for fu differences (c), dog corrected for fu differences (d), rat corrected for both KBF and fu differences (e), and dog corrected for both KBF and fu differences (f). Circles, acids; squares, bases; triangles, neutrals; diamonds, zwitterions.

- Most accurate prediction: 1. Direct correlation with the dog renal clearance with PPB and KBF corrections ($R^2 = 0.84$)



2. Input parameter for human PK prediction^{1,2}

- [Elimination] Microsome

- CL_{int,ini Rat} → CL_{int,rat} (in vitro)
 - CL_{h,ratinvivo} → CL_{int,rat} (in vivo)
- } ⇒ CL_{int,rat,iviv ratio}
 ⇒ human hepatic CL CL_{h,fromRativivc}
 ⇒ human bioavailability F_h (CL_{h,fromRativivc})

$$CL_{h,fromRativivc} = \frac{Q_{H,B} \cdot fu_B \cdot \boxed{CL_{u,int,H}} \times \boxed{CL_{int,rat,IVIV ratio}}}{Q_{H,B} + fu_B \cdot \boxed{CL_{u,int,H}} \times \boxed{CL_{int,rat,IVIV ratio}}}$$

$$CL_{int,rat,IVIV ratio} = \frac{\boxed{CL_{int,rat} (in vivo observed)}}{\boxed{CL_{int,rat} (in vitro predicted)}} < - CL_{h,rat} (in vivo observed)$$

$$\boxed{CL_{h,rat} (in vivo observed)} = \frac{Q_{H, rat} \cdot f_{ura_t} \cdot \boxed{CL_{int, rat}}}{Q_{H, rat} + f_{ur_{at}} \cdot \boxed{CL_{int, rat}}}$$

$$\boxed{CL_{int, rat}} = \boxed{CL_{int ini rat}} \cdot MPPGL_{rat} \cdot Liver wt_{rat}$$



[Elimination]

- Hepatic Clearance (CLh)
Microsome
 - CL int_ini Rat (ul/min/mg protein)
 - CLh_ratinvivo (ml/min/kg)

- Renal Clearance (CLr)
 - CLr_Rat (ml/min/kg)
 - CLr_Dog (ml/min/kg)

Dallphin-AtoM(ver 0.8.5)

Basic Info | Absorption | Distribution | Elimination | Final Parameters | Concentration Prediction | About

Hepatic Clearance(CLh)

☒ Microsome

CL int_ini : ☒ Rat ul/min/mg protein CLh_ratinvivo ml/min/kg

Human ul/min/mg protein

microsome protein concentration g/mL

fu_mic

CLh L/hr **CLh_fromRativivc L/hr** CLh_PS L/hr

☐ Hepatocyte

CL int_ini : Human ul/min/10⁶ cell

V_{cell} / V_{incubation}

fu_hep

CLh L/hr

Renal Clearance(CLr)

☒ CLr_Rat ml/min/kg

☒ CLr_Dog ml/min/kg

CLr (Rat to human prediction) L/hr

CLr (Dog to human prediction) L/hr



2. Input parameter for human PK prediction [Distribution]

- Mouse, rat, monkey, dog 총 4종의 정보 입력 가능
- 각 종 별 Weight
- Model structure – 1 compartment vs. 2 compartment
- Model 을 통해 구한 각 종 별 V_{ss} 또는 Q , V_{ss} (V_c , V_p)
- [Prediction] 을 통해 Allometric scaling 수행 =>
70 kg-human Q , V_c , V_p , V_{ss} 구함

- $V_c \rightarrow K_a$
$$k_{a,eq} = \frac{P_m S}{V_c}$$

P_m : drug permeability across intestinal mucosa (Caco-2 Papp)
 S : absorptive surface area



[Distribution]

- Animal data available ? YES
- One-compartment model
- Two-compartment model

Dalphin-AtoMVer 0.8.4

Basic Info | Absorption | Distribution | Elimination | Final Parameters | Concentration Prediction | About

Animal (in vivo) I.V. PK data in ≥ 2 species available? ☒ Yes ☐ No

Allometric approach



Enter distribution data in animals Two-Compartment Model

Species	Weight	Q	Vc	Vp	Vss
<input checked="" type="checkbox"/> Mouse	20 g	0.8 L/hr/kg	0.5 L/kg	1.0 L/kg	1.5 L/kg
<input checked="" type="checkbox"/> Rat	200 g	0.8 L/hr/kg	0.5 L/kg	1.0 L/kg	1.5 L/kg
<input checked="" type="checkbox"/> Monkey	3.0 kg	0.8 L/hr/kg	0.5 L/kg	1.0 L/kg	1.5 L/kg
<input checked="" type="checkbox"/> Dog	10.0 kg	0.8 L/hr/kg	0.5 L/kg	1.0 L/kg	1.5 L/kg

Prediction

Allometric Scaling Result

Human WT: 70kg Q: 0.47 L/hr/kg VC: 0.5 L/kg VP: 1.0 L/kg Vss: 1.5 L/kg



Dalphin-AtoMVer 0.8.4

Basic Info | Absorption | Distribution | Elimination | Final Parameters | Concentration Prediction | About

Animal (in vivo) I.V. PK data in ≥ 2 species available? ☒ Yes ☐ No

Allometric approach


Enter distribution data in animals One-Compartment Model

Species	Weight	Vss
<input checked="" type="checkbox"/> Mouse	20 g 0.5 L/kg	
<input checked="" type="checkbox"/> Rat	200 g 0.5 L/kg	
<input checked="" type="checkbox"/> Monkey	3.0 kg 0.5 L/kg	
<input checked="" type="checkbox"/> Dog	10.0 kg 0.5 L/kg	

Prediction

Allometric Scaling Result

Human WT: 70kg Vss: 0.5 L/kg



[Prediction]

- Allometric scaling result

Volume of distribution

- Allometric scaling as choice of method to predict human Vd
- Plasma protein binding should be considered³
- Multiple methods available for prediction of volume of distribution^{3,4,6}
 - ✓ Simple allometric scaling
 - ✓ Allometry with correction factors
 - ✓ Average fraction unbound tissue
 - ✓ Proportionality (dog-human)
 - ✓ QSAR



Volume of distribution^{3,4}

- Simple allometric scaling
- Allometry with correction factors*
- Average fraction unbound tissue*
- Proportionality (dog-human)*
- QSAR

*Pertains plasma protein binding data as an essential element

$$\log_{10} VD = a \cdot \log_{10} \text{body weight}_{(\text{kg})} + b$$

$$VD_{\text{free}} = \frac{VD_{\text{total}}}{f_u}$$

Table 4. Methods for prediction of volume of distribution (V_d).

Method	Equation	Comment *	Ref.
Simple AS	$V = a(W)^b$	(93) The prediction of V_d is well predicted equally with using two species in AS	[108]
Average fraction unbound in tissue ¹	$V = V_{\text{Plasma}}(1 + R_{E/I}) + f_u \cdot V_P \left(\frac{V_E}{V_P} - \frac{V_R \cdot f_u}{\alpha_R} \right)$	(94) It is useful to analyze and predict an alteration in apparent V_d then identify the cause of alteration. It is particularly useful for drugs with low V_d (<15 L or 0.2 L/kg)	[119]
Proportionality	$V_{\text{human,pred}} = \frac{V_{\text{animal}} \cdot f_{u,\text{human}}}{f_{u,\text{animal}}}$	(95) It is assumed that the volume of distribution at a steady state of free drug is identical between species	[120]
One species AS	$V_{\text{human,pred}} = -0.35V_{\text{rat}}^{0.91}$	(96) Statistical modeling is applied in this model	[121]
QSAR	$\begin{aligned} \log(V_{d_{ss,\text{human}}}) = & 0.1859 \\ & \cdot \log(V_{d_{ss,\text{rat}}}) \times \log(V_{d_{ss,\text{rat}}}) \\ & - 0.3887 \\ & \cdot \log(V_{d_{ss,\text{rat}}}) \times \log(\text{MW}) \\ & + 0.3089 \\ & \cdot \log(V_{d_{ss,\text{dog}}}) \times \log(\text{MW}) \\ & + 0.003306 \cdot \log(\text{MW}) \times c \log P \\ & + 1.71 \end{aligned}$	(97) $V_{d_{ss, \text{human}}}$ (mL/kg) is predicted by QSAR modeling with quadratic term descriptors	[122]

* Each comment corresponds to all the equations within each major section of the table defined by horizontal lines.


¹ Where V_d is apparent volume of distribution, V_{plasma} is plasma volume, V_E is extracellular space minus the plasma, V_R is physical volume into which the drug distributes minus the extracellular space, f_u is the fraction unbound in plasma, and $R_{E/I}$ is the ratio of distributed albumin in the extravascular space to that in the intravascular space. It is 1.4. α_R equals to C_u/C_R where C_u is unbound drug concentration at distribution equilibrium and C_R is concentration in V_R .

3. Allometry^{5,6}

- “Study of size and its consequences”
- Based on the assumption that there are anatomical, physiological, and biochemical similarities among animals which can be described by simple mathematical models
- Choice of method for projection to human PK
- Possible with in vivo IV PK data from 2 species, but ≥ 3 species provide more accurate prediction
- Plasma protein binding data in two or more species and human should be emphasized
- Completed after each species modeling to predict human PK parameters, which is used for simulations
- Different allometric approaches available

For which PK parameter prediction?

- Human Vd
- Human CL



General allometric approaches^{3,4,5}

- Simple allometry $Y = a(BW)^b$ $\log Y = \log(a) + b \cdot \log(BW)$
- Allometry with standard correction factors for Vd – PPB (plasma protein binding)

$$\log(VD) = \log(a) + b \cdot \log(BW)$$

$$VD_{\text{free}} = \frac{VD_{\text{total}}}{f_u}$$

- Allometry with standard correction factors for CL – MPL (product of maximum life-span), brain weight

$$\text{Clearance} = aBW^b$$

$$\text{Clearance} \times \text{MPL} = aBW^b$$

$$\text{Clearance} \times \text{BrainWeight} = aBW^b$$

cf) Correction factors for renally/biliary excreted drugs³ – GFR/bile flow, UDPGT

$$\text{Clearance} / \text{GFR} = aBW^b$$

$$\text{Cl} \times \text{Bile_flow} = aBW^b$$

$$\text{Cl} \times \text{UDPGT} = aBW^b$$



Different allometric approaches for CL¹⁰

- Simple allometry
- Rule of exponents
- Correction factor – MLP, brain weight
- Two-term method
- Multiexponential
- Normalized



Different allometric approaches for CL^{4,11}

Table 3. Methods for prediction of clearance (CL) using allometric scaling (AS).

Method	Equation	Comments *	Ref.
Simple AS	$CL = a(W)^b$ (74)	Select a proper equation by the rule of exponent (ROE)	-
AS with MLP ¹	$CL \cdot MLP = a(W)^b$ (75)	W and BW represent body and brain weight, respectively	-
AS with BW	$CL \cdot BW = a(W)^b$ (76)		[89]
Rule of exponent	If the exponent is 0.55 to 0.7, then use the simple AS, Equation (74) If the exponent is 0.71 to 1, then use the MLP, Equation (75) If the exponent is more than 1, then use the BW, Equation (76)		[90]
Two-term method	$CL = \theta(W)^a \cdot (BW)^b$ (77)	θ is a constant, which is determined by multiple regression analysis	[91]
Multieponential	$CL_{human} = aW^b + \left[\left(\frac{1 - \frac{3}{2}b}{1 - \frac{1}{2}b} \right) aW^{0.9} \right]$ (78)	The unit of CL is mL/min	[92]
Normalized AS	$CL_{animal} \frac{CL_{int,human}}{CL_{int,animal}} = a(W)^b$ (79)	CL_{int} refers the unbound CL_{int} in microsomes or hepatocytes in species and humans	[93]

Method	Equation	Comments *	Ref.
One species AS	$CL_{human} = CL_{animal} \cdot \left(\frac{W_{human}}{W_{animal}} \right)^b$ (80)		The exponent b is a constant 0.75, which is physiologically relevant value (eg, blood flow, filtration, etc) [94,95]
One species AS	$CL_{pred} = 0.152 \cdot CL_{rat} \cdot \left(\frac{W_{human}}{W_{rat}} \right)$ (81)		
One species AS	$CL_{pred} = 0.41 \cdot CL_{dog} \cdot \left(\frac{W_{human}}{W_{dog}} \right)$ (82)		Predict the CL of bound drug
One species AS	$CL_{pred} = 0.407 \cdot CL_{monkey} \cdot \left(\frac{W_{human}}{W_{monkey}} \right)$ (83)		[90]
Two species AS	$CL_{pred} = a_{rat-dog} \cdot W_{human}^{0.628}$ (84)		Predict the CL of bound drug
Two species AS	$CL_{pred} = a_{rat-monkey} \cdot W_{human}^{0.650}$ (85)		
Hepatic method	liver $CL_{pred} = CL_{animal} \cdot \left(\frac{Q_{H,human}}{Q_{H,animal}} \right)$ (86)		[96]
FCIM ²	$CL = 33.35 \times \left(\frac{a}{Rf_u} \right)^{0.77}$ (87)		Rf_u is the f_u ratio between rats and humans and a is the coefficient form AS The unit of CL is mL/min [97]
QSAR ³	$\log CL_{pred} = 0.433 \cdot \log(CL_{rat}) + 1.0 \cdot \log(CL_{dog}) - 0.00627 \cdot MW + 0.189 \cdot Ha - 0.00111 \cdot \log(CL_{dog}) \cdot MW + 0.0000144 \cdot MW^2 - 0.0004 \cdot MW \cdot Ha - 0.707$ (88)		The unit of ([1] i

* Each comment corresponds to all the equations within each major section of the table defining horizontal lines.

¹ The maximum therapeutic potential (MLP) is calculated by the equation: $MLP = \frac{100 \cdot 0.001 \cdot \text{mg} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}}{CL}$

² Fraction unbound intrapart correction method

³ Quantitative structure activity relationship (QSAR) consist of physicochemical properties, such as molecular weight (MW), partition coefficient (LogP), and number of hydrogen-bond donors (HBD).

(98) The unit of observed and predicted oral CL value is mL/min/kg.



2019 KSCPT Spring Meeting Pre-course Workshop

Theories and Practices for Human Pharmacokinetics Prediction Using

Simple allometry⁵


- $Y = a(BW)^b$
- Log-log transformation
- Dallphin's choice for V_d (exponents fixed)
- Easily done with Excel linear regression
- When a midpoint species (e.g. dog) is the source of the error, the change is primarily in the intercept rather than the slope
- Fixed exponents...

Simple allometry: the log-log transformation



The scaling exponents


- $Y = a(BW)^b$
- Scaling exponent (b) (i.e. the slope) defines the type of scaling relationship




Allometry with correction factors for CL

- Simple allometry vs. corrected allometry
- The rule of exponents provides an idea of which standard correction factor to select^{7,8}

Simple vs. corrected allometry (MLP): Cefazolin clearance in man



Simple allometry
Predicted: 141 mL/min
Actual: 61 mL/min
Error: 131%



Allometry with MLP as a correcting factor
Predicted: 50.55mL/min
Actual: 61mL/min
Error:17.1%

The rule of exponents to predict human CL

1. $0.55 \leq b < 0.71$: no correction factor is necessary
2. $0.71 \leq b < 1.00$ **MLP** should be incorporated into scaling method
3. $B > 1.00$ **Brain weight** should be incorporated into the scaling method


Methods	% Mean absolute error (MAE)
Simple allometry	106
CL x MLP	40
CL x brain Weight	49
Rule of exponents	25

How to use Dallphin allometric approach?

- 각 종별 PK modeling*을 선행적으로 수행해야함
- Model structures: 기본 1-comp or 2-compartment model
- 모든 종에서 동일한 model structure 를 사용

*PK modeling: General workflow


1. Model structure 정하기
2. Model optimization
3. Model diagnosis – GOF, individual plots
4. Model validation – VPC , (bootstrap for human data)
5. PK parameter estimation & extrapolation



Dallphin allometric approach (Vd, Q)

- For 2 compartment model:
 $V_{ss} = V_c + V_p$, Q
- Exponents fixed to 1 and 0.75
 $V = a(WT)^1$
 $Q = a'(WT)^{0.75}$

Two-compartment open model, IV injection.



Dallphin-AtoM(ver 0.8.4)

Basic Info | Absorption | Distribution | Elimination | Final Parameters | Concentration Prediction | About

Animal (in vivo) I.V. PK data in ≥ 2 species available? ☒ Yes ☐ No

Allometric approach



Enter distribution data in animals Two-Compartment Model

Species	Weight	Q	Vc	Vp	Vss
<input checked="" type="checkbox"/> Mouse	20 g	0.8 L/hr/kg	0.5 L/kg	1.0 L/kg	1.5 L/kg
<input checked="" type="checkbox"/> Rat	200 g	0.8 L/hr/kg	0.5 L/kg	1.0 L/kg	1.5 L/kg
<input checked="" type="checkbox"/> Monkey	3.0 kg	0.8 L/hr/kg	0.8 L/kg	0.5 L/kg	1.3 L/kg
<input checked="" type="checkbox"/> Dog	10.0 kg	0.6 L/hr/kg	1 L/kg	1.0 L/kg	2.0 L/kg

Prediction

Allometric Scaling Result

Human WT : 70kg Q : 0.37 L/hr/kg VC : 0.98 L/kg VP : 0.96 L/kg Vss : 1.94 L/kg



Dallphin allometric approach (Vd, Q)

- For 1 compartment model:

Vss only (Q, Vc, Vp)

- Exponents fixed to 1

$$V = a(WT)^1$$

$$Q = a'(WT)^{0.75}$$

One-compartment open model, IV injection.



Dallphin-AtoM(ver 0.8.4)

Basic Info Absorption Distribution Elimination Final Parameters Concentration Prediction About

Animal (in vivo) I.V. PK data in ≥ 2 species available? ☒ Yes ☐ No

Allometric approach


Enter distribution data in animals One-Compartment Model

Species	Weight	Vss
<input checked="" type="checkbox"/> Mouse	20 g	0.5 L/kg
<input checked="" type="checkbox"/> Rat	200 g	0.5 L/kg
<input checked="" type="checkbox"/> Monkey	3.0 kg	0.5 L/kg
<input checked="" type="checkbox"/> Dog	10.0 kg	0.5 L/kg

Prediction

Allometric Scaling Result

Human WT : 70kg Vss : 0.5 L/kg



*Note: when using 1-compartment model, in order to predict Ka,

Vss was used instead of Vc (as in 2-compartment model)¹³,

2019 KSCPT Spring Meeting Pre-course Workshop

which may provide potential discrepancy in predicting human Ka.


Theories and Practices for Human Pharmacokinetics Prediction Using Non-clinical Information

Discovery
Modeling & Simulation
Higher PTS
(Price to Stockist)


Dallphin's allometric approach example (Vd)

- Excel 데이터분석의 “회귀분석”과 같은 결과
- Natural scale 그래프로 제시, 상수는 0 을 사용 ((0,0) 을 지나가는 직선)
- $V_c = 0.9833(WT)$ ($R^2 = 0.9967$), $V_{ss} = 1.9420(WT)$ ($R^2 = 0.9902$)
- 다음 버전에는 scaling exponent 를 예측하도록 update 될 예정

	A	B	C	D	E	F	G	H	I
1	요약 출력		Vc						
2									
3	회귀분석 통계량								
4	다중 상관계수	0.996788	R2						
5	결정계수	0.996788							
6	조정된 결정계수	0.746788							
7	표준 오차	0.291426							
8	관측수	5							
9									
10	분산 분석								
11		자유도	제곱합	제곱 평균	F 비	유의한 F			
12	회귀	1	105.4304	105.4304	1241.396	5.03E-05			
13	잔차	4	0.339716	0.084929					
14	계	5	105.7701						
15									
16	계수	표준 오차	t 통계량	P-값	하위 95%	상위 95%	하위 95.0%	상위 95.0%	
17	Y 절편	0	#N/A	#N/A	#N/A	#N/A	#N/A	#N/A	
18	X 1	0.983307	0.027908	35.2345	3.87E-06	0.905821	1.060793	0.905821	1.060793
19	theta								



			Vss						
1	요약 출력		Vss						
2									
3	회귀분석 통계량								
4	다중 상관	0.995108							
5	결정계수	0.990239	R2						
6	조정된 결	0.740239							
7	표준 오차	1.006699							
8	관측수	5							
9									
10	분산 분석								
11		자유도	제곱합	제곱 평균	F 비	유의한 F			
12	회귀	1	411.2471	411.2471	405.7924	0.000267			
13	잔차	4	4.053769	1.013442					
14	계	5	415.3009						
15									
16	계수	표준 오차	t 통계량	P-값	하위 95%	상위 95%	하위 95.0%	상위 95.0%	
17	Y 절편	0	#N/A	#N/A	#N/A	#N/A	#N/A	#N/A	
18	X 1	1.942038	0.096406	20.1429	3.58E-05	1.674371	2.209705	1.674371	2.209705
19	theta								



4. Examples


1. Simple allometry using monkey data
2. Simple allometry using 4 species data (mouse, rat, monkey, dog)
3. Allometry with correction factor (brain weight)
4. Modified allometry with correction factor



Simple allometry example (1)

- 실제 수행된 모델링 프로젝트에서의 simple allometry
- Scaling 3-kg monkey to 70-kg human for mAb⁹
- Scaling CL (non-specific CL), Vc, Vp, Q, and Vmax, Km (target-specific CL)

	monkey	Unit	x 3 kg monkey	Unit	Human	allometry				
CL		L/h/kg		l/h		•CLh = CLm*(WTh/WTm)^0.85				
V1		l/kg		l		•V1 = V1m*(WTh/WTm)^1				
V2		l/kg		l		•V2 = V2m*(WTh/WTm)^1				
Q		l/h/kg		l/h		•Q h = CLm*(WTh/WTm)^0.85				
Vmax		mg/h/kg		mg/h		•Vmax = Vmaxmm*(WTh/WTm)^0.85				
Km		mg/l		mg/l		•Km 그대로				



$$\frac{dX_1}{dt} = -\frac{CL}{V_1} \times X_1 - \frac{CL_d}{V_1} \times X_1 - \frac{X_1}{V_1} \times \frac{V_{max}}{K_m + \frac{X_1}{V_1}} + \frac{CL_d}{V_2} \times X_2$$

$$\frac{dX_2}{dt} = \frac{CL_d}{V_1} \times X_1 - \frac{CL_d}{V_2} \times X_2$$


Fig. 1 Two-compartment nonlinear PK model










Simple allometry example (2)

	kg	Vss (L/kg)	vc (L/kg)	vc (L)	vp (L/kg)	vp (L)	Q (L/h/kg)	Q(L/h)
	0			0		0		0
mouse	0.02	1.5	0.5	0.01	1	0.02	0.8	0.016
rat	0.2	1.5	0.5	0.1	1	0.2	0.8	0.16
monkey	3	1.3	0.8	2.4	0.5	1.5	0.8	2.4
dog	10	2	1	10	1	10	0.6	6
exp	70	1.84	1.17	81.96	0.67	46.69	0.63	44.09
dallphine		1.94	0.98		0.96		0.37	

Two-compartment open model, IV injection.





- 4 Species PK data
- Each species PK parameters estimated from 2-compartment model
- $V_d = a(WT)^b$, $Q = a(WT)^b$
- a , b estimated from regression line
- Extrapolation to predict PK in a 70-kg human



Allometry with correction factor example (3)

L/h	CL1	CL3	CL5	CL10
mouse				
rat				
monkey				
dog				
human				

BRAIN WT*					
KG	L/h	CL1*Br WT	CL3*Br WT	CL5*Br WT	CL10*Br W
0.00036	mouse				
0.0018	rat				
0.09	monkey				
0.08	dog				
1.4	human				



- $CL = a(BW)^b$ vs. $CL \times Brain\ Weight = a(BW)^b$




Modified allometry with correction factor example (4)

- 실제 수행된 모델링 프로젝트에서의 modified allometry
- Brain weight 를 correction factor 로 사용하여 human CL 예측 후, 용량에 따른 CL 변화를 수식으로 설명함


$$CL \times Brain\ weight = a \cdot f(Dose) \cdot (BW)^b$$

CL: allometry with standard correction factor (brain weight) (1)



- Brain weight 을 standard correction factor로 사용하여 CL 에 대해 각각의 용량 (1, 3, 5, 10 mg/kg)에 따른 interspecies scaling (모든 용량군의 R²~0.99) 을 수행하여 CL 를 구함 (CL_point estimate by dose)
- Clearance x BrainWeight = aBW^b

CL: allometry with standard correction factor (brain weight) (2)



- 앞서 Mouse 에서 얻은 용량에 따른 CL 수식을 기반으로 이를 interspecies scaling 에 적용함, 400 mg 이하에서 수식을 적용하였으며, 400 mg 초과시 CL 는 일정하다고 가정함 -400 mg 이하: $CL \times BrainWeight = a \cdot (CL_{max} \cdot (1 - DOSE / (DOSE + CL50))) \cdot BW^b$
(CL_{max}= , CL50= , a= , b = 1 *상수 a 와 b 는 모델링을 통해 산출한 값임)
- 400 mg 초과: CL 는 용량에 따라 일정하다고 가정함
- 따라서, 용량에 따른 CL 변화를 BW 에 따라 나타낼 수 있으며, 용량과 BW 정보가 있다면 이젠 CL 산출이 가능해짐
- 이를 기반으로 70kg 사람에서의 CL 를 산출한 결과는 위와 같음(CL_equation by dose), 이를 앞의 각각 용량별 interspecies scaling 결과(CL_point estimate by dose)와 비교하였음

CL prediction: Allometry vs. in vitro data¹¹

- Allometry vs. IVIVE methods
- Allometry: fu intercept method and two-species-based allometry (rat-dog) were best
- IVIVE: Using microsomes (incorporating both plasma and microsomal binding) and hepatocytes (not incorporating binding) resulted in better prediction

⇒ Allometry slightly more accurate than predictions based on in vitro data

⇒ Dallphin uses only in vitro data for CL prediction

- 다음 버전에는 allometry(in vivo data)를 통해 CL 예측이 가능하도록 update 될 예정



Final parameters

- Hepatic CL: CL_h_fromRativivc
- Renal CL:
 - CL_r (Rat to human prediction)
 - CL_r (Dog to human prediction)
- V_{ss}
- V_c ➔ K_a
- Q
- F_h (CL_h_fromRativivc)

Dallphin-AtoM(ver 0.8.5)

Basic Info Absorption Distribution Elimination **Final Parameters** Concentration Prediction About

CL : 30.37 L/h

Hepatic CL(Microsome)

☐ CL_h 80.7 L/h

☒ CL_h_fromRativivc 12.2 L/h

☐ CL_h_PS 17.1 L/h

☐ CL_h_PS(logD) 24.5 L/h

Renal CL

☒ CL_r (Rat to human prediction) 18.0 L/h

☐ CL_r (Dog to human prediction) 12.8 L/h

☐ CL_r (User input) 0.0 L/h

V_{ss} : 1.94 L/kg

V_c : 0.98 L/kg

Q : 0.37 L/hr/kg

K_a : 2.72 /hr

F : 0.84048

F_h(Microsome)

☐ F_h(CL_h) 0.07

☒ F_h(CL_h_fromRativivc) 0.85

☐ F_h(CL_h_PS) 0.80

☐ F_h(CL_h_PS(logD)) 0.71


F_a

☒ F_a_Caco 0.97

☐ F_a_PSA 0.02

References

1. Certara Simcyp workshop material 2017
2. Paine, Stuart W., et al. "Prediction of human renal clearance from preclinical species for a diverse set of drugs that exhibit both active secretion and net reabsorption." *Drug Metabolism and Disposition* 39.6 (2011): 1008–1013.
3. Obach, R. Scott, et al. "The prediction of human pharmacokinetic parameters from preclinical and in vitro metabolism data." *Journal of Pharmacology and Experimental Therapeutics* 283.1 (1997): 46–58.
4. Choi, Go-Wun, Yong-Bok Lee, and Hea-Young Cho. "Interpretation of Non-Clinical Data for Prediction of Human Pharmacokinetic Parameters: In Vitro-In Vivo Extrapolation and Allometric Scaling." *Pharmaceutics* 11.4 (2019): 168.
5. <https://www.slideserve.com/oren/allometric-scaling-to-predict-pharmacokinetic-and-pharmacodynamic-parameters-in-man>
6. Ward, Keith W., and Brian R. Smith. "A comprehensive quantitative and qualitative evaluation of extrapolation of intravenous pharmacokinetic parameters from rat, dog, and monkey to humans. I. Clearance." *Drug Metabolism and Disposition* 32.6 (2004): 603–611.
7. Mahmood, Iftekhar, and J. D. Balian. "Interspecies scaling: predicting clearance of drugs in humans. Three different approaches." *Xenobiotica* 26.9 (1996): 887–895.
8. Nagilla, Rakesh, and Keith W. Ward. "A comprehensive analysis of the role of correction factors in the allometric predictivity of clearance from rat, dog, and monkey to humans." *Journal of pharmaceutical sciences* 93.10 (2004): 2522–2534.
9. Kamath, Amrita V., et al. "Preclinical pharmacokinetics of MEHD7945A, a novel EGFR/HER3 dual-action antibody, and prediction of its human pharmacokinetics and efficacious clinical dose." *Cancer chemotherapy and pharmacology* 69.4 (2012): 1063–1069.
10. Goteti, Kosalaram, C. Edwin Garner, and Iftekhar Mahmood. "Prediction of human drug clearance from two species: a comparison of several allometric methods." *Journal of pharmaceutical sciences* 99.3 (2010): 1601–1613.
11. Ring, Barbara J., et al. "PhRMA CPCDC initiative on predictive models of human pharmacokinetics, part 3: comparative assesment of prediction methods of human clearance." *Journal of pharmaceutical sciences* 100.10 (2011): 4090–4110.
12. Poulin, Patrick, and Frank-Peter Theil. "Prediction of pharmacokinetics prior to in vivo studies. 1. Mechanism-based prediction of volume of distribution." *Journal of pharmaceutical sciences* 91.1 (2002): 129–156.
13. Usansky, Helen H., and Patrick J. Sinko. "Estimating human drug oral absorption kinetics from Caco-2 permeability using an absorption-disposition model: model development and evaluation and derivation of analytical solutions for k_a and F_a ." *Journal of pharmacology and experimental therapeutics* 314.1 (2005): 391–399.



경청해 주셔서 감사합니다.

주요 *in vivo* 정보와
이를 활용한 인간 약동학 예측 이론

