

Hicksian aggregation and a world full of derivatives

Felipe Del Canto, 19 de Agosto de 2019

1. Taylor is that you?

To tackle the problem of aggregation across goods, I am considering an agent with utility function $u : \mathbb{R}_{++}^{n+m} \rightarrow \mathbb{R}$ that solves the following optimization problem

$$\begin{aligned} \max_{\mathbf{x}, \mathbf{z}} \quad & u(\mathbf{x}, \mathbf{z}) \\ \text{s.a.} \quad & \mathbf{p}\mathbf{x} + \mathbf{q}\mathbf{z} = y. \end{aligned}$$

With solution $\mathbf{x}(\mathbf{p}, \mathbf{q}, y)$. Assume we have a rule to aggregate the consumption bundle \mathbf{x} using a function $g : \mathbb{R}^n \rightarrow \mathbb{R}$ and call $X := g(x)$. The Hicks aggregation problem seeks to find $P := f(\mathbf{p})$ and $U : \mathbb{R}^{1+m} \rightarrow \mathbb{R}$ such that the function

$$\begin{aligned} X(f(\mathbf{p}), \mathbf{q}, y) &:= \arg \max_{X, \mathbf{z}} U(X, \mathbf{z}) \\ \text{s.a.} \quad & PX + \mathbf{q}\mathbf{z} = y \end{aligned}$$

satisfies

$$X(f(\mathbf{p}), \mathbf{q}, y) = g(\mathbf{x}(\mathbf{p}, \mathbf{q}, y)).$$

Typically this is not possible except in very restrictive cases. Suppose the model imposes some function U to the agent and some price index f . We would like to measure the error in the estimation of the aggregate demand $X = g(x)$. Note that the Lagrangians of both problems are

$$\mathcal{L}_d = u(\mathbf{x}, \mathbf{z}) + \lambda_d(y - \mathbf{p}\mathbf{x} - \mathbf{q}\mathbf{z}), \tag{1}$$

$$\mathcal{L}_a = U(X, \mathbf{z}) + \lambda_a(y - f(\mathbf{p})X - \mathbf{q}\mathbf{z}), \tag{2}$$

And the corresponding first order conditions are

$$\begin{cases} \nabla_{\mathbf{x}} u - \lambda_d \mathbf{p} = \mathbf{0}, \\ \nabla_{\mathbf{z}} u - \lambda_d \mathbf{q} = \mathbf{0} \end{cases} \quad (3)$$

$$\begin{cases} \frac{\partial U}{\partial x} - \lambda_a f(\mathbf{p}) = 0, \\ \nabla_{\mathbf{z}} U - \lambda_a \mathbf{q} = \mathbf{0} \end{cases} \quad (4)$$

We are interested in the difference

$$\left| X(f(\mathbf{p}), \mathbf{q}, y) - g(\mathbf{x}(\mathbf{p}, \mathbf{q}, y)) \right|.$$

Assuming the investigator has access to both aggregate and disaggregate data in consumption and a set of prices $\mathbf{p}_0, \mathbf{q}_0$ and income y_0 . It is possible to approximate each function using a Taylor expansion around $(\mathbf{p}_0, \mathbf{q}_0, y_0)$. Explicitly

$$\begin{aligned} X(f(\mathbf{p}), \mathbf{q}, y) &= X(f(\mathbf{p}_0), \mathbf{q}_0, y_0) + \frac{\partial X}{\partial P} \nabla f(\mathbf{p} - \mathbf{p}_0) + \nabla_q X(\mathbf{q} - \mathbf{q}_0) + \frac{\partial X}{\partial m}(m - m_0) + \mathcal{O}(2) \\ g(x(\mathbf{p}, \mathbf{q}, m)) &= g(x(\mathbf{p}_0, \mathbf{q}_0, m_0)) + \nabla g \cdot D_{\mathbf{x}} \left((\mathbf{p}, \mathbf{q}, m) - (\mathbf{p}_0, \mathbf{q}_0, m_0) \right) + \mathcal{O}(2) \end{aligned}$$

The derivatives of each function could in principle be obtained by implicit differentiation of equations (3) and (4). In particular, the derivatives of X and \mathbf{x} with respect to m can be obtained using the envelope theorem.

2. In other news...

The Schechtman y Escudero paper seems promising. Theorem 1.1 provides bounds on the MPC, but they depend on the solution of another (though deterministic) problem. Besides, the bounds work on a finite horizon problem although I think it may be enough if we take t large enough. Again, I have more questions than answers in this point.