

Homework 1

● Graded

Student

周洛君

Total Points

52 / 60 pts

Question 1

Questions 0 and 1

8 / 8 pts

✓ + 2 pts (Question 0) Page selection

✓ + 6 pts Correct

+ 4 pts A minor mistake (including no appropriate explanation/step)

+ 2 pts A major mistake or two minor mistakes

+ 0 pts Totally wrong or empty

Question 2

Question 2

12 / 12 pts

✓ + 12 pts Correct

+ 10 pts A minor mistake (including no appropriate explanation/step in 2.2)

+ 8 pts A major mistake or two minor mistakes

+ 0 pts Totally wrong or empty

Question 3

Question 3

6 / 6 pts

✓ + 6 pts Correct

+ 4 pts A minor mistake

+ 2 pts A major mistake or two minor mistakes

+ 0 pts Totally wrong or empty

Question 4

Question 4

12 / 16 pts

+ 16 pts Correct

+ 14 pts A minor mistake

✓ + 12 pts A major mistake or two minor mistakes

+ 10 pts A major mistake and other mistakes

+ 8 pts Two major mistakes

+ 4 pts Some reasonable efforts

+ 0 pts Totally wrong or empty

1 major mistake: this is the answer of 4-2.

Question 5

Question 5

6 / 6 pts

✓ + 6 pts Correct

+ 4 pts A minor mistake

+ 2 pts A major mistake or two minor mistakes

+ 0 pts Totally wrong or empty

Question 6

Question 6

4 / 6 pts

+ 6 pts Correct

✓ + 4 pts A minor mistake

+ 2 pts A major mistake or two minor mistakes

+ 0 pts Totally wrong or empty

Question 7

Question 7

4 / 6 pts

+ 6 pts Correct

✓ + 4 pts A minor mistake

+ 2 pts A major mistake or two minor mistakes

+ 0 pts Totally wrong or empty

Question assigned to the following page: [1](#)

Digital System Design and Lab: HW1

Lo Chun, Chou
R13922136

March 4, 2025

1

Since the following operation is correct:

$$024 + 043 + 013 + 033 = 201$$

Let base $= k \in \mathbb{Z}^+$, $k > 4$

We can formulate the following equations:

$$4 + 3 + 3 + 3 \pmod{k} = 13 \pmod{k} = 1 \quad (1)$$

$$2 + 4 + 1 + 3 + c_1 \pmod{k} = 10 + c_1 \pmod{k} = 0 \quad (2)$$

$$0 + 0 + 0 + 0 + c_2 \pmod{k} = c_2 \pmod{k} = 2 \quad (3)$$

Where c_1, c_2 are the carry-out generated by the addition of the previous digits.

By (1), and $k > 4$, we knew that the possible values of k are 6, 12.

Case 1: $k = 6$

Suppose $k = 6$, we have:

$$4 + 3 + 3 + 3 \pmod{6} = 13 \pmod{6} = 1 \quad (1)$$

$$2 + 4 + 1 + 3 + c_1 \pmod{6} = 10 + c_1 \pmod{6} = 0 \quad (2)$$

$$0 + 0 + 0 + 0 + c_2 \pmod{6} = c_2 \pmod{6} = 2 \quad (3)$$

By (1), we knew that $c_1 = 2$, thus equation (2) holds since $10 + 2 = 12 \equiv 0 \pmod{6}$. Similarly, we knew that $c_2 = 2$, thus equation (3) holds since $0 + 2 = 2 \equiv 2 \pmod{6}$.

Therefore, $k = 6$ is a valid solution.

Question assigned to the following page: [1](#)

Case 2: $k = 12$

Suppose $k = 12$, we have:

$$4 + 3 + 3 + 3 \pmod{12} = 13 \pmod{12} = 1 \quad (1)$$

$$2 + 4 + 1 + 3 + c_1 \pmod{12} = 10 + c_1 \pmod{12} = 0 \quad (2)$$

$$0 + 0 + 0 + 0 + c_2 \pmod{12} = c_2 \pmod{12} = 2 \quad (3)$$

By (1), we knew that $c_1 = 1$ since $\lfloor 13 \div 12 \rfloor = 1$, however, this contradicts with equation (2) since $10 + 1 = 11 \not\equiv 0 \pmod{12}$.

Therefore, $k = 12$ is not a valid solution.

The only possible base is $k = 6$. □

Question assigned to the following page: [2](#)

2

(1)

Decimal	weighted code			
	8	4	-2	-1
0	0	0	0	0
1	0	1	1	1
2	0	1	1	0
3	0	1	0	1
4	0	1	0	0
5	1	0	1	1
6	1	0	1	0
7	1	0	0	1
8	1	0	0	0
9	1	1	1	1

(2)

Decimal (d)	$9 - d$	weighted code for $9 - d$			
		8	4	-2	-1
0	9	1	1	1	1
1	8	1	0	0	0
2	7	1	0	0	1
3	6	1	0	1	0
4	5	1	0	1	1
5	4	0	1	0	0
6	3	0	1	0	1
7	2	0	1	1	0
8	1	0	1	1	1
9	0	0	0	0	0

By observing the table, and by the fact that $9 - d$ is the complement of d , we can see that the weighted code for $9 - d$ is the complement of the weighted code for d .

Which means that to obtain the code of $9 - d$, we could simply flip the bits of the code of d .

Question assigned to the following page: [3](#)

3

Find the complement of the function by only DeMorgan's laws and involution law:

$$F(A, B, C, D) = AB'C + (A' + B + D)(ABD' + B')$$

As required, we can only use the below properties:

$$\begin{aligned}(AB)' &= A' + B' \\ (A + B)' &= A'B' \\ (A')' &= A\end{aligned}$$

$$\begin{aligned}F'(A, B, C, D) &= (AB'C)' \cdot [(A' + B + D) \cdot (ABD' + B')] \\ &= [(AB')' + C'] \cdot [(A' + B + D)' + (ABD' + B')'] \\ &= (A' + B + C') \cdot [(A' + B)' D' + (ABD')' B] \\ &= (A' + B + C') \cdot [AB'D' + ((AB)' + D) B] \\ &= (A' + B + C') \cdot [AB'D' + (A' + B' + D) B] \\ &= (A' + B + C') \cdot (AB'D' + A'B + BD)\end{aligned}$$

Question assigned to the following page: [4](#)

4


(1)

The switching algebra expression is:

$$\begin{aligned} & D \cdot \{[(A' + B') \cdot C] + (A \cdot C')\} \\ &= D \cdot (A'C + B'C + AC') \\ &= A'CD + B'CD + AC'D \end{aligned}$$

1


(2)



(3)

We can convert the original expression into SOP form:

$$\begin{aligned} & A'CD + B'CD + AC'D \\ &= (A'C + B'C + AC') \cdot D \\ &= [(A' + B')C + AC'] \cdot D \\ &= (A' + B' + C') \cdot (A + C) \cdot D \end{aligned}$$



Question assigned to the following page: [5](#)

5

Obtain sum of product form:

$$\begin{aligned}
& (A + B + C + D)(A' + B' + C + D')(A' + C)(A + D)(B + C + D) \\
&= \{[A + (B + C + D)](B + C + D)\}(A' + B' + C + D')(A' + C)(A + D) \\
&= [A(B + C + D) + (B + C + D)(B + C + D)](A' + B' + C + D')(A' + C)(A + D) \\
&= [AB + AC + AD + (B + C + D)](A' + B' + C + D')(A' + C)(A + D) \\
&= (B + C + D)(A' + B' + C + D')(A' + C)(A + D) \\
&= (B + C + D)[A' + (B' + C + D')C](A + D) \\
&= (B + C + D)[A'D + A(B' + C + D')C] \\
&= (B + C + D)[A'D + AB'C + AC + ACD'] \\
&= (B + C + D)[A'D + AB'C + AC] \\
&= (B + C + D)[A'D + AC] \\
&= A'BD + A'CD + A'D + ABC + AC + ACD \\
&= A'D + AC \quad \square
\end{aligned}$$

Question assigned to the following page: [6](#)

6

Obtain product of sums form:

$$\begin{aligned} & BCD + C'D' + B'C'D + CD \\ &= C'D' + B'C'D + CD \\ &= (C' + D)(C + D') + B'(C + D)' \\ &= [(C' + D) + B'][(C + D')' + (C' + D)] \\ &= (B' + C' + D)(C'D + C' + D) \\ &= (B' + C' + D)(C' + D) \\ &= B'(C' + D) + (C' + D)(C' + D) \\ &= B'(C' + D) + (C' + D) \\ &= C' + D \quad \square \end{aligned}$$

Question assigned to the following page: [Z](#)

7

Inputs			Output
A	B	C	
0	0	0	0
1	0	0	0
0	1	0	0
0	0	1	0
1	1	0	1
0	1	1	1
1	0	1	1
1	1	1	1

From the table we can form the sum of products expression:

$$\begin{aligned}
 & ABC' + A'BC + AB'C + ABC \\
 = & ABC' + AB'C + BC \quad \square
 \end{aligned}$$