# Near-field singularity extraction for rooftop basis functions on uniform rectangular grids

## Contents


1 Introduction 2

2 Setting: cells, rooftops and moments 2

3 Singularity extraction on the touching cells 6

4 Reduction to two dimensions 7

5 Mixed antiderivatives 10

6 Closed forms for the touching cells 13

A The mixed antiderivatives $G_{mn}^{(p)}$ 15

B The touching moments 18

# 1 Introduction

This note continues the bachelor thesis [1]. There the Galerkin matrix of the electric field integral equation (EFIE) with rooftop basis functions on a uniform rectangular grid is assembled from three Toeplitz generators. Their entries are four-dimensional integrals of the Green's function

$$g(R) = \frac{\mathrm{e}^{-\mathrm{i}kR}}{R}, \qquad R = \|\boldsymbol{r} - \boldsymbol{r}'\|, \tag{1}$$

over pairs of cells, weighted by the linear profiles of the rooftops (the factor $1/(4\pi)$ is kept outside). The implementation in `src` integrates them with a tensor Gauss–Legendre (GL) rule of order $q$ and extracts $1/R$ on the coincident cell ([1], Entry *singex* and Appendix).

Cells that share an edge or only a vertex meet $R = 0$ as well, on the boundary of the integration domain. There the GL error does not decrease with the order: the generator entries keep a relative error of about $10^{-3}$ for cells of $\lambda/10$ at every order between 3 and 16, and extraction on the coincident cell alone changes nothing measurable. The note therefore extracts the non-smooth terms of (1),

$$\frac{1}{R}, \qquad -\frac{k^2}{2}R, \qquad \frac{k^4}{24}R^3,$$

on all nine touching cells and gives the closed forms of every static integral this needs.

**Three terms are enough.** With the three terms the entries reach $10^{-13}$–$10^{-14}$ at the order 12–16, and the computed current is at the rounding floor of double precision. A fourth term ($R^5$) would act below that floor, so no further extraction is needed.

**The way to the closed forms.** Section 2 expresses the three generators through seven cell-pair moments, so that only integrals over pairs of cells with linear weights are left. Section 3 splits the Green's function into the three extracted terms and a smooth remainder; the remainder stays with the GL rule, and the extracted terms leave the static moments with the kernels $R^{-1}$, $R$ and $R^3$. Section 4 reduces these four-dimensional integrals to two dimensions, and Section 5 gives the antiderivatives of the resulting integrands. Section 6 puts them together: every static moment of a touching cell pair is a short sum of antiderivative values. The note is written as a manual to the implementation; all closed forms are listed in Appendices A and B in the form of the thesis appendix.

**Conventions.** The time convention is $\mathrm{e}^{\mathrm{i}\omega t}$, and $\operatorname{arsinh} x = \log\bigl(x + \sqrt{x^2+1}\bigr)$; the logarithms of the thesis appendix are $\log\bigl((\sqrt{a^2+b^2}+a)/b\bigr) = \operatorname{arsinh}(a/b)$. The notation follows the thesis [1].

# 2 Setting: cells, rooftops and moments

**Definition 2.1 (Uniform grid).** Let $a, b > 0$ be the cell sizes in the $x$ and $y$ directions. For $(i, j) \in \mathbb{Z}^2$ the cell is

$$\Omega_{ij} = [ia, (i+1)a] \times [jb, (j+1)b] \subset \mathbb{R}^2.$$

A point $\boldsymbol{r} = (x, y) \in \Omega_{ij}$ has the local coordinates

$$\xi(\boldsymbol{r}) = \frac{x}{a} - i \in [0, 1], \qquad \eta(\boldsymbol{r}) = \frac{y}{b} - j \in [0, 1].$$

For an offset $\boldsymbol{d} = (d_x, d_y) \in \mathbb{Z}^2$ we write $\Omega_{\boldsymbol{d}} = \Omega_{d_x d_y}$. The two cells $\Omega_{\boldsymbol{0}}$ and $\Omega_{\boldsymbol{d}}$ share at least one point if and only if $\boldsymbol{d} \in \mathcal{T} = \{-1, 0, 1\}^2$. We call these offsets *touching* (Figure 1): $\boldsymbol{d} = \boldsymbol{0}$ is the coincident cell, the four offsets $(\pm 1, 0)$, $(0, \pm 1)$ share an edge and the four offsets $(\pm 1, \pm 1)$ a vertex.

**Definition 2.2 (Rooftop functions).** The $x$-directed rooftop $\boldsymbol{f}^x_{ij}$ is supported on $\Omega_{ij} \cup \Omega_{i+1,j}$ and

$$\boldsymbol{f}^x_{ij}(\boldsymbol{r}) = \hat{\boldsymbol{x}} \cdot \begin{cases} \xi(\boldsymbol{r}), & \boldsymbol{r} \in \Omega_{ij} \quad \text{(rising wedge)}, \\ 1 - \xi(\boldsymbol{r}), & \boldsymbol{r} \in \Omega_{i+1,j} \quad \text{(falling wedge)}. \end{cases}$$

Its normal component on the common edge is one, and its divergence is $+1/a$ on $\Omega_{ij}$ and $-1/a$ on $\Omega_{i+1,j}$. The $y$-directed rooftop $\boldsymbol{f}^y_{ij}$ is defined in the same way with $\eta$, $b$ and $\Omega_{i,j+1}$. These are the rooftops of [1] and of `src` (unit height on the edge).





Figure 1: Left: the reference cell $\Omega_{\boldsymbol{0}}$ and its eight touching neighbours $\Omega_{\boldsymbol{d}}$, labelled by the offset $\boldsymbol{d} \in \mathcal{T}$. The points with $R = 0$ that the two cells have in common are drawn in red: all of $\Omega_{\boldsymbol{0}}$ for the coincident cell, an edge for an edge neighbour and a vertex for a vertex neighbour. Right: the $x$-directed rooftop $\boldsymbol{f}^x_{ij}$, split into its rising wedge $\xi$ on $\Omega_{ij}$ and its falling wedge $1 - \xi$ on $\Omega_{i+1,j}$.

**Entry 2.1 (Galerkin entries).** With (1), the Galerkin entry of the EFIE between rooftops $\boldsymbol{f}_m$ and $\boldsymbol{f}_n$ is [1]

$$Z_{mn} = \frac{\mathrm{i}Z_0}{4\pi k}\Big(k^2 \iint \boldsymbol{f}_m(\boldsymbol{r}) \cdot \boldsymbol{f}_n(\boldsymbol{r}')\, g(R)\, \mathrm{d}A'\, \mathrm{d}A - \iint \nabla \cdot \boldsymbol{f}_m(\boldsymbol{r})\, \nabla' \cdot \boldsymbol{f}_n(\boldsymbol{r}')\, g(R)\, \mathrm{d}A'\, \mathrm{d}A\Big). \tag{2}$$

By translation invariance the entries depend only on the offset of the two rooftops. They are generated by

$$A^{xx}(\boldsymbol{d}) = \iint \boldsymbol{f}^x_{\boldsymbol{0}} \cdot \boldsymbol{f}^x_{\boldsymbol{d}}\, g, \qquad A^{yy}(\boldsymbol{d}) = \iint \boldsymbol{f}^y_{\boldsymbol{0}} \cdot \boldsymbol{f}^y_{\boldsymbol{d}}\, g, \qquad \Phi(\boldsymbol{d}) = \int_{\Omega_{\boldsymbol{0}}} \int_{\Omega_{\boldsymbol{d}}} g.$$

The divergences are piecewise constant, so the scalar-potential parts of all blocks are finite differences of $\Phi$; for example $\iint \nabla\!\cdot\boldsymbol{f}^{x}_{\mathbf{0}}\,\nabla'\!\cdot\boldsymbol{f}^{x}_{\boldsymbol{d}}\,g = a^{-2}\big(2\Phi(\boldsymbol{d})-\Phi(\boldsymbol{d}+\boldsymbol{e}_x)-\Phi(\boldsymbol{d}-\boldsymbol{e}_x)\big)$. This is how `src` assembles the three Toeplitz generators.

**Definition 2.3 (Cell-pair moments).** Let $K:(0,\infty)\to\mathbb{C}$ satisfy $K(\|\cdot\|)\in L^1_{\mathrm{loc}}(\mathbb{R}^2)$. For $\boldsymbol{d}\in\mathbb{Z}^2$ and $\alpha,\beta\in\{0,1\}$ define

$$M^{[K]}_{\alpha\beta}(\boldsymbol{d}) = \frac{1}{(ab)^2}\int_{\Omega_{\mathbf{0}}}\int_{\Omega_{\boldsymbol{d}}} \xi(\boldsymbol{r})^{\alpha}\,\xi(\boldsymbol{r}')^{\beta}\,K\big(\|\boldsymbol{r}-\boldsymbol{r}'\|\big)\,\mathrm{d}A(\boldsymbol{r}')\,\mathrm{d}A(\boldsymbol{r}),$$
$$N^{[K]}_{\alpha\beta}(\boldsymbol{d}) = \frac{1}{(ab)^2}\int_{\Omega_{\mathbf{0}}}\int_{\Omega_{\boldsymbol{d}}} \eta(\boldsymbol{r})^{\alpha}\,\eta(\boldsymbol{r}')^{\beta}\,K\big(\|\boldsymbol{r}-\boldsymbol{r}'\|\big)\,\mathrm{d}A(\boldsymbol{r}')\,\mathrm{d}A(\boldsymbol{r}),$$

where $\xi(\boldsymbol{r}')$ and $\eta(\boldsymbol{r}')$ are the local coordinates in $\Omega_{\boldsymbol{d}}$. For $K(R)=R^p$ we write $M^{(p)}_{\alpha\beta}(\boldsymbol{d})$ and $N^{(p)}_{\alpha\beta}(\boldsymbol{d})$. Since $N_{00}=M_{00}$, there are seven moments, ordered in `src` as $M_{00}$, $M_{10}$, $M_{01}$, $M_{11}$, $N_{10}$, $N_{01}$, $N_{11}$.

The moments carry all the geometry. The next lemma lists their symmetries.

**Lemma 2.1 (Symmetries of the moments).** Let $K$ be admissible, $\boldsymbol{d}=(d_x,d_y)\in\mathbb{Z}^2$ and $\alpha,\beta\in\{0,1\}$. Then:

(i) $M^{[K]}_{\alpha\beta}(d_x,-d_y) = M^{[K]}_{\alpha\beta}(d_x,d_y)$;

(ii) reflection in $x$:

$$\begin{aligned}
M^{[K]}_{00}(-d_x,d_y) &= M^{[K]}_{00}(\boldsymbol{d}),\\
M^{[K]}_{10}(-d_x,d_y) &= \big(M^{[K]}_{00}-M^{[K]}_{10}\big)(\boldsymbol{d}),\\
M^{[K]}_{01}(-d_x,d_y) &= \big(M^{[K]}_{00}-M^{[K]}_{01}\big)(\boldsymbol{d}),\\
M^{[K]}_{11}(-d_x,d_y) &= \big(M^{[K]}_{00}-M^{[K]}_{10}-M^{[K]}_{01}+M^{[K]}_{11}\big)(\boldsymbol{d});
\end{aligned}$$

(iii) $N^{[K]}_{\alpha\beta}(d_x,d_y;a,b) = M^{[K]}_{\alpha\beta}(d_y,d_x;b,a)$ (the cell sizes written explicitly);

(iv) $M^{[K]}_{\alpha\beta}(\boldsymbol{d}) = M^{[K]}_{\beta\alpha}(-\boldsymbol{d})$.

*Proof.* The kernel depends only on $\|\boldsymbol{r}-\boldsymbol{r}'\|$, which is invariant under reflections, translations and the interchange of $\boldsymbol{r}$ and $\boldsymbol{r}'$.

(i) The reflection $y\mapsto b-y$ maps $\Omega_{\mathbf{0}}$ onto itself and $\Omega_{(d_x,-d_y)}$ onto $\Omega_{(d_x,d_y)}$. It leaves both local $x$-coordinates unchanged.

(ii) The reflection $x\mapsto a-x$ maps $\Omega_{\mathbf{0}}$ onto itself with $\xi\mapsto 1-\xi$, and $\Omega_{(-d_x,d_y)}$ onto $\Omega_{(d_x,d_y)}$ with $\xi'\mapsto 1-\xi'$. Indeed, $[-d_xa,(1-d_x)a]$ goes to $[d_xa,(d_x+1)a]$ and $x'/a+d_x\mapsto 1-(x'/a+d_x)$. Expand $(1-\xi)^{\alpha}(1-\xi')^{\beta}$.

(iii) Exchange the roles of $x$ and $y$.

(iv) Interchange $\boldsymbol{r}\leftrightarrow\boldsymbol{r}'$ and translate by $-\boldsymbol{d}$; the cell $\Omega_{\boldsymbol{d}}$ becomes $\Omega_{\mathbf{0}}$ and $\Omega_{\mathbf{0}}$ becomes $\Omega_{-\boldsymbol{d}}$.

□

**Proposition 2.1 (Generators from moments).** For $\boldsymbol{d}=(d_x,d_y)$ with $d_x\ge 0$,

$$\begin{aligned}
\frac{A^{xx}(\boldsymbol{d})}{(ab)^2} &= M_{11}(\boldsymbol{d}) + \big(M_{00}-M_{10}-M_{01}+M_{11}\big)(\boldsymbol{d})\\
&\quad + \big(M_{10}-M_{11}\big)(\boldsymbol{d}+\boldsymbol{e}_x) + \big(M_{01}-M_{11}\big)(\boldsymbol{d}-\boldsymbol{e}_x),
\end{aligned}\tag{3}$$

with $M = M^{[g]}$. For $d_x = 0$ the last term is to be read as

$$\big(M_{01} - M_{11}\big)(-1, d_y) = \big(M_{10} - M_{11}\big)(1, d_y).$$

Furthermore $\Phi(\boldsymbol{d}) = (ab)^2 M_{00}^{[g]}(\boldsymbol{d})$, and $A^{yy}$ is given by (3) with $N$ and $\boldsymbol{e}_y$ in place of $M$ and $\boldsymbol{e}_x$.

*Proof.* Split both rooftops into their wedges, as in Definition (2.2). The reference rooftop $\boldsymbol{f}_{\boldsymbol{0}}^x$ has the rising wedge $\xi$ on $\Omega_{\boldsymbol{0}}$ and the falling wedge $1-\xi$ on $\Omega_{\boldsymbol{e}_x}$. The rooftop $\boldsymbol{f}_{\boldsymbol{d}}^x$ has the rising wedge $\xi'$ on $\Omega_{\boldsymbol{d}}$ and the falling wedge $1-\xi'$ on $\Omega_{\boldsymbol{d}+\boldsymbol{e}_x}$. Both are $x$-directed, so the dot product is the product of the profiles. The four wedge pairs of Figure 2 give

| pair | cells | offset | weight | moment |
|---|---|---|---|---|
| 1, rising–rising | $\Omega_{\boldsymbol{0}}$, $\Omega_{\boldsymbol{d}}$ | $\boldsymbol{d}$ | $\xi\xi'$ | $M_{11}$ |
| 2, falling–falling | $\Omega_{\boldsymbol{e}_x}$, $\Omega_{\boldsymbol{d}+\boldsymbol{e}_x}$ | $\boldsymbol{d}$ | $(1-\xi)(1-\xi')$ | $M_{00} - M_{10} - M_{01} + M_{11}$ |
| 3, rising–falling | $\Omega_{\boldsymbol{0}}$, $\Omega_{\boldsymbol{d}+\boldsymbol{e}_x}$ | $\boldsymbol{d}+\boldsymbol{e}_x$ | $\xi(1-\xi')$ | $M_{10} - M_{11}$ |
| 4, falling–rising | $\Omega_{\boldsymbol{e}_x}$, $\Omega_{\boldsymbol{d}}$ | $\boldsymbol{d}-\boldsymbol{e}_x$ | $(1-\xi)\xi'$ | $M_{01} - M_{11}$ |

with the moments taken at the offset of the pair (for pair 4 after the translation by $-\boldsymbol{e}_x$). The normalization $(ab)^{-2}$ in Definition (2.3) accounts for the factor $(ab)^2$. For $d_x = 0$ the offset $\boldsymbol{d} - \boldsymbol{e}_x = (-1, d_y)$ is negative. Lemma (2.1)(ii) gives

$$\begin{aligned}(M_{01} - M_{11})(-1, d_y) &= (M_{00} - M_{01})(1, d_y) - (M_{00} - M_{10} - M_{01} + M_{11})(1, d_y)\\ &= (M_{10} - M_{11})(1, d_y).\end{aligned}$$

The statement on $\Phi$ is the definition with $\alpha = \beta = 0$. The $y$-direction is identical. □





Figure 2: The four wedge pairs of the rooftops $\boldsymbol{f}_{\boldsymbol{0}}^x$ and $\boldsymbol{f}_{\boldsymbol{d}}^x$ in Proposition (2.1), drawn for $\boldsymbol{d} = (2, 2)$, numbered as in its proof. Every pair is a pair of cells at one offset, weighted by a product of two linear profiles.

The generators of all three Toeplitz blocks need the seven moments on the offsets $d_x \geq -1$, $d_y \geq 0$, and by Lemma (2.1) on $d_x, d_y \geq 0$ only.

# 3 Singularity extraction on the touching cells

**Proposition 3.1 (Non-smooth terms of the Green's function).** For $R \geq 0$,

$$g(R) = \frac{\mathrm{e}^{-\mathrm{i}kR}}{R} = \sum_{j=0}^{\infty} \frac{(-\mathrm{i}k)^j}{j!} R^{j-1} = \sum_{\ell=0}^{\infty} \frac{(-1)^\ell k^{2\ell}}{(2\ell)!} R^{2\ell-1} \; - \; \mathrm{i} \sum_{\ell=0}^{\infty} \frac{(-1)^\ell k^{2\ell+1}}{(2\ell+1)!} R^{2\ell}. \quad (4)$$

The series converge absolutely and locally uniformly for $R > 0$; the second one is an entire function of $R^2$. As functions of $\boldsymbol{z} = \boldsymbol{r} - \boldsymbol{r}' \in \mathbb{R}^2$ the even powers $R^{2\ell} = \|\boldsymbol{z}\|^{2\ell}$ are polynomials, while the odd powers $R^{2\ell-1}$ are not smooth at $\boldsymbol{z} = \boldsymbol{0}$.

*Proof.* Insert the exponential series and separate the even and odd $j$; this is Proposition *greensing* of [1]. □

**Definition 3.1 (Extraction of order $N$).** For $N \in \mathbb{N}$ let

$$S_N(R) = \sum_{\ell=0}^{N-1} \frac{(-1)^\ell k^{2\ell}}{(2\ell)!} R^{2\ell-1} = \frac{1}{R} - \frac{k^2}{2} R + \frac{k^4}{24} R^3 - \dots, \qquad g_N = g - S_N. \quad (5)$$

Table 1 lists the orders: $N = 1$ is the extraction of `src`, $N = 2$ the two-term extraction of [1] (Entry *singex*), there written as

$$g = g_{\mathrm{s}} + \frac{1}{4\pi} R^{-1} - \frac{k^2}{8\pi} R,$$

and $N = 3$ the one of this note. For every $N \geq 1$, $g_N$ extends continuously to $R = 0$ with $g_N(0) = -\mathrm{i}k$.

**Lemma 3.1 (Regularity of the remainder).** For $N \geq 1$ the function $\boldsymbol{z} \mapsto g_N(\|\boldsymbol{z}\|)$ is real-analytic on $\mathbb{R}^2 \setminus \{\boldsymbol{0}\}$ and belongs to $C^{2N-2,1}$ on every bounded set: its derivatives up to the order $2N-2$ are continuous, and those of the order $2N-2$ are Lipschitz. It is not of class $C^{2N-1}$ unless $k = 0$.

*Proof.* By (4),

$$g_N(R) = E(R^2) + R^{2N-1} F(R^2),$$

with the entire functions

$$E(s) = -\mathrm{i} \sum_{\ell \geq 0} \frac{(-1)^\ell k^{2\ell+1}}{(2\ell+1)!} s^\ell, \qquad F(s) = \sum_{\ell \geq N} \frac{(-1)^\ell k^{2\ell}}{(2\ell)!} s^{\ell-N}.$$

The function $E(\|\boldsymbol{z}\|^2)$ is real-analytic on $\mathbb{R}^2$.

The function $\|\boldsymbol{z}\|^{2N-1} = \left(\|\boldsymbol{z}\|^2\right)^{N-1} \|\boldsymbol{z}\|$ is smooth away from the origin and positively homogeneous of degree $2N-1$. Hence its derivatives of order $j \leq 2N-2$ are homogeneous of degree $2N-1-j \geq 1$. They are therefore continuous at the origin (with the value zero), and those of order $2N-2$, being homogeneous of degree one, are Lipschitz. The derivatives of order $2N-1$ are homogeneous of degree zero and not constant, so they are bounded but discontinuous at the origin.

Multiplication by the analytic factor $F(\|\boldsymbol{z}\|^2)$ preserves the class $C^{2N-2,1}$ on bounded sets. Since $F(0) = (-1)^N k^{2N}/(2N)! \neq 0$ for $k \neq 0$, the term $F(0)\,\|\boldsymbol{z}\|^{2N-1}$ is present and the class is sharp. □

On a touching cell pair the integrand of a moment meets $\boldsymbol{z} = \mathbf{0}$, in the interior of the four-dimensional domain for the coincident cell and on its boundary for a shared edge or vertex. A tensor GL rule integrates polynomials exactly, so its error on such a pair is governed by the regularity of Lemma (3.1) (Table 1).

Table 1: The extraction orders and the regularity of the remainder on a touching cell pair.

| $N$ | extracted $S_N$ | used in | remainder $g_N$ |
|---|---|---|---|
| 0 | — | | unbounded |
| 1 | $1/R$ | `src` | Lipschitz |
| 2 | $1/R - k^2R/2$ | [1] | $C^{2,1}$ |
| 3 | $1/R - k^2R/2 + k^4R^3/24$ | this note | $C^{4,1}$ |

The error then decreases algebraically in the order $q$, and the faster, the larger $N$. The lemma is not a quadrature error estimate; for that theory see [2]. On the cell pairs that do not touch, $g$ is real-analytic on a neighbourhood of the domain, and the GL error decreases exponentially.

**Corollary 3.1 (Exact decomposition of the touching moments).** For every $N \geq 1$, $\boldsymbol{d} \in \mathcal{T}$ and $\alpha, \beta \in \{0, 1\}$,

$$M^{[g]}_{\alpha\beta}(\boldsymbol{d}) = M^{[g_N]}_{\alpha\beta}(\boldsymbol{d}) + \sum_{\ell=0}^{N-1} \frac{(-1)^\ell k^{2\ell}}{(2\ell)!}\, M^{(2\ell-1)}_{\alpha\beta}(\boldsymbol{d}), \tag{6}$$

and likewise for the $N$-moments. The first term is evaluated with GL; the static moments $M^{(p)}_{\alpha\beta}(\boldsymbol{d})$, $p \in \{-1, 1, 3\}$, are given in closed form by Theorem (6.1).

*Proof.* The moments are linear in $K$, and every term is absolutely integrable: the kernels $R^{2\ell-1}$ with $\ell \geq 0$ have $2\ell - 1 \geq -1 > -2$, and $g_N$ is bounded. □

**What has to be computed.** By Lemma (2.1), the static moments of the four offsets

$$\boldsymbol{d} \in \{(0,0),\ (1,0),\ (0,1),\ (1,1)\}$$

determine all nine touching offsets. These are the coincident cell, the two kinds of shared edge and the shared vertex. For $N = 3$ we need, for each of them, the four $x$-moments $M_{00}, M_{10}, M_{01}, M_{11}$ and the three $y$-moments, with $p \in \{-1, 1, 3\}$. The $y$-moments follow from the $x$-moments by Lemma (2.1)(iii).

# 4 Reduction to two dimensions

The integrand of a moment depends on the four coordinates only through the two differences $x - x'$ and $y - y'$, and the weights are polynomials. Integrating out the "position along the diagonal" in each direction leaves a two-dimensional integral with piecewise polynomial weights. The coincident-cell transformation of [1] (Proposition *singinteval*, regions I–IV), which follows [3], is the special case $\boldsymbol{d} = \mathbf{0}$.

**Lemma 4.1 (One-dimensional correlation).** Let $\varphi, \psi \in L^\infty(0,1)$, $\delta \in \mathbb{R}$, and let $h : \mathbb{R} \to \mathbb{C}$ be locally integrable. Then

$$\int_0^1\!\!\int_0^1 \varphi(\xi)\,\psi(\tau)\,h(\xi-\tau-\delta)\,\mathrm{d}\tau\,\mathrm{d}\xi = \int_{-1}^{1} W_{\varphi\psi}(\nu)\,h(\nu-\delta)\,\mathrm{d}\nu,$$
$$W_{\varphi\psi}(\nu) = \int_{\max(0,\nu)}^{\min(1,1+\nu)} \varphi(\xi)\,\psi(\xi-\nu)\,\mathrm{d}\xi, \tag{7}$$

and both integrals converge absolutely.

*Proof.* The linear map $(\xi,\tau) \mapsto (\xi,\nu)$, $\nu = \xi - \tau$, has determinant one. It maps the unit square onto the parallelogram (Figure 3)

$$\{(\xi,\nu) :\ -1 \le \nu \le 1,\ \max(0,\nu) \le \xi \le \min(1,1+\nu)\},$$

because $0 \le \tau = \xi - \nu \le 1$ is equivalent to $\nu \le \xi \le 1+\nu$.

By Tonelli's theorem applied to $|\varphi|\,|\psi|\,|h|$, the left-hand side is absolutely convergent, since

$$\int_0^1\!\!\int_0^1 |\varphi(\xi)\,\psi(\tau)\,h(\xi-\tau-\delta)|\,\mathrm{d}\tau\,\mathrm{d}\xi \le \|\varphi\|_\infty\,\|\psi\|_\infty \int_{-1}^{1} (1-|\nu|)\,|h(\nu-\delta)|\,\mathrm{d}\nu < \infty.$$

Fubini's theorem then allows the integration in $\xi$ first, for fixed $\nu$, which gives $W_{\varphi\psi}(\nu)$. □

**Corollary 4.1 (Weights of the rooftop moments).** For $\varphi(\xi) = \xi^\alpha$ and $\psi(\tau) = \tau^\beta$, $\alpha, \beta \in \{0,1\}$, write $W_{\alpha\beta} = W_{\varphi\psi}$. On $I_- = [-1,0]$ and $I_+ = [0,1]$ these are the polynomials

| | $\nu \in I_-$ | $\nu \in I_+$ |
|---|---|---|
| $W_{00}$ | $1+\nu$ | $1-\nu$ |
| $W_{10}$ | $\frac{1}{2}(1+\nu)^2$ | $\frac{1}{2}(1-\nu^2)$ |
| $W_{01}$ | $\frac{1}{2}(1-\nu^2)$ | $\frac{1}{2}(1-\nu)^2$ |
| $W_{11}$ | $\frac{1}{6}(1+\nu)^2(2-\nu)$ | $\frac{1}{6}(2-3\nu+\nu^3)$ |

(8)

They are continuous on $[-1,1]$, $W_{00}$ and $W_{11}$ are even, and $W_{01}(\nu) = W_{10}(-\nu)$.

*Proof.* For $\nu \in I_+$ the limits in (7) are $\nu \le \xi \le 1$:

$$W_{00} = \int_\nu^1 \mathrm{d}\xi = 1-\nu, \qquad W_{10} = \int_\nu^1 \xi\,\mathrm{d}\xi = \frac{1}{2}(1-\nu^2),$$
$$W_{01} = \int_\nu^1 (\xi-\nu)\,\mathrm{d}\xi = \frac{1}{2}(1-\nu)^2, \qquad W_{11} = \int_\nu^1 \xi(\xi-\nu)\,\mathrm{d}\xi = \frac{1}{6}\big(2-3\nu+\nu^3\big).$$

For $\nu \in I_-$ they are $0 \le \xi \le 1+\nu$:

$$W_{00} = \int_0^{1+\nu} \mathrm{d}\xi = 1+\nu, \qquad W_{10} = \int_0^{1+\nu} \xi \,\mathrm{d}\xi = \frac{1}{2}(1+\nu)^2,$$

$$W_{01} = \int_0^{1+\nu} (\xi-\nu) \,\mathrm{d}\xi = \frac{1}{2}(1-\nu^2), \qquad W_{11} = \int_0^{1+\nu} \xi(\xi-\nu) \,\mathrm{d}\xi = \frac{1}{6}(1+\nu)^2(2-\nu).$$

At $\nu = 0$ both sides give the values 1, 1/2, 1/2 and 1/3, so the weights are continuous. For the symmetries, the substitution $s = \xi + \nu$ in (7) gives $W_{\varphi\psi}(-\nu) = W_{\psi\varphi}(\nu)$. Hence $W_{01}(\nu) = W_{10}(-\nu)$, and $W_{00}$, $W_{11}$ are even; for $W_{11}$ compare

$$\frac{1}{6}(1-\nu)^2(2+\nu) = \frac{1}{6}\big(2 - 3\nu + \nu^3\big).$$

□





Figure 3: Left: the substitution $\nu = \xi - \tau$ of Lemma (4.1) maps the unit square onto a parallelogram; the line $\xi - \tau = \nu$ (red) becomes the horizontal segment over which $W_{\varphi\psi}(\nu)$ integrates. The two triangles $\nu \in I_-$ and $\nu \in I_+$ have different limits in $\xi$, which is why the weights change their polynomial form at $\nu = 0$. Right: the weights (8).

**Theorem 4.1 (Two-dimensional representation).** Let $p > -2$, $\boldsymbol{d} = (d_x, d_y) \in \mathbb{Z}^2$ and $\alpha, \beta \in \{0, 1\}$, and put

$$\rho_{\boldsymbol{d}}(\nu, \eta) = \sqrt{a^2(\nu - d_x)^2 + b^2(\eta - d_y)^2}.$$

Then

$$\begin{aligned} M^{(p)}_{\alpha\beta}(\boldsymbol{d}) &= \int_{-1}^{1}\int_{-1}^{1} W_{\alpha\beta}(\nu)\, W_{00}(\eta)\, \rho_{\boldsymbol{d}}(\nu,\eta)^p \,\mathrm{d}\eta \,\mathrm{d}\nu, \\ N^{(p)}_{\alpha\beta}(\boldsymbol{d}) &= \int_{-1}^{1}\int_{-1}^{1} W_{00}(\nu)\, W_{\alpha\beta}(\eta)\, \rho_{\boldsymbol{d}}(\nu,\eta)^p \,\mathrm{d}\eta \,\mathrm{d}\nu. \end{aligned} \tag{9}$$

The same holds for a general admissible kernel $K$, with $K(\rho_{\boldsymbol{d}})$ in place of $\rho^p_{\boldsymbol{d}}$.

*Proof.* Parametrize $\boldsymbol{r} = (a\xi_1, b\eta_1) \in \Omega_{\boldsymbol{0}}$ and $\boldsymbol{r}' = \big(a(d_x + \xi_2),\, b(d_y + \eta_2)\big) \in \Omega_{\boldsymbol{d}}$ with $\xi_1, \eta_1, \xi_2, \eta_2 \in [0, 1]$. Then $\mathrm{d}A'\,\mathrm{d}A = (ab)^2\,\mathrm{d}\eta_2\,\mathrm{d}\xi_2\,\mathrm{d}\eta_1\,\mathrm{d}\xi_1$, which cancels the normalization of Definition (2.3), $\xi(\boldsymbol{r}) = \xi_1$, $\xi(\boldsymbol{r}') = \xi_2$, and

$$\|\boldsymbol{r} - \boldsymbol{r}'\| = \sqrt{a^2(\xi_1 - \xi_2 - d_x)^2 + b^2(\eta_1 - \eta_2 - d_y)^2}.$$

*Absolute convergence.* The function $\boldsymbol{z} \mapsto \|\boldsymbol{z}\|^p$ is locally integrable on $\mathbb{R}^2$ for $p > -2$ (polar coordinates). Lemma (4.1), applied in each direction to the nonnegative kernel by Tonelli's theorem, gives

$$\iiiint |\cdot| = \int_{-1}^{1}\int_{-1}^{1} (1-|\nu|)(1-|\eta|)\,\rho_{\boldsymbol{d}}^{p} \le \int_{-1}^{1}\int_{-1}^{1} \rho_{\boldsymbol{d}}^{p} < \infty.$$

*Reduction.* By Fubini's theorem, fix $(\eta_1, \eta_2)$ with $\eta_1 - \eta_2 - d_y \neq 0$ (almost every pair) and apply Lemma (4.1) in $(\xi_1, \xi_2)$ with $\varphi(\xi) = \xi^{\alpha}$, $\psi(\tau) = \tau^{\beta}$, $\delta = d_x$ and

$$h(s) = \big(a^2 s^2 + b^2(\eta_1 - \eta_2 - d_y)^2\big)^{p/2}.$$

This gives the inner integral $\int_{-1}^{1} W_{\alpha\beta}(\nu)\, h(\nu - d_x)\,\mathrm{d}\nu$. Applying the lemma once more in $(\eta_1, \eta_2)$, with the weights 1 and 1, gives the factor $W_{00}(\eta)$.

The $N$-moments are the same with the roles of the two directions exchanged, and the general kernel changes nothing in the argument. □

**Remark.** For $\boldsymbol{d} \in \mathcal{T}$ the singular point of $\rho_{\boldsymbol{d}}^{p}$, $p < 0$, is $(\nu, \eta) = (d_x, d_y)$. It is the centre, a midpoint of an edge or a corner of $[-1, 1]^2$ (Figure 5). In every case it lies on the lines $\nu \in \{-1, 0, 1\}$ and $\eta \in \{-1, 0, 1\}$, where the weights change their polynomial form. This is the structural fact behind the closed forms of Section 6.

# 5 Mixed antiderivatives

The antiderivatives below are functions of two lengths, which are written as $a$ and $b$ like in the thesis appendix. This does not clash with the cell sizes: the touching moments evaluate them only at the points $(a, b)$, $(2a, b)$, $(a, 2b)$ and $(2a, 2b)$ (Section 6).

**Definition 5.1 (Mixed antiderivatives).** Let $Q = [0, \infty)^2$ be the closed first quadrant and $Q^{\circ}$ its interior. For $p > -2$, $m, n \in \mathbb{N}_0$ and $(a, b) \in Q$ put

$$G_{mn}^{(p)}(a, b) = \int_{0}^{a}\int_{0}^{b} u^m v^n \big(u^2 + v^2\big)^{p/2}\,\mathrm{d}v\,\mathrm{d}u. \tag{10}$$

For a function $F$ on $Q$ and $0 \le a_0 \le a_1$, $0 \le b_0 \le b_1$, the corner difference is

$$\Delta F\,[a_0, a_1] \times [b_0, b_1] = F(a_1, b_1) - F(a_0, b_1) - F(a_1, b_0) + F(a_0, b_0).$$

**Proposition 5.1 (Properties of $G$).** Let $p > -2$ and $m, n \in \mathbb{N}_0$.

(i) $G_{mn}^{(p)}$ is continuous on $Q$ and vanishes on both axes: $G_{mn}^{(p)}(a, 0) = G_{mn}^{(p)}(0, b) = 0$.

(ii) For every rectangle $[a_0, a_1] \times [b_0, b_1] \subset Q$,

$$\int_{a_0}^{a_1}\int_{b_0}^{b_1} u^m v^n \big(u^2 + v^2\big)^{p/2}\,\mathrm{d}v\,\mathrm{d}u = \Delta G_{mn}^{(p)}\,[a_0, a_1] \times [b_0, b_1]. \tag{11}$$

(iii) $\partial_a \partial_b G_{mn}^{(p)} = a^m b^n \big(a^2 + b^2\big)^{p/2}$ on $Q^{\circ}$, and $G_{mn}^{(p)}$ is the only function in $C(Q) \cap C^2(Q^{\circ})$ with this property that vanishes on both axes.





Figure 4: Left: the corner difference (11). The integral over the shaded rectangle is the alternating sum of $G^{(p)}_{mn}$ at its corners, each value being the integral over $[0,a]\times[0,b]$ (dashed). The circles are level lines of $\sqrt{u^2+v^2}$ around the singular point (red). Right: a piece that touches the singular point. Three of its corners lie on the axes, where $G^{(p)}_{mn}$ vanishes, and only $G^{(p)}_{mn}(a,b)$ remains.

(iv) $G^{(p)}_{mn}(\lambda a,\lambda b)=\lambda^{m+n+p+2}\,G^{(p)}_{mn}(a,b)$ for $\lambda>0$.

*Proof.* (i) The integrand is locally integrable on $Q$ because $p>-2$; continuity follows by dominated convergence, and the integral over a rectangle with a side of length zero is zero.

(ii) This is inclusion–exclusion (Figure 4): the rectangle $[a_0,a_1]\times[b_0,b_1]$ is $[0,a_1]\times[0,b_1]$ minus $[0,a_0]\times[0,b_1]$ and $[0,a_1]\times[0,b_0]$, plus their overlap $[0,a_0]\times[0,b_0]$.

(iii) The derivative follows from the fundamental theorem of calculus in each variable. If $D$ is the difference of two such functions, $\partial_a\partial_b D=0$ gives $D(a,b)=e(a)+c(b)$ on $Q^\circ$; letting $b\to0^+$ and $a\to0^+$ with $D=0$ on the axes shows that $e$ and $c$ are constants with $e+c=D(0,0)=0$.

(iv) Substitute $u=\lambda u'$ and $v=\lambda v'$ in (10). □

**Lemma 5.1 (One-dimensional integrals).** Let $a\ge0$, $b>0$, $m\in\mathbb{N}_0$ and $q\in\{-1,1,3,5\}$, and let $R=\sqrt{a^2+b^2}$. The integrals

$$K^{(q)}_m(a)=\int_0^a u^m\bigl(u^2+b^2\bigr)^{q/2}\,\mathrm{d}u$$

are

$$K^{(q)}_0=\begin{cases}\operatorname{arsinh}\dfrac{a}{b}, & q=-1,\\[2mm] \dfrac12\Bigl(aR+b^2\operatorname{arsinh}\dfrac{a}{b}\Bigr), & q=1,\\[2mm] \dfrac18\Bigl(2aR^3+3b^2aR+3b^4\operatorname{arsinh}\dfrac{a}{b}\Bigr), & q=3,\\[2mm] \dfrac{1}{48}\Bigl(8aR^5+10b^2aR^3+15b^4aR+15b^6\operatorname{arsinh}\dfrac{a}{b}\Bigr), & q=5,\end{cases}\tag{12}$$

$$K^{(q)}_1=\frac{R^{q+2}-b^{q+2}}{q+2},\tag{13}$$

$$K^{(q)}_m=\frac{a^{m-1}R^{q+2}-(m-1)b^2K^{(q)}_{m-2}}{m+q+1},\qquad m\ge2,\tag{14}$$

and, for the logarithmic terms produced by the inner integrations,

$$\int_0^a u^m \operatorname{arsinh}\frac{b}{u}\,\mathrm{d}u = \frac{1}{m+1}\Big(a^{m+1}\operatorname{arsinh}\frac{b}{a} + b\,K_m^{(-1)}(a)\Big). \tag{15}$$

The same formulas with $a$ and $b$ exchanged give the inner integrals $\int_0^b v^n(u^2+v^2)^{p/2}\,\mathrm{d}v$.

*Proof.* *Base cases.* They follow from

$$\frac{\mathrm{d}}{\mathrm{d}u}\operatorname{arsinh}\frac{u}{b} = \big(u^2+b^2\big)^{-1/2}, \qquad \frac{\mathrm{d}}{\mathrm{d}u}\big(u^2+b^2\big)^{(q+2)/2} = (q+2)\,u\,\big(u^2+b^2\big)^{q/2},$$
$$\frac{\mathrm{d}}{\mathrm{d}u}\Big(u\big(u^2+b^2\big)^{q/2}\Big) = (q+1)\big(u^2+b^2\big)^{q/2} - qb^2\big(u^2+b^2\big)^{(q-2)/2};$$

the first one gives $K_0^{(-1)}$ and the second one (13). The last one gives the recursion

$$K_0^{(q)} = \frac{aR^q + qb^2K_0^{(q-2)}}{q+1},$$

which produces (12) from $K_0^{(-1)}$.

*Recursion.* For $m \geq 2$ integrate from 0 to $a$

$$\frac{\mathrm{d}}{\mathrm{d}u}\big(u^{m-1}(u^2+b^2)^{(q+2)/2}\big) = (m-1)b^2u^{m-2}(u^2+b^2)^{q/2} + (m+q+1)u^m(u^2+b^2)^{q/2}.$$

*Logarithmic terms.* Integrate by parts with

$$\frac{\mathrm{d}}{\mathrm{d}u}\operatorname{arsinh}\frac{b}{u} = -\frac{b}{u\sqrt{u^2+b^2}};$$

the boundary term $u^{m+1}\operatorname{arsinh}(b/u)$ vanishes as $u \to 0^+$. □

**Proposition 5.2 (Closed forms).** For $p \in \{-1,1,3\}$, $m \in \{0,1,2,3\}$, $n \in \{0,1\}$ and $(a,b) \in Q^\circ$,

$$G_{mn}^{(p)}(a,b) = \frac{1}{D}\left(P_0 + P_1\sqrt{a^2+b^2} + P_2\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right) + P_3\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)\right), \tag{16}$$

with an integer $D$ and homogeneous integer polynomials $P_i(a,b)$, listed in Appendix A. These are all the antiderivatives the moments need; any other one follows from the symmetry $G_{nm}^{(p)}(b,a) = G_{mn}^{(p)}(a,b)$.

*Proof.* *Construction.* Integrate first in $v$ and then in $u$ with Lemma (5.1). Every step produces only terms of the form (16).

*Verification.* Each formula of Appendix A is continuous on $Q$, vanishes on the axes and satisfies $\partial_a\partial_b F = a^m b^n(a^2+b^2)^{p/2}$; these properties were checked by exact symbolic computation (SymPy). Hence $F = G_{mn}^{(p)}$ by Proposition (5.1)(iii). All formulas also agree with 30-digit adaptive quadrature of (10) within $2\cdot 10^{-25}$. □

**Example.** For $p=-1$, (15) gives directly

$$G_{00}^{(-1)}(a,b) = \int_0^a \operatorname{arsinh}\frac{b}{u}\,\mathrm{d}u = a\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) + b\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right),$$

the classical potential integral of a uniformly charged rectangle at its corner.

# 6 Closed forms for the touching cells

Theorem (4.1) turns a touching moment into an integral over $[-1,1]^2$. In each direction the square splits into $I_- = [-1,0]$ and $I_+ = [0,1]$, where the weights are polynomials. Each of the four pieces becomes a rectangle of $Q$ with the singular point at the origin (Figure 5), and Proposition (5.1)(ii) evaluates it exactly.

**Lemma 6.1 (The polynomial pieces).** Let $d \in \{0,1\}$ and $s \in \{-,+\}$. On $\nu \in I_s$ the quantity $\nu - d$ has a constant sign $\sigma$, and $u = \sigma a(\nu - d) \geq 0$ ranges over $[0,a]$ on both pieces if $d = 0$ ($\sigma = \mp 1$ on $I_\mp$), and over $[a,2a]$ on $I_-$ and $[0,a]$ on $I_+$ if $d = 1$ ($\sigma = -1$ on both). The singular value $\nu = d$ is $u = 0$. Every weight of (8) is on each piece a polynomial in $u$ of degree at most three, obtained by substituting $\nu = \sigma u/a + d$ (Table 2). The $y$-direction is the same with $v$, $b$ and $d_y$.

*Proof.* $\nu - d \in [-1,0]$ or $[0,1]$ for $d = 0$, and $\nu - d \in [-2,-1]$ or $[-1,0]$ for $d = 1$. The polynomials of Table 2 are the expansions of (8). □





Figure 5: The integration regions of the four touching configurations ($a = b$). Top: the domain $[-1,1]^2$ of Theorem (4.1), split into the pieces $I_s \times I_t$, labelled $s\,t$; the singular point $(\nu,\eta) = (d_x,d_y)$ (red) is the centre, the midpoint of an edge or a corner, and the circles are level lines of $\rho_{\boldsymbol{d}}$. Bottom: the same pieces in the variables $u = \sigma_s a(\nu - d_x)$, $v = \sigma_t b(\eta - d_y)$ of Lemma (6.1). In a direction with $d = 0$ both pieces fold onto $[0,a]$; with $d = 1$ they lie side by side on $[0,a]$ and $[a,2a]$. The singular point is the origin, a corner of every piece that contains it.

**Theorem 6.1 (Touching moments).** Let $p \in \{-1,1,3\}$, $\boldsymbol{d} = (d_x,d_y) \in \{0,1\}^2$ and $\alpha,\beta \in \{0,1\}$. On the pieces $s,t \in \{-,+\}$ write, with Table 2,

$$W_{\alpha\beta}\big|_{I_s} = \sum_{m=0}^{3} c_m^s\, u^m, \qquad W_{00}\big|_{I_t} = \sum_{n=0}^{1} e_n^t\, v^n,$$

and let $A_s \times B_t$ be the image of the piece, with $A_s$ equal to $[0,a]$ or $[a,2a]$ according to $d_x$ and $B_t$ equal to $[0,b]$ or $[b,2b]$ according to $d_y$. Then

$$M_{\alpha\beta}^{(p)}(\boldsymbol{d}) = \frac{1}{ab} \sum_{s,t\in\{-,+\}} \sum_{m=0}^{3} \sum_{n=0}^{1} c_m^s\, e_n^t\, \Delta G_{mn}^{(p)}\, A_s \times B_t, \tag{17}$$

Table 2: The weights (8) on the pieces of Lemma (6.1) ($x$-direction).

| $d_x$ | 0 | 0 | 1 | 1 |
|---|---|---|---|---|
| piece | $I_-$, $\sigma=-1$ | $I_+$, $\sigma=+1$ | $I_-$, $\sigma=-1$ | $I_+$, $\sigma=-1$ |
| $u$ | $[0,\,a]$ | $[0,\,a]$ | $[a,\,2a]$ | $[0,\,a]$ |
| $W_{00}$ | $\dfrac{a-u}{a}$ | $\dfrac{a-u}{a}$ | $\dfrac{2a-u}{a}$ | $\dfrac{u}{a}$ |
| $W_{10}$ | $\dfrac{(a-u)^2}{2a^2}$ | $\dfrac{(a-u)\,(a+u)}{2a^2}$ | $\dfrac{(2a-u)^2}{2a^2}$ | $\dfrac{u\,(2a-u)}{2a^2}$ |
| $W_{01}$ | $\dfrac{(a-u)\,(a+u)}{2a^2}$ | $\dfrac{(a-u)^2}{2a^2}$ | $\dfrac{u\,(2a-u)}{2a^2}$ | $\dfrac{u^2}{2a^2}$ |
| $W_{11}$ | $\dfrac{(2a+u)\,(a-u)^2}{6a^3}$ | $\dfrac{(2a+u)\,(a-u)^2}{6a^3}$ | $\dfrac{(a+u)\,(2a-u)^2}{6a^3}$ | $\dfrac{u^2\,(3a-u)}{6a^3}$ |

where, because $G^{(p)}_{mn}$ vanishes on the axes,

$$\begin{aligned}
\Delta G\,[0,a]\times[0,b] &= G(a,b),\\
\Delta G\,[a,2a]\times[0,b] &= G(2a,b)-G(a,b),\\
\Delta G\,[0,a]\times[b,2b] &= G(a,2b)-G(a,b),\\
\Delta G\,[a,2a]\times[b,2b] &= G(2a,2b)-G(2a,b)-G(a,2b)+G(a,b).
\end{aligned}\tag{18}$$

Every touching moment is therefore an exact, finite combination of $G^{(p)}_{mn}$ at the four points $(a,b)$, $(2a,b)$, $(a,2b)$, $(2a,2b)$. The $N$-moments follow from Lemma (2.1)(iii) with $a$ and $b$ exchanged.

*Proof.* By Theorem (4.1), $M^{(p)}_{\alpha\beta}(\boldsymbol{d})$ is the sum over the four pieces $I_s\times I_t$. On a piece substitute $u=\sigma_s a(\nu-d_x)$ and $v=\sigma_t b(\eta-d_y)$: then $\mathrm{d}\nu\,\mathrm{d}\eta = \mathrm{d}u\,\mathrm{d}v/(ab)$, $\rho_{\boldsymbol{d}}=\sqrt{u^2+v^2}$, the weights become the polynomials of Lemma (6.1), and the piece is $\int_{A_s}\int_{B_t}\sum_{m,n}c^s_m e^t_n u^m v^n (u^2+v^2)^{p/2}\,\mathrm{d}v\,\mathrm{d}u$. Apply (11) term by term; the corners on the axes contribute zero by Proposition (5.1)(i). Each piece is absolutely integrable ($p>-2$), so no limit is involved. □

**Remark.** The same corner-sum construction works for every offset, but for distant cells it gives a sum of large, nearly cancelling terms, whereas GL converges exponentially there. It is meant for the touching offsets only.

**Corollary 6.1 (Homogeneity).** $M^{(p)}_{\alpha\beta}(\boldsymbol{d};\lambda a,\lambda b)=\lambda^p\,M^{(p)}_{\alpha\beta}(\boldsymbol{d};a,b)$ for $\lambda>0$, so the static moments of a grid depend, up to the factor $a^p$, on the aspect ratio $b/a$ only.

*Proof.* In (9), $\rho_{\boldsymbol{d}}$ scales by $\lambda$ and the weights do not depend on $a$ and $b$. □

**Proposition 6.1 (Agreement with the coincident-cell formulas of the thesis).** Consider the functions `type1`, `type2`, `type3` of `src` and $\mathcal{I}^{(1)}$ of the appendix of [1]. Then

$$\begin{aligned}
M^{(-1)}_{00}(\mathbf{0}) &= \texttt{type3}(a,b), & M^{(-1)}_{10}(\mathbf{0})-M^{(-1)}_{11}(\mathbf{0}) &= \texttt{type2}(a,b),\\
M^{(-1)}_{11}(\mathbf{0}) &= \texttt{type1}(a,b), & (ab)^2 M^{(1)}_{00}(\mathbf{0}) &= \mathcal{I}^{(1)}[1],\\
& & (ab)^2 M^{(1)}_{11}(\mathbf{0}) &= \mathcal{I}^{(1)}[\lambda_1\lambda_1'].
\end{aligned}$$

The first one reads

$$M_{00}^{(-1)}(\mathbf{0}) = \frac{1}{3a^2b^2}\Bigg(2a^3 + 2b^3 - 2\sqrt{a^2+b^2}\,\left(a^2+b^2\right) + 6a^2b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) + 6ab^2\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg).$$

*Proof.* By Corollary (6.1) it suffices to take $a = 1$ and keep $b > 0$ symbolic. The difference of the two sides, written with (17) and the cited formula, simplifies to zero exactly (SymPy). □

**Remark (thesis appendix).** The values of the formulas in the appendix of [1] are correct, but three labels are misprinted: the integrands of the $\mathcal{I}^{(1)}$ entries are written with $1/R$ instead of $R$; the last $\mathcal{I}^{(1)}$ entry repeats the label $[\lambda_1(1-\lambda_1')]$ instead of $[\lambda_2(1-\lambda_2')]$; and the integrand of $\mathcal{I}^{(-1)}[\lambda_2(1-\lambda_2')]$ contains $(u+\lambda_2')$ instead of $(v+\lambda_2')$.

# A The mixed antiderivatives $G_{mn}^{(p)}$

This appendix lists the closed forms (16) of the antiderivatives (10) for $p \in \{-1, 1, 3\}$, $m \le 3$ and $n \le 1$, which are the ones Theorem (6.1) needs. Any other one follows from $G_{nm}^{(p)}(b,a) = G_{mn}^{(p)}(a,b)$, and on the axes $G_{mn}^{(p)} = 0$.

**Kernel $R^{-1}$.**

$$G_{00}^{(-1)}(a,b) = a\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) + b\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)$$

$$G_{01}^{(-1)}(a,b) = \frac{1}{2}\left(-a^2 + a\sqrt{a^2+b^2} + b^2\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\right)$$

$$G_{10}^{(-1)}(a,b) = \frac{1}{2}\left(-b^2 + b\sqrt{a^2+b^2} + a^2\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)\right)$$

$$G_{11}^{(-1)}(a,b) = \frac{1}{3}\left(-a^3 - b^3 + \sqrt{a^2+b^2}\,\left(a^2+b^2\right)\right)$$

$$G_{20}^{(-1)}(a,b) = \frac{1}{6}\left(ab\sqrt{a^2+b^2} + 2a^3\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) - b^3\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\right)$$

$$G_{21}^{(-1)}(a,b) = \frac{1}{8}\left(-2a^4 + a\sqrt{a^2+b^2}\,\left(2a^2+b^2\right) - b^4\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\right)$$

$$G_{30}^{(-1)}(a,b) = \frac{1}{12}\left(2b^4 + b\sqrt{a^2+b^2}\,\left(a^2-2b^2\right) + 3a^4\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)\right)$$

$$G_{31}^{(-1)}(a,b) = \frac{1}{15}\left( -3a^5 + 2b^5 + \sqrt{a^2+b^2}\left(3a^4 + a^2b^2 - 2b^4\right)\right)$$

**Kernel $R^1$.**

$$G_{00}^{(1)}(a,b) = \frac{1}{6}\left(2ab\sqrt{a^2+b^2} + a^3\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) + b^3\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\right)$$

$$G_{01}^{(1)}(a,b) = \frac{1}{24}\left( -2a^4 + a\sqrt{a^2+b^2}\left(2a^2+5b^2\right) + 3b^4\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\right)$$

$$G_{10}^{(1)}(a,b) = \frac{1}{24}\left( -2b^4 + b\sqrt{a^2+b^2}\left(5a^2+2b^2\right) + 3a^4\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)\right)$$

$$G_{11}^{(1)}(a,b) = \frac{1}{15}\left( -a^5 - b^5 + \sqrt{a^2+b^2}\left(a^4+2a^2b^2+b^4\right)\right)$$

$$G_{20}^{(1)}(a,b) = \frac{1}{40}\left(ab\sqrt{a^2+b^2}\left(6a^2+b^2\right) + 4a^5\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) - b^5\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\right)$$

$$G_{21}^{(1)}(a,b) = \frac{1}{144}\left( -8a^6 + a\sqrt{a^2+b^2}\left(8a^4+14a^2b^2+3b^4\right) - 3b^6\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\right)$$

$$G_{30}^{(1)}(a,b) = \frac{1}{180}\left(4b^6 + b\sqrt{a^2+b^2}\left(21a^4+2a^2b^2-4b^4\right) + 15a^6\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)\right)$$

$$G_{31}^{(1)}(a,b) = \frac{1}{105}\left( -5a^7 + 2b^7 + \sqrt{a^2+b^2}\left(5a^6+8a^4b^2+a^2b^4-2b^6\right)\right)$$

**Kernel $R^3$.**

$$G_{00}^{(3)}(a,b) = \frac{1}{40}\left(7ab\sqrt{a^2+b^2}\left(a^2+b^2\right) + 3a^5\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)\right.$$

$$+ 3b^5 \log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)$$

$$G_{01}^{(3)}(a,b) = \frac{1}{240}\Bigg(-8a^6 + a\sqrt{a^2+b^2}\left(8a^4+26a^2b^2+33b^4\right) + 15b^6 \log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)$$

$$G_{10}^{(3)}(a,b) = \frac{1}{240}\Bigg(-8b^6 + b\sqrt{a^2+b^2}\left(33a^4+26a^2b^2+8b^4\right) + 15a^6 \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)\Bigg)$$

$$G_{11}^{(3)}(a,b) = \frac{1}{35}\left(-a^7-b^7+\sqrt{a^2+b^2}\left(a^6+3a^4b^2+3a^2b^4+b^6\right)\right)$$

$$G_{20}^{(3)}(a,b) = \frac{1}{336}\Bigg(ab\sqrt{a^2+b^2}\left(38a^4+26a^2b^2+3b^4\right) + 18a^7 \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) - 3b^7 \log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)$$

$$G_{21}^{(3)}(a,b) = \frac{1}{1920}\Bigg(-48a^8 + a\sqrt{a^2+b^2}\left(48a^6+136a^4b^2+118a^2b^4+15b^6\right) - 15b^8 \log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)$$

$$G_{30}^{(3)}(a,b) = \frac{1}{2240}\Bigg(16b^8 + b\sqrt{a^2+b^2}\left(215a^6+134a^4b^2+8a^2b^4-16b^6\right) + 105a^8 \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)\Bigg)$$

$$G_{31}^{(3)}(a,b) = \frac{1}{315}\Bigg(-7a^9+2b^9 + \sqrt{a^2+b^2}\left(7a^8+19a^6b^2+15a^4b^4+a^2b^6-2b^8\right)\Bigg)$$

# B The touching moments

This appendix lists the $x$-moments $M^{(p)}_{\alpha\beta}(\boldsymbol{d})$ of the cell $a \times b$ for $p \in \{-1, 1, 3\}$, $\alpha, \beta \in \{0, 1\}$ and the touching offsets $\boldsymbol{d} \in \{(0,0), (1,0), (0,1), (1,1)\}$. They are the corner sums (17) expanded with SymPy, and each one was checked against its corner sum to 30 digits. The $N$-moments are $N^{(p)}_{\alpha\beta}(d_x, d_y; a, b) = M^{(p)}_{\alpha\beta}(d_y, d_x; b, a)$, and the negative offsets follow from Lemma (2.1)(i) and (ii). For evaluation in double precision the corner sum (17) is the better form; the expanded formulas are meant for reading and checking.

## Kernel $R^{-1}$

**Coincident cell, $\boldsymbol{d} = (0,0)$.**

$$M^{(-1)}_{00}(0,0) = \frac{1}{3a^2b^2}\Bigg(2a^3 + 2b^3 - 2\sqrt{a^2+b^2}\,\left(a^2+b^2\right) + 6a^2b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) + 6ab^2\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)$$

$$M^{(-1)}_{10}(0,0) = \frac{1}{3a^2b^2}\Bigg(a^3 + b^3 - \sqrt{a^2+b^2}\,\left(a^2+b^2\right) + 3a^2b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) + 3ab^2\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)$$

$$M^{(-1)}_{01}(0,0) = \frac{1}{3a^2b^2}\Bigg(a^3 + b^3 - \sqrt{a^2+b^2}\,\left(a^2+b^2\right) + 3a^2b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) + 3ab^2\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)$$

$$M^{(-1)}_{11}(0,0) = \frac{1}{90a^4b^2}\Bigg(12a^5 + 30a^2b^3 + 2b^5 - \sqrt{a^2+b^2}\,\left(12a^4 + 29a^2b^2 + 2b^4\right) + 45a^4b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) + 60a^3b^2\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)$$

**Shared edge parallel to $y$, $\boldsymbol{d} = (1,0)$.**

$$M^{(-1)}_{00}(1,0) = \frac{1}{3a^2b^2}\Bigg(6a^3 - b^3 + 2\sqrt{a^2+b^2}\,\left(a^2+b^2\right) - \sqrt{4a^2+b^2}\,\left(4a^2+b^2\right) + 12a^2b\log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right)$$

$$
\begin{aligned}
&- 6a^2 b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) - 6ab^2 \log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right) \\
&+ 6ab^2 \log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)
\end{aligned}
$$

$$
\begin{aligned}
M_{10}^{(-1)}(1,0) = \frac{1}{24a^3b^2}\Bigg(&20a^4 - 8ab^3 + 2a\sqrt{a^2+b^2}\left(6a^2+11b^2\right) \\
&- 2a\sqrt{4a^2+b^2}\left(8a^2+7b^2\right) \\
&+ 64a^3 b \log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right) \\
&- 40a^3 b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) \\
&- 2b^2\left(24a^2-b^2\right)\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right) \\
&+ b^2\left(48a^2-b^2\right)\log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)
\end{aligned}
$$

$$
\begin{aligned}
M_{01}^{(-1)}(1,0) = \frac{1}{24a^3b^2}\Bigg(&28a^4 + 2a\sqrt{a^2+b^2}\left(2a^2-3b^2\right) \\
&- 2a\sqrt{4a^2+b^2}\left(8a^2-3b^2\right) \\
&+ 32a^3 b \log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right) - 8a^3 b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) \\
&- 2b^4 \log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right) + b^4 \log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)
\end{aligned}
$$

$$
\begin{aligned}
M_{11}^{(-1)}(1,0) = \frac{1}{360a^4b^2}\Bigg(&180a^5 - 4b^5 + 2\sqrt{a^2+b^2}\left(54a^4+13a^2b^2+4b^4\right) \\
&- 2\sqrt{4a^2+b^2}\left(72a^4+11a^2b^2+2b^4\right) \\
&+ 480a^4 b \log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right) \\
&- 300a^4 b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) \\
&- 30ab^2\left(8a^2+b^2\right)\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right) \\
&+ 15ab^2\left(16a^2+b^2\right)\log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)
\end{aligned}
$$

**Shared edge parallel to $x$, $\boldsymbol{d} = (0,1)$.**

$$
\begin{aligned}
M_{00}^{(-1)}(0,1) = \frac{1}{3a^2b^2}\Bigg( & -a^3 + 6b^3 + 2\sqrt{a^2+b^2}\left(a^2+b^2\right) \\
& - \sqrt{a^2+4b^2}\left(a^2+4b^2\right) - 6a^2b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) \\
& + 6a^2b\log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right) \\
& + 12ab^2\log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right) - 6ab^2\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)
\end{aligned}
$$

$$
\begin{aligned}
M_{10}^{(-1)}(0,1) = \frac{1}{6a^2b^2}\Bigg( & -a^3 + 6b^3 + 2\sqrt{a^2+b^2}\left(a^2+b^2\right) \\
& - \sqrt{a^2+4b^2}\left(a^2+4b^2\right) - 6a^2b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) \\
& + 6a^2b\log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right) \\
& + 12ab^2\log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right) - 6ab^2\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)
\end{aligned}
$$

$$
\begin{aligned}
M_{01}^{(-1)}(0,1) = \frac{1}{6a^2b^2}\Bigg( & -a^3 + 6b^3 + 2\sqrt{a^2+b^2}\left(a^2+b^2\right) \\
& - \sqrt{a^2+4b^2}\left(a^2+4b^2\right) - 6a^2b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) \\
& + 6a^2b\log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right) \\
& + 12ab^2\log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right) - 6ab^2\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)
\end{aligned}
$$

$$
\begin{aligned}
M_{11}^{(-1)}(0,1) = \frac{1}{90a^4b^2}\Bigg( & -6a^5 + 90a^2b^3 + 30b^5 \\
& + \sqrt{a^2+b^2}\left(12a^4 + 29a^2b^2 + 2b^4\right) \\
& - 2\sqrt{a^2+4b^2}\left(3a^4 + 29a^2b^2 + 8b^4\right) \\
& - 45a^4b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) \\
& + 45a^4b\log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right)
\end{aligned}
$$

$$+ 120a^3b^2 \log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right)$$
$$- 60a^3b^2 \log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)$$

**Shared vertex, $\boldsymbol{d} = (1,1)$.**

$$M_{00}^{(-1)}(1,1) = \frac{1}{3a^2b^2}\Bigg(-3a^3-3b^3-6\sqrt{a^2+b^2}\left(a^2+b^2\right)$$
$$+\sqrt{a^2+4b^2}\left(a^2+4b^2\right)+\sqrt{4a^2+b^2}\left(4a^2+b^2\right)$$
$$- 12a^2b\log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right)$$
$$+ 18a^2b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)$$
$$- 6a^2b\log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right)$$
$$- 12ab^2\log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right)$$
$$+ 18ab^2\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)$$
$$- 6ab^2\log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)$$

$$M_{10}^{(-1)}(1,1) = \frac{1}{24a^3b^2}\Bigg(-10a^4-24ab^3-2a\sqrt{a^2+b^2}\left(14a^2+39b^2\right)$$
$$+2a\sqrt{a^2+4b^2}\left(3a^2+22b^2\right)$$
$$+2a\sqrt{4a^2+b^2}\left(8a^2+7b^2\right)$$
$$- 64a^3b\log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right)$$
$$+ 104a^3b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)$$
$$- 40a^3b\log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right)$$
$$- 16b^2\left(6a^2-b^2\right)\log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right)$$

$$
\begin{aligned}
&+ 2b^2\left(72a^2 - 5b^2\right)\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\\
&- b^2\left(48a^2 - b^2\right)\log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)
\end{aligned}
$$

$$
\begin{aligned}
M_{01}^{(-1)}(1,1) = \frac{1}{24a^3b^2}\Bigg(&- 14a^4 - 10a\sqrt{a^2+b^2}\left(2a^2 - 3b^2\right)\\
&+ 2a\sqrt{a^2+4b^2}\left(a^2 - 6b^2\right)\\
&+ 2a\sqrt{4a^2+b^2}\left(8a^2 - 3b^2\right)\\
&- 32a^3b\log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right)\\
&+ 40a^3b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)\\
&- 8a^3b\log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right)\\
&- 16b^4\log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right) + 10b^4\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\\
&- b^4\log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)
\end{aligned}
$$

$$
\begin{aligned}
M_{11}^{(-1)}(1,1) = \frac{1}{360a^4b^2}\Bigg(&- 90a^5 - 60b^5 - 6\sqrt{a^2+b^2}\left(42a^4 + 19a^2b^2 + 12b^4\right)\\
&+ 2\sqrt{a^2+4b^2}\left(27a^4 + 26a^2b^2 + 32b^4\right)\\
&+ 2\sqrt{4a^2+b^2}\left(72a^4 + 11a^2b^2 + 2b^4\right)\\
&- 480a^4b\log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right)\\
&+ 780a^4b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)\\
&- 300a^4b\log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right)\\
&- 240ab^2\left(2a^2+b^2\right)\log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right)\\
&+ 30ab^2\left(24a^2+5b^2\right)\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)
\end{aligned}
$$

$$- 15ab^2\left(16a^2+b^2\right)\log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)$$

## Kernel $R^1$

**Coincident cell, $\boldsymbol{d}=(0,0)$.**

$$M_{00}^{(1)}(0,0)=\frac{1}{30a^2b^2}\Bigg(2a^5+2b^5-2\sqrt{a^2+b^2}\left(a^4-3a^2b^2+b^4\right)$$
$$+5a^4b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)+5ab^4\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)$$

$$M_{10}^{(1)}(0,0)=\frac{1}{60a^2b^2}\Bigg(2a^5+2b^5-2\sqrt{a^2+b^2}\left(a^4-3a^2b^2+b^4\right)$$
$$+5a^4b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)+5ab^4\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)$$

$$M_{01}^{(1)}(0,0)=\frac{1}{60a^2b^2}\Bigg(2a^5+2b^5-2\sqrt{a^2+b^2}\left(a^4-3a^2b^2+b^4\right)$$
$$+5a^4b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)+5ab^4\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)$$

$$M_{11}^{(1)}(0,0)=\frac{1}{3780a^4b^2}\Bigg(36a^7+126a^2b^5+8b^7$$
$$-\sqrt{a^2+b^2}\left(36a^6-165a^4b^2+122a^2b^4+8b^6\right)$$
$$+105a^6b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)$$
$$+210a^3b^4\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)$$

**Shared edge parallel to $y$, $\boldsymbol{d}=(1,0)$.**

$$M_{00}^{(1)}(1,0)=\frac{1}{30a^2b^2}\Bigg(30a^5-b^5+2\sqrt{a^2+b^2}\left(a^4-3a^2b^2+b^4\right)$$
$$-\sqrt{4a^2+b^2}\left(16a^4-12a^2b^2+b^4\right)$$
$$+40a^4b\log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right)-5a^4b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)$$
$$-5ab^4\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)+5ab^4\log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)$$

$$
\begin{aligned}
M_{10}^{(1)}(1,0) = \frac{1}{720a^3b^2}\Bigg( & 224a^6 - 24ab^5 \\
& + 2a\sqrt{a^2+b^2}\left(16a^4 - 56a^2b^2 + 33b^4\right) \\
& - 2a\sqrt{4a^2+b^2}\left(64a^4 - 80a^2b^2 + 21b^4\right) \\
& + 384a^5 b \log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right) \\
& - 84a^5 b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) \\
& - 6b^4\left(20a^2 - b^2\right)\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right) \\
& + 3b^4\left(40a^2 - b^2\right)\log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)
\end{aligned}
$$

$$
\begin{aligned}
M_{01}^{(1)}(1,0) = \frac{1}{720a^3b^2}\Bigg( & 496a^6 + 2a\sqrt{a^2+b^2}\left(8a^4 - 16a^2b^2 - 9b^4\right) \\
& - 2a\sqrt{4a^2+b^2}\left(128a^4 - 64a^2b^2 - 9b^4\right) \\
& + 576a^5 b \log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right) \\
& - 36a^5 b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) - 6b^6 \log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right) \\
& + 3b^6 \log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)
\end{aligned}
$$

$$
\begin{aligned}
M_{11}^{(1)}(1,0) = \frac{1}{15120a^4b^2}\Bigg( & 3360a^7 - 16b^7 \\
& + 2\sqrt{a^2+b^2}\left(240a^6 - 666a^4b^2 + 55a^2b^4 + 16b^6\right) \\
& - 2\sqrt{4a^2+b^2}\left(960a^6 - 960a^4b^2 + 47a^2b^4 + 8b^6\right) \\
& + 5376a^6 b \log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right) \\
& - 1176a^6 b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) \\
& - 42ab^4\left(20a^2 + 3b^2\right)\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)
\end{aligned}
$$

$$+ 21ab^4\left(40a^2+3b^2\right)\log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)$$

**Shared edge parallel to $x$, $\boldsymbol{d}=(0,1)$.**

$$\begin{aligned}
M_{00}^{(1)}(0,1) = \frac{1}{30a^2b^2}\Bigg( & -a^5+30b^5+2\sqrt{a^2+b^2}\left(a^4-3a^2b^2+b^4\right)\\
& -\sqrt{a^2+4b^2}\left(a^4-12a^2b^2+16b^4\right)\\
& -5a^4b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)+5a^4b\log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right)\\
& +40ab^4\log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right)-5ab^4\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)\\
M_{10}^{(1)}(0,1) = \frac{1}{60a^2b^2}\Bigg( & -a^5+30b^5+2\sqrt{a^2+b^2}\left(a^4-3a^2b^2+b^4\right)\\
& -\sqrt{a^2+4b^2}\left(a^4-12a^2b^2+16b^4\right)\\
& -5a^4b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)+5a^4b\log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right)\\
& +40ab^4\log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right)-5ab^4\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)\\
M_{01}^{(1)}(0,1) = \frac{1}{60a^2b^2}\Bigg( & -a^5+30b^5+2\sqrt{a^2+b^2}\left(a^4-3a^2b^2+b^4\right)\\
& -\sqrt{a^2+4b^2}\left(a^4-12a^2b^2+16b^4\right)\\
& -5a^4b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)+5a^4b\log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right)\\
& +40ab^4\log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right)-5ab^4\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)\\
M_{11}^{(1)}(0,1) = \frac{1}{3780a^4b^2}\Bigg( & -18a^7+1890a^2b^5+504b^7\\
& +\sqrt{a^2+b^2}\left(36a^6-165a^4b^2+122a^2b^4+8b^6\right)\\
& -2\sqrt{a^2+4b^2}\left(9a^6-165a^4b^2+488a^2b^4+128b^6\right)\\
& -105a^6b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)
\end{aligned}$$

$$
\begin{aligned}
&+ 105a^6 b \log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right) \\
&+ 1680a^3b^4 \log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right) \\
&- 210a^3b^4 \log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)
\end{aligned}
$$

**Shared vertex, $\boldsymbol{d} = (1,1)$.**

$$
\begin{aligned}
M_{00}^{(1)}(1,1) = \frac{1}{30a^2b^2}\Bigg( &- 15a^5 - 15b^5 - 18\sqrt{a^2+b^2}\left(a^4 - 3a^2b^2 + b^4\right) \\
&+ \sqrt{a^2+4b^2}\left(a^4 - 12a^2b^2 + 16b^4\right) \\
&+ \sqrt{4a^2+b^2}\left(16a^4 - 12a^2b^2 + b^4\right) \\
&- 40a^4 b \log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right) \\
&+ 45a^4 b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) \\
&- 5a^4 b \log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right) \\
&- 40ab^4 \log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right) \\
&+ 45ab^4 \log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right) \\
&- 5ab^4 \log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)
\end{aligned}
$$

$$
\begin{aligned}
M_{10}^{(1)}(1,1) = \frac{1}{720a^3b^2}\Bigg( &- 112a^6 - 360ab^5 \\
&- 2a\sqrt{a^2+b^2}\left(80a^4 - 376a^2b^2 + 369b^4\right) \\
&+ 16a\sqrt{a^2+4b^2}\left(a^4 - 14a^2b^2 + 33b^4\right) \\
&+ 2a\sqrt{4a^2+b^2}\left(64a^4 - 80a^2b^2 + 21b^4\right) \\
&- 384a^5 b \log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right)
\end{aligned}
$$

$$
\begin{aligned}
&+ 468a^5 b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)\\
&- 84a^5 b \log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right)\\
&- 192b^4\left(5a^2-b^2\right)\log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right)\\
&+ 6b^4\left(180a^2-17b^2\right)\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\\
&- 3b^4\left(40a^2-b^2\right)\log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)
\end{aligned}
$$

$$
\begin{aligned}
M_{01}^{(1)}(1,1) = \frac{1}{720a^3b^2}\Bigg( &- 248a^6 - 34a\sqrt{a^2+b^2}\left(8a^4-16a^2b^2-9b^4\right)\\
&+ 8a\sqrt{a^2+4b^2}\left(a^4-8a^2b^2-18b^4\right)\\
&+ 2a\sqrt{4a^2+b^2}\left(128a^4-64a^2b^2-9b^4\right)\\
&- 576a^5 b\log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right)\\
&+ 612a^5 b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)\\
&- 36a^5 b\log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right)\\
&- 192b^6\log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right)\\
&+ 102b^6\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right) - 3b^6\log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)
\end{aligned}
$$

$$
\begin{aligned}
M_{11}^{(1)}(1,1) = \frac{1}{15120a^4b^2}\Bigg( &- 1680a^7 - 1008b^7 - 6\sqrt{a^2+b^2}\big(400a^6-1502a^4b^2\\
&+ 269a^2b^4+176b^6\big)\\
&+ 8\sqrt{a^2+4b^2}\left(30a^6-333a^4b^2+110a^2b^4+128b^6\right)\\
&+ 2\sqrt{4a^2+b^2}\left(960a^6-960a^4b^2+47a^2b^4+8b^6\right)\\
&- 5376a^6 b\log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right)
\end{aligned}
$$

$$
\begin{aligned}
&+ 6552a^6 b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)\\
&- 1176a^6 b \log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right)\\
&- 1344ab^4\left(5a^2+3b^2\right)\log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right)\\
&+ 126ab^4\left(60a^2+17b^2\right)\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\\
&- 21ab^4\left(40a^2+3b^2\right)\log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)
\end{aligned}
$$

## Kernel $R^3$

**Coincident cell, $\boldsymbol{d} = (0,0)$.**

$$
\begin{aligned}
M_{00}^{(3)}(0,0) = \frac{1}{420a^2b^2}\Bigg(&8a^7+8b^7\\
&-\sqrt{a^2+b^2}\left(8a^6-25a^4b^2-25a^2b^4+8b^6\right)\\
&+21a^6b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)+21ab^6\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)
\end{aligned}
$$

$$
\begin{aligned}
M_{10}^{(3)}(0,0) = \frac{1}{840a^2b^2}\Bigg(&8a^7+8b^7\\
&-\sqrt{a^2+b^2}\left(8a^6-25a^4b^2-25a^2b^4+8b^6\right)\\
&+21a^6b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)+21ab^6\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)
\end{aligned}
$$

$$
\begin{aligned}
M_{01}^{(3)}(0,0) = \frac{1}{840a^2b^2}\Bigg(&8a^7+8b^7\\
&-\sqrt{a^2+b^2}\left(8a^6-25a^4b^2-25a^2b^4+8b^6\right)\\
&+21a^6b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)+21ab^6\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)
\end{aligned}
$$

$$
\begin{aligned}
M_{11}^{(3)}(0,0) = \frac{1}{30240a^4b^2}\Bigg(&64a^9+288a^2b^7+16b^9-\sqrt{a^2+b^2}\big(64a^8\\
&-275a^6b^2-390a^4b^4+280a^2b^6+16b^8\big)
\end{aligned}
$$

$$+ 189a^8 b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)$$
$$+ 504a^3 b^6 \log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)$$

**Shared edge parallel to $y$, $\boldsymbol{d} = (1,0)$.**

$$M_{00}^{(3)}(1,0) = \frac{1}{420a^2b^2}\Bigg(504a^7 - 4b^7$$
$$+ \sqrt{a^2+b^2}\left(8a^6 - 25a^4b^2 - 25a^2b^4 + 8b^6\right)$$
$$- 2\sqrt{4a^2+b^2}\left(128a^6 - 100a^4b^2 - 25a^2b^4 + 2b^6\right)$$
$$+ 672a^6 b \log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right)$$
$$- 21a^6 b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) - 21ab^6 \log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)$$
$$+ 21ab^6 \log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)$$

$$M_{10}^{(3)}(1,0) = \frac{1}{13440a^3b^2}\Bigg(3936a^8 - 128ab^7$$
$$+ 2a\sqrt{a^2+b^2}\left(80a^6 - 272a^4b^2 - 310a^2b^4 + 177b^6\right)$$
$$- 2a\sqrt{4a^2+b^2}\big(1024a^6 - 1152a^4b^2 - 440a^2b^4$$
$$+ 113b^6\big) + 6144a^7 b \log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right)$$
$$- 432a^7 b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)$$
$$- 6b^6\left(112a^2 - 5b^2\right)\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)$$
$$+ 3b^6\left(224a^2 - 5b^2\right)\log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)$$

$$M_{01}^{(3)}(1,0) = \frac{1}{13440a^3b^2}\Bigg(12192a^8$$
$$+ 2a\sqrt{a^2+b^2}\left(48a^6 - 128a^4b^2 - 90a^2b^4 - 49b^6\right)$$

$$
\begin{aligned}
&- 2a\sqrt{4a^2+b^2}\big(3072a^6 - 2048a^4b^2 - 360a^2b^4 \\
&- 49b^6\big) + 15360a^7 b \log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right) \\
&- 240a^7 b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) - 30b^8 \log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right) \\
&+ 15b^8 \log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)
\end{aligned}
$$

$$
\begin{aligned}
M_{11}^{(3)}(1,0) = \frac{1}{120960a^4b^2}\Bigg(&27552a^9 - 32b^9 + 2\sqrt{a^2+b^2}\big(560a^8 - 1702a^6b^2 \\
&- 1590a^4b^4 + 119a^2b^6 + 32b^8\big) \\
&- 2\sqrt{4a^2+b^2}\big(7168a^8 - 7232a^6b^2 - 2280a^4b^4 \\
&+ 103a^2b^6 + 16b^8\big) + 41472a^8 b \log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right) \\
&- 2916a^8 b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) \\
&- 18ab^6\left(112a^2+15b^2\right)\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right) \\
&+ 9ab^6\left(224a^2+15b^2\right)\log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)
\end{aligned}
$$

**Shared edge parallel to $x$, $\boldsymbol{d} = (0,1)$.**

$$
\begin{aligned}
M_{00}^{(3)}(0,1) = \frac{1}{420a^2b^2}\Bigg(&- 4a^7 + 504b^7 \\
&+ \sqrt{a^2+b^2}\left(8a^6 - 25a^4b^2 - 25a^2b^4 + 8b^6\right) \\
&- 2\sqrt{a^2+4b^2}\left(2a^6 - 25a^4b^2 - 100a^2b^4 + 128b^6\right) \\
&- 21a^6 b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) \\
&+ 21a^6 b \log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right) \\
&+ 672ab^6 \log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right)
\end{aligned}
$$

$$
\begin{aligned}
&\quad - 21ab^6 \log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)\\
M_{10}^{(3)}(0,1) = \frac{1}{840a^2b^2}\Bigg(&- 4a^7 + 504b^7\\
&+ \sqrt{a^2+b^2}\left(8a^6 - 25a^4b^2 - 25a^2b^4 + 8b^6\right)\\
&- 2\sqrt{a^2+4b^2}\left(2a^6 - 25a^4b^2 - 100a^2b^4 + 128b^6\right)\\
&- 21a^6b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)\\
&+ 21a^6b \log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right)\\
&+ 672ab^6 \log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right)\\
&- 21ab^6 \log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)\\
M_{01}^{(3)}(0,1) = \frac{1}{840a^2b^2}\Bigg(&- 4a^7 + 504b^7\\
&+ \sqrt{a^2+b^2}\left(8a^6 - 25a^4b^2 - 25a^2b^4 + 8b^6\right)\\
&- 2\sqrt{a^2+4b^2}\left(2a^6 - 25a^4b^2 - 100a^2b^4 + 128b^6\right)\\
&- 21a^6b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)\\
&+ 21a^6b \log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right)\\
&+ 672ab^6 \log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right)\\
&- 21ab^6 \log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)\\
M_{11}^{(3)}(0,1) = \frac{1}{30240a^4b^2}\Bigg(&- 32a^9 + 18144a^2b^7 + 4080b^9 + \sqrt{a^2+b^2}\big(64a^8\\
&- 275a^6b^2 - 390a^4b^4 + 280a^2b^6 + 16b^8\big)\\
&- 2\sqrt{a^2+4b^2}\big(16a^8 - 275a^6b^2 - 1560a^4b^4
\end{aligned}
$$

$$
\begin{aligned}
&+ 4480a^2b^6 + 1024b^8\big) - 189a^8b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)\\
&+ 189a^8b\log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right)\\
&+ 16128a^3b^6\log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right)\\
&- 504a^3b^6\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\Bigg)
\end{aligned}
$$

**Shared vertex, $\boldsymbol{d} = (1,1)$.**

$$
\begin{aligned}
M_{00}^{(3)}(1,1) = \frac{1}{420a^2b^2}\Bigg(&- 252a^7 - 252b^7\\
&- 33\sqrt{a^2+b^2}\left(8a^6 - 25a^4b^2 - 25a^2b^4 + 8b^6\right)\\
&+ 2\sqrt{a^2+4b^2}\left(2a^6 - 25a^4b^2 - 100a^2b^4 + 128b^6\right)\\
&+ 2\sqrt{4a^2+b^2}\left(128a^6 - 100a^4b^2 - 25a^2b^4 + 2b^6\right)\\
&- 672a^6b\log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right)\\
&+ 693a^6b\log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right)\\
&- 21a^6b\log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right)\\
&- 672ab^6\log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right)\\
&+ 693ab^6\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right)\\
&- 21ab^6\log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)
\end{aligned}
$$

$$
\begin{aligned}
M_{10}^{(3)}(1,1) = \frac{1}{13440a^3b^2}\Bigg(&- 1968a^8 - 8064ab^7 - 2a\sqrt{a^2+b^2}\big(1104a^6\\
&- 4880a^4b^2 - 7350a^2b^4 + 7409b^6\big)\\
&+ 16a\sqrt{a^2+4b^2}\left(5a^6 - 68a^4b^2 - 310a^2b^4 + 708b^6\right)
\end{aligned}
$$

$$
\begin{aligned}
&+ 2a\sqrt{4a^2+b^2}\big(1024a^6 - 1152a^4b^2 - 440a^2b^4 \\
&+ 113b^6\big) - 6144a^7 b \log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right) \\
&+ 6576a^7 b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) \\
&- 432a^7 b \log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right) \\
&- 768b^6\left(28a^2 - 5b^2\right)\log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right) \\
&+ 6b^6\left(3696a^2 - 325b^2\right)\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right) \\
&- 3b^6\left(224a^2 - 5b^2\right)\log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)
\end{aligned}
$$

$$
\begin{aligned}
M_{01}^{(3)}(1,1) = \frac{1}{13440a^3b^2}\Bigg( &- 6096a^8 \\
&- 130a\sqrt{a^2+b^2}\left(48a^6 - 128a^4b^2 - 90a^2b^4 - 49b^6\right) \\
&+ 16a\sqrt{a^2+4b^2}\left(3a^6 - 32a^4b^2 - 90a^2b^4 - 196b^6\right) \\
&+ 2a\sqrt{4a^2+b^2}\big(3072a^6 - 2048a^4b^2 - 360a^2b^4 \\
&- 49b^6\big) - 15360a^7 b \log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right) \\
&+ 15600a^7 b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) \\
&- 240a^7 b \log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right) \\
&- 3840b^8 \log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right) \\
&+ 1950b^8 \log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right) \\
&- 15b^8 \log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)
\end{aligned}
$$

$$
M_{11}^{(3)}(1,1) = \frac{1}{120960a^4b^2}\Bigg( - 13776a^9 - 8160b^9 - 6\sqrt{a^2+b^2}\big(2576a^8
$$

$$
\begin{aligned}
&- 10210a^6b^2 - 12690a^4b^4 + 2237a^2b^6 + 1376b^8\big) \\
&+ 8\sqrt{a^2+4b^2}\big(70a^8 - 851a^6b^2 - 3180a^4b^4 \\
&+ 952a^2b^6 + 1024b^8\big) + 2\sqrt{4a^2+b^2}\big(7168a^8 \\
&- 7232a^6b^2 - 2280a^4b^4 + 103a^2b^6 + 16b^8\big) \\
&- 41472a^8 b \log\left(\frac{\sqrt{4a^2+b^2}+b}{2a}\right) \\
&+ 44388a^8 b \log\left(\frac{\sqrt{a^2+b^2}+b}{a}\right) \\
&- 2916a^8 b \log\left(\frac{\sqrt{a^2+4b^2}+2b}{a}\right) \\
&- 2304ab^6\left(28a^2+15b^2\right)\log\left(\frac{\sqrt{a^2+4b^2}+a}{2b}\right) \\
&+ 54ab^6\left(1232a^2+325b^2\right)\log\left(\frac{\sqrt{a^2+b^2}+a}{b}\right) \\
&- 9ab^6\left(224a^2+15b^2\right)\log\left(\frac{\sqrt{4a^2+b^2}+2a}{b}\right)\Bigg)
\end{aligned}
$$

# References


[1] T. Macháček, "Method of moments for analysis of planar radiators," Bachelor thesis, Czech Technical University in Prague, Faculty of Electrical Engineering, 2025, supervisor M. Čapek.

[2] S. A. Sauter and C. Schwab, *Boundary Element Methods*, ser. Springer Series in Computational Mathematics. Berlin, Heidelberg: Springer, 2011, vol. 39.

[3] S. Bao, D. Wang, Y. Mo, S. Hu, J. Gu, and W. Tang, "Fully analytical evaluation of singular integrals with RWG and rooftop basis functions," *IEEE Journal on Multiscale and Multiphysics Computational Techniques*, vol. 5, pp. 217–226, 2020.