

Appunti di intro QFT

Università degli Studi di Torino

Anno Accademico 2026/2027

Prof. Boglione, appunti di Paolo Rocchietti March¹

¹paolo.rocchiettimarc@edu.unito.it

Indice

1	Lesson 1, 23/09	3
2	Lesson 2, 24/09	3
2.1	Natural units	3
2.2	Klein-Gordon Equation	3
2.3	Symmetries and conservations	4

1 Lesson 1, 23/09

PRIMA LEZIONE SU TABLET

2 Lesson 2, 24/09

2.1 Natural units

From this point onwards, natural units will be used:

$$\hbar = c = 1$$

Note: this messes up dimensionality, but, conveniently, forces mass and energy to have the same units

$$E = mc^2 \xrightarrow{c=1} E = m,$$

and length and time to have the same dimensionality

$$x = ct \xrightarrow{c=1} [x] = [t].$$

To convert back into more familiar units the constants are, respectively,

$$c = 299.792.458 \text{ m/s}$$

$$\hbar c \approx 0.197 \text{ GeV}\cdot\text{fm}$$

2.2 Klein-Gordon Equation

The relativistic quantum mechanics' fundamental equation is

$$(\square + m^2)\psi(x) = 0$$

where

$$\square = \partial^\mu \partial_\mu = \frac{\partial^2}{\partial t^2} - \nabla^2$$

and $\psi(x)$ is still the Schrödinger's wavefunction.

It will now be shown that this equation **does not** conserve probability, using a continuity equation.

Regular quantum mechanics states

$$\frac{\partial \rho}{\partial t} + \nabla \cdot \mathbf{j} = 0,$$

where

$$\rho = \langle \psi_t | \psi_t \rangle \quad \mathbf{j} = \frac{i}{m} (\psi^* \nabla \psi - \psi \nabla \psi^*)$$

Consider the Klein-Gordon equation left-multiplied by $\psi^*(x)$ and its conjugate

$$\begin{cases} \psi^*(x) (\square + m^2) \psi(x) = 0 \\ \psi(x) (\square + m^2) \psi^*(x) = 0 \end{cases}$$

their difference is

$$\psi^*(\square + m^2)\psi - \psi(\square + m^2)\psi^* = 0$$

$$\partial^\mu (\psi^* \partial_\mu \psi - \psi \partial_\mu \psi^*) = 0$$

$$\partial^\mu J_\mu = 0$$

this is a 4D continuity equation and J^μ is a Noether current.

Expanding the equation yields

$$\frac{\partial}{\partial t} \left(\psi^* \frac{\partial \psi}{\partial t} - \psi \frac{\partial \psi^*}{\partial t} \right) - \nabla \cdot (\psi^* \nabla \psi - \psi \nabla \psi^*) = 0$$

the second term is proportional to the quantum probability current, but it is missing an i ! Moreover, the first term, representing ρ , is **not definite positive**, so it cannot represent the norm of the state $|\psi_t\rangle$!

This proves that it is **impossible** to make QM relativistic using wavefunctions and retaining the probabilistic approach.

Paul Dirac used another approach, he wanted to decrease the order of space derivative (instead of increasing the time derivative order, like K-G): this result was better, but for free particles it yielded a double solution with $E < 0$ states, violating a fundamental QM theorem!

For this reason, he devised the concept of the *Dirac sea*: these $E < 0$ states must all be occupied (because they cannot be observed), and a uniform charge can be factored out of Maxwell's equations with proper boundary conditions; when one of these particles in the sea is excited, the "hole" left behind can be perceived as a "positive electron": he predicted the existence of *antiparticles*.

This interpretation is still classical, but it begins to convey the concept that a finite number of particles is *not* possible for relativistic QM, so the concept of wavefunctions must be left behind in favor of field theories.

2.3 Symmetries and conservations

To build a theory, it is necessary to start with principles and symmetry: **Lorentz-invariance** to guarantee the conservation of distance between points (this is also useful to define a metric topology!).

Why symmetry? Noether theorem states that for every 4-current that respects a continuity equation

$$\partial_\mu J^\mu = 0$$

there is a conserved Noether charge

$$Q := \int d^3x J^0, \text{ so that } \frac{dQ}{dt} = 0$$

2.3.1 Lorentz-Poincaré group

The Lorentz invariant is

$$s^2 = t^2 - x_i x^i = t'^2 - x'_i x'^i = x_\mu x^\mu :$$

the Lorentz group is the set of all transformations $x^\mu \rightarrow x'^\mu$ over the Minkowski space that conserve s^2 ; in matricial representation the elements are

$$x'^\mu = \Lambda^\mu_\nu x^\nu$$

the invariant is conserved **if and only if**

$$\boxed{g_{\mu\nu} \Lambda^\mu_\rho \Lambda^\nu_\sigma = g_{\rho\sigma} .}$$

The covariant quantities transform with the inverse matrix

$$x'_\mu = \Lambda_\nu^\mu x_\nu$$

Observation 1 Since $g_{\mu\nu}$ is a symmetric matrix, this equation encompasses 10 constraints on Λ^μ_ν , leaving 3 degrees of freedom for rotations and 3 for boosts.

Naming convention Consider the determinant of the invariance condition:

$$\det(\Lambda^T g \Lambda) = \det(g) \leftrightarrow \det(\Lambda)^2 = 1 \leftrightarrow \det(\Lambda) = \pm 1$$

Transformations with $\det(\Lambda) = 1$ are called *proper Lorentz transformations*.

Transformations with $\det(\Lambda) = -1$ are called *improper Lorentz transformations*.

Naming convention Consider the (0,0) component of the invariance condition:

$$g_{\mu\nu} \Lambda^\mu{}_0 \Lambda^\nu{}_0 = 1 \leftrightarrow (\Lambda_{00})^2 - \Lambda^i{}_0 g_{ii} \Lambda^i{}_0 = 1 \leftrightarrow |\Lambda^0{}_0|^2 \geq 1$$

Because this term regulates how the transformation relates x^0 and x'^0 , transformations with $\Lambda^0{}_0 \geq 1$ are called *orthochronous Lorentz transformations*, while those with $\Lambda^0{}_0 \leq -1$ are called *non-orthochronous Lorentz transformations*.

These 2 divisions split the Lorentz group into 4 parts

- Proper and orthochronous $L_+^\uparrow$: restricted Lorentz group, sometimes referred to as "Lorentz Group";
- Improper and orthochronous $L_-^\uparrow$: spacial inversion;
- Proper and non-orthochronous $L_+^\downarrow$: time inversion;
- Improper and non-orthochronous $L_-^\downarrow$: total inversion of spacetime.

Note The only subgroup is $L_+^\uparrow$, as it is the only one that contains the identity matrix $\mathbb{I}$.