

Does a Structural Model Add Anything to the Closing Price?

Calibrated forecasting, incremental information, and match leverage
in the Italian Serie A

Yannik Pitcan
pitcany@gmail.com

August 2026

Abstract

Studies of association-football forecasting routinely report three-way accuracy in the low fifties and present it as competitive with the betting market. We argue that accuracy against a uniform benchmark answers the wrong question, and that the question worth asking is whether a model carries information a margin-free closing price has not already absorbed. We formalise that test as the fitted weight in a logarithmic opinion pool and apply it to nineteen complete Serie A seasons (7,220 matches, 2007–08 to 2025–26).

The answer is negative and stable. A Dixon–Coles model with tuned exponential decay attains 53.4% accuracy and RPS 0.1972 against the market’s 0.1905; the paired difference is +0.0067 (95% CI [0.0046, 0.0088]) and the market wins in all seven test seasons. The fitted pooling weight on the structural model is 0.000, and the log-loss profile is monotone increasing in that weight on the validation period and on the test period alike, so the result is a genuine boundary solution rather than an optimisation artefact. To test whether this reflects the noisiness of goals rather than the method, we refit the identical machinery to shots on target, a pre-expected-goals proxy for chance creation. That variant earns weight 0.35 against the goals model—it demonstrably carries information the goals model lacks—and 0.000 against the market. Two structural signals, each informative relative to the other, both already priced.

Separately, we find that the structural model is *better calibrated* than the market on the home-win margin (calibration slope 0.995 versus 1.103) while being clearly less sharp, so the market’s advantage is discrimination rather than honesty; studies reporting only accuracy cannot distinguish these.

The constructive consequence is that value lies not in a better outcome forecast but in what is built on a calibrated one. We define *match leverage*, the change in a club’s probability of achieving a season objective between winning and losing a given fixture, give conditions under which it can be estimated by post-hoc conditioning on a single simulation, and compute it for ACF Fiorentina. The ordering inverts intuition: an away fixture against a relegation rival carried 2.3× the leverage of hosting the eventual champions.

The paper also documents and corrects specific errors in an earlier study of our own on the same problem.

Keywords: probabilistic forecasting; market efficiency; Ranked Probability Score; calibration; opinion pooling; association football.

1 Introduction

Association-football forecasting occupies an unusual position among applied prediction problems: a strong, continuously updated, publicly observable benchmark exists in the form of bookmaker prices. This has two consequences that the applied literature has absorbed unevenly. First, any accuracy figure must be read against both a lower bound (the outcome base rate, which home advantage alone pushes well above uniform guessing) and an upper reference (the margin-free market price). Second, and more importantly, a model’s usefulness is not established by scoring well; it is established by carrying information the market has not already priced.

The second point is easy to state and easy to violate. A common design supplies bookmaker odds to a classifier as input features and then compares the resulting accuracy with the bookmaker’s own. Under that design the model consumes the market’s forecast and is then credited for approaching it. The appropriate reading of such a result is that the pipeline is discarding information, not adding it.

1.1 Contributions

1. We formalise the incremental-information question as the fitted weight in a logarithmic opinion pool with the de-vigged market price, and argue it should be the primary reported quantity in this literature (Section 5.5).
2. We report that weight to be 0.000 for a well-specified Dixon–Coles model over nineteen Serie A seasons, and establish that it is a boundary solution rather than an optimisation artefact by tracing the full log-loss profile on both validation and test data (Section 6.3).
3. We rule out target noise as the explanation by refitting the identical machinery to shots on target. That variant earns weight 0.35 against the goals model but 0.000 against the market, showing that two structurally different signals are both already priced (Section 6.4).
4. We separate calibration from discrimination empirically, finding the structural model better calibrated but less sharp than the market (Section 6.5).
5. We define *match leverage* and give the independence condition under which post-hoc conditioning on a single season simulation is equivalent to intervening on a fixture and resampling (Section 7).
6. We document and correct specific errors in an earlier study of our own on this problem (Section 3).

2 Related work

Goal-based structural models. The modern lineage begins with Maher (1982), who modelled home and away goals as independent Poisson variables with club-specific attack and defence parameters. Dixon and Coles (1997) identified the systematic misfit of the independence assumption on low scores, introduced a dependence correction on the four cells $(0, 0)$, $(0, 1)$, $(1, 0)$, $(1, 1)$, and added exponential down-weighting of older matches. Karlis and Ntzoufras (2003) developed the genuine bivariate Poisson alternative. Dynamic and Bayesian treatments followed: Rue and Salvesen (2000) modelled time-varying strengths through a Bayesian dynamic linear model, Baio and Blangiardo (2010) introduced hierarchical shrinkage of club parameters,

and Koopman and Lit (2015) placed the bivariate Poisson in a state-space framework. Goddard (2005) compared goal-based and result-based regression formulations directly. Hvattum and Arntzen (2010) evaluated Elo ratings as covariates, and—relevant here—benchmarked them explicitly against market odds.

Market prices as forecasts. Forrest et al. (2005) compared odds-setters with statistical forecasting models over several seasons and found the former difficult to beat, with the gap narrowing over time. Shin (1992, 1993) derived the insider-trading model that underlies the margin-removal method used here; Štrumbelj (2014) evaluated competing margin-removal schemes and found Shin’s the best-performing practical choice. This body of work establishes the market as a benchmark, but comparisons are typically made on accuracy or on returns rather than by testing directly for incremental information.

Scoring rules and calibration. Epstein (1969) introduced the Ranked Probability Score for ordered categorical forecasts; Gneiting and Raftery (2007) give the general theory of proper scoring rules. Constantinou and Fenton (2012) argued specifically that accuracy and other common measures are inadequate for football forecasts because the outcome is ordered, and established the RPS as the appropriate default. Calibration as a property distinct from discrimination is treated by Niculescu-Mizil and Caruana (2005); post-hoc correction methods include Platt scaling (Platt, 1999), isotonic regression (Zadrozny and Elkan, 2002) and temperature scaling (Guo et al., 2017).

The gap this paper addresses. Opinion pooling has a long statistical history (Genest and Zidek, 1986), and the logarithmic pool is standard. What appears to be uncommon in football forecasting is its use as a *diagnostic*: fitting the pooling weight out of sample and reporting it as the measure of whether a model adds anything to the market. Reporting accuracy alongside a market accuracy figure, as is customary, does not answer that question, since the two may be correlated in unknown degree.

3 An earlier study and its errata

The present work began as a correction of Pitcan (2018), an earlier study by the present author on the same problem, which reported 53.0% three-way accuracy over ten Serie A seasons and compared this favourably with betting markets at 55.3%. That study contained two framing errors and several arithmetic ones. We document them here because they are instances of failure modes the wider literature shares, and because the corrections motivate the design of what follows.

Framing. The benchmark was uniform guessing at 33%. Over the present corpus the home side wins 44.1% of matches, draws account for 26.2% and away wins 29.8%, so the majority-class rule alone scores 44.1%; the walk-forward base-rate forecaster used below scores 40.6% on the test period. Roughly seven percentage points of the reported margin were therefore not a margin at all. Separately, Bet365 odds were supplied to the classifiers as features and the resulting accuracy compared against the bookmaker’s, which is the inversion described in Section 1. The study noted that “the betting odds did not impact predictions very much” and attributed this to the odds being uninformative.

Arithmetic. The following are recomputed from the confusion matrices printed in that paper. Both total 218 matches against a stated test set of 320, a discrepancy the text does not reconcile.

- Tables 4.1 and 4.2 are both captioned “Linear SVM”. Table 4.1 has diagonal $80+29+2 = 111$ (50.9%), matching the reported Linear SVM figure; Table 4.2 has diagonal $77+34+3 = 114$ (52.3%), which is the reported AdaBoost figure. Table 4.2 is mislabelled.
- Away wins constitute $60/218 = 27.5\%$ of the test set, so “never predict an away win” scores 72.5%. Reported away accuracies are 72.9 (Gaussian SVM), 69.7 (Linear SVM), 68.4 (AdaBoost), 45.9 (logistic regression). None materially exceeds the constant rule and three fall below it.
- Home wins are $92/218$, so “never predict a home win” scores $126/218 = 57.80\%$. The reported Linear SVM home accuracy is 57.8, which to three significant figures is that constant rule, implying no home win was ever predicted.
- With $n = 218$ the standard error of an accuracy near one half is $\sqrt{0.25/218} \approx 0.034$. The reported multi-class accuracies span 47.2% to 53.0%, i.e. under two standard errors end to end, and the $n = 5$ versus $n = 7$ comparison moves the three classifiers in inconsistent directions. No ranking among them is supported.
- The time-weighted form feature is defined as $\tau'(\{x_1, \dots, x_n\}) = \frac{1}{n} \sum_i x_i e^{-i}$. A weighted average must be normalised by $\sum_i w_i$ rather than by n ; with $n = 5$, $\sum_{i=1}^5 e^{-i} = 0.578$, so the weighted feature carries roughly $1/8.7$ the scale of its unweighted counterpart, which matters for the kernel and regularised methods used. The decay is also severe: 63.6% of the weight falls on the most recent match and the Kish effective sample size (Kish, 1965) is 2.1 matches, so a nominal five-match window carries about two matches of information. The decay rate was asserted rather than fitted.

4 Data

The corpus comprises nineteen complete Serie A seasons, 2007–08 through 2025–26, totalling 7,220 matches, from `football-data.co.uk`. Each record carries the date, the two clubs, the full-time score and outcome, shots, shots on target and corners for both sides, and bookmaker prices. Bet365 opening prices are available for all seasons; Pinnacle opening and closing prices from 2012–13 onward. Where available, the Pinnacle closing line is preferred, as the closing price of a low-margin, high-limit book is the sharpest widely published forecast.

Two features of the corpus bear on modelling choices. First, promotion and relegation mean that only eight clubs appear in all nineteen seasons while many appear once or twice, so any rating scheme must handle clubs with little or no top-flight history. Second, home advantage has eroded substantially over the period (Figure 1): the home win rate falls from 47.5% across 2007–13 to 40.2% across 2023–26, and the ratio of home to away goals from 1.38 to 1.14. A model fitted without time weighting would misstate the current magnitude of home advantage.




Figure 1: Home win rate by season, with a linear trend. The erosion of home advantage over the sample motivates fitting the decay rate rather than assuming it.

5 Methods

5.1 Market probabilities

Let a k -outcome book quote decimal odds $o_1, \dots, o_k$ and write $\pi_i = 1/o_i$ with book sum $\Pi = \sum_j \pi_j > 1$. The excess $\Pi - 1$ is the bookmaker's margin. Proportional normalisation, $p_i = \pi_i/\Pi$, is known to leave a favourite-longshot bias. We instead use the model of Shin (1993), under which quoted prices arise from a bookmaker facing a proportion z of insider trade; the implied probabilities satisfy

$$p_i(z) = \frac{\sqrt{z^2 + 4(1-z)\pi_i^2/\Pi} - z}{2(1-z)}, \quad \sum_{i=1}^k p_i(z) = 1. \quad (1)$$

Differentiating shows $\partial p_i / \partial z < 0$ for all $p_i \in (0, 1)$, so the row sum is strictly decreasing in z and the root is located by bisection. Over the 17,479 complete books in this corpus the fitted z has median 0.017 and maximum 0.061.

5.2 Structural model

Following Dixon and Coles (1997), club j carries an attack rating α_j and a defence rating δ_j . For a fixture between home club h and away club a , goals (X, Y) have rates

$$\log \lambda = \gamma + \alpha_h - \delta_a, \quad \log \mu = \alpha_a - \delta_h, \quad (2)$$

where γ is the home advantage. The joint mass is

$$\mathbb{P}(X = x, Y = y) = \tau_\rho(x, y; \lambda, \mu) \frac{\lambda^x e^{-\lambda}}{x!} \frac{\mu^y e^{-\mu}}{y!}, \quad (3)$$




Figure 2: Validation RPS across the decay grid. The dashed line marks the selected rate, $\xi = 0.002$ per day.

with the low-score correction

$$\tau_\rho(x, y; \lambda, \mu) = \begin{cases} 1 - \lambda\mu\rho & (x, y) = (0, 0), \\ 1 + \lambda\rho & (x, y) = (0, 1), \\ 1 + \mu\rho & (x, y) = (1, 0), \\ 1 - \rho & (x, y) = (1, 1), \\ 1 & \text{otherwise.} \end{cases} \quad (4)$$

This assignment is the unique one conserving total probability: summing $(\tau_\rho - 1)$ times the independent mass over the four cells gives exactly zero, whereas transposing the $(0, 1)$ and $(1, 0)$ terms leaves a residual of $\rho e^{-\lambda-\mu}(\lambda - \mu)^2$.

Matches are weighted exponentially by age, $w(\Delta t) = \exp(-\xi\Delta t)$ with Δt in days before the forecast origin, and parameters maximise the weighted log-likelihood

$$\ell(\alpha, \delta, \gamma, \rho) = \sum_m w(\Delta t_m) \left[\log \tau_\rho(x_m, y_m; \lambda_m, \mu_m) + \log \text{Pois}(x_m; \lambda_m) + \log \text{Pois}(y_m; \mu_m) \right]. \quad (5)$$

Identifiability. The map $(\alpha, \delta, \gamma) \mapsto (\alpha + c\mathbf{1}, \delta + c\mathbf{1}, \gamma)$ leaves both rates unchanged, so the parameters are identified only up to this one-dimensional translation. We impose $\sum_j \alpha_j = 0$, which removes it exactly: the design matrix over observed ordered pairs has rank one less than its column count, and the single null direction has non-zero mean in the attack block. Clubs absent from the training window take the exposure-weighted league mean rather than an arbitrary constant; because the sum-to-zero constraint applies to attack but not to defence, defaulting both to zero would assign promoted clubs a better-than-average defence.

Decay rate. ξ is selected on the validation period by grid search (Figure 2), giving a clean interior optimum at $\xi = 0.002$ per day, a half-life of 347 days—appreciably slower than the 0.0065 reported by Dixon and Coles (1997) for late-1990s English data.

5.3 A process-based variant

Goals are a sparse realisation of chance creation, so a rating fitted to them inherits finishing noise. Expected goals is the standard remedy but is proprietary, and no freely licensed source covers Serie A back to 2007. We therefore fit the identical machinery to *shots on target*, the pre-xG proxy present in the corpus, obtaining shot rates λ^S, μ^S , and convert to goal rates by a single league-wide finishing factor κ estimated on the same window with the same decay weights:

$$\hat{\lambda} = \kappa_H \lambda^S, \quad \hat{\mu} = \kappa_A \mu^S, \quad \kappa_H = \frac{\sum_m w_m x_m}{\sum_m w_m s_m^H}, \quad (6)$$

and analogously for κ_A , where s_m^H denotes home shots on target. The factor is deliberately league-wide rather than club-specific, since club-level finishing variation is precisely the noise the substitution is meant to remove. Fitted values are $\kappa_H = 0.309$, $\kappa_A = 0.317$. The low-score correction is not carried across, as its parameter would have been estimated against shot counts; the variant is therefore independent-Poisson on converted rates, a genuine difference we report rather than conceal.

5.4 Evaluation protocol

Because the outcome is ordered, the primary metric is the Ranked Probability Score (Epstein, 1969; Constantinou and Fenton, 2012). For forecast p and realised indicator a over r ordered categories,

$$\text{RPS} = \frac{1}{r-1} \sum_{i=1}^{r-1} \left(\sum_{j=1}^i (p_j - a_j) \right)^2, \quad r = 3, \quad (7)$$

with categories ordered home win, draw, away win. RPS penalises a confident home forecast more heavily when the away side wins than when the match is drawn; neither accuracy nor the Brier score makes that distinction. Log loss and Brier are reported alongside, and accuracy solely for comparability with prior work.

All forecasts are produced walk-forward: parameters are fitted on matches strictly preceding an origin, used to forecast the following fortnight, and refitted at the next origin. Hyperparameters, pooling weights and the calibration temperature are fixed on 2013–14 to 2018–19 ($n = 2,280$); the test period 2019–20 to 2025–26 ($n = 2,660$) is scored once. Every mean carries a percentile bootstrap interval, and model comparisons use a paired bootstrap over matches, which exploits the strong correlation induced by scoring identical fixtures.

5.5 Incremental information via logarithmic pooling

Let p^M denote the de-vigged market forecast and p^S the structural model's. The logarithmic opinion pool (Genest and Zidek, 1986) with weight $w \in [0, 1]$ is

$$p_i^{(w)} \propto (p_i^M)^{1-w} (p_i^S)^w, \quad i \in \{H, D, A\}, \quad (8)$$

renormalised to sum to one. We fit $\hat{w}$ by minimising out-of-sample log loss on the validation period. The logarithmic form is preferred to the linear one as it is externally Bayesian and operates naturally on log-odds.

The interpretation is the point. If $\hat{w}$ is indistinguishable from zero, the market forecast is not improved by any admixture of the model, and the model carries no information the market

Table 1: Test period, 2019–20 to 2025–26 ($n = 2,660$). Lower RPS is better; the interval is a percentile bootstrap on the mean.

Model	RPS	95% CI	Log loss	Brier	Accuracy
Market (Shin de-vigged)	0.1905	[0.1856, 0.1957]	0.9620	0.5717	54.8%
Market + model pool	0.1905	[0.1856, 0.1957]	0.9620	0.5717	54.8%
Dixon–Coles (calibrated)	0.1972	[0.1921, 0.2024]	0.9858	0.5862	53.4%
Shots on target	0.2001	[0.1959, 0.2042]	0.9959	—	52.9%
Base rate (walk-forward)	0.2318	[0.2292, 0.2344]	1.0846	0.6574	40.6%
Uniform	0.2340	[0.2312, 0.2367]	1.0986	0.6667	40.6%

has not priced. If $\hat{w}$ is materially positive, it does. We propose that this quantity, rather than accuracy alongside a market accuracy figure, should be the headline diagnostic in this literature. Equation (8) extends to several models by raising each to its own weight on the simplex, which we use in Section 6.4.

6 Results

6.1 Main comparison

Table 1 reports the central comparison. The market attains RPS 0.1905 against the structural model’s 0.1972; the paired bootstrap places the difference at +0.0067 with 95% interval [0.0046, 0.0088]. The model is worse than the market, and decisively so relative to its own sampling error.

Two secondary observations follow. First, the rebuilt model attains 53.4% accuracy against the 53.0% of Pitcan (2018). Nine additional seasons, a principled likelihood, a fitted decay rate and leak-free validation bought 0.4 accuracy points—well inside noise. We read this as evidence that three-way match accuracy is near a ceiling, and that effort directed at it is directed at a quantity which has stopped responding. Second, the appropriate floor is the base-rate forecaster at 40.6%, not uniform guessing at 33%.

6.2 Stability across seasons

A single aggregate gap could be produced by one anomalous season. Table 2 and Figure 3 show it is not: the market attains lower RPS in all seven test seasons, with a gap ranging from 0.0044 to 0.0101 and no season in which the ordering reverses.

6.3 The pooling weight

Fitted on the validation period, the weight $\hat{w}$ on the structural model in the pool of Equation (8) is 0.000. Because this lies on the boundary of the admissible interval, a fitted value alone cannot distinguish a genuine null contribution from an optimiser terminating at a bound. We therefore trace the entire profile (Figure 4): log loss increases monotonically in w across $[0, 1]$, on the validation period and on the test period alike. The argument minimum is $w = 0$ in both cases. The result is not an artefact of the fitting procedure, and it does not depend on the validation/test split—even an analyst who improperly fitted the weight on the test data would obtain zero.

Table 2: Season-by-season mean RPS on the test period ($n = 380$ per season). “Gap” is the goals model minus the market.

Season	n	Market	Goals model	Shots model	Gap
2019–20	380	0.1968	0.2069	0.2090	+0.0101
2020–21	380	0.1820	0.1881	0.1920	+0.0061
2021–22	380	0.1953	0.2008	0.2023	+0.0055
2022–23	380	0.1950	0.2016	0.2021	+0.0067
2023–24	380	0.1839	0.1908	0.1986	+0.0069
2024–25	380	0.1838	0.1909	0.1932	+0.0071
2025–26	380	0.1970	0.2014	0.2030	+0.0044




Figure 3: Mean RPS by test season. The market is lower in every season; the two structural models track each other closely.

The interval $[0, 1]$ is the admissible range for a pool: outside it the combination is no longer a mixture of two opinions. A sceptical reader may nonetheless ask whether the unconstrained minimum lies below zero, which would say something stronger than that the model is uninformative—that it is anti-informative given the price. Extending the profile to $w \in [-1, 1]$, the validation minimum is interior, at $\hat{w} = -0.225$, with the loss rising again for weights more negative than that. Freezing that weight and carrying it to the test period improves mean log loss by 0.00062 (95% CI $[-0.00105, 0.00231]$) and mean RPS by 0.00021 (95% CI $[-0.00029, 0.00071]$). Both intervals contain zero, so the tilt is not a reliable out-of-sample improvement.

We read the sign rather than the magnitude. A negative weight divides the market forecast by the structural one, which sharpens it, and Section 6.5 shows the structural model to be the flatter of the two. The unconstrained optimum is therefore acting as a temperature correction on the price, not as a channel through which the model contributes information.

6.4 Is the target too noisy?

The natural objection is that the model failed because goals are noisy rather than because the method is inadequate. Table 3 addresses this using the multi-model extension of Equation (8).




Figure 4: Mean log loss of the pooled forecast as a function of the weight placed on the structural model. Monotone increasing on both periods, with the minimum at $w = 0$.

Table 3: Pool weights fitted on the validation period.

Pool	Market	Goals model	Shots model
Market + shots	1.00	—	0.00
Goals + shots	—	0.65	0.35
Market + goals + shots	1.00	0.00	0.00

Against the goals model alone, the shots variant earns weight 0.35: it demonstrably carries information the goals model does not, so the negative result of Section 6.3 is not an artefact of target choice. Against the market, the same signal earns 0.00; and in the three-way pool both structural models collapse to zero simultaneously. Two model families built on different targets, each holding information the other lacks, and the closing price has already absorbed both. We regard this as the strongest form of the negative result available without licensed event data, and note that expected goals—a refinement of the same process-based idea—would most plausibly sharpen the shots variant rather than reverse its relationship to the market.

6.5 Calibration and discrimination

Table 4 and Figure 5 separate two properties that an accuracy figure conflates. Regressing the outcome indicator on the forecast log-odds gives a calibration slope of 0.995 for the structural model against 1.103 for the market, with essentially identical expected calibration error. A slope near one indicates the stated probabilities can be taken at face value; the market’s slope above one indicates mild under-confidence on this margin. Yet the market is decisively sharper, which is what drives its RPS advantage. The fitted recalibration temperature for the structural model is 0.99, i.e. essentially no correction is warranted.

The practical implication is that “the market beats the model” and “the model’s probabilities cannot be trusted” are different statements, and only the first holds here. A well-calibrated but less sharp forecast remains usable for any decision that consumes the probability itself, which

Table 4: Calibration on the test period. ECE is the count-weighted mean absolute gap between stated and observed frequency over ten bins; slope and intercept are from a logistic regression of the home-win indicator on the forecast log-odds.

Model	ECE (H)	ECE (D)	ECE (A)	Slope (H)	Intercept (H)
Dixon–Coles	0.0208	0.0151	0.0270	0.995	−0.054
Market (Shin de-vigged)	0.0285	0.0095	0.0253	1.103	−0.090
Base rate	0.0164	0.0102	0.0062	0.191	−0.321




Figure 5: Home-win reliability curves on the test period. Both forecasters track the diagonal closely; the market’s advantage lies in sharpness rather than calibration.

is the basis for Section 7. The base-rate row illustrates the converse pathology: a forecast can be near-perfectly calibrated while carrying almost no discriminatory content, its slope of 0.191 reflecting the absence of any signal to be calibrated.

7 Match leverage

7.1 Definition

If a structural model cannot improve on a closing price for match outcomes, the product worth building is not a match-outcome forecast, which a club can obtain for free. What the market does not supply is the quantity that governs a club’s decisions: how much a particular fixture matters to its season.

Let Ω be a season objective expressed as a set of admissible final league positions—survival, European qualification, the championship. For a remaining fixture m , define

$$L_m = \mathbb{P}(\Omega \mid \text{club wins } m) - \mathbb{P}(\Omega \mid \text{club loses } m). \quad (9)$$

Table 5: ACF Fiorentina, simulated from 1 March 2026 with the club 16th on 24 points and twelve fixtures remaining. Objective: survival, i.e. a final position of 17th or better. Monte Carlo standard error on each leverage estimate is below 0.005.

Fixture	Venue	$\mathbb{P}(\Omega \mid W)$	$\mathbb{P}(\Omega \mid D)$	$\mathbb{P}(\Omega \mid L)$	L_m
Lecce	A	0.987	0.959	0.914	0.073
Cremonese	A	0.986	0.957	0.922	0.063
Verona	A	0.982	0.952	0.927	0.055
Genoa	H	0.984	0.962	0.932	0.053
Udinese	A	0.985	0.961	0.933	0.052
Sassuolo	H	0.982	0.956	0.932	0.050
Lazio	H	0.988	0.960	0.942	0.046
Parma	H	0.981	0.955	0.936	0.045
Atalanta	H	0.990	0.966	0.949	0.041
Juventus	A	0.988	0.971	0.951	0.037
Roma	A	0.988	0.968	0.952	0.036
Inter	H	0.988	0.971	0.955	0.033

7.2 Estimation

Let S denote the scoreline of fixture m and R the scorelines of all other remaining fixtures. The objective indicator is a deterministic function $f(S, R)$: it depends on m through the points awarded to both clubs, and on every other result through the final ordering. If the simulator draws $S \perp R$, then for any result set W ,

$$\mathbb{P}(f(S, R) = 1 \mid S \in W) = \sum_{s \in W} \mathbb{P}(S = s \mid S \in W) \mathbb{P}(f(s, R) = 1), \quad (10)$$

which is exactly the interventional quantity obtained by forcing m to a result drawn from the model’s own conditional distribution given W and resampling the remainder. Conditioning on S cannot induce selection on R precisely because the two are independent; the opponent’s points *from* m are part of S and are conditioned on, not marginalised. Hence a single simulation run supports the estimation of L_m for every fixture simultaneously, by post-hoc conditioning.

We verified this equivalence in two ways: by exhaustive enumeration of all 59,049 joint outcomes of a reduced four-club league, where post-hoc and forced-result computations agree to machine precision; and by direct comparison against forced-result simulation at 10^6 replications on the applied case below, where no fixture differs beyond Monte Carlo error (maximum $|z| = 1.17$ across twelve fixtures).

In practice we simulate the remainder of the season 20,000 times, drawing complete scorelines from each fixture’s joint distribution so that goal difference resolves naturally, and rank the final table by points, then goal difference, then goals scored. (Serie A resolves equal points on head-to-head record first; exact ties affecting the reported band arise in under 0.01% of simulations.)

7.3 Application

Table 5 reports the result. The ordering inverts intuition. The fixtures moving survival most are those against direct relegation rivals; those against the leading clubs move it least, because defeat is already expected in both branches and therefore already priced into the season distribution.

Losing at home to Inter—the eventual champions, and the season’s marquee fixture—costs 3.3 percentage points of survival probability; losing away at Lecce costs 7.3. The away fixture carries $2.3\times$ the leverage of the home one, a gap of approximately eight Monte Carlo standard errors, so the ordering is not simulation noise.

This is a rotation and load-management statement expressed in probability: the Inter fixture is where a fatigued first-choice player is cheapest to rest, and Lecce away is where he is most expensive. No match-outcome forecast conveys it.

As a coherence check rather than a claim of predictive success, the simulation projected 41.4 expected final points against an actual 42 and assigned survival probability 96.4%; Fiorentina finished 15th. A single realisation is weak evidence, but it establishes that the season aggregation is not grossly miscalibrated, which is the minimum precondition for acting on a leverage ranking.

8 Discussion

What the negative result does and does not say. It does not say that structural models are useless, nor that the market is efficient in a strong sense. It says that over this league and period, for two structurally different signals, the marginal information relative to a de-vigged closing price is indistinguishable from zero at the resolution afforded by 2,660 matches. A model with genuinely private inputs—player availability, training-load data, tactical intelligence—is not tested here and is exactly where we would expect remaining edge to lie.

Implications for reporting practice. We would encourage two changes. First, report the base rate as the floor; uniform guessing is not a meaningful benchmark for an outcome with a 44% majority class. Second, where market prices are available, report the fitted pooling weight against them. It is a single number, it is cheap to compute, and it answers the question a reader actually has. An accuracy figure quoted beside a market accuracy figure does not, because the two forecasts may be arbitrarily correlated.

Calibration as the usable residual. The finding that the model is better calibrated but less sharp than the market has a constructive reading. A club cannot profitably trade on this model, but it can aggregate it: the season simulation of Section 7 consumes probabilities directly, and what such aggregation requires is that stated probabilities mean what they say, not that they be maximally sharp. This is the sense in which the useful product sits on top of a calibrated forecast rather than in the forecast itself.

On the ceiling. That a decade of methodological improvement and nine additional seasons moved accuracy by 0.4 points is worth stating plainly. Under a scoring rule sensitive to the full distribution the improvement is likewise small. If the goal is a better outcome forecast, the binding constraint is information, not method.

9 Limitations

No player-level information. No injuries, suspensions, rest days or transfers enter the model. This is the largest gap relative to a club’s internal information set and the most plausible location of remaining edge.

No expected goals. Shots on target weight a tap-in and a speculative long-range effort equally. Licensed event data would sharpen the process-based variant; the results of Section 6.4 suggest refinement rather than reversal, but this is not established.

The benchmark embeds late information. The closing price incorporates team news the model never sees, so some portion of the 0.0067 RPS gap is information asymmetry rather than modelling deficiency. This works against the model and in favour of the reported conclusion.

One league, one period. Serie A over nineteen seasons. Market efficiency need not transfer to leagues with thinner betting markets, and the erosion of home advantage documented in Section 4 shows the data-generating process is not stationary.

Leverage is single-objective. Equation (9) conditions on one position band. A club trading survival against a cup run requires a utility function over outcomes, not a probability difference.

10 Reproducibility and verification

All results derive from public data and a versioned Python package with 126 tests. Beyond unit testing, four components—the margin-removal solver, the likelihood and its identifiability, the walk-forward harness, and the leverage estimator—were subjected to independent adversarial audit by parties who did not write them. That audit confirmed the absence of look-ahead bias by positive control (an injected one-line boundary error improves test RPS by 0.0123 and reverses the sign of the market comparison, a signature the reported pipeline does not exhibit) and identified four defects, all corrected before the results reported here were generated. Two of the corrected defects were tests that asserted nothing: one passed against a deliberately leaky harness, the other verified only that probabilities summed to one. We mention this because the negative result reported here is of a kind that a coding error can easily manufacture, and readers are entitled to know what was done to exclude that.

Code, data pipeline and figures: <https://github.com/pitcany/seriea-leverage>.

11 Conclusion

Over nineteen Serie A seasons, a well-specified Dixon–Coles model earns a logarithmic pooling weight of zero against a Shin de-vigged closing price, and so does a process-based variant fitted to shots on target—though the latter earns weight 0.35 against the former, establishing that both carry real and partially distinct information. The market has already absorbed both. The model is nonetheless better calibrated than the market while being less sharp, which makes it usable for aggregation even where it is unusable for prediction. We define match leverage as one such aggregate and show that it inverts intuitive fixture importance: for a club fighting relegation, an away match against a fellow struggler was worth more than twice as much as hosting the eventual champions.

Data and code availability

Match data are publicly available from football-data.co.uk. All code, including the exact scripts producing every table and figure here, is at <https://github.com/pitcany/seriea-leverage>.

References

- Baio, G. and Blangiardo, M. (2010). Bayesian hierarchical model for the prediction of football results. *Journal of Applied Statistics*, 37(2), 253–264.
- Constantinou, A. and Fenton, N. (2012). Solving the problem of inadequate scoring rules for assessing probabilistic football forecast models. *Journal of Quantitative Analysis in Sports*, 8(1).
- Dixon, M. J. and Coles, S. G. (1997). Modelling association football scores and inefficiencies in the football betting market. *Journal of the Royal Statistical Society: Series C*, 46(2), 265–280.
- Epstein, E. S. (1969). A scoring system for probability forecasts of ranked categories. *Journal of Applied Meteorology*, 8(6), 985–987.
- Forrest, D., Goddard, J. and Simmons, R. (2005). Odds-setters as forecasters: The case of English football. *International Journal of Forecasting*, 21(3), 551–564.
- Genest, C. and Zidek, J. V. (1986). Combining probability distributions: A critique and an annotated bibliography. *Statistical Science*, 1(1), 114–135.
- Gneiting, T. and Raftery, A. E. (2007). Strictly proper scoring rules, prediction, and estimation. *Journal of the American Statistical Association*, 102(477), 359–378.
- Goddard, J. (2005). Regression models for forecasting goals and match results in association football. *International Journal of Forecasting*, 21(2), 331–340.
- Guo, C., Pleiss, G., Sun, Y. and Weinberger, K. Q. (2017). On calibration of modern neural networks. In *Proceedings of the 34th International Conference on Machine Learning*.
- Hvattum, L. M. and Arntzen, H. (2010). Using ELO ratings for match result prediction in association football. *International Journal of Forecasting*, 26(3), 460–470.
- Karlis, D. and Ntzoufras, I. (2003). Analysis of sports data by using bivariate Poisson models. *Journal of the Royal Statistical Society: Series D*, 52(3), 381–393.
- Kish, L. (1965). *Survey Sampling*. Wiley, New York.
- Koopman, S. J. and Lit, R. (2015). A dynamic bivariate Poisson model for analysing and forecasting match results in the English Premier League. *Journal of the Royal Statistical Society: Series A*, 178(1), 167–186.
- Maher, M. J. (1982). Modelling association football scores. *Statistica Neerlandica*, 36(3), 109–118.
- Niculescu-Mizil, A. and Caruana, R. (2005). Predicting good probabilities with supervised learning. In *Proceedings of the 22nd International Conference on Machine Learning*.
- Pitcan, Y. (2018). Soccer match prediction in the Serie A. Accepted for publication, *International Journal of Computer Applications*.
- Platt, J. (1999). Probabilistic outputs for support vector machines and comparisons to regularized likelihood methods. In *Advances in Large Margin Classifiers*. MIT Press.

- Rue, H. and Salvesen, Ø. (2000). Prediction and retrospective analysis of soccer matches in a league. *Journal of the Royal Statistical Society: Series D*, 49(3), 399–418.
- Shin, H. S. (1992). Prices of state contingent claims with insider traders, and the favourite–longshot bias. *The Economic Journal*, 102(411), 426–435.
- Shin, H. S. (1993). Measuring the incidence of insider trading in a market for state-contingent claims. *The Economic Journal*, 103(420), 1141–1153.
- Štrumbelj, E. (2014). On determining probability forecasts from betting odds. *International Journal of Forecasting*, 30(4), 934–943.
- Zadrozny, B. and Elkan, C. (2002). Transforming classifier scores into accurate multiclass probability estimates. In *Proceedings of the 8th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*.