Beyond Points Accuracy: Survival Forecasts and Fixture Stakes in Serie A

Yannik Pitcan

Revised 20 September 2026

Abstract

Does a better forecast of final points establish a better forecast of survival? We examine that distinction and, separately, the survival stakes of avoiding defeat versus turning a draw into a win. Using 7,220 Serie A fixtures from 2007/08–2025/26, we compare goals-based Dixon–Coles forecasts, independent-Poisson shots-on-target ratings, and a documented closing-price benchmark. Parameters and combinations are selected on 2013/14–2018/19; 2,660 subsequent matches form a historical evaluation period. We evaluate final-points and survival forecasts for all clubs at three dates across seven seasons, using the applicable head-to-head and playoff rules. On March 1, structural-model points mean absolute error is 3.48, versus 5.45 for a league-average Poisson comparator: difference -1.97 (95% season-bootstrap interval [-2.62, -1.40]). Survival Brier scores are 0.0552 and 0.0536, with difference +0.0016 ([-0.0058, +0.0090]). Points gains have intervals excluding zero at all three dates; survival gains remain unresolved. An exploratory breakdown uses the bottom eight clubs at each forecast origin. For match forecasts, calibrated goals have ranked probability score 0.19721, versus 0.19037 for the closing market. For Fiorentina against Lecce, the simulated survival difference between a draw and a loss is 4.82 percentage points, versus 3.12 between a win and a draw. Reweighting draws toward wins and losses yields a break-even ratio of added win probability to added loss probability of 1.54 (approximate 95% simulation-only interval [1.08, 2.19]). This illustrates how fixture stakes depend on the type of probability change, without identifying a tactical or rotation effect. The historical forecast comparison and the hypothetical probability trade-off answer different questions; neither establishes a useful real-world policy. The contribution connects outcome-specific forecast evaluation to an application of established conditional-importance methods. Revised analyses are retrospective, and prospective forecast and action validation remain necessary.

1 Question, application and prior work

Clubs fighting relegation need probabilities of survival, while forecast improvement is often summarized through match scores or final points. We ask whether a transparent model that improves final-points predictions also demonstrates better survival forecasts. Survival depends on the club's position relative to other teams, their joint results and league rules; a better points prediction alone does not establish that these probabilities improve.

We connect that forecast-evaluation question to a separate scenario question: when does avoiding defeat have more survival value than turning a draw into a win? A single win-versus-loss importance score does not distinguish these two consequences. We retain both and derive the probability trade-offs needed for a hypothetical change to improve survival. Historical forecasting performance and the modeled consequences of assumed probability changes are different forms of evidence. Neither identifies how an actual tactic changes match outcomes.

Our application is survival planning in Serie A, where a match's relevance depends on rival results, the remaining schedule and league rules. We compare the model with a league-average baseline for season forecasts and separately with closing betting prices for individual matches.

The underlying methods are established. Dixon and Coles (1997) introduced a low-score dependence adjustment and time weighting for football scores. Egidi et al. (2018) combine historical information and bookmaker odds in a Bayesian score model. Reade et al. (2020) compare statistical and betting-market forecasts, including forecast-encompassing analysis of scoreline probabilities. Wheatcroft (2021) forecasts match statistics and finds information beyond odds with a different rating and combination procedure. A negative result for our shots model cannot overturn that evidence or generalise to expected goals.

Our match-importance definition is also established. Scarf and Shi (2008) develop conditional tournament importance and discuss effort allocation. Link and de Lorenzo (2016) describe the difference in the probability of a season consequence conditional on winning versus losing and its earlier antecedents; they study its association with player activity and pacing. de Lorenzo et al. (2016) explicitly incorporate win and draw importance. We apply this established framework to the asymmetry of Serie A survival stakes and do not claim a new importance metric or an optimal tactical rule.

This paper contributes a retrospective comparison of points and survival forecast performance, an exploratory breakdown by standings at the forecast origin, and a documented match-level evaluation of goals, shots, pooling and recalibration. The Fiorentina example separately distinguishes avoiding defeat from turning a draw into a win, reports simulation uncertainty, and makes their hypothetical probability trade-off explicit. Its contribution is the integration and evaluation of these components, not a new forecasting family or causal decision rule.

2 Data, timing and evaluation protocol

Football-Data provides results, shots on target and bookmaker prices (Football-Data, 2026). The local snapshot contains 19 complete seasons, 2007/08–2025/26, each with 380 fixtures: 7,220 matches overall. Date, season and home/away club identify each fixture. We check duplicate fixtures, self-matches, score/result agreement and season completeness. One match lacks complete shots-on-target data and is excluded from shots fitting.

Model forecasts use only matches dated strictly before their forecast origin. Origins advance by 14 days; the training window covers the preceding 1,460 days. The cap is a substantive modeling choice: at decay 0.002/day the oldest retained observation still has weight about 0.054. A forecast can therefore precede kickoff by up to 13 calendar days. Closing prices have later information, including potentially lineups and injuries. This is an operational comparison with the eventual close, not a comparison at identical information times.

The 2,280 fixtures from 2013/14–2018/19 select the goals-model decay, temperature and forecast-combination parameters. The 2,660 fixtures from 2019/20–2025/26 evaluate the selected forecasts. Forecast origins still refit club ratings as new past matches become available; combination parameters remain fixed. Revisions were developed after inspecting the historical test period. We retain the chronological split but do not describe the revised analyses as an untouched or preregistered holdout.

2.1 Closing-price provenance

We use Pinnacle closing prices before 23 July 2025, then Bet365 closing prices; average closing prices are an explicitly labelled fallback if needed. Football-Data warns that Pinnacle prices became systematically stale from 23 July 2025. We ingest closing columns directly and never replace missing closing prices

with pre-closing snapshots. A source and timing label accompanies every forecast. In the evaluated sample, all earlier seasons use Pinnacle closing prices and all 380 fixtures in 2025/26 use Bet365 closing prices.

Shin de-vigging (Shin, 1992) converts complete three-outcome price books to normalised probabilities. The market remains a forecast, not ground truth. Changing bookmaker across seasons, the de-vigging assumption and unmatched forecast times constrain interpretation. The six-season comparison excluding 2025/26 and a 2025/26 average-closing sensitivity are reported in the reproduction outputs.

3 Forecasts and statistical comparisons

3.1 Goals and shots models

For home club i and away club j,

$$\lambda_{ij} = \exp(\gamma + a_i - d_j), \qquad \mu_{ij} = \exp(a_j - d_i), \qquad \sum_i a_i = 0.$$

The goals model multiplies the independent Poisson mass by

$$\tau(x, y) = \begin{cases} 1 - \lambda\mu\rho, & x = y = 0, \\ 1 + \lambda\rho, & x = 0, y = 1, \\ 1 + \mu\rho, & x = 1, y = 0, \\ 1 - \rho, & x = y = 1, \\ 1, & \text{otherwise.} \end{cases}$$

We maximise the weighted log likelihood with weights $\exp(-\xi\Delta t)$, where Δt is measured in days. The validation grid is $\xi \in \{0, .002, .004, .0065, .009, .012, .018\}$ and selects 0.002. The original value 0.0065 used half-week units in Dixon and Coles (1997); it corresponds to about 0.001857/day, not 0.0065/day. Our selected daily rate has a half-life of approximately 346.6 days.

Probability support must hold at every possible score, even unobserved low scores. We therefore optimise a fraction $\eta \in [-.999999, .999999]$ of a rate-dependent admissible interval. Let

$$B = \exp\{\max(\gamma, 0) + \max_i a_i - \min_i d_i\}, \quad C = \exp\{\gamma + 2(\max_i a_i - \min_i d_i)\}.$$

We set $\rho = \eta/B$ for negative η and $\rho = \eta \min(1, 1/C)$ otherwise. These conservative bounds guarantee positive low-score corrections for every pairing of the fitted ratings and the stated fallback ratings. They may constrain the likelihood more than checking only observed pairings. Prediction rejects invalid support rather than clipping negative probabilities. Optimiser convergence and minimum forecast tau are retained with the forecasts.

The shots model fits independent Poisson rates to home and away shots on target: ρ is fixed to zero during fitting, as well as in the converted goal grid. Venue-specific league conversion factors are the weighted totals of goals divided by weighted totals of shots on target in the same training window. The shots model shares the selected goals decay for a controlled comparison; we do not claim that this is the best possible shots specification. Changing both target and dependence structure means this is not an experiment that isolates target noise.

Both models use attack zero for an unseen club (the unweighted mean under the identifiability constraint) and exposure-weighted mean defence. The shots model uses the same rate-extraction path, preventing a different unseen-club fallback. Score grids include 0–12 goals per side and are renormalised after truncation. The season model uses raw score grids; the separately temperature-calibrated three-way goals forecast is used only in match evaluation.

3.2 Pooling is a restricted comparison

For market probabilities m and structural probabilities s, a logarithmic pool is $p_k \propto m_k^{1-w} s_k^w$. We select $w \in [0, 1]$ on validation log loss, explicitly evaluating both endpoints. Multi-model weights use a nonnegative simplex grid with spacing 0.05.

A zero weight establishes only that the structural forecast was not selected within this family on validation data. A positive weight can compensate for calibration error without adding match-specific information: even uniform probabilities can soften an overconfident forecast. Conversely, a useful signal can receive zero weight if its mapping into probabilities is misspecified. A negative coefficient generally changes match-specific relative odds and is not equivalent to a single temperature.

To distinguish a simple recalibration gain from a gain using structural forecasts, we fit two nested multinomial-logit combinations. With away as the reference class, for $k \in \{H, D\}$,

$$\log \frac{p_k}{p_A} = \alpha_k + \beta_M \log \frac{m_k}{m_A} + \beta_G \log \frac{g_k}{g_A} + \beta_S \log \frac{s_k}{s_A}.$$

The market-only model fixes $\beta_G = \beta_S = 0$. Both versions estimate the same two class intercepts and market coefficient on validation data. Bounds are $[-3, 3]$ for intercepts, $[0, 5]$ for β_M and $[-2, 2]$ for structural coefficients; no test-set tuning is performed. Their held-out score difference is the relevant comparison. This is still a restricted combination model, not a general test of informational completeness.

3.3 Scores, uncertainty and calibration

The primary match score is the ranked probability score (RPS), using H–D–A ordering and division by two; lower values are better. We also report log loss, the sum-of-squares multiclass Brier score and accuracy. A uniform forecast's argmax ties select H in the software: its displayed accuracy therefore equals the home-win frequency, not the expected accuracy of random guessing.

Every comparison scores the same fixtures. We provide paired match-bootstrap and paired season-bootstrap intervals using 10,000 resamples. Season resampling retains dependence among matches in a season, but only seven clusters are available. Neither procedure propagates model-estimation or specification uncertainty. Intervals for the additional comparisons are descriptive, without a multiplicity correction or an equivalence margin.

For each outcome we report equal-width-bin calibration, bin counts, descriptive Wilson intervals, and calibration intercept/slope estimates with Fisher intervals. The latter assume independent observations and are not a clustered test of superiority. Macro one-versus-rest AUC describes discrimination; entropy describes concentration. Neither is interchangeable with calibration.

4 Match-forecast results

The closing benchmark outperforms the calibrated goals model by +0.00684 RPS, with a season-bootstrap interval of [+0.00596, +0.00805]. The market has lower RPS in 7 of seven seasons. The nonnegative market/goals and market/shots pools select the market alone. This supports a scoped negative result about those pools, not the conclusion that all structural information is priced or that improved forecasting is impossible.

The validation goals/shots weights are 0.60/0.40. Relative to raw goals, that pool changes test RPS by -0.00055 with a season-bootstrap interval of [-0.00122, +0.00024]. The comparison quantifies the gain rather than inferring information from a positive pooling weight.

Table 1: Historical test forecasts, 2019/20–2025/26 ($n = 2,660$). RPS, log loss and Brier score: lower is better.

Forecast	RPS	Log loss	Brier	Accuracy
Market + structural logits	0.18974	0.9591	0.5700	54.6%
Market recalibrated	0.18985	0.9594	0.5702	54.8%
Market	0.19037	0.9614	0.5714	54.7%
Goals + shots	0.19667	0.9834	0.5852	53.6%
Goals calibrated	0.19721	0.9858	0.5862	53.4%
Goals raw	0.19722	0.9857	0.5862	53.4%
Shots	0.19871	0.9905	0.5900	53.2%

Adding goals and shots log odds to market recalibration changes test RPS by -0.00012 (season-bootstrap interval [-0.00040, +0.00012]) and log loss by -0.00026 (interval [-0.00116, +0.00045]). This result is assessed against the recalibrated market, not just the raw prices. The RPS interval includes zero, so these data do not establish a stable improvement from the added structural forecasts.

Calibration diagnostics are in the accompanying tables and reliability figure. A slope near one for one class, or a fitted temperature near one, cannot establish global calibration. Sparse extreme-probability bins have visibly large uncertainty. We therefore make no claim that the structural forecast is generally better calibrated or that calibration diagnostics establish the reliability of its joint season distribution.

5 Season simulation and retrospective validation

Each simulation independently draws a scoreline for every remaining league fixture from the raw goals model fitted at the decision date. Ratings are frozen throughout the remaining schedule. Final totals combine completed results with these draws. This excludes future transfers, injuries, tactical changes, parameter uncertainty and dependence induced by latent team-strength updates.

Points-tied groups are ordered by their head-to-head points, head-to-head goal difference, overall goal difference, goals scored and drawing lots. From 2022/23, ties at the title and survival boundaries trigger playoffs after the mini-table selects the participants. The 2022/23 rule uses one neutral match. From 2023/24, the title uses one match at the better-ranked club and survival uses two legs, with the better-ranked club hosting the return (FIGC, 2022, 2023a; Lega Serie A, 2025). Equal aggregate goals go directly to penalties: there is no extra time or away-goals rule. Playoff scoring uses the same frozen ratings; penalty shootouts are modeled as 50/50. We assume the scheduled home venue for title playoffs, excluding exceptional public-order venue changes.

5.1 Evaluation design

We reconstruct forecasts for all 20 clubs on January 1, March 1 and May 1 in each season from 2019/20 through 2025/26: 140 club-season observations per horizon, 420 overall. Dates and comparisons were fixed for this revision after the historical outcomes had been inspected. This is exploratory validation, not a prospective trial.

For every origin we draw 10,000 seasons and evaluate survival Brier score, final-points mean absolute error, empirical continuous ranked probability score (CRPS) and central 90% point-interval coverage. The comparator uses league-wide, time-weighted home and away goal rates from the identical training window. It preserves the current table and remaining fixtures but assigns no club-specific strengths. It is a transparent baseline, not a claim to be the strongest alternative.




Figure 1: Top: proper-score difference from the closing benchmark by season. Bottom: selected forecasts on the historical test sample with paired season-bootstrap intervals for their RPS difference from the market. Intervals are conditional on fitted forecasts and based on seven seasons.

The future fixture list is taken from the completed dataset, with realised schedule dates; no future scores or closing prices enter the forecasts. The season model has no time-varying future ratings, so future fixture dates do not affect its rate estimates. Known points deductions are applied at the origin. For Juventus in 2022/23 we use the 15-point sanction after January 20, its removal after April 20, and the eventual 10-point sanction after May 22 (FIGC, 2023b; CONI, 2023; FIGC, 2023c). Future legal changes are not anticipated. Final points include the actual sanction; actual survival labels include Verona's win over Spezia in the June 2023 playoff. Errors can therefore include unanticipated administrative changes.

The structural model improves points MAE and CRPS at every evaluated horizon; the paired season intervals for those improvements exclude zero. In March, points MAE is 3.48 versus 5.45, a difference of -1.97 points (interval [-2.62, -1.40]). Survival Brier differences include zero at all three horizons. Better final-points forecasts therefore do not establish better survival probabilities. For March forecasts, the structural-minus-baseline survival Brier difference is +0.0016 with season-bootstrap interval [-0.0058, +0.0090]. March point-interval coverage is 91.4%. Coverage at this horizon is at least 90%; this finite sample does not establish calibration across horizons or subgroups. Repeated dates for the same club-season are not independent replications; inference resamples whole seasons. Seven seasons,




Figure 2: Raw goals and market reliability for all three outcomes. Point size encodes bin count; vertical intervals are descriptive 95% Wilson intervals under an independent-binomial approximation. Count strips show the number of matches in each bin. These are not tests accounting for season dependence.

containing only 21 actual relegations, sharply limit calibration assessment. Predictive interval coverage also tests the frozen-rate simulation assumptions, rather than just Monte Carlo precision.

5.2 Exploratory breakdown by standings at the forecast origin

To check whether the points gains occur only away from the relegation contest, we separately evaluate the lowest eight clubs and the other twelve at each origin. Eligibility uses current points after known deductions, then current goal difference, goals scored and alphabetical club name to resolve ties. This deterministic subgroup ordering is distinct from the final-ranking rules used by the simulator. It uses only results strictly before the origin; no final standings or survival outcomes determine membership. The groups contain 56 and 84 club-season observations per date, respectively. The choice of eight and this breakdown were made after inspecting the historical study, so the results are exploratory. We retain the all-club analysis and resample the same seven seasons within each subgroup.

Table 2: All-club retrospective season evaluation. Each row contains 140 club-season forecasts. Survival uses binary Brier score; point coverage is for a nominal central 90% interval.

Date	Model	Brier	Pts MAE	Pts CRPS	Coverage
Jan 1	Dixon-Coles	0.0882	5.26	3.73	84.3%
Jan 1	League Poisson	0.0904	8.66	6.47	65.0%
Mar 1	Dixon-Coles	0.0552	3.48	2.43	91.4%
Mar 1	League Poisson	0.0536	5.45	3.88	78.6%
May 1	Dixon-Coles	0.0256	2.33	1.64	95.0%
May 1	League Poisson	0.0277	2.95	2.08	87.9%

Table 3: Exploratory origin-defined subgroup comparisons. Differences are Dixon–Coles minus league-average Poisson; negative values favor Dixon–Coles. Brackets are paired 95% season-bootstrap intervals. n counts club-season observations at each date, not independent seasons.

Date	Origin group	n	Points MAE difference	Survival Brier difference
Jan 1	Bottom eight	56	-3.12 [-3.99, -2.15]	-0.008487 [-0.054905, +0.037179]
Jan 1	Other twelve	84	-3.58 [-4.96, -2.15]	+0.001936 [-0.000902, +0.005099]
Mar 1	Bottom eight	56	-2.13 [-2.74, -1.58]	+0.004079 [-0.014325, +0.022588]
Mar 1	Other twelve	84	-1.87 [-2.61, -1.16]	-0.000050 [-0.000095, -0.000016]
May 1	Bottom eight	56	-0.75 [-1.22, -0.42]	-0.005193 [-0.021654, +0.015922]
May 1	Other twelve	84	-0.54 [-1.11, +0.01]	-0.000004 [-0.000012, +0.000000]

In the bottom eight on March 1, points MAE is 3.24 versus 5.36, a difference of -2.13 (interval [-2.74, -1.58]). Survival Brier is 0.1379 versus 0.1338, a difference of +0.0041 ([-0.0143, +0.0226]). Bottom-eight points gains have intervals excluding zero at all three dates; their survival-score differences include zero at every date. The other twelve also show points gains in January and March, although their May interval includes zero. Their March survival Brier improvement is very small in absolute terms (Table 3). Thus the evidence for points improvement is not confined to the other twelve clubs, while better survival forecasts for the bottom eight remain unestablished. This breakdown does not identify which modeling assumptions cause the remaining survival errors or establish equivalence between the forecasts.

6 Fiorentina: avoiding defeat or chasing victory?

The preceding analysis evaluates historical forecast quality. The following illustration asks a separate question about the consequences of assumed probability changes; it does not explain the historical forecast-score gaps. At March 1, 2026, we fit the model using only prior results and simulate 20,000 season completions. Fiorentina's projected survival probability is 96.00%, compared with 96.41% under the earlier points/overall-goal-difference ranking applied to the same league draws. A survival playoff is triggered in 11.23% of simulations across the league. That is a rule-trigger rate, not a prediction-error rate. Expected final points are 41.40. The later observed finish is an illustration, not evidence that one probabilistic forecast was calibrated.

For fixture m and survival event S, define

$$q_{mr} = P(S \mid R_m = r), \qquad L_m = q_{mW} - q_{mL}.$$

These quantities are extracted from one joint simulation, retaining conditional sample counts and Monte Carlo standard errors. The largest fixture importance is Lecce (7.94 percentage points). Inter's

importance is 3.49 percentage points. To distinguish the two improvements involving a draw, define

$$U_m = q_{mW} - q_{mD}, \qquad A_m = q_{mD} - q_{mL}, \qquad L_m = U_m + A_m.$$

Here U_m is the survival gain from a win rather than a draw, and A_m is the gain from a draw rather than a loss. Both include the associated opponent points and conditional scorelines; neither isolates the causal value of a league point. Table 4 shows the three fixtures with highest total importance, Lazio as an illustrative contrasting profile, and Inter. These examples were selected retrospectively; all 12 fixtures are reported in the accompanying results.

Table 4: Fiorentina's conditional survival stakes at March 1, 2026. U and A are percentage-point differences; A/U is dimensionless. Brackets contain approximate 95% simulation-only intervals (normal for differences, Fieller for ratios), conditional on the fitted model. They exclude parameter and specification uncertainty.

Opponent	Venue	Win over draw U	Draw over loss A	Break-even A/U
Lecce	A	3.12 [2.52, 3.71]	4.82 [3.78, 5.86]	1.54 [1.08, 2.19]
Cremonese	A	2.83 [2.22, 3.44]	4.21 [3.22, 5.21]	1.49 [0.99, 2.21]
Verona	A	3.31 [2.63, 4.00]	2.74 [1.67, 3.81]	0.83 [0.45, 1.36]
Lazio	H	3.12 [2.52, 3.73]	1.99 [1.20, 2.79]	0.64 [0.34, 1.05]
Inter	H	1.76 [1.09, 2.43]	1.73 [1.07, 2.38]	0.98 [0.47, 2.04]

Against Lecce, avoiding defeat has a survival difference of 4.82 percentage points, versus 3.12 for turning a draw into a win. Against Lazio, the corresponding differences are 1.99 and 3.12. This distinction describes the consequences of results; it does not establish which changes a team can achieve.

Fixture importance is not the expected cost of resting a player. If a hypothetical intervention changes only the W/D/L probabilities from p_r to p'_r while preserving conditional scorelines, all other fixtures and the playoff model, then

$$\Delta P(S) = \sum_{r \in \{W,D,L\}} (p'_r - p_r) q_{mr}.$$

A shift of δ probability specifically from win to loss has cost δL_m; shifts involving draws need the draw conditional as well. In particular, consider moving probability $a \geq 0$ from draws to wins and $b \geq 0$ from draws to losses, with $a + b \leq p_D$. Then

$$p'_W = p_W + a, \quad p'_D = p_D - a - b, \quad p'_L = p_L + b, \qquad \Delta P(S) = aU_m - bA_m.$$

For $U_m > 0$ and $b > 0$, survival improves when $a/b > A_m/U_m$. Lecce's estimated break-even ratio is 1.54 (interval [1.08, 2.19]), compared with 0.64 ([0.34, 1.05]) for Lazio. Thus the estimated probability trade-off varies by fixture. These intervals do not establish a precise ordering of all fixtures or account for uncertainty in a proposed action's effects.

The same hypothetical shift changes expected league points by $2a - b$. For illustration, moving five percentage points of probability from draws to wins and another five to losses adds 0.05 expected points. It changes simulated survival by -0.085 percentage points against Lecce (interval [-0.156, -0.014]), and +0.057 against Lazio ([-0.006, +0.119]), whose interval includes zero. The effects are small; they illustrate that an expected-points improvement need not improve modeled survival. The shifts are feasible in the empirical outcome distribution of each displayed fixture, but are assumed scenarios rather than estimated tactics. Each calculation changes one fixture separately; the effects cannot generally be added across simultaneous changes. Appendix B describes the simulation intervals, including the shared draw conditional.

Actual rotation can also alter score margins, later fatigue and other fixtures. No player-level causal model is estimated here, so the table identifies fixtures for further analysis rather than prescribes a lineup.

Under the independent-fixture, fixed-parameter simulator, conditioning on an observed simulated result is equivalent to drawing the focal scoreline from its conditional distribution and resimulating all other fixtures and independent playoff randomness. It is not equivalent to fixing an arbitrary scoreline, such as 1–0, and does not generally survive conditioning on shared latent strengths.

7 Limitations and implications

The revised empirical claims are tied to the implemented models, forecast timings and evaluation periods. Shots on target are an imperfect chance-quality proxy; a shared decay and Poisson count model do not exhaust their predictive potential. The closing benchmark is stronger in information timing than a fortnightly structural forecast. A wider market/structural combination class may perform differently.

The season analysis adds evidence beyond a single case, but remains retrospective, small at the season level, and dependent on fixed-rate assumptions. Improved survival forecasting against a league-average baseline would not alone demonstrate practical benefit from player rotation. Prospective validation should archive forecasts before outcomes, compare stronger same-time baselines, and estimate intervention effects independently of fixture importance. The accompanying study protocol specifies these steps.

Competition rules can change, and penalty strength and exceptional venue decisions are additional modeling assumptions. League points, top-six position and qualification for a particular European competition must not be equated; the latter also depends on cup and eligibility rules outside this study.

Reproducibility materials include the implementation, regression tests, forecast-level outputs, input hashes, exact dependency versions and scripts generating the reported tables and figures. Raw source files are held locally; public redistribution and the conference's data-sharing requirement need permission review. No independent audit or large simulation verification is claimed without a corresponding executable artifact.

8 Conclusion

The structural model improves final-points predictions against the league-average comparator at all three dates, while survival-score gains remain uncertain. Those results support evaluating forecasts against the specific season outcome of interest, rather than treating points accuracy as evidence of survival-probability quality. The origin-defined subgroup analysis retains that retrospective, exploratory interpretation.

Separately, the Fiorentina example shows why avoiding defeat and turning a draw into a win need different conditional values: a hypothetical probability shift can improve expected points while reducing modeled survival. Its effects are small and conditional on the simulator, and no tactic is shown to deliver the assumed change. The match-level comparison likewise establishes neither informational completeness of betting prices nor a decision-policy benefit. Prospective forecast validation and independently estimated action effects remain necessary.

A Reproduction and revision record

The repository's `README.md` lists the complete execution order. The validation decay sweep precedes both match experiments and the season study. Saved forecasts record origin, price provenance, convergence and minimum forecast tau. Tables and numeric manuscript macros are generated from saved reports, rather than independently transcribed.

This version replaces an invalid shots-count dependence fit with an independent-Poisson likelihood, unifies unseen-club rate handling, constrains goals-model probability support, replaces mixed pre-closing/closing prices, and applies season-specific competition rules. It also narrows the original information and decision claims, adds direct prior work, removes unsupported audit and equivalence claims, and labels new historical analyses as retrospective. The new numbers supersede the prior manuscript. The draw-versus-win extension derives conditional stakes, simulation-only intervals and probability-transfer scenarios from the saved Fiorentina conditional probabilities and sample counts; it introduces no new match forecasts or player-effect estimates. The subsequent points-versus-survival extension joins the saved all-club forecasts to standings reconstructed strictly at each origin, adds a retrospective bottom-eight breakdown, and leaves the fitted forecasts intact.

B Simulation uncertainty for conditional stakes

Let $v_r = \hat{q}_r(1 - \hat{q}_r)/n_r$. Conditional on the W/D/L sample counts, the independent season draws give

$$\mathrm{Var}(\hat{U}) = v_W + v_D, \qquad \mathrm{Var}(\hat{A}) = v_D + v_L, \qquad \mathrm{Cov}(\hat{U}, \hat{A}) = -v_D.$$

Normal intervals use these variances. Ratio intervals use the Fieller set

$$(\hat{A} - t\hat{U})^2 \le z_{.975}^2 \left\{ \mathrm{Var}(\hat{A}) - 2t\,\mathrm{Cov}(\hat{U}, \hat{A}) + t^2\,\mathrm{Var}(\hat{U}) \right\}.$$

If the denominator is poorly resolved, the set can be unbounded or disconnected; the accompanying output retains these cases rather than forcing a finite interval. The scenario variance is $a^2 v_W + (a + b)^2 v_D + b^2 v_L$. These are pointwise Monte Carlo approximations conditional on fitted assumptions, not season-bootstrap intervals or simultaneous comparisons. More simulations would not resolve uncertainty about the model or an actual action. The full fixture report includes counts, covariance, interval status, and feasibility checks using the empirical W/D/L frequencies.

References

Dixon, M. J. and Coles, S. G. (1997). Modelling association football scores and inefficiencies in the football betting market. *Applied Statistics*, 46(2), 265–280.

Shin, H. S. (1992). Prices of state contingent claims with insider traders, and the favourite-longshot bias. *Economic Journal*, 102(411), 426–435.

Egidi, L., Pauli, F. and Torelli, N. (2018). Combining historical data and bookmakers' odds in modelling football scores. `https://arxiv.org/abs/1802.08848`.

Reade, J. J., Singleton, C. and Vaughan Williams, L. (2020). Betting markets for English Premier League results and scorelines: evaluating a forecasting model. University of Reading Economics Discussion Paper 2020-03. `https://www.reading.ac.uk/web/files/economics/emdp202003.pdf`.

Wheatcroft, E. (2021). Forecasting football matches by predicting match statistics. *Journal of Sports Analytics*. https://doi.org/10.3233/JSA-200462.

Link, D. and de Lorenzo, M. F. (2016). Seasonal pacing—Match importance affects activity in professional soccer. *PLOS ONE*, 11(6), e0157127. https://doi.org/10.1371/journal.pone.0157127.

Scarf, P. A. and Shi, X. (2008). The importance of a match in a tournament. *Computers & Operations Research*, 35(7), 2406–2418. https://doi.org/10.1016/j.cor.2006.11.005.

de Lorenzo, M., O'Bree, B. and Grundy, I. (2016). Simplifying probabilistic match importance calculation in football. In R. Stefani and A. Schembri (eds.), *Proceedings of the 13th Australasian Conference on Mathematics and Computers in Sport*, pp. 157–162. ANZIAM MathSport, Melbourne, 11–13 July 2016. https://www.mathsportinternational.com/anziam/Mathsport%2013.pdf.

Football-Data (accessed September 2026). Data files and notes, including the Pinnacle stale-price warning. https://football-data.co.uk/data.

FIGC (2022). Comunicato Ufficiale 304/A: Serie A 2022/23 ranking rules. https://www.figc.it/media/173115/304-deroga-art-51-noif-determinazione-classifica-campionato-serie-a-ss-2022-2023.pdf.

FIGC (2023a). Federal Council decisions, 7 July 2023, Serie A ranking rules. https://www.figc.it/media/208310/comunicato-stampa-cf-7-luglio-2023.pdf.

Lega Serie A (2025). *Regolamento Generale Competizioni 2025/26*, p. 44, reproducing FIGC 331/A, 19 June 2025. https://img.legaseriea.it/vimages/68a58bc9/Regolamento%20Generale%20Competizioni%20Lega%20Serie%20A_2025-26.pdf.

FIGC (2023b). Plusvalenza investigation: Juventus handed 15-point penalty, 20 January 2023. https://www.figc.it/en/figc/news/plusvalenza-investigation-juventus-handed-15-point-penalty-kyqu0aru.

CONI (2023). Collegio di Garanzia dello Sport, decision deposited 20 April 2023, appeals 13–19/2023. https://www.coni.it/images/collegiodigaranzia/Dispositivo_decisione_-_Ricc._13_14_15_16_17_18_19-2023.pdf.

FIGC (2023c). Juventus deducted ten points, 22 May 2023. https://www.figc.it/en/figc/news/juventus-deducted-ten-points-hpb6xbpy.