

# Brownian motion and stochastic integrals

**Rémy Degenne (Univ. Lille, Inria, CNRS, Centrale Lille, CRISAL)**

David Ledvinka (Imperial College London)

Etienne Marion (ENS Lyon)

Peter Pfaffelhuber (Universität Freiburg)

Other contributors: Markus Himmel, Jonas Bayer, Lorenzo Luccioli,  
Alessio Rondelli, Jérémy Scanvic, Pietro Monticone


12/12/2025

# Table of Contents

## 1 Brownian motion

- ▶ Brownian motion
- ▶ The Kolmogorov-Chentsov theorem
- ▶ Stochastic integrals

# Brownian motion



## Brownian motion, mathematically

Real valued Brownian motion indexed by time  $t \in \mathbb{R}_+$ :

- A **stochastic process**  $(B_t)_{t \in \mathbb{R}_+} : \mathbb{R}_+ \rightarrow \Omega \rightarrow \mathbb{R}$  such with a **specific distribution**
- Almost surely **continuous paths**:  $t \mapsto B_t(\omega)$  (in Lean:  $(B \cdot \omega)$ ) is continuous for almost all  $\omega$

Alternatively: the increments  $B_t - B_s$  are Gaussian with mean 0 and variance  $t - s$ , independent of  $B_u$  for  $u \leq s$ , and  $B_0 = 0$ . + continuity.

## Genesis of the project

With Peter Pfaffelhuber, we formalized Kolmogorov's extension theorem in 2023.

- A family  $(P_J)$  of probability measures on  $E^J$  for finite  $J \subseteq T$ , is projective if for all  $J \subseteq I$ , the pushforward of  $P_I$  by restriction to  $E^J$  is  $P_J$ .
- **Extension theorem**: a projective family  $(P_J)$  defines a unique measure on  $E^T$  whose finite dimensional marginals are the  $P_J$ .

Finite dimensional distributions of a process characterize its distribution.

Then we wanted to go further, towards Brownian motion.

## Genesis of the project

With Peter Pfaffelhuber, we formalized Kolmogorov's extension theorem in 2023.

- A family  $(P_J)$  of probability measures on  $E^J$  for finite  $J \subseteq T$ , is projective if for all  $J \subseteq I$ , the pushforward of  $P_I$  by restriction to  $E^J$  is  $P_J$ .
- **Extension theorem**: a projective family  $(P_J)$  defines a unique measure on  $E^T$  whose finite dimensional marginals are the  $P_J$ .

Finite dimensional distributions of a process characterize its distribution.

Then we wanted to go further, towards Brownian motion.

But we had no time and the project was stuck for two years.

## Genesis of the project

With Peter Pfaffelhuber, we formalized Kolmogorov's extension theorem in 2023.

- A family  $(P_J)$  of probability measures on  $E^J$  for finite  $J \subseteq T$ , is projective if for all  $J \subseteq I$ , the pushforward of  $P_I$  by restriction to  $E^J$  is  $P_J$ .
- **Extension theorem**: a projective family  $(P_J)$  defines a unique measure on  $E^T$  whose finite dimensional marginals are the  $P_J$ .

Finite dimensional distributions of a process characterize its distribution.

Then we wanted to go further, towards Brownian motion.

But we had no time and the project was stuck for two years.

We decided to turn it into a collaborative project.

First construct a “pre-Brownian”, with the right distributions but not necessarily continuous paths.

- Kolmogorov extension: sufficient to build a projective family of measures.
- For the Brownian motion, finite dimensional distributions are **multivariate Gaussians** with mean 0 and covariance  $\text{cov}(B_{t_i}, B_{t_j}) = \min(t_i, t_j)$ .

Then get a continuous modification.

- Build multivariate Gaussians (only Gaussians on  $\mathbb{R}$  in Matlab at the time).
- prove that family is projective
- prove the extension theorem to get a process with the right finite distributions.
- use the **Kolmogorov-Chentsov continuity theorem** to get a modification with continuous paths.



# *Inria* Project organization

## Preparation phase

- Choose a general proof strategy
- Identify references (textbooks, articles)
- Write a detailed blueprint

## Collaboration/formalization phase

- Zulip channel for discussions
- Public project on GitHub, with continuous integration
- Contributors make pull requests

PR phase (ongoing): move everything to Mathlib. 60+ pull requests so far.



## Project organization

### Preparation phase (3+ weeks, 2 contributors)

- Choose a general proof strategy
- Identify references (textbooks, articles)
- Write a detailed blueprint

### Collaboration/formalization phase (3 weeks, 9 contributors)

- Zulip channel for discussions
- Public project on GitHub, with continuous integration
- Contributors make pull requests

PR phase (ongoing): move everything to Mathlib. 60+ pull requests so far.

# *Inria* **Blueprint**

Developed by Patrick Massot for the Sphere Eversion project.

→ Let's look at it!

Writing a blueprint:

- detailed definitions, many small lemmas
- for people who don't know the full proof
- from Mathlib to the goal
- the dependency graph is very important

**Bottleneck of the project!**

## What did we formalize?

- Gaussian measures (in Banach spaces, Hilbert spaces, Euclidean spaces)
- Finite distributions of stochastic processes, modifications, indistinguishability
- Kolmogorov extension theorem
- Kolmogorov–Chentsov continuity theorem [Krätschmer & Urusov, 2023]
  - Covering numbers, chaining
- Putting it all together: real valued Brownian motion on  $\mathbb{R}_+$ , Wiener measure

**Brownian motion on  $\mathbb{R}_+$ :** a stochastic process  $(B_t)_{t \in \mathbb{R}_+}$  such that

- for  $(t_1, \dots, t_n) \in \mathbb{R}_+^n$ ,  $B_{t_1}, \dots, B_{t_n}$  are jointly Gaussian with mean 0 and covariance  $\text{cov}(B_{t_i}, B_{t_j}) = \min(t_i, t_j)$ ,
- $B$  has continuous paths.

# Table of Contents

## 2 The Kolmogorov-Chentsov theorem

- ▶ Brownian motion
- ▶ The Kolmogorov-Chentsov theorem
- ▶ Stochastic integrals

Modification:


- $(X_t)_{t \in T}$  and  $(Y_t)_{t \in T}$  are modifications of each other if for all  $t \in T$ ,  $X_t = Y_t$  almost surely.
- $\rightarrow$  Same distribution.
- Not necessarily indistinguishable:  $\mathbb{P}(\forall t \in T, X_t = Y_t)$  may be  $< 1$ .

**Kolmogorov-Chentsov theorem:** sufficient conditions for a process to have a modification with continuous paths.

We followed the proof from

[Krätschmer and Urusov, A Kolmogorov–Chentsov type theorem on general metric spaces with applications to limit theorems for Banach-valued processes, Journal of Theoretical Probability, 2023]

# Assumptions of the theorem



$T$ ,  $E$  extended pseudo-metric spaces with metrics  $d_T$ ,  $d_E$ .  $(X_t)_{t \in T}$  with values in  $E$ .

## Assumptions:

- $T$  has bounded covering number with exponent  $d > 0$ : for some  $C > 0$  and all  $\varepsilon > 0$ ,  $N_\varepsilon(T) \leq C\varepsilon^{-d}$
- Kolmogorov condition for exponents  $(p, q)$ , with  $q > 0$  and  $p > 0$ :  $(X_s, X_t)$  measurable for the Borel  $\sigma$ -algebra on  $E \times E$ , and

$$\exists M, \forall s, t \in T, \quad \mathbb{E}[d_E(X_s, X_t)^p] \leq M d_T(s, t)^q.$$

## The Kolmogorov-Chentsov theorem

$T$ ,  $E$  extended pseudo-metric spaces with metrics  $d_T$ ,  $d_E$ .  $(X_t)_{t \in T}$  with values in  $E$ .

### Theorem (Kolmogorov-Chentsov, Krätschmer & Urusov, 2023)

*Suppose that  $T$  has bounded covering number with exponent  $d > 0$  and that  $(X_t)_{t \in T}$  satisfies the Kolmogorov condition for exponents  $(p, q)$ , with  $q > d$  and  $p > 0$ . Then for all  $\beta \in (0, (q - d)/p)$  there exists a finite constant  $L$  such that for every countable subset  $T' \subseteq T$ ,*

$$\mathbb{E} \left[ \sup_{s, t \in T'} \frac{d_E(X_s, X_t)^p}{d_T(s, t)^{\beta p}} \right] \leq L.$$

*If furthermore  $E$  is complete, and  $T$  is second-countable, then the process  $X$  has a modification with Hölder continuous paths of exponent  $\beta$  for all  $\beta \in (0, (q - d)/p)$ .*



It's useful to have definitions focused on random variables

- `HasLaw X P Q`:  $X$  has law  $P$  on  $(\Omega, Q)$ .
- Could be `Q.map X = P`, but `HasLaw` also contains `AE measurable X Q`, to allow composition.
- `HasGaussianLaw`, random variable version of `IsGaussian`.

Constants are hard on paper, easy in Lean.

1. Constant improved in Lemma 1
2. error in lemma 8 where it changes a value
3. quick to fix!

Sometimes you don't use the theorem, but actually the particular proof.

Important in the K.C. theorem: the modification is built from limits.

$E$  not second-countable:

- Measurability of  $d_E(X_s, X_t)$  needs measurability of pairs  $(X_s, X_t)$ .
- If we build 2 modifications  $Y^{(1)}, Y^{(2)}$ , and want to combine them, we want measurability of  $(Y_s^{(1)}, Y_t^{(2)})$ .
- The theorem as stated does not give that. Our formal theorem does:

`IsLimitOfIndicator Y X P U`

Easy to track where assumptions are used.

$T$  and  $E$  not true metric spaces but pseudo-metric spaces (distance can be 0 between different points).

- The paper we follow has metric spaces
- We used that in a first formalization
- Then: change into pseudo-metric, see what breaks
- Many minor breaks. Two places needs thought.
- Focus on those two: zulip discussions. Adaptation of the proof, thanks to Sébastien Gouëzel.

## Did we formalize the right thing?

The computer checks the proofs, but not the definitions

If it looks like a duck, swims like a duck, and quacks like a duck, then it probably is a duck.

- $B$  has stationary increments: for all  $s \leq t \in \mathbb{R}_+$ ,  $B_t - B_s$  has law  $N(0, t - s)$ ,
- $B$  has independent increments,
- a process with independent increments and such that  $X_t$  has law  $N(0, t)$  for all  $t \in \mathbb{R}_+$  has the same distribution as  $B$ ,
- for  $c > 0$ , the process  $B_{ct}/\sqrt{c}$  is a Brownian motion,
- for  $t_0 \in \mathbb{R}_+$ , the process  $B_{t_0+t} - B_{t_0}$  is a Brownian motion,
- $tB_{1/t}$  is a Brownian motion,
- $B_t$  tends to 0 almost surely as  $t$  tends to 0.

- ▶ Brownian motion
- ▶ The Kolmogorov-Chentsov theorem
- ▶ Stochastic integrals

Goal: Stochastic integrals

$$\int X dB$$

- Project ongoing, accepting contributions.
- Change in organization: blueprint and code in parallel. Both collaboratively.
- Goal: Itô's lemma for integrals against cadlag semimartingales.
- Some steps: Début theorem, Doob-Meyer decomposition, quadratic variation, square integrable martingales...
- Recent progress: right-continuous filtrations, local properties, local martingales.

- We formalized Brownian motion in Lean: Gaussian measures, Kolmogorov extension, Kolmogorov-Chentsov theorem.
- We are PRing everything to Mathlib.
- You can contribute to the stochastic integral project!

- We formalized Brownian motion in Lean: Gaussian measures, Kolmogorov extension, Kolmogorov-Chentsov theorem.
- We are PRing everything to Mathlib.
- You can contribute to the stochastic integral project!

### Announcements:

- Conference **Lean Together 2026**: virtually via Zoom, 19-23 Jan 2026
- Mathematics in Lean session at the **International Congress on Mathematical Software 2026** (ICMS 2026), Waterloo, Ontario, Canada, 20-23 July 2026



- We formalized Brownian motion in Lean: Gaussian measures, Kolmogorov extension, Kolmogorov-Chentsov theorem.
- We are P<sub>R</sub>ing everything to Mathlib.
- You can contribute to the stochastic integral project!

### Announcements:

- Conference **Lean Together 2026**: virtually via Zoom, 19-23 Jan 2026
- Mathematics in Lean session at the **International Congress on Mathematical Software 2026** (ICMS 2026), Waterloo, Ontario, Canada, 20-23 July 2026

# Thank you!