

Addendum: "Note on the weak convergence of hyperplane α -quantile functionals and their continuity in the Skorokhod J1 topology"

P. M. Sparago

ARTICLE HISTORY

Compiled May 11, 2026

- (1) In the arguments for the proof of Th. 1.1, as is standard, we always assume that the variance-covariance matrix of a $\mathbb{R}^d$ -Brownian motion with nontrivial covariance structure is nondegenerate (hence, strictly positive-definite). Indeed, if the variance-covariance matrix is degenerate, the finite-dimensional distributions of the process do not have a Gaussian density, hence the process is not a *bona fide* Brownian motion in $\mathbb{R}^d$, but takes values only in a linear subspace.
- (2) Typo: Th. 2.7 should have s in the arguments $\inf_{s \leq t}(\dots)$ and $\sup_{s \leq t}(\dots)$, not t .
- (3) If $\gamma = 0$, Th. 1.1 is automatically true since the choice forces $x \mapsto (M_{t,\alpha}(x), \tau_{M_{t,\alpha}}(x)) = (0, 0)$, so that the case $\gamma = 0$ is implicitly ignored in the proofs. For clarity: Prop. 3.2, Prop. 3.6 and Le. 3.5 should read " $\gamma \in \mathbb{R}^d / \{0\}$ " instead of " $\gamma \in \mathbb{R}^d$ ".