

Visvesvaraya National Institute of Technology, Nagpur
Department of Mathematics

Assignment: Probability Theory(PT-1)

Topics: D.F., P.M.F., P.D.F.

Notations: DF-distribution function, PMF-probability mass function, PDF-probability density function

1. Do the following functions define DFs?

$$(i) F(x) = \begin{cases} 0 & \text{if } x < 0 \\ x & \text{if } 0 \leq x < \frac{1}{2} \\ 1 & \text{if } x \geq \frac{1}{2} \end{cases} \quad (ii) F(x) = \begin{cases} 0 & \text{if } x \leq 1 \\ 1 - \frac{1}{x} & \text{if } x > 1 \end{cases} \quad (iii) F(x) = \begin{cases} 0 & \text{if } x < 0 \\ 1 - e^{-x} & \text{if } x \geq 0 \end{cases}$$

2. Does the function $f_{\theta}(x) = \begin{cases} \theta^2 x e^{-\theta x} & \text{if } x > 0 \\ 0 & \text{if } x \leq 0 \end{cases}$ where $\theta > 0$, define a PDF? Find the DF associated with $f_{\theta}(x)$; if X is RV with PDF $f_{\theta}(x)$, find $P\{X \geq 1\}$.

3. Are the following functions distribution functions? IF so, find the corresponding density functions.

(i) $F(x) = 0$ if $x < -\theta$, $= \frac{1}{2} \left(\frac{x}{\theta} + 1 \right)$ if $|x| \leq \theta$, and 1 for $x > \theta$ where $\theta > 0$.

(ii) $F(x) = 0$ if $x < 0$, and $= 1 - e^{-x^2}$ if $x \geq 0$.

(iii) $F(x) = 0$ if $x < 0$, and $= 1 - (1 + x)e^{-x}$ if $x \geq 0$.

(iv) $F(x) = 0$ if $x < 1$, and $= \frac{(x-1)^2}{8}$ if $1 \leq x < 3$, and 1 for $x \geq 3$.

4. Determine the constant A in the following function, so that the function is PMF and find the DF $F(x)$ in each case. Plot PMF and DF.

$$(i) f(x) = A \left(\frac{1}{4} \right)^x, \quad x = 1, 2, 3, 4, \dots \quad (ii) f(x) = \begin{cases} A \binom{6}{x} \left(\frac{1}{4} \right)^x \left(\frac{3}{4} \right)^{6-x}, & x = 0, 1, 2, 3, 4, 5, 6 \\ 0 & \text{otherwise} \end{cases}$$

$$(iii) f(x) = \begin{cases} A \frac{3^x}{x!}, & x = 0, 1, 2, 3, \dots \\ 0 & \text{otherwise} \end{cases} \quad (iv) f(x) = \begin{cases} A \left(\frac{2}{3} \right)^{x-1}, & x = 1, 2, 3, \dots \\ 0 & \text{otherwise} \end{cases}$$

5. Determine the constant A in the following function, so that the function is PDF and find the DF $F(x)$ in each case. Plot PDF and DF.

$$(i) f(x) = \begin{cases} A[2 - |2 - x|], & 0 \leq x \leq 4 \\ 0 & \text{otherwise} \end{cases} \quad (ii) f(x) = \begin{cases} Ax^{-\frac{1}{2}}, & 0 < x < 1 \\ 0 & \text{otherwise} \end{cases}$$

$$(iii) f(x) = \begin{cases} A \frac{1}{\sqrt{4 - x^2}}, & |x| < 2 \\ 0 & \text{otherwise} \end{cases} \quad (iv) f(x) = \begin{cases} A \frac{(\log x)^2}{x} e^{-\frac{(\log x)^2}{2}}, & x > 0 \\ 0 & \text{otherwise} \end{cases}$$

$$(v) f(x) = Ae^{-|x|}, \quad -\infty < x < \infty \quad (vi) f(x) = \begin{cases} Ax^2, & 1 \leq x \leq 2 \\ Ax & 2 < x < 3 \\ 0 & \text{otherwise} \end{cases}$$

$$(vii) f(x) = Ae^{-\frac{(x-2)^2}{2}}, \quad -\infty < x < \infty.$$

6. A box contains good and defective items. If an item drawn is good, we assign the number 1 to the drawing, otherwise the number 0. Let p be the probability of the drawing at random a good item. Then find the DF $F(x)$.
7. Suppose that $P(X \geq x)$ is given for a RV X (of the continuous type) for all x . Find the PDF in each of the following cases:
- (i) $P(X \geq x) = 1$ if $x < 0$, and $= (1 + \frac{x}{\lambda})^{-\lambda}$ if $x \geq 0$, $\lambda > 0$ is a constant.
- (ii) $P(X \geq x) = 1$ if $x < x_0$, and $= (\frac{x_0}{x})^\alpha$ if $x \geq x_0$, $x_0 > 0$ and $\alpha > 0$ are constants.

8. Suppose one is told that the time, one has to wait (in mins.) for a bus on a certain bus stop is a random phenomenon, with the probability function specified by the PDF $f(x)$, given by

$$(i) f(x) = \begin{cases} 4x - 2x^2 - 1 & \text{for } 0 \leq x \leq 2 \\ 0 & \text{otherwise} \end{cases} \quad (ii) f(x) = \begin{cases} 4x - 2x^2 & \text{for } 0 \leq x \leq 2 \\ 0 & \text{otherwise} \end{cases}$$

$$(iii) f(x) = \begin{cases} \frac{3}{8}(4x - 2x^2) & \text{for } 0 \leq x \leq 2 \\ 0 & \text{otherwise} \end{cases}.$$

Is it possible for $f(x)$ to be a PDF, if so find the probability that one has to wait for a bus is (a) more than 1 min. (b) less than 1 min.

9. The length of time (in minutes) that a certain young lady speaks on the telephone is found to be a random phenomenon with the probability function specified by the PDF $f(\cdot)$, given by

$$f(x) = \begin{cases} Ae^{-\frac{x}{5}}, & x > 0 \\ 0 & x \leq 0 \end{cases}.$$

(i) Find the value of A that makes f a PDF. (ii) Show that the telephone conversation will last more than $a + b$ minutes given that it has lasted at least a minutes is equal to the unconditional probability that it will last more than b minutes.

10. The number of times a certain equipment operates before having to be discarded is found to be a random phenomenon with the probability function specified by the PMF $p(\cdot)$, given by

$$p(x) = \begin{cases} A \left(\frac{1}{3}\right)^x, & x = 0, 1, 2, 3, \dots \\ 0 & \text{otherwise} \end{cases}.$$

(i) Find the value of A that makes $p(\cdot)$ the PMF. (ii) What is the probability that equipment will operate more than 15 times given that it has operated more than 5 times.

11. For a certain random variable X the PDF $f(\cdot)$ is given by $f(x) = \frac{k}{1+x^2}$, $-\infty < x < \infty$.

(i) Find the value of k that makes $f(\cdot)$ the PDF and the DF $F(x)$ of X . (ii) Find $P\left(\frac{1}{3} < X^2 < 1\right)$.

12. For a certain random variable(RV) X the DF F is given by

$$F(x) = \begin{cases} 0 & \text{if } x \leq 1 \\ k(x-1)^4 & \text{if } 1 < x \leq 3 \\ 1 & \text{if } x > 3 \end{cases}.$$

(i) Find the value of k which makes F a DF. (ii) Find the PDF $f(x)$ and $P\{1.5 \leq X \leq 3.5\}$.