

#### 

### SALES



# My Name is Piyusha.

I have done a project in SQL named "Pizza Sales".
In this I have tried to solve some questions related to pizza sales by using SQL

Queries.

### RETRIEVE THE TOTAL NUMBER OF ORDERS PLACED?

SELECT

COUNT(\*) As total\_orders

FROM

orders;

total\_orders 21350





### CALCULATE THE TOTAL REVENUE GENERATED FROM PIZZA SALES?

#### SELECT

```
ROUND(SUM(order_details.quantity * pizzas.price),
2) AS total_revenue
```

#### FROM

```
order_details
JOIN
```

pizzas ON pizzas.pizza\_id = order\_details.pizza\_id;

#### total\_revenue ▶ 817860.05





#### IDENTIFY THE HIGHEST-PRICED PIZZA?

	name	price
<b>&gt;</b>	The Greek Pizza	35.95



## IDENTIFY THE MOST COMMON PIZZA SIZE ORDERED?

	size	order_count
<b>&gt;</b>	L	18526
	M	15385
	S	14137
	XL	544
	XXL	28



# LIST THE TOP 5 MOST ORDERED PIZZA TYPES ALONG WITH THEIR QUANTITIES?

```
SELECT
    pizza_types.name, SUM(order_details.quantity) AS quantity
FROM
    pizza_types
        JOIN
    pizzas ON pizza_types.pizza_type_id = pizzas.pizza_type_id
        JOIN
    order_details ON order_details.pizza_id = pizzas.pizza_id
GROUP BY pizza_types.name
ORDER BY quantity DESC
LIMIT 5;
```

	name	quantity
Þ	The Classic Deluxe Pizza	2453
	The Barbecue Chicken Pizza	2432
	The Hawaiian Pizza	2422
	The Pepperoni Pizza	2418
	The Thai Chicken Pizza	2371





## JOIN THE NECESSARY TABLES TO FIND THE TOTAL QUANTITY OF EACH PIZZA CATEGORY ORDERED?

```
SELECT
    pizza_types.category,
    SUM(order_details.quantity) AS quantity
FROM
    pizza_types
        JOIN
    pizzas ON pizza_types.pizza_type_id = pizzas.pizza_type_id
        JOIN
    order_details ON pizzas.pizza_id = order_details.pizza_id
GROUP BY pizza_types.category
ORDER BY quantity DESC;
```

	category	quantity
١	Classic	14888
	Supreme	11987
	Veggie	11649
	Chicken	11050





### DETERMINE THE DISTRIBUTION OF ORDERS BY HOUR OF THE DAY?

```
SELECT
```

HOUR(order\_time) AS hour, COUNT(order\_id) AS order\_count
FROM

orders

GROUP BY HOUR(order\_time);

	hour	order_count
<b>&gt;</b>	11	1231
	12	2520
	13	2455
	14	1472
	15	1468
	16	1920
	17	2336
	18	2399
	19	2009
	20	1642
	21	1198
	22	663
	23	28
	10	8
	9	1





### JOIN RELEVANT TABLES TO FIND THE CATEGORY-WISE DISTRIBUTION OF PIZZAS?

```
category, COUNT(name)
FROM
pizza_types
GROUP BY category;
```

	category	count(name)
Þ	Chicken	6
	Classic	8
	Supreme	9
	Veggie	9



### GROUP THE ORDERS BY DATE AND CALCULATE THE AVERAGE NUMBER OF PIZZAS ORDERED PER DAY?

```
SELECT
    ROUND(AVG(quantity), 0)
FROM
    (SELECT
         orders.order_date, SUM(order_details.quantity) AS quantity
    FROM
         orders
    JOIN order_details ON orders.order_id = order_details.order_id
    GROUP BY orders.order_date) AS order_quantity;
```

round(avg(quantity),0)

138





### DETERMINE THE TOP 3 MOST ORDERED PIZZA TYPES BASED ON REVENUE?

```
SELECT
    pizza_types.name,
    SUM(order_details.quantity * pizzas.price) AS revenue
FROM
    pizza_types
        JOIN
    pizzas ON pizza_types.pizza_type_id = pizzas.pizza_type_id
        JOIN
    order_details ON order_details.pizza_id = pizzas.pizza_id
GROUP BY pizza_types.name
ORDER BY revenue DESC
LIMIT 3;
```

name	revenue
The Thai Chicken Pizza	43434.25
The Barbecue Chicken Pizza	42768
The California Chicken Pizza	41409.5





### CALCULATE THE PERCENTAGE CONTRIBUTION OF EACH PIZZA TYPE TO TOTAL REVENUE?

```
SELECT
    pizza_types.category,
    ROUND(SUM(order_details.quantity * pizzas.price) / (SELECT
                    ROUND(SUM(order_details.quantity * pizzas.price),
                                2) AS total revenue
                FROM
                    order details
                        JOIN
                    pizzas ON pizzas.pizza_id = order_details.pizza_id) * 100,
            2) AS revenue
FROM
    pizza types
        JOIN
    pizzas ON pizza_types.pizza_type_id = pizzas.pizza_type_id
        JOIN
    order details ON order details.pizza id = pizzas.pizza id
GROUP BY pizza_types.category
```

	category	revenue
١	Classic	26.91
	Supreme	25.46
	Chicken	23.96
	Veggie	23.68





#### ANALYZE THE CUMULATIVE REVENUE GENERATED OVER TIME?

SELECT order\_date,

SUM(revenue) over(order by order\_date) AS cum\_revenue

FROM

(SELECT orders.order\_date,

SUM(order\_details.quantity \* pizzas.price) AS revenue

FROM order\_details JOIN pizzas

ON order\_details.pizza\_id = pizzas.pizza\_id

JOIN orders

ON orders.order\_id = order\_details.order\_id

GROUP BY orders.order\_date) AS sales;

	order_date	cum_revenue
Þ	2015-01-01	2713.85000000000004
	2015-01-02	5445.75
	2015-01-03	8108.15
	2015-01-04	9863.6
	2015-01-05	11929.55
	2015-01-06	14358.5
	2015-01-07	16560.7
	2015-01-08	19399.05
	2015-01-09	21526.4



# DETERMINATION ORDERED PIZ REVENUE CA

### DETERMINE THE TOP 3 MOST ORDERED PIZZA TYPES BASED ON REVENUE FOR EACH PIZZA CATEGORY?

(SELECT category, name, revenue,
rank() over(partition by category ORDER BY revenue DESC ) AS rn
FROM
(SELECT pizza\_types.category, pizza\_types.name,
SUM((order\_details.quantity) \* pizzas.price) AS revenue
FROM pizza\_types JOIN pizzas
ON pizza\_types.pizza\_type\_id = pizzas.pizza\_type\_id
JOIN order\_details
ON order\_details.pizza\_id = pizzas.pizza\_id
GROUP BY pizza\_types.category, pizza\_types.name) AS a) AS b
WHERE rn <= 3;</pre>

	name	revenue
Þ	The Thai Chicken Pizza	43434.25
	The Barbecue Chicken Pizza	42768
	The California Chicken Pizza	41409.5
	The Classic Deluxe Pizza	38180.5
	The Hawaiian Pizza	32273.25
	The Pepperoni Pizza	30161.75
	The Spicy Italian Pizza	34831.25



