

Solution

Section 1.3 - Quadratic Equations

Exercise

Solve: $x^2 = -25$

Solution

$$x = \pm\sqrt{-25}$$
$$= \pm 5i$$

Exercise

Solve: $x^2 = 49$

Solution

$$x = \pm 7$$

Exercise

Solve: $9x^2 = 100$

Solution

$$x^2 = \frac{100}{9}$$
$$x = \pm\sqrt{\frac{100}{9}}$$
$$= \pm \frac{10}{3}$$

Exercise

Solve: $4x^2 + 25 = 0$

Solution

$$4x^2 = -25$$
$$x^2 = -\frac{25}{4}$$
$$x = \pm\sqrt{-\frac{25}{4}}$$
$$= \pm \frac{5}{2}i$$

Exercise

Solve: $5x^2 + 35 = 0$

Solution

$$5x^2 = -35$$

$$x^2 = -7$$

$$\underline{x = \pm i\sqrt{7}}$$

Exercise

Solve: $5x^2 - 45 = 0$

Solution

$$5x^2 = 45$$

$$x = \frac{45}{5}$$

$$x^2 = 9$$

$$\underline{x = \pm 3}$$

Exercise

Solve: $(x - 4)^2 = 12$

Solution

$$x - 4 = \pm\sqrt{12}$$

$$x = 4 \pm \sqrt{12} \quad \sqrt{12} = \sqrt{4(3)} = 2\sqrt{3}$$

$$\underline{x = 4 \pm 2\sqrt{3}}$$

Exercise

Solve: $(x + 3)^2 = -16$

Solution

$$x + 3 = \pm\sqrt{-16}$$

$$\underline{x = -3 \pm 4i}$$

Exercise

Solve: $(x-2)^2 = -20$

Solution

$$x-2 = \pm\sqrt{-20}$$

$$x = \underline{2 \pm 4i\sqrt{5}} \quad |$$

Exercise

Solve: $(4x+1)^2 = 20$

Solution

$$4x+1 = \pm\sqrt{20}$$

$$4x = -1 \pm 2\sqrt{5}$$

$$x = \underline{\frac{-1 \pm 2\sqrt{5}}{4}} \quad |$$

Exercise

Solve $x^2 - 6x = -7$

Solution

$$x = \frac{-(-6) \pm \sqrt{(-6)^2 - 4(1)(7)}}{2(1)}$$

$$= \frac{6 \pm \sqrt{8}}{2}$$

$$= \frac{6 \pm 2\sqrt{2}}{2}$$

$$= \frac{2(3 \pm \sqrt{2})}{2}$$

$$= \underline{3 \pm \sqrt{2}} \quad |$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve $-6x^2 = 3x + 2$

Solution

$$6x^2 + 3x + 2 = 0$$

$$\begin{aligned}
 x &= \frac{-3 \pm \sqrt{3^2 - 4(6)(2)}}{2(6)} \\
 &= \frac{-3 \pm \sqrt{-39}}{12} \\
 &= \frac{-3}{12} \pm i \frac{\sqrt{39}}{12} \\
 &= \underline{-\frac{1}{4} \pm i \frac{\sqrt{39}}{12}}
 \end{aligned}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $3x^2 + 2x = 7$

Solution

$$3x^2 + 2x - 7 = 0 \quad \Rightarrow \quad \mathbf{a = 3, \quad b = 2, \quad c = -7}$$

$$x = \frac{-2 \pm \sqrt{4 - 4(3)(-7)}}{2(3)}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$= \frac{-2 \pm \sqrt{88}}{6}$$

$$= \frac{-2 \pm \sqrt{4(22)}}{6}$$

$$= \frac{-2 \pm 2\sqrt{22}}{6}$$

$$= \frac{2(-1 \pm \sqrt{22})}{6}$$

$$= \underline{-\frac{1 \pm \sqrt{22}}{3}}$$

$$x = -\frac{1}{3} \pm \frac{\sqrt{22}}{3}$$

Exercise

$3x^2 + 6 = 10x$

Solution

$$3x^2 - 10x + 6 = 0$$

$$x = \frac{-(-10) \pm \sqrt{(-10)^2 - 4(3)(6)}}{2(3)}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$\begin{aligned}
&= \frac{10 \pm \sqrt{100 - 72}}{6} \\
&= \frac{10}{6} \pm \frac{\sqrt{28}}{6} \\
&= \frac{5}{3} \pm \frac{2\sqrt{7}}{6} \\
&= \frac{5}{3} \pm \frac{\sqrt{7}}{3}
\end{aligned}$$

Exercise

Solve: $5x^2 + 2 = x$

Solution

$$5x^2 - x + 2 = 0$$

$$\begin{aligned}
x &= \frac{1 \pm \sqrt{1 - 40}}{10} \\
&= \frac{1 \pm \sqrt{-39}}{10} \\
&= \frac{1 \pm i\sqrt{39}}{10} \\
&= \frac{1}{10} \pm i\frac{\sqrt{39}}{10}
\end{aligned}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $5x^2 = 2x - 3$

Solution

$$5x^2 - 2x + 3 = 0$$

$$\begin{aligned}
x &= \frac{-(-2) \pm \sqrt{(-2)^2 - 4(5)(3)}}{2(5)} \\
&= \frac{2 \pm \sqrt{4 - 60}}{10} \\
&= \frac{2 \pm \sqrt{-56}}{10} \\
&= \frac{2 \pm i\sqrt{4(14)}}{10}
\end{aligned}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$\begin{aligned}
&= \frac{2 \pm i2\sqrt{14}}{10} \\
&= \frac{2}{10} \pm i \frac{2\sqrt{14}}{10} \\
&= \frac{1}{5} \pm i \frac{\sqrt{14}}{5}
\end{aligned}$$

Exercise

Solve: $x^2 + 8x + 15 = 0$

Solution

$$x = \frac{-8 \pm \sqrt{8^2 - 4(1)(15)}}{2(1)}$$

$$= \frac{-8 \pm \sqrt{64 - 60}}{2}$$

$$= \frac{-8 \pm \sqrt{4}}{2}$$

$$= \frac{-8 \pm 2}{2}$$

$$= \left\{ \begin{aligned} \frac{-8+2}{2} &= \frac{-6}{2} = -3 \\ \frac{-8-2}{2} &= \frac{-10}{2} = -5 \end{aligned} \right\}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $x^2 + 5x + 2 = 0$

Solution

$$x = \frac{-5 \pm \sqrt{5^2 - 4(1)(2)}}{2(1)}$$

$$= \frac{-5 \pm \sqrt{25 - 8}}{2}$$

$$= \frac{-5 \pm \sqrt{17}}{2}$$

$$= \frac{-5}{2} \pm \frac{\sqrt{17}}{2}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $x^2 + x - 12 = 0$

Solution

$$\begin{aligned}x &= \frac{-1 \pm \sqrt{1+48}}{2} \\&= \frac{-1 \pm 7}{2} \\&= \begin{cases} \frac{-1-7}{2} = -4 \\ \frac{-1+7}{2} = 3 \end{cases}\end{aligned}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $x^2 - 2x - 15 = 0$

Solution

$$\begin{aligned}x &= \frac{2 \pm \sqrt{4+60}}{2} \\&= \frac{2 \pm 8}{2} \\&= \begin{cases} \frac{2+8}{2} = 5 \\ \frac{2-8}{2} = -3 \end{cases}\end{aligned}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $x^2 - 4x - 45 = 0$

Solution

$$\begin{aligned}x &= \frac{4 \pm \sqrt{16+180}}{2} \\&= \frac{4 \pm \sqrt{196}}{2} \\&= \frac{4 \pm 14}{2} \\&= \begin{cases} \frac{4+14}{2} = 9 \\ \frac{4-14}{2} = -5 \end{cases}\end{aligned}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $x^2 - 6x - 10 = 0$

Solution

$$x^2 - 6x = 10$$

$$x^2 - 6x + \left(\frac{-6}{2}\right)^2 = 10 + \left(\frac{-6}{2}\right)^2$$

$$x^2 - 2(3)x + (3)^2 = 10 + 9$$

$$(x - 3)^2 = 19$$

$$x - 3 = \pm\sqrt{19}$$

$$\underline{x = 3 \pm \sqrt{19}}$$

Exercise

Solve: $2x^2 + 3x - 4 = 0$

Solution

$$x^2 + \frac{3}{2}x = 2$$

$$x^2 + \frac{3}{2}x + \left(\frac{1}{2} \frac{3}{2}\right)^2 = 2 + \left(\frac{1}{2} \frac{3}{2}\right)^2$$

$$x^2 + \frac{3}{2}x + \left(\frac{3}{4}\right)^2 = 2 + \frac{9}{16}$$

$$\left(x + \frac{3}{4}\right)^2 = \frac{41}{16}$$

$$x + \frac{3}{4} = \pm\sqrt{\frac{41}{16}}$$

$$\underline{x = -\frac{3}{4} \pm \frac{\sqrt{41}}{4}}$$

Exercise

Solve $x^2 - x + 8 = 0$

Solution

$$x = \frac{1 \pm \sqrt{1 - 32}}{2}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$= \frac{1 \pm \sqrt{-31}}{2}$$

$$= \frac{1 \pm i\sqrt{31}}{2}$$

Exercise

Solve $2x^2 - 13x = 1$

Solution

$$2x^2 - 13x - 1 = 0$$

$$x = \frac{13 \pm \sqrt{169 + 8}}{4}$$

$$= \frac{13 \pm \sqrt{169 + 8}}{4}$$

$$= \frac{13 \pm \sqrt{177}}{4}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve $r^2 + 3r - 3 = 0$

Solution

$$r = \frac{-3 \pm \sqrt{3^2 - 4(1)(-3)}}{2(1)}$$

$$= \frac{-3 \pm \sqrt{9 + 12}}{2}$$

$$= \frac{-3 \pm \sqrt{21}}{2}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $x^3 + 8 = 0$

Solution

$$(x + 2)(x^2 - 2x + 4) = 0$$

$$a^3 + b^3 = (a + b)(a^2 - ab + b^2)$$

$$x + 2 = 0$$

$$x^2 - 2x + 4 = 0$$

$$x = -2$$

$$\begin{aligned} x &= \frac{-(-2) \pm \sqrt{(-2)^2 - 4(1)(4)}}{2(1)} \\ &= \frac{2 \pm \sqrt{-12}}{2} \\ &= \frac{2 \pm 2i\sqrt{3}}{2} \\ &= \frac{2(1 \pm i\sqrt{3})}{2} \\ &= 1 \pm i\sqrt{3} \end{aligned}$$

The solution set is $\{-2, 1 \pm i\sqrt{3}\}$

Exercise

Solve: $4x^2 - 12x + 9 = 0$

Solution

$$\begin{aligned} x &= \frac{12 \pm \sqrt{144 - 144}}{8} \\ &= \frac{12}{8} \\ &= \frac{3}{2} \end{aligned}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $9x^2 - 30x + 25 = 0$

Solution

$$\begin{aligned} x &= \frac{30 \pm \sqrt{900 - 900}}{18} \\ &= \frac{30}{18} \\ &= \frac{5}{3} \end{aligned}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $x^2 - 14x + 49 = 0$

Solution

$$\begin{aligned}
 x &= \frac{14 \pm \sqrt{196 - 196}}{2} \\
 &= \frac{14}{2} \\
 &= \underline{7}
 \end{aligned}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $x^2 - 8x + 16 = 0$

Solution

$$\begin{aligned}
 x &= \frac{8 \pm \sqrt{64 - 64}}{2} \\
 &= \frac{8}{2} \\
 &= \underline{4}
 \end{aligned}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $x^2 + 6x + 13 = 0$

Solution

$$\begin{aligned}
 x &= \frac{-6 \pm \sqrt{36 - 52}}{2} \\
 &= \frac{-6 \pm \sqrt{-16}}{2} \\
 &= \frac{-6 \pm 4i}{2} \\
 &= \underline{-3 \pm 2i}
 \end{aligned}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $2x^2 - 2x + 13 = 0$

Solution

$$\begin{aligned}
 x &= \frac{2 \pm \sqrt{4 - 104}}{4} \\
 &= \frac{2 \pm \sqrt{-100}}{4}
 \end{aligned}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$= \frac{2 \pm 10i}{4}$$

$$= \frac{1}{2} \pm \frac{5}{2}i$$

Exercise

Solve: $x^2 + 2x + 29 = 0$

Solution

$$x = \frac{-2 \pm \sqrt{4 - 116}}{2}$$

$$= \frac{-2 \pm \sqrt{-112}}{2}$$

$$= \frac{-2 \pm 4i\sqrt{7}}{2}$$

$$= -1 \pm 2i\sqrt{7}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $4x^2 + 4x + 13 = 0$

Solution

$$x = \frac{-4 \pm \sqrt{16 - 16(13)}}{8}$$

$$= \frac{-4 \pm 4\sqrt{-12}}{8}$$

$$= \frac{-4 \pm 8i\sqrt{3}}{8}$$

$$= -\frac{1}{2} \pm i\sqrt{3}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $x^2 - 2x + 26 = 0$

Solution

$$x = \frac{2 \pm \sqrt{4 - 4(26)}}{2}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$= \frac{2 \pm 2\sqrt{-25}}{2}$$

$$= \underline{1 \pm 5i}$$

Exercise

Solve: $9x^2 - 4x + 20 = 0$

Solution

$$x = \frac{4 \pm \sqrt{16 - 16(45)}}{18}$$

$$= \frac{4 \pm 4\sqrt{-44}}{18}$$

$$= \frac{-4 \pm 8i\sqrt{11}}{8}$$

$$= \underline{-\frac{1}{2} \pm i\sqrt{11}}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $x^2 + 6x + 21 = 0$

Solution

$$x = \frac{-6 \pm \sqrt{36 - 84}}{2}$$

$$= \frac{-6 \pm \sqrt{-48}}{2}$$

$$= \frac{-6 \pm 4i\sqrt{3}}{2}$$

$$= \underline{-3 \pm 2i\sqrt{3}}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $9x^2 - 12x - 49 = 0$

Solution

$$x = \frac{12 \pm \sqrt{2^4 3^2 - 2^2 3^2 7^2}}{18}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$\begin{aligned}
&= \frac{12 \pm 6\sqrt{4-49}}{18} \\
&= \frac{12 \pm 6\sqrt{-45}}{18} \\
&= \frac{12 \pm 18i\sqrt{5}}{18} \\
&= \underline{\underline{\frac{2}{3} \pm i\sqrt{5}}}
\end{aligned}$$

Exercise

Solve: $x(x-3) = 18$

Solution

$$x^2 - 3x - 18 = 0$$

$$x = \frac{3 \pm \sqrt{9+72}}{2}$$

$$= \frac{3 \pm 9}{2}$$

$$= \left\{ \begin{array}{l} \frac{3+9}{2} = 6 \\ \frac{3-9}{2} = -3 \end{array} \right.$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $x(x-4) - 21 = 0$

Solution

$$x^2 - 4x - 21 = 0$$

$$x = \frac{4 \pm \sqrt{16+84}}{2}$$

$$= \frac{4 \pm 10}{2}$$

$$= \left\{ \begin{array}{l} \frac{4+10}{2} = 7 \\ \frac{4-10}{2} = -3 \end{array} \right.$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $(x-1)(x+4) = 14$

Solution

$$x^2 + 3x - 18 = 0$$

$$x = \frac{-3 \pm \sqrt{9 + 72}}{2}$$

$$= \frac{-3 \pm 9}{2}$$

$$= \left\{ \begin{array}{l} \frac{-3+9}{2} = 3 \\ \frac{-3-9}{2} = -6 \end{array} \right.$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $(x-3)(x+8) = -30$

Solution

$$x^2 + 5x + 6 = 0$$

$$x = \frac{-5 \pm \sqrt{25 - 24}}{2}$$

$$= \frac{-5 \pm 1}{2}$$

$$= \left\{ \begin{array}{l} \frac{-5+1}{2} = -2 \\ \frac{-5-1}{2} = -3 \end{array} \right.$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $x(x+8) = 16(x-1)$

Solution

$$x^2 + 8x = 16x - 16$$

$$x^2 - 8x + 16 = 0$$

$$x = \frac{8 \pm \sqrt{64 - 64}}{2}$$

$$= \frac{8}{2}$$

$$= 4$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $x(x+9) = 4(2x+5)$

Solution

$$x^2 + 9x = 8x + 20$$

$$x^2 + x - 20 = 0$$

$$x = \frac{-1 \pm \sqrt{1+80}}{2}$$

$$= \frac{-1 \pm 9}{2}$$

$$= \left\{ \begin{array}{l} \frac{-1+9}{2} = 4 \\ \frac{-1-9}{2} = -5 \end{array} \right.$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $(x+1)^2 = 2(x+3)$

Solution

$$x^2 + 2x + 1 = 2x + 6$$

$$x^2 = 5$$

$$x = \pm\sqrt{5}$$

Exercise

Solve: $(x+1)^2 - 5(x+2) = 3x+7$

Solution

$$x^2 + 2x + 1 - 5x - 10 = 3x + 7$$

$$x^2 - 6x - 16 = 0$$

$$x = \frac{6 \pm \sqrt{36+64}}{2}$$

$$= \frac{6 \pm 10}{2}$$

$$= \left\{ \begin{array}{l} \frac{6+10}{2} = 8 \\ \frac{6-10}{2} = -2 \end{array} \right.$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $x(8x+1) = 3x^2 - 2x + 2$

Solution

$$8x^2 + x = 3x^2 - 2x + 2$$

$$5x^2 + 3x - 2 = 0$$

$$x = \frac{-3 \pm \sqrt{9 + 40}}{10}$$

$$= \frac{-3 \pm 7}{2}$$

$$= \left\{ \begin{array}{l} \frac{-3+7}{10} = \underline{\underline{\frac{2}{5}}} \\ \frac{-3-7}{10} = \underline{\underline{-1}} \end{array} \right.$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

Solve: $x^2 + 6x - 7 = 0$

Solution

$$1 + 6 - 7 = 0 \quad a + b + c = 0$$

$$\underline{x = 1, -7} \quad x_1 = 1, \quad x_2 = \frac{c}{a}$$

Exercise

Solve: $x^2 - 6x - 7 = 0$

Solution

$$1 - (-6) - 7 = 0 \quad a - b + c = 0$$

$$\underline{x = -1, 7} \quad x_1 = -1, \quad x_2 = -\frac{c}{a}$$

Exercise

Solve: $3x^2 + 4x - 7 = 0$

Solution

$$3 + 4 - 7 = 0 \quad a + b + c = 0$$

$$\underline{x = 1, -\frac{7}{3}} \quad x_1 = 1, \quad x_2 = \frac{c}{a}$$

Exercise

Solve: $3x^2 - 4x - 7 = 0$

Solution

$$3 - (-4) - 7 = 0 \quad a - b + c = 0$$

$$\underline{x = -1, \frac{7}{3}} \quad x_1 = -1, \quad x_2 = -\frac{c}{a}$$

Exercise

Solve: $3x^2 - x - 2 = 0$

Solution

$$3 - 1 - 2 = 0 \quad a + b + c = 0$$

$$\underline{x = 1, -\frac{2}{3}} \quad x_1 = 1, \quad x_2 = \frac{c}{a}$$

Exercise

Solve: $3x^2 + x - 2 = 0$

Solution

$$3 - 1 - 2 = 0 \quad a - b + c = 0$$

$$\underline{x = -1, \frac{2}{3}} \quad x_1 = -1, \quad x_2 = -\frac{c}{a}$$

Exercise

Solve: $2x^2 + 3x - 5 = 0$

Solution

$$2 + 3 - 5 = 0 \quad a + b + c = 0$$

$$\underline{x = 1, -\frac{5}{2}} \quad x_1 = 1, \quad x_2 = \frac{c}{a}$$

Exercise

Solve: $2x^2 - 3x - 5 = 0$

Solution

$$2 - (-3) - 5 = 0 \quad a + b + c = 0$$

$$\underline{x = -1, \frac{5}{2}} \quad x_1 = -1, \quad x_2 = -\frac{c}{a}$$

Exercise

$$\text{Solve: } x^2 - 3x - 4 = 0$$

Solution

$$1 - (-3) - 4 = 0 \quad a - b + c = 0$$

$$\underline{x = -1, 4} \quad x_1 = -1, \quad x_2 = -\frac{c}{a}$$

Exercise

$$\text{Solve: } x^2 + 3x - 4 = 0$$

Solution

$$1 + 3 - 4 = 0 \quad a + b + c = 0$$

$$\underline{x = 1, -4} \quad x_1 = 1, \quad x_2 = \frac{c}{a}$$

Exercise

$$\text{Solve: } x^2 + 2x + 1 = 0$$

Solution

$$1 - 2 + 1 = 0 \quad a - b + c = 0$$

$$\underline{x = -1, -1} \quad x_1 = -1, \quad x_2 = -\frac{c}{a}$$

Exercise

$$\text{Solve: } 4x^2 - x - 5 = 0$$

Solution

$$4 - (-1) - 5 = 0 \quad a - b + c = 0$$

$$\underline{x = -1, \frac{5}{4}} \quad x_1 = -1, \quad x_2 = -\frac{c}{a}$$

Exercise

Solve for the specified variable $A = \frac{\pi d^2}{4}$, for d

Solution

$$\frac{4}{\pi} A = \frac{4}{\pi} \frac{\pi d^2}{4}$$

$$4A = 4 \frac{\pi d^2}{4}$$

$$\frac{4A}{\pi} = d^2$$

$$4A = \pi d^2$$

$$d^2 = \frac{4A}{\pi}$$

$$\frac{4A}{\pi} = \frac{\pi}{\pi} d^2$$

$$d = \pm \sqrt{\frac{4A}{\pi}}$$

$$= \pm 2 \frac{\sqrt{A}}{\sqrt{\pi}}$$

$$= \pm 2 \frac{\sqrt{A}}{\sqrt{\pi}} \frac{\sqrt{\pi}}{\sqrt{\pi}}$$

$$= \pm \frac{2\sqrt{\pi A}}{\pi}$$

Exercise

Solve for the specified variable $rt^2 - st - k = 0$ ($r \neq 0$), for t

Solution

$$t = \frac{-(-s) \pm \sqrt{(-s)^2 - 4(r)(-k)}}{2(r)}$$

$$t = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$t = \frac{s \pm \sqrt{s^2 + 4rk}}{2r}$$

Exercise

A rectangular park is 6 miles long and 2 miles wide. How long is a pedestrian route that runs diagonally across the park?

Solution

$$d^2 = 6^2 + 2^2$$

$$d^2 = 40$$

$$d = \sqrt{40}$$

$$\approx 6.32 \text{ miles}$$

Exercise

What is the width of a 25-inch television set whose height is 15 inches?

Solution

$$w^2 + 15^2 = 25^2$$

$$w^2 = 25^2 - 15^2$$

$$w = \sqrt{625 - 225}$$

$$= 20 \text{ in}$$

Exercise

The length of a rectangular sign is 3 feet longer than the width. If the sign's area is 54 square feet, find its length and width.

Solution

$$\ell = w + 3$$

$$\text{Area} = \ell w = 54$$

$$(w + 3)w = 54$$

$$w^2 + 3w - 54 = 0$$

$$w = \frac{-3 \pm \sqrt{9 + 216}}{2}$$

$$= \frac{-3 \pm 15}{2}$$

$$= \begin{cases} \frac{-3 - 15}{2} = -9 \\ \frac{-3 + 15}{2} = 6 \end{cases}$$

$$\ell = 6 + 3 = 9$$

\therefore the length of sign is 6 feet and width is 3 feet.

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exercise

A rectangular parking lot has a length that is 3 yards greater than the width. The area of the parking lot is 180 square yards, find the length and the width.

Solution

$$\ell = w + 3$$

$$\text{Area} = \ell w = 180$$

$$(w + 3)w = 180$$

$$w^2 + 3w - 180 = 0$$

$$\begin{aligned} w &= \frac{-3 \pm \sqrt{9 + 720}}{2} \\ &= \frac{-3 \pm 27}{2} \\ &= \begin{cases} \frac{-3 - 27}{2} = -\cancel{15} \\ \frac{-3 + 27}{2} = \underline{12} \end{cases} \end{aligned}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$\ell = 12 + 3 = \underline{15}$$

\therefore the length of sign is **15 feet** and width is **12 feet**.

Exercise

Each side of a square is lengthened by 3 inches. The area of this new, larger square is 64 square inches. Find the length of a side of the original square.

Solution

The new length of each side of a square is $= x + 3$

$$A = (x + 3)^2 = 64$$

$$x + 3 = \pm 8$$

$$x = -3 \pm 8$$

$$= \begin{cases} -3 - 8 = -\cancel{11} \\ -3 + 8 = \underline{5} \end{cases}$$

\therefore the length of the original square side is **5 inches**.

Exercise

Each side of a square is lengthened by 2 inches. The area of this new, larger square is 36 square inches. Find the length of a side of the original square.

Solution

The new length of each side of a square is $= x + 2$

$$A = (x + 2)^2 = 36$$

$$x + 2 = \pm 6$$

$$x = -2 \pm 6$$

$$= \begin{cases} -2 - 6 = -\cancel{8} \\ -2 + 6 = \underline{4} \end{cases}$$

\therefore the length of the original square side is **4 inches**.

Exercise

One number is 5 greater than another. The product of the numbers is 36. Find the numbers.

Solution

$$n = m + 5$$

$$P = mn = 36$$

$$m(m + 5) = 36$$

$$m^2 + 5m - 36 = 0$$

$$m = \frac{-5 \pm \sqrt{25 + 144}}{2}$$

$$= \frac{-5 \pm 13}{2}$$

$$= \left\{ \begin{array}{l} \frac{-5 - 13}{2} = -9 \\ \frac{-5 + 13}{2} = 4 \end{array} \right.$$

$$n = -9 + 5 = -4$$

$$n = 4 + 5 = 9$$

\therefore The numbers are 4 & 9 or -4 & -9

Exercise

One number is 6 less than another. The product of the numbers is 72. Find the numbers.

Solution

$$n = m - 6$$

$$P = mn = 72$$

$$m(m - 6) = 72$$

$$m^2 - 6m - 72 = 0$$

$$m = \frac{6 \pm \sqrt{36 + 288}}{2}$$

$$= \frac{6 \pm 18}{2}$$

$$= \left\{ \begin{array}{l} \frac{6 - 18}{2} = -6 \\ \frac{6 + 18}{2} = 12 \end{array} \right.$$

$$n = -6 - 6 = -12$$

$$n = 12 - 6 = 6$$

\therefore The numbers are 6 & 12 or -6 & -12

Exercise

A vacant rectangular lot is being turned into a community vegetable garden measuring 15 meters by 12 meters. A path of uniform width is to surround the garden. If the area of the garden and path combined is 378 square meters, find the width of the path.

Solution

$$\text{Area} = (15 + 2x)(12 + 2x)$$

$$378 = (15 + 2x)(12 + 2x)$$

$$378 = 180 + 30x + 24x + 4x^2$$

$$0 = 180 + 54x + 4x^2 - 378$$

$$0 = 4x^2 + 54x - 198$$

$$4x^2 + 54x - 198 = 0$$


$$x = \frac{-(54) \pm \sqrt{(54)^2 - 4(4)(-198)}}{2(4)}$$

$$= \frac{-54 \pm \sqrt{6084}}{8}$$

$$= \frac{-54 \pm 78}{8}$$

$$= \left\{ \begin{array}{l} \frac{-54 + 78}{8} = 3 \\ \frac{-54 - 78}{8} = -16.5 \end{array} \right.$$

\therefore the width of the path is 3 meters.



Exercise

A pool measuring 10 m by 20 m is surrounded by a path of uniform width. If the area of the pool and the path combined is 600 m^2 , what is the width of the path?

Solution

$$A = lw$$

$$600 = (20 + 2x)(10 + 2x)$$

$$600 = 200 + 40x + 20x + 4x^2$$


$$0 = -600 + 200 + 60x + 4x^2$$

$$0 = -400 + 60x + 4x^2$$

$$0 = -100 + 15x + x^2$$

$$x^2 + 15x - 100 = 0$$

$$x = \frac{-15 \pm \sqrt{15^2 + 400}}{2(1)}$$



$$= \frac{-15 \pm \sqrt{625}}{2}$$

$$= \begin{cases} \frac{-15-25}{2} = -20 \\ \frac{-15+25}{2} = 5 \end{cases}$$

∴ The width of the path is **5 m**

Exercise

You put in flower bed measuring 10 feet by 12 feet. You plan to surround the bed with uniform border of low-growing plants.

- Write a polynomial that describes the area of the uniform border that surrounds your flowers.
- The low growing plants surrounding the flower bed require 1 square foot each when mature. If you have 168 of these plants, how wide a strip around the flower bed should you prepare for the border?

Solution

$$a) \text{ Area} = 4x^2 + 2(12x) + 2(10x)$$

$$= 4x^2 + 44x$$

$$b) \quad A = 4x^2 + 44x = 168 \times 1$$

$$4x^2 + 44x - 168 = 0$$


$$x^2 + 11x - 42 = 0$$

$$x = \frac{-11 \pm \sqrt{121 + 168}}{2}$$

$$= \frac{-11 \pm 17}{2}$$

$$= \begin{cases} \frac{-11-17}{2} = -\cancel{14} \\ \frac{-11+17}{2} = 3 \end{cases}$$

∴ The width of the path is **3 feet**.



Exercise

A rectangular garden measures 80 feet by 60 feet. A large path of uniform width is to be added along both shorter sides and one longer side of the garden. The landscape designer doing the work wants to double the garden's area with the addition of this path. How wide should the path be?

Solution

$$\text{Total Area} = 2 \times (\text{area of the garden})$$

$$(80 + 2x)(60 + x) = 2(60)(80)$$

$$4800 + 200x + 2x^2 = 9600$$

$$2x^2 + 200x - 4800 = 0$$

$$x^2 + 100x - 2400 = 0$$


$$x = \frac{-100 \pm \sqrt{10,000 + 9,600}}{2}$$

$$= \frac{-100 \pm 10\sqrt{196}}{2}$$

$$= \frac{-100 \pm 140}{2}$$

$$= \begin{cases} \frac{-100 - 140}{2} = -\cancel{120} \\ \frac{-100 + 140}{2} = \underline{20} \end{cases}$$

\therefore the path should be **20 feet**.



Exercise

The length of a rectangular poster is 1 *foot* more than the width, and a diagonal of the poster is 5 *feet*. Find the length and the width.

Solution

Given: $\ell = w + 1$ $d = 5$

$$\ell^2 + w^2 = d^2$$

$$(w + 1)^2 + w^2 = 25$$

$$w^2 + 2w + 1 + w^2 = 25$$

$$2w^2 + 2w - 24 = 0$$

$$w^2 + w - 12 = 0$$

$$w = \underline{3}, \quad \cancel{4}$$

$$\ell = 3 + 1 = \underline{4}$$

\therefore The length is **4 feet** and the width is **3 feet**.

Exercise

One leg of a right triangle is 7 *cm* less than the length of the other leg. The length of the hypotenuse is 13 *cm*. find the lengths of the legs.

Solution

Given: $x = y - 7$ $d = 13$

$$x^2 + y^2 = d^2$$

$$(y - 7)^2 + y^2 = 169$$

$$y^2 - 14y + 49 + y^2 - 169 = 0$$

$$2y^2 - 14y - 120 = 0$$

$$y^2 - 7y - 60 = 0$$

$$y = \frac{7 \pm \sqrt{49 + 240}}{2}$$

$$= \frac{7 \pm \sqrt{289}}{2}$$

$$= \frac{7 \pm 17}{2}$$

$$= \begin{cases} \frac{7-17}{2} = -\cancel{5} \\ \frac{7+17}{2} = \underline{12} \end{cases}$$

$$\underline{y = 12}$$

$$x = 12 - 7 = \underline{5}$$

∴ The length of the leg: **5** & **12** cm.

Exercise

A tent with wires attached to help stabilize it, as shown below. The length of each wire is 8 feet greater than the distance from the ground to where it is attached to the tent.

The distance from the base of the tent to where the wire is anchored exceeds this height by 7 feet, Find the length of each wire used to stabilize the tent.

Solution


$$x^2 + (x + 7)^2 = (x + 8)^2$$

$$x^2 + x^2 + 14x + 49 = x^2 + 16x + 64$$

$$x^2 - 2x - 15 = 0$$

$$\underline{x = 5, \cancel{-3}}$$

∴ The length of each wire: **5** feet, **12** feet, and **13** feet.



Exercise

A boat is being pulled into a dock with a rope attached to the boat at water level. Where the boat is 12 ft from the dock, the length of the rope from the boat to the dock is 3 ft longer than twice the height of the dock above the water. Find the height of the dock.

Solution

$$(2h + 3)^2 = h^2 + 12^2$$

$$4h^2 + 12h + 9 = h^2 + 144$$

$$4h^2 + 12h + 9 - h^2 - 144 = 0$$


$$3h^2 + 12h - 135 = 0$$

$$h^2 + 4h - 45 = 0$$

$$(h + 9)(h - 5) = 0$$

$$h = -9, 5$$

Height = 5 ft.



Exercise

A piece of wire measuring 20 feet is attached to a telephone pole as a guy wire. The distance along the ground from the bottom of the pole to the end of the wire is 4 feet greater than the height where the wire is attached to the pole. How far up the pole does the guy wire reach?

Solution

$$(x + 4)^2 + x^2 = 20^2$$

$$x^2 + 8x + 16 + x^2 = 400$$

$$2x^2 + 8x - 384 = 0$$

$$x^2 + 4x - 192 = 0$$


$$x = \frac{-4 \pm \sqrt{16 + 768}}{2}$$

$$= \frac{-4 \pm \sqrt{784}}{2}$$

$$= \frac{-4 \pm 28}{2}$$

$$= \left\{ \begin{array}{l} \frac{-4 - 28}{2} = -16 \\ \frac{-4 + 28}{2} = 12 \end{array} \right.$$

\therefore the guy wire reaches the pole at 12 feet high.



Exercise

Logan and Cassidy leave a campsite, Logan biking due north and Cassidy biking due east. Logan bikes 7 km/h slower than Cassidy. After 4 hr, they are 68 km apart. Find the speed of each bicyclist.

Solution

$$4r^2 + [4(r - 7)]^2 = 68^2$$

$$16r^2 + 16(r^2 - 14r + 49) = 4624$$

$$16r^2 + 16r^2 - 224r + 784 = 4624$$

$$32r^2 - 224r + 784 - 4624 = 0$$

$$32r^2 - 224r - 3840 = 0$$

$$r^2 - 7r - 120 = 0$$

$$\rightarrow r = -8, 15$$

$$\Rightarrow \text{Cassidy's} = 15 \text{ km/h}$$

$$\Rightarrow \text{Logan's} = 8 \text{ km/h}$$



Exercise

Two trains leave a station at the same time. One train travels due west, and the other travels due south. The train traveling west travels 20 km/hr faster than the train traveling south. After 2 hr., the trains are 200 km apart. Find the speed of each train.

Solution

Given: $w = s + 20$ & $t = 2$

$$[2(s + 20)]^2 + (2s)^2 = 200^2$$

$$4(s^2 + 40s + 400) + 4s^2 = 40,000$$

$$s^2 + 40s + 400 + s^2 = 10,000$$

$$2s^2 + 40s + 9,600 = 0$$

$$s^2 + 20s + 4,800 = 0$$

$$\rightarrow s = \cancel{80}, 60$$

\therefore Speed of south train: 60 km/hr

Speed of west train: $60 + 20 = 80$ km/hr



Exercise

Towers are 1482 *feet* tall. How long would it take an object dropped from the top to reach the ground?

Given $s = t^2$

Solution

$$1482 = 16t^2$$

$$\frac{1482}{16} = t^2$$

$$t = \sqrt{\frac{1482}{16}}$$
$$= \frac{\sqrt{1482}}{4}$$
$$\approx 9.624 \text{ sec}$$

Exercise

The formula $P = 0.01A^2 + .05A + 107$ models a woman's normal Point systolic blood pressure, P , an age A . Use this formula to find the age, to the nearest year, of a woman whose normal systolic blood pressure is 115 mm Hg.

Solution

$$0.01A^2 + 0.05A + 107 = 115 \Rightarrow 0.01A^2 + 0.05A - 8 = 0$$

$$A = \frac{-0.05 \pm \sqrt{.05^2 - 4(.01)(-8)}}{2(.01)}$$

$$= \frac{-0.05 \pm \sqrt{.0025 + .32}}{.02}$$

$$= \frac{-0.05 \pm .567}{.02}$$

$$= \begin{cases} \frac{-0.05 - .567}{.02} = \cancel{31} \text{ (Not a Solution)} \\ \frac{-0.05 + .567}{.02} = 25.89 \approx 26 \end{cases}$$

Exercise

A rectangular piece of metal is 10 *in.* longer than it is wide. Squares with sides 2 *in.* long are cut from the four corners, and the flaps folded upward to form an open box. If the volume of the box is 832 in^3 , what were the original dimensions of the piece of metal?

Solution

$$l = w + 10$$

Bottom width: $w - 4$

Bottom length: $l - 4 = w + 10 - 4 = w + 6$

$$V = lwh = (w + 6)(w - 4)2$$

$$= 2(w^2 - 4w + 6w - 24)$$

$$= 2w^2 + 4w - 48$$

$$2w^2 + 4w - 48 = 832$$

$$2w^2 + 4w - 880 = 0$$

$$w^2 + 2w - 440 = 0$$

$$(w + 22)(w - 20) = 0$$

$$w + 22 = 0$$

$$w - 20 = 0$$

$$w = -22$$

$$w = 20$$

Width of the metal is 20 in by the length (20+10) 30 in.

Exercise

An astronaut on the moon throws a baseball upward. The astronaut is 6 ft., 6 in., tall, and the initial velocity of the ball is 30 ft/sec. The height s of the ball in feet is given by the equation

$$s = -2.7t^2 + 30t + 6.5$$

Where t is the number of seconds after the ball was thrown.

- After how many seconds is the ball 12 ft. above the moon's surface?
- How many seconds will it take for the ball to return to the surface?

Solution

- After how many seconds is the ball 12 ft above the moon's surface?

$$12 = -2.7t^2 + 30t + 6.5$$

$$0 = -2.7t^2 + 30t + 6.5 - 12$$

$$0 = -2.7t^2 + 30t - 5.5$$

$$t = \frac{-30 \pm \sqrt{(30)^2 - 4(-2.7)(-5.5)}}{2(-2.7)} \approx \frac{-30 \pm 29}{-5.4}$$

$$t \approx \frac{-30 - 29}{-5.4}$$

$$t \approx 10.9 \text{ sec}$$

$$t \approx \frac{-30 + 29}{-5.4}$$

$$t \approx .12 \text{ sec}$$

- How many seconds will it take for the ball to return to the surface?

$$0 = -2.7t^2 + 30t + 6.5$$

$$t = \frac{-30 \pm \sqrt{(30)^2 - 4(-2.7)(6.5)}}{2(-2.7)} \approx \frac{-30 \pm 31.15}{-5.4}$$

$$t \approx \frac{-30-31.15}{-5.4}$$

$$t \approx 11.32$$

$$t \approx \frac{-30+31.15}{-5.4}$$


$$t \approx -0.212$$

It will take 11.32 sec.

Exercise

The bar graph shows of SUVs (sport utility vehicles in the US, in *millions*. The quadratic equation

$S = .00579x^2 + .2579x + .9703$ models sales of SUVs from 1992 to 2003, where S represents sales in *millions*, and $x = 0$ represents 1992, $x = 1$ represents 1993 and so on.



- Use the model to determine sales in 2002 and 2003. Compare the results to the actual figures of 4.2 million and 4.4 million from the graph.
- According to the model, in what year do sales reach 3.5 million? Is the result accurate?

Solution

- For 2002 $\Rightarrow x = 10$

$$S = .00579(10)^2 + .2579(10) + .9703$$

$$\approx 4.1 \text{ million}$$

- For 2003 $\Rightarrow x = 11$

$$S = .00579(11)^2 + .2579(11) + .9703$$

$$\approx 4.5 \text{ million}$$

- $3.5 = .00579x^2 + .2579x + .9703$

$$0 = .00579x^2 + .2579x + .9703 - 3.5$$

$$0 = .00579x^2 + .2579x - 2.5297$$

$$x = \frac{-.2579 \pm \sqrt{(.2579)^2 - 4(.00579)(-2.5297)}}{2(.00579)}$$

$$= \frac{-.2579 \pm \sqrt{.1251}}{.01158}$$

$$x = \frac{-.2579 - .3537}{.01158}$$

$$x = \frac{-.2579 + .3537}{.01158}$$

$$x \approx -52.8$$

$$x \approx 8.3$$

According to the model, the number reached 3.5 million in the year 2000. The model closely matches the graph, so it is accurate

Exercise

Cynthia wants to buy a rug for a room that is 20 feet wide and 27 feet long. She wants to leave a uniform strip of floor around the rug. She can afford to buy 170 square feet of carpeting. What dimension should the rug have?

Solution

The area of the rug is:

$$(27 - 2x)(20 - 2x) = 170$$

$$540 - 54x - 40x + 4x^2 = 170$$

$$540 - 94x + 4x^2 - 170 = 0$$

$$4x^2 - 94x + 370 = 0$$



Solve for x .

$$\cancel{x = 18.5} \text{ or } x = 5$$

$$20 - 2x = 20 - 2(5) = 10$$

$$\text{and } 27 - 2x = 27 - 2(5) = 17$$

Therefore, the dimensions are: 10, 20 feet.



Exercise

Erik finds a piece of property in the shape of a right triangle. He finds that the longer leg is 20 m longer than twice the length of the shorter leg. The hypotenuse is 10 m longer than the length of the longer leg. Find the lengths of the sides of the triangular lot.

Solution

l : longer leg

s : shorter leg

Longer leg is 20 m longer than twice the length of the shorter leg

$$l = 2s + 20$$

The hypotenuse is 10 m longer than the length of the longer leg

$$h = l + 10$$

$$= 2s + 20 + 10$$

$$= 2s + 30$$

$$l^2 + s^2 = h^2$$

$$\begin{aligned}
(2s + 20)^2 + s^2 &= (2s + 30)^2 \\
4s^2 + 80s + 400 + s^2 &= 4s^2 + 120s + 900 \\
4s^2 + 80s + 400 + s^2 - 4s^2 - 120s - 900 &= 0 \\
s^2 - 40s - 500 &= 0 \\
(s + 10)(s - 50) &= 0 \\
s + 10 = 0 &\qquad s - 50 = 0 \\
s = -10 &\qquad s = 50
\end{aligned}$$

The shorter length is 50 *m*.

The longer length is $l = 2s + 20 = 2(50) + 20 = 120$

$$h = l + 10 = 120 + 10 = 130 \text{ m}$$

Exercise

An open box is made from a 10-*cm* by 20-*cm* piece of tin by cutting a square from each corner and folding up the edges. The area of the resulting base is 96 cm^2 . What is the length of the sides of the squares?

Solution

$$\begin{aligned}
\text{Area of the base} &= (10 - 2x)(20 - 2x) \\
&= 200 - 20x - 40x + 4x^2 \\
&= 4x^2 - 60x + 200
\end{aligned}$$


$$4x^2 - 60x + 200 = 96$$

$$4x^2 - 60x + 104 = 0$$

$$x^2 - 15x + 26 = 0$$

$$(x - 13)(x - 2) = 0$$

$$\begin{cases} x - 13 = 0 \rightarrow x = 13 \\ x - 2 = 0 \rightarrow x = 2 \end{cases} \Rightarrow x = 2 \text{ (only)}$$



Exercise

A square piece of cardboard is formed into a box by cutting out 3-*inch* squares from each of the corners and folding up the sides. If the volume of the box needs to be 126.75 cubic *inches*, what size square piece of cardboard is needed?

Solution

$$V = 3(x - 6)^2 = 126.75$$

$$(x - 6)^2 = 42.25$$


$$x - 6 = \sqrt{\frac{4225}{100}}$$

$$x = 6 + \frac{65}{10}$$

$$= 6 + \frac{13}{2}$$

$$= \frac{25}{2}$$

$$= \underline{12.5 \text{ in.}}$$



Exercise

You want to use 132 *feet* of chain-link fencing to enclose a rectangular region and subdivide the region into two smaller rectangular regions. If the total enclosed area is 576 *square feet*, find the dimensions of the enclosed region.

Solution

$$P = 2l + 3w = 132$$

$$l = \frac{1}{2}(132 - 3w)$$

$$A = lw = 576$$

$$\frac{w}{2}(132 - 3w) = 576$$

$$132w - 3w^2 = 1,152$$

$$3w^2 - 132w + 1,152 = 0$$

$$w^2 - 44w + 384 = 0$$

$$w = \frac{44 \pm \sqrt{1936 - 1536}}{2}$$

$$= \frac{44 \pm \sqrt{400}}{2}$$

$$= \frac{44 \pm 20}{2}$$


$$= \begin{cases} \frac{44 - 20}{2} = 12 \\ \frac{44 + 20}{2} = 32 \end{cases}$$

$$w = 12 \rightarrow l = \frac{1}{2}(132 - 36) = \underline{48}$$

$$w = 32 \rightarrow l = \frac{1}{2}(132 - 96) = \underline{18}$$

\therefore the dimensions: Length **48 feet**, width **12 feet**.

Or Length **18 feet**, width **32 feet**.



Exercise

How far is it from home plate to second base on a baseball diamond?


Solution

$$\begin{aligned}x^2 &= 90^2 + 90^2 \\&= 2(90^2)\end{aligned}$$

$$\underline{x = 90\sqrt{2}}$$

∴ The distance between home plate and second base is


$90\sqrt{2}$ feet



Exercise

Two equations can be used to track the position of a baseball t seconds after it is hit.

For instance, suppose $h = -16t^2 + 50t + 4.5$ gives the height, in *feet*, of a baseball t seconds after it is hit and $s = 103.9t$ gives the horizontal distance, in *feet*, of the ball from home plate t seconds after it is hit.



Use these equations to determine whether this particular baseball will clear a 10-foot fence positioned 360 feet from home plate.

Solution

$$s = 103.9t = 360$$

$$\begin{aligned}t &= \frac{360}{103.9} \\&= \frac{3600}{1039} \text{ sec} \quad \underline{\approx 3.46 \text{ sec}}\end{aligned}$$

$$\begin{aligned}h(3.46) &= -16(3.46)^2 + 50(3.46) + 4.5 \\&\quad \underline{\approx -14.05}\end{aligned}$$

Since the height is negative, then the ball hit the ground before the fence.

∴ The baseball will **not** clear the 10-foot fence.

Exercise

A ball is thrown downward with an initial velocity of 5 feet per second from the Golden Gate Bridge, which is 220 feet above the water. How long will it take for the ball to hit the water?

Solution

$$s(t) = -16t^2 - 5t + 220$$

$$s(t) = -\frac{1}{2}gt^2 + v_0t + s_0$$

$$-16t^2 - 5t + 220 = 0$$

$$t = \frac{5 \pm \sqrt{25 + 4(16)(220)}}{-32}$$

$$= \frac{5 \pm \sqrt{25 + 14,080}}{-32}$$

$$= \frac{-5 + \sqrt{14,105}}{32}$$

$$\therefore \text{It will take for the ball to hit the water } \frac{-5 + \sqrt{14,105}}{32} \approx 3.56 \text{ sec}$$

Exercise

A television screen measures 60 inches diagonally, and its aspect ratio is 16 to 9. This means that the ratio of the width of the screen to the height of the screen is 16 to 9.

Find the width and height of the screen.

Solution

$$(16x)^2 + (9x)^2 = 60^2$$

$$256x^2 + 81x^2 = 3600$$

$$337x^2 = 3600$$

$$x^2 = \frac{3600}{337}$$

$$x = \sqrt{\frac{3600}{337}}$$

$$= \frac{60}{\sqrt{337}} \text{ in.} \approx 3.268 \text{ in.}$$



$$\therefore \text{The width of TV is } 16 \times \frac{60}{\sqrt{337}} = \frac{960}{\sqrt{337}} \text{ in.} \approx 52 \text{ in.}$$

$$\text{The height of TV is } 9 \times \frac{60}{\sqrt{337}} = \frac{540}{\sqrt{337}} \text{ in.} \approx 29.4 \text{ in.}$$

Exercise

A company makes rectangular solid candy bars that measures 5 inches by 2 inches by 0.5 inch. Due to difficult financial times, the company has decided to keep the price of the candy bar fixed and reduce the volume of the bar by 20%. What should the dimensions be for the new candy bar if the company keeps the height at 0.5 inch and makes length of the candy bar 3 inches longer than the width?

Solution

The original volume is given:

$$V = 5 \times 2 \times \frac{1}{2}$$
$$= 5 \text{ in}^3$$

Reduction the volume of the bar by 20% which leave 80% of the new candy.

$$V_{\text{new}} = (.8)(5)$$
$$= 4 \text{ in}^3$$

$$V = lwh$$

$$4 = (w + 3)(w)\left(\frac{1}{2}\right)$$

$$w^2 + 3w = 8$$

$$w^2 + 3w - 8 = 0$$


$$w = \frac{-3 \pm \sqrt{9 + 32}}{2}$$

$$= \frac{-3 \pm \sqrt{41}}{2}$$

$$= \frac{-3 + \sqrt{41}}{2} \quad \text{and} \quad \cancel{\frac{-3 - \sqrt{41}}{2}} < 0$$

$$\therefore \text{The new width of the chocolate bar is } \frac{-3 + \sqrt{41}}{2} \approx 1.7 \text{ in.}$$

$$\text{The new length of the chocolate bar is } \frac{-3 + \sqrt{41}}{2} + 3 = \frac{3 + \sqrt{41}}{2} \approx 4.7 \text{ in.}$$



Exercise

A company makes rectangular solid candy bars that measures 5 inches by 2 inches by 0.5 inch. Due to difficult financial times, the company has decided to keep the price of the candy bar fixed and reduce the volume of the bar by 20%. What should the dimensions be for the new candy bar if the company keeps the height at 0.5 inch and makes length of the candy bar 2.5 times as long as its width?

Solution

The original volume is given:

$$V = 5 \times 2 \times \frac{1}{2}$$

$$= 5 \text{ in}^3$$

Reduction the volume of the bar by 20% which leave 80% of the new candy.

$$V_{\text{new}} = (.8)(5)$$

$$= 4 \text{ in}^3$$


$$V = lwh$$

$$4 = \left(\frac{3}{2}w\right)(w)\left(\frac{1}{2}\right)$$

$$3w^2 = 16$$

$$w^2 = \frac{16}{3}$$

$$w = \frac{4}{\sqrt{3}}$$



$$\therefore \text{The new width of the chocolate bar is } \frac{4\sqrt{3}}{3} \text{ in.}$$

$$\text{The new length of the chocolate bar is } 3\frac{4\sqrt{3}}{3} = 4\sqrt{3} \text{ in.}$$