

## **Solution**      **Section 1.6 – Surface Area**

### **Exercise**

Find the lateral (side) surface area of the cone generated by revolving the line segment  $y = \frac{x}{2}$ ,  $0 \leq x \leq 4$ , about the  $x$ -axis. Check your answer with the geometry formula

$$\text{Lateral surface area} = \frac{1}{2} \times \text{base circumference} \times \text{slant height}$$

### **Solution**

$$\frac{dy}{dx} = \frac{1}{2}$$

$$\sqrt{1 + \left(\frac{dy}{dx}\right)^2} = \sqrt{1 + \frac{1}{4}} = \frac{\sqrt{5}}{2}$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$


$$= 2\pi \int_0^4 \left(\frac{x}{2}\right) \frac{\sqrt{5}}{2} dx$$

$$= \frac{\pi\sqrt{5}}{2} \int_0^4 x dx$$

$$= \frac{\pi\sqrt{5}}{2} \frac{1}{2} x^2 \Big|_0^4$$

$$= \frac{\pi\sqrt{5}}{4} (4^2 - 0)$$

$$= \underline{4\pi\sqrt{5} \text{ unit}^2}$$



$$\text{base circumference} = 2\pi r = 2\pi(2) = 4\pi$$

$$\text{slant height} = \sqrt{4^2 + 2^2} = \sqrt{20} = 2\sqrt{5}$$

$$\text{Lateral surface area} = \frac{1}{2} \times \text{base circumference} \times \text{slant height}$$

$$= \frac{1}{2} \times (4\pi) \times (2\sqrt{5})$$

$$= \underline{4\pi\sqrt{5}}$$

### Exercise

Find the lateral surface area of the cone generated by revolving the line segment  $y = \frac{x}{2}$ ,  $0 \leq x \leq 4$ , about the y-axis. Check your answer with the geometry formula

$$\text{Lateral surface area} = \frac{1}{2} \times \text{base circumference} \times \text{slant height}$$

### Solution

$$y = \frac{x}{2} \Rightarrow x = 2y \rightarrow \begin{cases} x = 0 & \rightarrow y = 0 \\ x = 4 & \rightarrow y = 2 \end{cases}$$

$$\frac{dx}{dy} = 2$$

$$\sqrt{1 + \left(\frac{dy}{dx}\right)^2} = \sqrt{1 + 4} = \sqrt{5}$$

$$S = 2\pi \int_c^d x \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy$$


$$= 2\pi \int_0^2 2y \sqrt{5} dy$$

$$= 4\pi \sqrt{5} \int_0^2 y dy$$

$$= 4\pi \sqrt{5} \left. \frac{1}{2} y^2 \right|_0^2$$

$$= 2\pi \sqrt{5} (4 - 0)$$

$$= \underline{8\pi\sqrt{5} \text{ unit}^2}$$



$$\text{base circumference} = 2\pi(4) = 8\pi$$

$$\text{slant height} = \sqrt{4^2 + 2^2} = \sqrt{20} = 2\sqrt{5}$$

$$\text{Lateral surface area} = \frac{1}{2} \times \text{base circumference} \times \text{slant height}$$

$$= \frac{1}{2} \times (8\pi) \times (2\sqrt{5})$$

$$= \underline{8\pi\sqrt{5}}$$

### Exercise

Find the lateral surface area of the cone frustum generated by revolving the line segment

$y = \frac{x}{2} + \frac{1}{2}$ ,  $1 \leq x \leq 3$ , about the  $x$ -axis. Check your answer with the geometry formula

$$\text{Frustum surface area} = \pi(r_1 + r_2) \times \text{slant height}$$

### Solution

$$\frac{dy}{dx} = \frac{1}{2}$$

$$\sqrt{1 + \left(\frac{dy}{dx}\right)^2} = \sqrt{1 + \frac{1}{4}} = \frac{\sqrt{5}}{2}$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

$$= 2\pi \int_1^3 \left(\frac{x}{2} + \frac{1}{2}\right) \left(\frac{\sqrt{5}}{2}\right) dx$$

$$= \pi \frac{\sqrt{5}}{2} \int_1^3 (x+1) dx$$


$$= \pi \frac{\sqrt{5}}{2} \left[ \frac{1}{2}x^2 + x \right]_1^3$$

$$= \pi \frac{\sqrt{5}}{2} \left[ \frac{1}{2}(3)^2 + (3) - \left( \frac{1}{2}(1)^2 + (1) \right) \right]$$

$$= \pi \frac{\sqrt{5}}{2} \left[ \frac{9}{2} + 3 - \frac{3}{2} \right]$$

$$= \pi \frac{\sqrt{5}}{2} (6)$$

$$= \underline{3\pi\sqrt{5} \text{ unit}^2}$$



$$r_1 = \frac{1}{2} + \frac{1}{2} = 1 \quad r_2 = \frac{3}{2} + \frac{1}{2} = 2$$

$$\text{slant height} = \sqrt{(2-1)^2 + (3-1)^2} = \sqrt{5}$$

$$\text{Frustum surface area} = \pi(r_1 + r_2) \times \text{slant height}$$

$$= \pi(1+2)\sqrt{5}$$

$$= \underline{3\pi\sqrt{5}}$$

### Exercise

Find the lateral surface area of the cone frustum generated by revolving the line segment

$y = \frac{x}{2} + \frac{1}{2}$ ,  $1 \leq x \leq 3$ , about the  $y$ -axis. Check your answer with the geometry formula

$$\text{Frustum surface area} = \pi(r_1 + r_2) \times \text{slant height}$$

### Solution

$$y = \frac{x}{2} + \frac{1}{2} \rightarrow 2y = x + 1 \Rightarrow \boxed{x = 2y - 1}$$

$$\frac{dx}{dy} = 2$$

$$\sqrt{1 + \left(\frac{dx}{dy}\right)^2} = \sqrt{1 + 4} = \sqrt{5}$$

$$S = 2\pi \int_1^2 (2y - 1)(\sqrt{5}) dy$$

$$= 2\pi\sqrt{5} \int_1^2 (2y - 1) dy$$

$$= 2\pi\sqrt{5} \left[ y^2 - y \right]_1^2$$

$$= 2\pi\sqrt{5} \left[ (2)^2 - 2 - (1^2 - 1) \right]$$

$$= \underline{4\pi\sqrt{5} \text{ unit}^2}$$


$$r_1 = 1 \quad r_2 = 3$$

$$\text{slant height} = \sqrt{(2-1)^2 + (3-1)^2} = \sqrt{5}$$

$$\text{Frustum surface area} = \pi(r_1 + r_2) \times \text{slant height}$$

$$= \pi(1 + 3)\sqrt{5}$$

$$= \underline{4\pi\sqrt{5}}$$



### Exercise

Set up and evaluate the definite integral for the area of the surface generated by revolving the curve

$y = \frac{1}{3}x^3$  about the  $x$ -axis


### Solution

$$y = \frac{1}{3}x^3 \rightarrow y' = x^2$$

$$\sqrt{1+(y')^2} = \sqrt{1+x^4}$$

$$\begin{aligned} S &= 2\pi \int_0^3 \frac{1}{3}x^3 \sqrt{1+x^4} \, dx \\ &= \frac{\pi}{6} \int_0^3 (1+x^4)^{1/2} d(1+x^4) \\ &= \frac{\pi}{9} (1+x^4)^{3/2} \Big|_0^3 \\ &= \frac{\pi}{9} ((82)^{3/2} - 1) \\ &= \frac{\pi}{9} (82\sqrt{82} - 1) \end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} \, dx$$



### Exercise

Set up and evaluate the definite integral for the area of the surface generated by revolving the curve  $y = 2\sqrt{x}$  about the  $x$ -axis


### Solution

$$y = 2\sqrt{x} \rightarrow y' = \frac{1}{\sqrt{x}}$$

$$\sqrt{1+(y')^2} = \sqrt{1+\frac{1}{x}}$$

$$\begin{aligned} S &= 2\pi \int_4^9 2\sqrt{x} \sqrt{\frac{x+1}{x}} \, dx \\ &= 4\pi \int_4^9 (1+x)^{1/2} d(1+x) \\ &= \frac{8}{3} \pi (1+x)^{3/2} \Big|_4^9 \\ &= \frac{8}{3} \pi (10^{3/2} - 5^{3/2}) \approx 171.285 \end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} \, dx$$



### Exercise

Find the area of the surface generated by  $y = \frac{x^3}{9}$ ,  $0 \leq x \leq 2$ ,  $x$ -axis

### Solution

$$\frac{dy}{dx} = \frac{1}{3}x^2$$

$$\sqrt{1 + \left(\frac{dy}{dx}\right)^2} = \sqrt{1 + \frac{1}{9}x^4} = \frac{1}{3}\sqrt{9 + x^4}$$

$$S = 2\pi \int_0^2 \frac{x^3}{9} \frac{1}{3} \sqrt{9 + x^4} dx$$

$$= \frac{2\pi}{27} \int_0^2 x^3 \sqrt{9 + x^4} dx$$

$$= \frac{2\pi}{27} \int_9^{25} u^{1/2} \left(\frac{1}{4} du\right)$$

$$= \frac{\pi}{54} \int_9^{25} u^{1/2} du$$

$$= \frac{\pi}{54} \frac{2}{3} u^{3/2} \Big|_9^{25}$$



$$= \frac{\pi}{81} (25^{3/2} - 9^{3/2})$$

$$= \frac{98\pi}{81} \text{ unit}^2$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

$$u = 9 + x^4 \rightarrow du = 4x^3 dx \Rightarrow \frac{1}{4} du = x^3 dx$$

$$\rightarrow \begin{cases} x=2 & \rightarrow u=25 \\ x=0 & \rightarrow u=9 \end{cases}$$



### Exercise


Find the area of the surface generated by  $y = \sqrt{x+1}$ ,  $1 \leq x \leq 5$ ,  $x$ -axis

### Solution

$$y = \sqrt{x+1} = (x+1)^{1/2}$$

$$\frac{dy}{dx} = \frac{1}{2}(x+1)^{-1/2}$$

$$\begin{aligned}\sqrt{1 + \left(\frac{dy}{dx}\right)^2} &= \sqrt{1 + \frac{1}{4}(x+1)^{-1}} \\ &= \sqrt{1 + \frac{1}{4(x+1)}} \\ &= \sqrt{\frac{4x+4+1}{4(x+1)}} \\ &= \frac{1}{2}\sqrt{\frac{4x+5}{x+1}}\end{aligned}$$



$$S = 2\pi \int_1^5 \sqrt{x+1} \frac{1}{2} \frac{\sqrt{4x+5}}{\sqrt{x+1}} dx$$

$$= \pi \int_1^5 \sqrt{4x+5} dx$$

$$= \frac{\pi}{4} \int_1^5 (4x+5)^{1/2} d(4x+5)$$

$$= \frac{\pi}{6} (4x+5)^{3/2} \Big|_1^5$$

$$= \frac{\pi}{6} (25^{3/2} - 9^{3/2})$$

$$= \frac{\pi}{6} (98)$$

$$= \frac{49\pi}{3} \text{ unit}^2$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$


### Exercise

Find the area of the surface generated by  $y = \sqrt{2x - x^2}$ ,  $0.5 \leq x \leq 1.5$ ,  $x$ -axis

### Solution

$$\frac{dy}{dx} = \frac{1}{2}(2x - x^2)^{-1/2} (2 - 2x) = (1 - x)(2x - x^2)^{-1/2}$$

$$\begin{aligned}\sqrt{1 + \left(\frac{dy}{dx}\right)^2} &= \sqrt{1 + (1 - x)^2 (2x - x^2)^{-1}} \\ &= \sqrt{1 + \frac{1 - 2x + x^2}{2x - x^2}} \\ &= \sqrt{\frac{2x - x^2 + 1 - 2x + x^2}{2x - x^2}} \\ &= \sqrt{\frac{1}{2x - x^2}} \\ &= \frac{1}{\sqrt{2x - x^2}}\end{aligned}$$



$$\begin{aligned}S &= 2\pi \int_{.5}^{1.5} \sqrt{2x - x^2} \frac{1}{\sqrt{2x - x^2}} dx \\ &= 2\pi \int_{.5}^{1.5} dx \\ &= 2\pi x \Big|_{.5}^{1.5} = 2\pi(1.5 - .5) \\ &= \underline{2\pi \text{ unit}^2}\end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

### Exercise

Find the area of the surface generated by  $y = 3x + 4$ ,  $0 \leq x \leq 6$ , *revolved about x-axis*

### Solution

$$y' = 3$$

$$\begin{aligned}S &= 2\pi \int_0^6 (3x + 4) \sqrt{1 + 9} dx \\ &= 2\pi \sqrt{10} \left( \frac{3}{2}x^2 + 4x \right) \Big|_0^6 \\ &= 2\pi \sqrt{10} (54 + 24) \\ &= \underline{156\pi \sqrt{10} \text{ unit}^2}\end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$



### Exercise

Find the area of the surface generated by  $y = 12 - 3x$ ,  $1 \leq x \leq 3$ , revolved about  $x$ -axis

#### Solution

$$y' = -3$$

$$\begin{aligned} S &= 2\pi \int_1^3 (12 - 3x) \sqrt{1 + 9} \, dx \\ &= 2\pi \sqrt{10} \left( 12x - \frac{3}{2}x^2 \right) \Big|_1^3 \\ &= 2\pi \sqrt{10} \left( 36 - \frac{27}{2} - 12 + \frac{3}{2} \right) \\ &= \underline{24\pi \sqrt{10} \text{ unit}^2} \end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left( \frac{dy}{dx} \right)^2} \, dx$$

### Exercise

Find the area of the surface generated by  $y = 8\sqrt{x}$ ,  $9 \leq x \leq 20$ , revolved about  $x$ -axis

#### Solution

$$y' = \frac{4}{\sqrt{x}}$$

$$\begin{aligned} S &= 2\pi \int_9^{20} 8\sqrt{x} \sqrt{1 + \frac{16}{x}} \, dx \\ &= 16\pi \int_9^{20} \sqrt{x} \frac{\sqrt{x+16}}{\sqrt{x}} \, dx \\ &= 16\pi \int_9^{20} (x+16)^{1/2} \, d(x+16) \\ &= \frac{32\pi}{3} (x+16)^{3/2} \Big|_9^{20} \\ &= \frac{32\pi}{3} \left( (36)^{3/2} - (25)^{3/2} \right) \\ &= \frac{32\pi}{3} (216 - 125) \\ &= \underline{\frac{2912\pi}{3} \text{ unit}^2} \end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left( \frac{dy}{dx} \right)^2} \, dx$$

### Exercise

Find the area of the surface generated by  $y = x^3$ ,  $0 \leq x \leq 1$ , revolved about  $x$ -axis

#### Solution

$$y' = 3x^2$$

$$\begin{aligned} S &= 2\pi \int_0^1 x^3 \sqrt{1+9x^4} \, dx \\ &= \frac{\pi}{18} \int_0^1 (1+9x^4)^{1/2} d(1+9x^4) \\ &= \frac{\pi}{27} (1+9x^4)^{3/2} \Big|_0^1 \\ &= \frac{\pi}{27} \left( (10)^{3/2} - 1 \right) \\ &= \frac{\pi}{27} (10\sqrt{10} - 1) \text{ unit}^2 \end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left( \frac{dy}{dx} \right)^2} \, dx$$

### Exercise

Find the area of the surface generated by  $y = x^{3/2} - \frac{1}{3}x^{1/2}$ ,  $1 \leq x \leq 2$ , revolved about  $x$ -axis

### Solution

$$y' = \frac{3}{2}x^{1/2} - \frac{1}{6}x^{-1/2} = \frac{1}{2} \left( 3\sqrt{x} - \frac{1}{3\sqrt{x}} \right) = \frac{9x-1}{6\sqrt{x}}$$

$$\begin{aligned} S &= 2\pi \int_1^2 \left( x^{3/2} - \frac{1}{3}x^{1/2} \right) \sqrt{1 + \frac{(9x-1)^2}{36x}} \, dx \\ &= \frac{2}{3}\pi \int_1^2 \left( 3x^{3/2} - x^{1/2} \right) \frac{\sqrt{36x + 81x^2 - 18x + 1}}{6\sqrt{x}} \, dx \\ &= \frac{\pi}{9} \int_1^2 (3x-1) \sqrt{81x^2 + 18x + 1} \, dx \\ &= \frac{\pi}{9} \int_1^2 (3x-1) \sqrt{(9x+1)^2} \, dx \\ &= \frac{\pi}{9} \int_1^2 (3x-1)(9x+1) \, dx \\ &= \frac{\pi}{9} \int_1^2 (27x^2 - 6x - 1) \, dx \\ &= \frac{\pi}{9} \left( 9x^3 - 3x^2 - x \right) \Big|_1^2 = \frac{\pi}{9} (72 - 12 - 2 - 9 + 3 + 1) \\ &= \frac{53\pi}{9} \text{ unit}^2 \end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left( \frac{dy}{dx} \right)^2} \, dx$$

### Exercise

Find the area of the surface generated by  $y = \sqrt{4x+6}$ ,  $0 \leq x \leq 5$ , *revolved about x-axis*

#### Solution

$$y' = \frac{2}{\sqrt{4x+6}}$$

$$\begin{aligned} S &= 2\pi \int_0^5 \sqrt{4x+6} \sqrt{1 + \frac{4}{4x+6}} dx \\ &= 2\pi \int_0^5 \sqrt{4x+6} \sqrt{\frac{4x+6+4}{4x+6}} dx \\ &= 2\pi \int_0^5 (4x+10)^{1/2} dx \\ &= \frac{\pi}{2} \int_0^5 (4x+10)^{1/2} d(4x+10) \\ &= \frac{\pi}{3} (4x+10)^{3/2} \Big|_0^5 \\ &= \frac{\pi}{3} (30^{3/2} - 10^{3/2}) \\ &= \frac{\pi}{3} (30\sqrt{30} - 10\sqrt{10}) \\ &= \frac{10\pi\sqrt{10}}{3} (3\sqrt{3} - 1) \text{ unit}^2 \end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

### Exercise

Find the area of the surface generated by  $y = \frac{1}{4}(e^{2x} + e^{-2x})$ ,  $-2 \leq x \leq 2$ , *revolved about x-axis*

#### Solution

$$y' = \frac{1}{2}(e^{2x} - e^{-2x})$$

$$\begin{aligned} S &= 2\pi \int_{-2}^2 \frac{1}{4}(e^{2x} + e^{-2x}) \sqrt{1 + \frac{1}{4}(e^{2x} - e^{-2x})^2} dx \\ &= \frac{\pi}{2} \int_{-2}^2 (e^{2x} + e^{-2x}) \frac{1}{2} \sqrt{4 + e^{4x} - 2 + e^{-4x}} dx \\ &= \frac{\pi}{4} \int_{-2}^2 (e^{2x} + e^{-2x}) \sqrt{e^{4x} + 2 + e^{-4x}} dx \end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

$$\begin{aligned}
&= \frac{\pi}{4} \int_{-2}^2 \left( e^{2x} + e^{-2x} \right) \sqrt{\left( e^{2x} + e^{-2x} \right)^2} dx \\
&= \frac{\pi}{4} \int_{-2}^2 \left( e^{2x} + e^{-2x} \right)^2 dx \\
&= \frac{\pi}{4} \int_{-2}^2 \left( e^{4x} + 2 + e^{-4x} \right) dx \\
&= \frac{\pi}{4} \left( \frac{1}{4} e^{4x} + 2x - \frac{1}{4} e^{-4x} \right) \Big|_{-2}^2 \\
&= \frac{\pi}{4} \left( \frac{1}{4} e^8 + 4 - \frac{1}{4} e^{-8} - \frac{1}{4} e^{-8} + 4 + \frac{1}{4} e^8 \right) \\
&= \frac{\pi}{4} \left( \frac{1}{2} e^8 + 8 - \frac{1}{2} e^{-8} \right) \\
&= \frac{\pi}{8} \left( e^8 + 16 - e^{-8} \right) \text{ unit}^2
\end{aligned}$$

### Exercise

Find the area of the surface generated by  $y = \frac{1}{8}x^4 + \frac{1}{4x^2}$ ,  $1 \leq x \leq 2$ , revolved about  $x$ -axis

### Solution

$$y' = \frac{1}{2}x^3 - \frac{1}{2x^3} = \frac{1}{2} \frac{x^6 - 1}{x^3}$$

$$\begin{aligned}
S &= 2\pi \int_1^2 \left( \frac{1}{8}x^4 + \frac{1}{4x^2} \right) \sqrt{1 + \left( \frac{x^6 - 1}{2x^3} \right)^2} dx \\
&= \frac{\pi}{4} \int_1^2 \left( \frac{x^6 + 2}{x^2} \right) \sqrt{1 + \frac{x^{12} - 2x^6 + 1}{4x^6}} dx \\
&= \frac{\pi}{4} \int_1^2 \left( \frac{x^6 + 2}{x^2} \right) \sqrt{\frac{x^{12} + 2x^6 + 1}{4x^6}} dx \\
&= \frac{\pi}{4} \int_1^2 \left( \frac{x^6 + 2}{x^2} \right) \frac{\sqrt{(x^6 + 1)^2}}{2x^3} dx \\
&= \frac{\pi}{4} \int_1^2 \frac{(x^6 + 2)(x^6 + 1)}{2x^5} dx \\
&= \frac{\pi}{4} \int_1^2 \frac{x^{12} + 3x^6 + 2}{2x^5} dx
\end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left( \frac{dy}{dx} \right)^2} dx$$

$$\begin{aligned}
&= \frac{\pi}{8} \int_1^2 \left( x^7 + 3x + 2x^{-5} \right) dx \\
&= \frac{\pi}{8} \left( \frac{1}{8}x^8 + \frac{3}{2}x^2 - \frac{1}{2}x^{-4} \right) \Big|_1^2 \\
&= \frac{\pi}{8} \left( 32 + 6 - \frac{1}{32} - \frac{1}{8} - \frac{3}{2} + \frac{1}{2} \right) \\
&= \frac{\pi}{8} \left( 37 - \frac{5}{32} \right) \\
&= \underline{\underline{\frac{1179\pi}{256} \text{ unit}^2}}
\end{aligned}$$

### Exercise

Find the area of the surface generated by  $y = \frac{1}{3}x^3 + \frac{1}{4x}$ ,  $\frac{1}{2} \leq x \leq 2$ , revolved about  $x$ -axis

### Solution

$$y' = x^2 - \frac{1}{4x^2} = \frac{4x^4 - 1}{4x^2}$$

$$\begin{aligned}
S &= 2\pi \int_{1/2}^2 \left( \frac{1}{3}x^3 + \frac{1}{4x} \right) \sqrt{1 + \left( \frac{4x^4 - 1}{4x^2} \right)^2} dx \\
&= 2\pi \int_{1/2}^2 \left( \frac{4x^4 + 3}{12x} \right) \sqrt{1 + \frac{16x^8 - 8x^4 + 1}{16x^4}} dx \\
&= \frac{\pi}{6} \int_{1/2}^2 \left( \frac{4x^4 + 3}{x} \right) \sqrt{\frac{16x^8 + 8x^4 + 1}{16x^4}} dx \\
&= \frac{\pi}{6} \int_{1/2}^2 \left( \frac{4x^4 + 3}{x} \right) \frac{\sqrt{(4x^4 + 1)^2}}{4x^2} dx \\
&= \frac{\pi}{24} \int_{1/2}^2 \left( \frac{4x^4 + 3}{x^3} \right) (4x^4 + 1) dx \\
&= \frac{\pi}{24} \int_{1/2}^2 (4x + 3x^{-3})(4x^4 + 1) dx \\
&= \frac{\pi}{24} \int_{1/2}^2 (16x^5 + 16x + 3x^{-3}) dx \\
&= \frac{\pi}{24} \left( \frac{8}{3}x^6 + 8x^2 - \frac{3}{2}x^{-2} \right) \Big|_{1/2}^2 \\
&= \frac{\pi}{24} \left( \frac{512}{3} + 32 - \frac{3}{8} - \frac{1}{24} - 2 + 6 \right)
\end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left( \frac{dy}{dx} \right)^2} dx$$

$$\begin{aligned}
&= \frac{\pi}{24} \left( \frac{4086}{24} + 36 \right) \\
&= \frac{\pi}{24} \left( \frac{681}{4} + 36 \right) \\
&= \frac{\pi}{24} \left( \frac{825}{4} \right) \\
&= \frac{275\pi}{32} \text{ unit}^2
\end{aligned}$$

### Exercise

Find the area of the surface generated by  $y = \sqrt{5x - x^2}$ ,  $1 \leq x \leq 4$ , revolved about  $x$ -axis

### Solution

$$\begin{aligned}
y' &= \frac{5 - 2x}{2\sqrt{5x - x^2}} \\
1 + \left( \frac{dy}{dx} \right)^2 &= 1 + \frac{(5 - 2x)^2}{4(5x - x^2)} \\
&= \frac{20x - 4x^2 + 25 - 20x + 4x^2}{4(5x - x^2)} \\
&= \frac{25}{4(5x - x^2)}
\end{aligned}$$

$$\begin{aligned}
S &= 2\pi \int_1^4 \sqrt{5x - x^2} \sqrt{\frac{25}{4(5x - x^2)}} dx \\
&= 5\pi \int_1^4 dx \\
&= 5\pi x \Big|_1^4 \\
&= 15\pi \text{ unit}^2
\end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left( \frac{dy}{dx} \right)^2} dx$$

### Exercise


Set up an evaluate the definite integral for the area of the surface generated by revolving the curve about the  $x$ -axis

$$y = \frac{1}{6}x^3 + \frac{1}{2x}, \quad 1 \leq x \leq 2$$

### Solution


$$y' = \frac{1}{2}x^2 - \frac{1}{2x^2}$$

$$\begin{aligned}
 \sqrt{1+(y')^2} &= \sqrt{1+\frac{1}{4}x^4-\frac{1}{2}+\frac{1}{4x^4}} \\
 &= \sqrt{\frac{1}{4}x^4+\frac{1}{2}+\frac{1}{4x^4}} \\
 &= \sqrt{\left(\frac{1}{2}x^2+\frac{1}{2x^2}\right)^2} \\
 &= \frac{1}{2}x^2+\frac{1}{2x^2}
 \end{aligned}$$



$$\begin{aligned}
 S &= 2\pi \int_1^2 \left(\frac{1}{6}x^3 + \frac{1}{2x}\right) \left(\frac{1}{2}x^2 + \frac{1}{2x^2}\right) dx \\
 &= 2\pi \int_1^2 \left(\frac{x^5}{12} + \frac{x}{3} + \frac{1}{4x^3}\right) dx \\
 &= 2\pi \left(\frac{1}{72}x^6 + \frac{1}{6}x^2 - \frac{1}{8x^2}\right) \Big|_1^2 \\
 &= 2\pi \left(\frac{64}{72} + \frac{2}{3} - \frac{1}{32} - \frac{1}{72} - \frac{1}{6} + \frac{1}{8}\right) \\
 &= 2\pi \left(\frac{63}{72} + \frac{19}{32}\right) \\
 &= 2\pi \left(\frac{423}{288}\right) \\
 &= \frac{47\pi}{16}
 \end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$



### Exercise

Set up and evaluate the definite integral for the area of the surface generated by revolving the curve about the  $x$ -axis


$$y = \sqrt{4-x^2}, \quad -1 \leq x \leq 1$$

### Solution


$$\begin{aligned}
 y' &= \frac{-x}{\sqrt{4-x^2}} \\
 \sqrt{1+(y')^2} &= \sqrt{1+\frac{x^2}{4-x^2}} \\
 &= \sqrt{\frac{4}{4-x^2}}
 \end{aligned}$$

$$S = 2\pi \int_{-1}^1 \sqrt{4-x^2} \frac{2}{\sqrt{4-x^2}} dx$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$



$$\begin{aligned}
 &= 4\pi \int_{-1}^1 dx \\
 &= 4\pi x \Big|_{-1}^1 \\
 &= 8\pi
 \end{aligned}$$



### Exercise

Set up an evaluate the definite integral for the area of the surface generated by revolving the curve about

the  $x$ -axis  $y = \sqrt{9 - x^2}, \quad -2 \leq x \leq 2$

### Solution

$$y' = \frac{-x}{\sqrt{9 - x^2}}$$


$$\begin{aligned}
 \sqrt{1 + (y')^2} &= \sqrt{1 + \frac{x^2}{9 - x^2}} \\
 &= \sqrt{\frac{9}{9 - x^2}}
 \end{aligned}$$

$$S = 2\pi \int_{-2}^2 \sqrt{9 - x^2} \cdot \frac{3}{\sqrt{9 - x^2}} dx$$


$$= 6\pi \int_{-2}^2 dx$$

$$= 6\pi x \Big|_{-2}^2$$

$$= 24\pi$$



$$S = 2\pi \int_a^b y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$




### Exercise

Set up an evaluate the definite integral for the area of the surface generated by revolving the curve about the  $y$ -axis

### Solution

$$y' = \frac{1}{3}x^{-2/3} = \frac{1}{3x^{2/3}}$$

$$\begin{aligned}
 \sqrt{1 + (y')^2} &= \sqrt{1 + \frac{1}{9x^{4/3}}} \\
 &= \frac{\sqrt{9x^{4/3} + 1}}{3x^{2/3}}
 \end{aligned}$$






$$\begin{aligned}
 S &= 2\pi \int_1^8 x \frac{\sqrt{9x^{4/3}+1}}{3x^{2/3}} dx & S &= 2\pi \int_a^b x \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx \\
 &= \frac{2}{3}\pi \int_1^8 x^{1/3} \sqrt{9x^{4/3}+1} dx \\
 &= \frac{\pi}{18} \int_1^8 \left(9x^{4/3}+1\right)^{1/2} d\left(9x^{4/3}+1\right) \\
 &= \frac{\pi}{27} \left(9x^{4/3}+1\right)^{3/2} \Big|_1^8 \\
 &= \frac{\pi}{27} \left( \left(72(8)^{1/3}+1\right)^{3/2} - 10^{3/2} \right) \\
 &= \frac{\pi}{27} \left( 145\sqrt{145} - 10\sqrt{10} \right)
 \end{aligned}$$

### Exercise

Set up and evaluate the definite integral for the area of the surface generated by revolving the curve about the  $y$ -axis

#### Solution

$$\begin{aligned}
 y' &= -2x \\
 \sqrt{1+(y')^2} &= \sqrt{1+4x^2} \\
 S &= 2\pi \int_0^3 x \sqrt{1+4x^2} dx & S &= 2\pi \int_a^b x \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx \\
 &= \frac{\pi}{4} \int_0^3 \left(1+4x^2\right)^{1/2} d\left(1+4x^2\right) \\
 &= \frac{\pi}{6} \left(1+4x^2\right)^{3/2} \Big|_0^3 \\
 &= \frac{\pi}{6} \left( 37\sqrt{37} - 1 \right)
 \end{aligned}$$



### Exercise

Find the area of the surface generated by  $y = (3x)^{1/3}$ ;  $0 \leq x \leq \frac{8}{3}$  about  $y$ -axis

#### Solution

$$3x = y^3 \rightarrow x = \frac{1}{3}y^3 \Rightarrow x' = y^2 \quad \begin{cases} x=0 & \rightarrow y=0 \\ x=\frac{8}{3} & \rightarrow y=\left(3\frac{8}{3}\right)^{1/3} = 2 \end{cases}$$

$$\begin{aligned}
S &= 2\pi \int_0^2 \frac{1}{3} y^3 \sqrt{1+y^4} \, dy \\
&= \frac{\pi}{6} \int_0^2 (1+y^4)^{1/2} d(1+y^4) \\
&= \frac{\pi}{9} (1+y^4)^{3/2} \Big|_0^2 \\
&= \frac{\pi}{9} ((17)^{3/2} - 1) \\
&= \frac{\pi}{9} (17\sqrt{17} - 1)
\end{aligned}$$

$$S = 2\pi \int_c^d x \sqrt{1 + \left(\frac{dx}{dy}\right)^2} \, dy$$

### Exercise

Find the area of the surface generated of the curve  $y = 4x - 1$  between the points  $(1, 3)$  and  $(4, 15)$  about  $y$ -axis

#### Solution

$$y = 4x - 1 \rightarrow x = \frac{1}{4}(y + 1) \Rightarrow x' = \frac{1}{4}$$

$$\begin{aligned}
S &= 2\pi \int_3^{15} \frac{1}{4}(y+1) \sqrt{1 + \frac{1}{16}} \, dy \\
&= \frac{\pi}{2} \int_3^{15} (y+1) \sqrt{\frac{17}{16}} \, dy \\
&= \frac{\pi\sqrt{17}}{8} \left( \frac{1}{2}y^2 + y \right) \Big|_3^{15} \\
&= \frac{\pi\sqrt{17}}{8} \left( \frac{225}{2} + 15 - \frac{9}{2} - 3 \right) \\
&= \frac{\pi\sqrt{17}}{8} (120) \\
&= 15\pi\sqrt{17}
\end{aligned}$$

$$S = 2\pi \int_c^d x \sqrt{1 + \left(\frac{dx}{dy}\right)^2} \, dy$$

### Exercise

Find the area of the surface generated of the curve  $y = \frac{1}{2} \ln(2x + \sqrt{4x^2 - 1})$  between the points  $(\frac{1}{2}, 0)$  and  $(\frac{17}{16}, \ln 2)$  about  $y$ -axis

#### Solution

$$2y = \ln(2x + \sqrt{4x^2 - 1}) \rightarrow (2x + \sqrt{4x^2 - 1})^2 = (e^{2y})^2$$

$$4x^2 + 4x\sqrt{4x^2 - 1} + 4x^2 - 1 = e^{4y}$$

$$4x\left(2x + \sqrt{4x^2 - 1}\right) = e^{4y} + 1$$

$$2x + \sqrt{4x^2 - 1} = e^{2y}$$

$$4x\left(e^{2y}\right) = e^{4y} + 1$$

$$x = \frac{e^{4y} + 1}{4e^{2y}} = \frac{1}{4}\left(e^{2y} + e^{-2y}\right)$$

$$x' = \frac{1}{2}\left(e^{2y} - e^{-2y}\right)$$

$$S = 2\pi \int_0^{\ln 2} \frac{1}{4}\left(e^{2y} + e^{-2y}\right) \sqrt{1 + \frac{1}{4}\left(e^{2y} - e^{-2y}\right)^2} dy$$

$$S = 2\pi \int_c^d x \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy$$

$$= \frac{\pi}{4} \int_0^{\ln 2} \left(e^{2y} + e^{-2y}\right) \sqrt{4 + e^{4y} - 2 + e^{-4y}} dy$$

$$= \frac{\pi}{4} \int_0^{\ln 2} \left(e^{2y} + e^{-2y}\right) \sqrt{\left(e^{2y} + e^{-2y}\right)^2} dy$$

$$= \frac{\pi}{4} \int_0^{\ln 2} \left(e^{2y} + e^{-2y}\right)^2 dy$$

$$= \frac{\pi}{4} \int_0^{\ln 2} \left(e^{4y} + 2 + e^{-4y}\right) dy$$

$$= \frac{\pi}{4} \left( \frac{1}{4} e^{4y} + 2y - \frac{1}{4} e^{-4y} \right) \Big|_0^{\ln 2}$$

$$= \frac{\pi}{4} \left( \frac{1}{4} e^{4 \ln 2} + 2 \ln 2 - \frac{1}{4} e^{-4 \ln 2} - \frac{1}{4} + \frac{1}{4} \right)$$

$$= \frac{\pi}{4} \left( \frac{1}{4} e^{\ln 2^4} + 2 \ln 2 - \frac{1}{4} e^{\ln 2^{-4}} \right)$$

$$= \frac{\pi}{4} \left( \frac{1}{4} 2^4 + 2 \ln 2 - \frac{1}{4} 2^{-4} \right)$$

$$= \frac{\pi}{4} \left( 4 + 2 \ln 2 - \frac{1}{64} \right)$$

$$= \frac{\pi}{4} \left( \frac{255}{64} + 2 \ln 2 \right) \text{ unit}^2$$

### Exercise

Find the area of the surface generated by  $x = \sqrt{12y - y^2}$ ;  $2 \leq y \leq 10$  about y-axis

### Solution

$$x' = \frac{6 - y}{\sqrt{12y - y^2}}$$

$$\begin{aligned} S &= 2\pi \int_2^{10} \sqrt{12y - y^2} \sqrt{1 + \frac{(6 - y)^2}{12y - y^2}} dy \\ &= 2\pi \int_2^{10} \sqrt{12y - y^2 + 36 - 12y + y^2} dy \\ &= 12\pi \int_2^{10} dy \\ &= 12\pi y \Big|_2^{10} \\ &= \underline{96\pi \text{ unit}^2} \end{aligned}$$

$$S = 2\pi \int_c^d x \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy$$

### Exercise

Find the area of the surface generated by  $x = 4y^{3/2} - \frac{1}{12}y^{1/2}$ ;  $1 \leq y \leq 4$  about y-axis

### Solution

$$x' = 6y^{1/2} - \frac{1}{24\sqrt{y}} = \frac{144y - 1}{24\sqrt{y}}$$

$$\begin{aligned} S &= 2\pi \int_1^4 \left(4y^{3/2} - \frac{1}{12}y^{1/2}\right) \sqrt{1 + \frac{(144y - 1)^2}{576y}} dy \\ &= 2\pi \int_1^4 \left(4y^{3/2} - \frac{1}{12}y^{1/2}\right) \sqrt{\frac{576y + (144y)^2 - 288y + 1}{576y}} dy \\ &= \frac{\pi}{12} \int_1^4 \left(4y^{3/2} - \frac{1}{12}y^{1/2}\right) \frac{1}{\sqrt{y}} \sqrt{(144y + 1)^2} dy \\ &= \frac{\pi}{144} \int_1^4 (48y - 1)(144y + 1) dy \\ &= \frac{\pi}{144} \int_1^4 (6,912y^2 - 96y - 1) dy \\ &= \frac{\pi}{144} \left(2304y^3 - 48y^2 - y\right) \Big|_1^4 \end{aligned}$$

$$S = 2\pi \int_c^d x \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy$$

$$\begin{aligned}
 &= \frac{\pi}{144} (147,456 - 768 - 4 - 2304 + 48 + 1) \\
 &= \frac{144,429\pi}{144} \\
 &= \frac{48,143 \pi}{48} \text{ unit}^2
 \end{aligned}$$

### Exercise

Set up and evaluate the definite integral for the area of the surface generated by revolving the curve about the  $y$ -axis

$$y = 1 - \frac{1}{4}x^2, \quad 0 \leq x \leq 2$$

### Solution

$$y' = -\frac{1}{2}x$$

$$\begin{aligned}
 \sqrt{1 + (y')^2} &= \sqrt{1 + \frac{x^2}{4}} \\
 &= \frac{1}{2}\sqrt{4 + x^2}
 \end{aligned}$$



$$S = 2\pi \int_0^2 x \frac{\sqrt{4 + x^2}}{2} dx$$

$$= \frac{\pi}{2} \int_0^2 (4 + x^2)^{1/2} d(4 + x^2)$$

$$= \frac{\pi}{3} (4 + x^2)^{3/2} \Big|_0^2$$

$$= \frac{\pi}{3} (8^{3/2} - 4^{3/2})$$

$$= \frac{\pi}{3} (16\sqrt{2} - 8) \approx 15.318 \text{ unit}^2$$



### Exercise

Set up and evaluate the definite integral for the area of the surface generated by revolving the curve about the  $y$ -axis

$$y = \frac{1}{2}x + 3, \quad 1 \leq x \leq 5$$


### Solution

$$y' = \frac{1}{2}$$


$$\sqrt{1 + (y')^2} = \sqrt{1 + \frac{1}{4}} = \frac{\sqrt{5}}{2}$$

$$S = \pi \sqrt{5} \int_1^5 x dx$$

$$S = 2\pi \int_a^b x \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$



$$\begin{aligned}
&= \pi\sqrt{5}\left(\frac{1}{2}x^2\right)\Big|_1^5 \\
&= \frac{\sqrt{5}}{2}\pi(25-1) \\
&= \underline{12\pi\sqrt{5}} \quad \text{unit}^2
\end{aligned}$$



### Exercise

A right circular cone is generated by revolving the region bounded by  $y = \frac{3}{4}x$ ,  $y = 3$ , and  $x = 0$  about the  $y$ -axis. Find the lateral surface area of the cone.

#### Solution

$$y' = \frac{3}{4}$$

$$\sqrt{1+(y')^2} = \sqrt{1+\frac{9}{16}} = \frac{5}{4}$$

$$y = 3 = \frac{3}{4}x \Rightarrow \underline{x = 4}$$

$$S = \frac{5\pi}{2} \int_0^4 x \, dx$$

$$= \frac{5\pi}{4} x^2 \Big|_0^4$$

$$= \underline{20\pi} \quad \text{unit}^2$$

$$S = 2\pi \int_a^b x \sqrt{1+\left(\frac{dy}{dx}\right)^2} \, dx$$

### Exercise

A right circular cone is generated by revolving the region bounded by  $y = \frac{h}{r}x$ ,  $y = h$ , and  $x = 0$  about the  $y$ -axis. Verify that the lateral surface area of the cone is  $S = \pi r \sqrt{r^2 + h^2}$

#### Solution

$$y' = \frac{h}{r}$$

$$\sqrt{1+(y')^2} = \sqrt{1+\frac{h^2}{r^2}}$$

$$= \frac{\sqrt{r^2 + h^2}}{r}$$

$$y = h = \frac{h}{r}x \Rightarrow \underline{x = r}$$

$$\begin{aligned}
 S &= 2\pi \int_0^r x \frac{\sqrt{r^2 + h^2}}{r} dx \\
 &= \frac{\pi \sqrt{r^2 + h^2}}{r} \left( x^2 \right) \Big|_0^r \\
 &= \pi r \sqrt{r^2 + h^2}
 \end{aligned}$$

$$S = 2\pi \int_a^b x \sqrt{1 + \left( \frac{dy}{dx} \right)^2} dx$$

### Exercise

Find the area of the zone of a sphere formed by revolving the graph of  $y = \sqrt{9 - x^2}$ ,  $0 \leq x \leq 2$ , about the  $y$ -axis

#### Solution

$$y' = \frac{-x}{\sqrt{9 - x^2}}$$

$$\sqrt{1 + (y')^2} = \sqrt{1 + \frac{x^2}{9 - x^2}} = \frac{3}{\sqrt{9 - x^2}}$$

$$\begin{aligned}
 S &= 2\pi \int_0^2 x \frac{3}{\sqrt{9 - x^2}} dx \\
 &= -3\pi \int_0^2 (9 - x^2)^{-1/2} d(9 - x^2) \\
 &= -6\pi (9 - x^2)^{1/2} \Big|_0^2 \\
 &= -6\pi (\sqrt{5} - 3) \\
 &= 6\pi (3 - \sqrt{5})
 \end{aligned}$$

$$S = 2\pi \int_a^b x \sqrt{1 + \left( \frac{dy}{dx} \right)^2} dx$$

### Exercise

Find the area of the zone of a sphere formed by revolving the graph of  $y = \sqrt{r^2 - x^2}$ ,  $0 \leq x \leq a$ , about the  $y$ -axis. Assume that  $a < r$ .

#### Solution

$$y' = \frac{-x}{\sqrt{r^2 - x^2}}$$

$$\sqrt{1 + (y')^2} = \sqrt{1 + \frac{x^2}{r^2 - x^2}} = \frac{r}{\sqrt{r^2 - x^2}}$$

$$\begin{aligned}
S &= 2\pi \int_0^a x \frac{r}{\sqrt{r^2 - x^2}} dx & S &= 2\pi \int_a^b x \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx \\
&= -\pi r \int_0^a (r^2 - x^2)^{-1/2} d(r^2 - x^2) \\
&= -2\pi r \sqrt{r^2 - x^2} \Big|_0^a \\
&= -2\pi r \left( \sqrt{r^2 - a^2} - r \right) \\
&= \underline{2\pi r \left( r - \sqrt{r^2 - a^2} \right)}
\end{aligned}$$

### Exercise

Find the area of the surface generated by the curve  $y = 1 + \sqrt{1 - x^2}$  between the points  $(1, 1)$  and  $\left(\frac{\sqrt{3}}{1}, \frac{3}{2}\right)$  about y-axis

### Solution

$$\begin{aligned}
\left(\sqrt{1 - x^2}\right)^2 &= (y - 1)^2 \Rightarrow 1 - x^2 = y^2 - 2y + 1 \\
x &= \sqrt{2y - y^2} \rightarrow x' = \frac{1 - y}{\sqrt{2y - y^2}} \\
S &= 2\pi \int_1^{3/2} \sqrt{2y - y^2} \sqrt{1 + \frac{(1 - y)^2}{2y - y^2}} dy & S &= 2\pi \int_c^d x \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy \\
&= 2\pi \int_1^{3/2} \sqrt{2y - y^2 + 1 - 2y + y^2} dy \\
&= 2\pi \int_1^{3/2} dy \\
&= 2\pi y \Big|_1^{3/2} \\
&= \underline{\pi \text{ unit}^2}
\end{aligned}$$



### Exercise

Find the area of the surface generated by  $x = 2\sqrt{4-y}$   $0 \leq y \leq \frac{15}{4}$ ,  $y$ -axis


### Solution

$$\frac{dy}{dx} = 2 \cdot \frac{1}{2} (4-y)^{-1/2} (-1) = \frac{-1}{\sqrt{4-y}}$$

$$\begin{aligned}\sqrt{1 + \left(\frac{dy}{dx}\right)^2} &= \sqrt{1 + \frac{1}{4-y}} \\ &= \sqrt{\frac{4-y+1}{4-y}} \\ &= \sqrt{\frac{5-y}{4-y}}\end{aligned}$$

$$\begin{aligned}S &= 2\pi \int_0^{15/4} 2\sqrt{4-y} \frac{\sqrt{5-y}}{\sqrt{4-y}} dy \\ &= 4\pi \int_0^{15/4} \sqrt{5-y} dy \\ &= 4\pi \int_0^{15/4} (5-y)^{1/2} (-d(5-y)) \\ &= -4\pi \frac{2}{3} (5-y)^{3/2} \Big|_0^{15/4} \\ &= -\frac{8\pi}{3} \left[ \left(5 - \frac{15}{4}\right)^{3/2} - (5-0)^{3/2} \right] \\ &= -\frac{8\pi}{3} \left[ \left(\frac{5}{4}\right)^{3/2} - 5^{3/2} \right] \\ &= -\frac{8\pi}{3} \left[ \frac{5\sqrt{5}}{8} - 5\sqrt{5} \right] \\ &= -\frac{8\pi}{3} 5\sqrt{5} \left( \frac{1}{8} - 1 \right) \\ &= -\frac{8\pi}{3} 5\sqrt{5} \left( -\frac{7}{8} \right) \\ &= \underline{\underline{\frac{35\pi\sqrt{5}}{3} \text{ unit}^2}}\end{aligned}$$

$$d(5-y) = -dy$$



### Exercise


Find the area of the surface generated by  $x = \sqrt{2y-1}$   $\frac{5}{8} \leq y \leq 1$ ,  $y$ -axis

### Solution

$$\frac{dy}{dx} = \frac{1}{2}(2y-1)^{-1/2}(2) = \frac{1}{\sqrt{2y-1}}$$

$$\begin{aligned}\sqrt{1 + \left(\frac{dy}{dx}\right)^2} &= \sqrt{1 + \frac{1}{2y-1}} \\ &= \sqrt{\frac{2y}{2y-1}}\end{aligned}$$

$$\begin{aligned}S &= 2\pi \int_{5/8}^1 \sqrt{2y-1} \frac{\sqrt{2y}}{\sqrt{2y-1}} dy \\ &= 2\pi \int_{5/8}^1 \sqrt{2y} dy & u = 2y \rightarrow du = 2dy \\ &= 2\pi \int_{5/8}^1 u^{1/2} \left(\frac{1}{2} du\right) \\ &= \pi \int_{5/8}^1 u^{1/2} du \\ &= \frac{2\pi}{3} (2y)^{3/2} \Big|_{5/8}^1 \\ &= \frac{2\pi}{3} \left( (2)^{3/2} - \left(\frac{5}{4}\right)^{3/2} \right) \\ &= \frac{2\pi}{3} \left( 2\sqrt{2} - \frac{5\sqrt{5}}{8} \right) \\ &= \frac{2\pi}{3} \left( \frac{16\sqrt{2} - 5\sqrt{5}}{8} \right) \\ &= \frac{\pi}{12} (16\sqrt{2} - 5\sqrt{5}) \text{ unit}^2\end{aligned}$$



### Exercise

$y = \frac{1}{3}(x^2 + 2)^{3/2}$ ,  $0 \leq x \leq \sqrt{2}$ ;  $y$ -axis (Hint: Express  $ds = \sqrt{dx^2 + dy^2}$  in terms of  $dx$ , and evaluate the integral  $S = \int 2\pi x ds$  with appropriate limits.)

### Solution

$$dy = \frac{1}{3} \frac{3}{2} (x^2 + 2)^{1/2} (2x) dx = x \sqrt{x^2 + 2} dx$$

$$\begin{aligned} ds &= \sqrt{dx^2 + \left( x \sqrt{x^2 + 2} dx \right)^2} \\ &= \sqrt{dx^2 + x^2 (x^2 + 2) dx^2} \\ &= \sqrt{1 + x^4 + 2x^2} dx \\ &= \sqrt{(1 + x^2)^2} dx \\ &= (1 + x^2) dx \end{aligned}$$

$$S = \int 2\pi x ds$$

$$= 2\pi \int_0^{\sqrt{2}} x(1 + x^2) dx$$

$$d(1 + x^2) = 2x dx$$

$$= \pi \int_0^{\sqrt{2}} (1 + x^2) d(1 + x^2)$$

$$= \pi \frac{1}{2} u^2 \Big|_1^3$$

$$= \frac{\pi}{2} (3^2 - 1^2)$$

$$= \frac{\pi}{2} (8)$$

$$= \underline{4\pi \text{ unit}^2}$$


### Exercise

Find the area of the surface generated by revolving the curve  $x = \frac{1}{2}(e^y + e^{-y})$ ,  $0 \leq y \leq \ln 2$ , about y-axis

### Solution

$$\begin{aligned} S &= 2\pi \int_0^{\ln 2} \frac{1}{2}(e^y + e^{-y}) \sqrt{1 + \left( \frac{e^y - e^{-y}}{2} \right)^2} dy \\ &= \pi \int_0^{\ln 2} (e^y + e^{-y}) \sqrt{1 + \frac{e^{2y} + e^{-2y} - 2}{4}} dy \end{aligned}$$

$$\begin{aligned}
&= \pi \int_0^{\ln 2} (e^y + e^{-y}) \sqrt{\frac{4 + e^{2y} + e^{-2y} - 2}{4}} dy \\
&= \frac{\pi}{2} \int_0^{\ln 2} (e^y + e^{-y}) \sqrt{e^{2y} + e^{-2y} + 2} dy \\
&= \frac{\pi}{2} \int_0^{\ln 2} (e^y + e^{-y}) \sqrt{(e^y + e^{-y})^2} dy \\
&= \frac{\pi}{2} \int_0^{\ln 2} (e^y + e^{-y})^2 dy \\
&= \frac{\pi}{2} \int_0^{\ln 2} (e^{2y} + e^{-2y} + 2) dy \\
&= \frac{\pi}{2} \left[ \frac{1}{2} e^{2y} - \frac{1}{2} e^{-2y} + 2y \right]_0^{\ln 2} \\
&= \frac{\pi}{2} \left[ \left( \frac{1}{2} e^{2 \ln 2} - \frac{1}{2} e^{-2 \ln 2} + 2 \ln 2 \right) - \left( \frac{1}{2} e^0 - \frac{1}{2} e^0 + 0 \right) \right] \\
&= \frac{\pi}{2} \left( \frac{1}{2} \cdot 4 - \frac{1}{2} \cdot \frac{1}{4} + 2 \ln 2 \right) \\
&= \frac{\pi}{2} \left( \frac{15}{8} + 2 \ln 2 \right) \text{ unit}^2
\end{aligned}$$




## Exercise

Did you know that if you can cut a spherical loaf of bread into slices of equal width, each slice will have the same amount of crust? To see why, suppose the semicircle  $y = \sqrt{r^2 - x^2}$  shown here is revolved about the  $x$ -axis to generate a sphere. Let **AB** be an arc of the semicircle that lies above an interval of length  $h$  on the  $x$ -axis. Show that the area swept out by **AB** does not depend on the location of the interval. (It does depend on the length of the interval.)

### Solution

$$\begin{aligned}
\frac{dy}{dx} &= \frac{-2x}{2\sqrt{r^2 - x^2}} = \frac{-x}{\sqrt{r^2 - x^2}} \\
\sqrt{1 + \left(\frac{dy}{dx}\right)^2} &= \sqrt{1 + \frac{x^2}{r^2 - x^2}} \\
&= \sqrt{\frac{r^2}{r^2 - x^2}}
\end{aligned}$$



$$\begin{aligned}
 S &= 2\pi \int_a^{a+h} \sqrt{r^2 - x^2} \frac{r}{\sqrt{r^2 - x^2}} dx \\
 &= 2\pi r \int_a^{a+h} dx \\
 &= 2\pi r x \Big|_a^{a+h} \\
 &= 2\pi r(a+h-a) \\
 &= \underline{2\pi rh \text{ unit}^2}
 \end{aligned}$$

### Example

The curved surface of a funnel is generated by revolving the graph of  $y = f(x) = x^3 + \frac{1}{12x}$  on the interval  $[1, 2]$  about the  $x$ -axis. Approximately what volume of paint is needed to cover the outside of the funnel with a layer of paint 0.05 cm thick? Assume that  $x$  and  $y$  measured in centimeters.


### Solution

$$f'(x) = 3x^2 - \frac{1}{12x^2}$$

$$1 + f'(x)^2 = 1 + \left(3x^2 - \frac{1}{12x^2}\right)^2$$

$$\begin{aligned}
 &= 1 + 9x^4 - \frac{1}{2} + \frac{1}{144x^4} \\
 &= 9x^4 + \frac{1}{2} + \frac{1}{144x^4} \\
 &= \left(3x^2 + \frac{1}{12x^2}\right)^2
 \end{aligned}$$

$$\begin{aligned}
 S &= 2\pi \int_1^2 \left(x^3 + \frac{1}{12x}\right) \sqrt{\left(3x^2 + \frac{1}{12x^2}\right)^2} dx \\
 &= 2\pi \int_1^2 \left(x^3 + \frac{1}{12x}\right) \left(3x^2 + \frac{1}{12x^2}\right) dx \\
 &= 2\pi \int_1^2 \left(3x^5 + \frac{x}{3} + \frac{1}{144}x^{-3}\right) dx \\
 &= 2\pi \left(\frac{1}{2}x^6 + \frac{1}{6}x^2 - \frac{1}{288}x^{-2}\right) \Big|_1^2
 \end{aligned}$$



$$\begin{aligned}
&= 2\pi \left( 32 + \frac{2}{3} - \frac{1}{1152} - \frac{1}{2} - \frac{1}{6} + \frac{1}{288} \right) \\
&= 2\pi \left( \frac{36864 + 768 - 1 - 576 - 192 + 4}{1152} \right) \\
&= \underline{\underline{\frac{12,289}{192} \pi \text{ cm}^2}}
\end{aligned}$$

Because the paint layer is  $0.05 \text{ cm}$  thick, the approximate volume of paint needed is

$$= \left( \frac{12,289}{192} \pi \text{ cm}^2 \right) (0.05 \text{ cm}) \approx \underline{\underline{10.1 \text{ cm}^3}}$$

### Exercise

When the circle  $x^2 + (y - a)^2 = r^2$  on the interval  $[-r, r]$  is revolved about the  $x$ -axis, the result is the surface of a torus, where  $0 < r < a$ . Show that the surface area of the torus is  $S = 4\pi^2 ar$ .

### Solution

$$\begin{aligned}
x^2 + (y - a)^2 = r^2 &\Rightarrow (y - a)^2 = r^2 - x^2 \\
&y = a \pm \sqrt{r^2 - x^2}
\end{aligned}$$

$$f(x) = a + \sqrt{r^2 - x^2}$$

$$\begin{aligned}
1 + f'(x)^2 &= 1 + \left( \frac{-x}{\sqrt{r^2 - x^2}} \right)^2 \\
&= 1 + \frac{x^2}{r^2 - x^2} \\
&= \frac{r^2}{r^2 - x^2}
\end{aligned}$$

$$\begin{aligned}
S_1 &= 2\pi \int_{-r}^r \left( a + \sqrt{r^2 - x^2} \right) \frac{r}{\sqrt{r^2 - x^2}} dx \\
&= 4\pi \int_0^r \left( \frac{ar}{\sqrt{r^2 - x^2}} + r \right) dx \\
&= 4\pi \left[ ar \sin^{-1} \left( \frac{x}{r} \right) + rx \right]_0^r \\
&= 4\pi \left( ar \frac{\pi}{2} + r^2 \right) \\
&= \underline{\underline{2\pi^2 ar + 4\pi r^2}}
\end{aligned}$$

$$\begin{aligned}
S_2 &= 2\pi \int_{-r}^r \left( a - \sqrt{r^2 - x^2} \right) \frac{r}{\sqrt{r^2 - x^2}} dx \\
&= 4\pi \int_0^r \left( \frac{ar}{\sqrt{r^2 - x^2}} - r \right) dx \\
&= 4\pi \left[ ar \sin^{-1} \left( \frac{x}{r} \right) - rx \right]_0^r \\
&= 4\pi \left( ar \frac{\pi}{2} - r^2 \right) \\
&= \underline{2\pi^2 ar - 4\pi r^2} \\
S &= 2\pi^2 ar + 4\pi r^2 + 2\pi^2 ar - 4\pi r^2 = \underline{4\pi^2 ar \text{ unit}^2}
\end{aligned}$$

### Exercise

A 1.5-mm layer of paint is applied to one side. Find the approximate volume of paint needed of the spherical zone generated when the curve  $y = \sqrt{8x - x^2}$  on the interval  $[1, 7]$  is revolved about the  $x$ -axis. Assume  $x$  and  $y$  are in *meters*.

### Solution

$$\begin{aligned}
y' &= \frac{4-x}{\sqrt{8x-x^2}} \\
S &= 2\pi \int_1^7 \sqrt{8x-x^2} \sqrt{1 + \frac{(4-x)^2}{8x-x^2}} dx \\
&= 2\pi \int_1^7 \sqrt{8x-x^2} \frac{\sqrt{8x-x^2+16-8x+x^2}}{\sqrt{8x-x^2}} dx \\
&= 2\pi \int_1^7 \sqrt{16} dx \\
&= 8\pi x \Big|_1^7 \\
&= \underline{48\pi \text{ m}^2}
\end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left( \frac{dy}{dx} \right)^2} dx$$

The volume of paint required to cover the surface to a thickness 0.0015 *m* is

$$\begin{aligned}
V &= 48\pi(0.0015) \approx \underline{0.226195 \text{ m}^3} & 1 \text{ m}^3 &= 264.172052 \text{ gal} \\
&= 0.226195 \times 264.172052 \approx \underline{59.75 \text{ gal}}
\end{aligned}$$

### Exercise

A 1.5-*mm* layer of paint is applied to one side. Find the approximate volume of paint needed of the spherical zone generated when the upper portion of the circle  $x^2 + y^2 = 100$  on the interval  $[-8, 8]$  is revolved about the  $x$ -axis. Assume  $x$  and  $y$  are in *meters*.

### Solution

$$\begin{aligned}y &= \sqrt{100 - x^2} \Rightarrow y' = \frac{-x}{\sqrt{100 - x^2}} \\S &= 2\pi \int_{-8}^8 \sqrt{100 - x^2} \sqrt{1 + \frac{x^2}{100 - x^2}} dx \\&= 2\pi \int_{-8}^8 \sqrt{100 - x^2} \frac{\sqrt{100 - x^2 + x^2}}{\sqrt{100 - x^2}} dx \\&= 20\pi \int_{-8}^8 dx \\&= 20\pi x \Big|_{-8}^8 \\&= \underline{320\pi \text{ m}^2}\end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

The volume of paint required to cover the surface to a thickness 0.0015 *m* is


$$\begin{aligned}V &= 320\pi(0.0015) \approx \underline{1.507965 \text{ m}^3} & 1 \text{ m}^3 &= 264.172052 \text{ gal} \\&= 1.507965 \times 264.172052 \approx \underline{398.36 \text{ gal}}\end{aligned}$$

### Exercise

Suppose a sphere of radius  $r$  is sliced by two horizontal planes  $h$  units apart. Show that the surface area of the resulting zone on the sphere is  $2\pi h$ , independent of the location of the cutting planes.

### Solution

$$\begin{aligned}f(x) &= \sqrt{r^2 - x^2} \\1 + f'(x)^2 &= 1 + \left(\frac{-x}{\sqrt{r^2 - x^2}}\right)^2 \\&= 1 + \frac{x^2}{r^2 - x^2} \\&= \frac{r^2}{r^2 - x^2}\end{aligned}$$





$$\begin{aligned}
 S &= 2\pi \int_a^{a+h} \sqrt{r^2 - x^2} \frac{r}{\sqrt{r^2 - x^2}} dx \\
 &= 2\pi r x \Big|_a^{a+h} \\
 &= 2\pi r (a+h-a) \\
 &= \underline{2\pi r h \text{ unit}^2}
 \end{aligned}$$


$$S = 2\pi \int_a^b f(x) \sqrt{1 + f'(x)^2} dx$$

### Exercise

An ornamental light bulb is designed by revolving the graph of  $y = \frac{1}{3}x^{1/2} - x^{3/2}$ ,  $0 \leq x \leq \frac{1}{3}$  about the  $x$ -axis, where  $x$  and  $y$  are measured in *feet*. Find the surface area of the bulb and use the result to approximate the amount of glass needed to make the bulb.  
(Assume that the glass is 0.015 *inch* thick)

### Solution

$$\begin{aligned}
 y' &= \frac{1}{6}x^{-1/2} - \frac{3}{2}x^{1/2} \\
 \sqrt{1 + (y')^2} &= \sqrt{1 + \frac{1}{36}x^{-1} - \frac{1}{2} + \frac{9}{4}x} \\
 &= \frac{1}{6}\sqrt{x^{-1} + 18 + 81x} \\
 &= \frac{1}{6}\sqrt{\left(x^{-1/2} + 9x^{1/2}\right)^2} \\
 &= \frac{1}{6}\left(x^{-1/2} + 9x^{1/2}\right) \\
 S &= 2\pi \frac{1}{6} \int_0^{1/3} \left(\frac{1}{3}x^{1/2} - x^{3/2}\right)\left(x^{-1/2} + 9x^{1/2}\right) dx \\
 &= \frac{\pi}{3} \int_0^{1/3} \left(\frac{1}{3} + 2x - 9x^2\right) dx \\
 &= \frac{\pi}{3} \left(\frac{1}{3}x + x^2 - 3x^3\right) \Big|_0^{1/3} \\
 &= \frac{\pi}{3} \left(\frac{1}{9} + \frac{1}{9} - \frac{1}{9}\right) \\
 &= \underline{\frac{\pi}{27}} \approx 0.1164 \text{ ft}^2 \approx 16.8 \text{ in}^2
 \end{aligned}$$



$$\text{Amount of glass needed: } V = \frac{\pi}{2} \left(\frac{0.015}{12}\right) \approx 0.00015 \text{ ft}^3 \approx \underline{0.25 \text{ in}^3}$$