

Solution **Section 2.10 – Applications**

Exercise

Solve using the Laplace transform: $y' + y = te^t$, $y(0) = -2$

Solution

$$\mathcal{L}(y' + y) = \mathcal{L}(te^t)$$

$$\mathcal{L}(y') + \mathcal{L}(y) = \mathcal{L}(te^t)$$

$$sY(s) - y(0) + Y(s) = \frac{1}{(s-1)^2} \quad y(0) = -2$$

$$(s+1)Y(s) + 2 = \frac{1}{(s-1)^2}$$

$$(s+1)Y(s) = \frac{1}{(s-1)^2} - 2$$

$$Y(s) = \frac{1}{(s+1)(s-1)^2} - \frac{2}{s+1}$$

$$\frac{1}{(s+1)(s-1)^2} = \frac{A}{s+1} + \frac{B}{s-1} + \frac{C}{(s-1)^2}$$

$$1 = (A+B)s^2 + (C-2A)s + A - B + C$$

$$\begin{cases} A+B=0 \\ C-2A=0 \\ A-B+C=1 \end{cases} \Rightarrow A = \frac{1}{4} \quad B = -\frac{1}{4} \quad C = \frac{1}{2}$$

$$Y(s) = \frac{1}{4} \frac{1}{s+1} - \frac{1}{4} \frac{1}{s-1} + \frac{1}{2} \frac{1}{(s-1)^2} - \frac{2}{s+1}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{-\frac{7}{4} \frac{1}{s+1} - \frac{1}{4} \frac{1}{s-1} + \frac{1}{2} \frac{1}{(s-1)^2}\right\}$$

$$y(t) = \underline{-\frac{7}{4}e^{-t} - \frac{1}{4}e^t + \frac{1}{2}te^t}$$

Exercise

Solve using the Laplace transform: $y' - y = 2\cos 5t$, $y(0) = 0$

Solution

$$\mathcal{L}\{y' - y\} = \mathcal{L}\{2\cos 5t\}$$

$$sY(s) - y(0) - Y(s) = \frac{2s}{s^2 + 25} \quad y(0) = 0$$

$$(s-1)Y(s) = \frac{2s}{s^2 + 25}$$

$$Y(s) = \frac{2s}{(s-1)(s^2 + 25)} = \frac{A}{s-1} + \frac{Bs+C}{s^2 + 25}$$

$$2s = As^2 + 25A + Bs^2 - Bs + Cs - C$$

$$\begin{cases} A+B=0 \\ -B+C=2 \\ 25A-C=0 \end{cases} \Rightarrow A = \frac{1}{13} \quad B = -\frac{1}{13} \quad C = \frac{25}{13}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{1}{13}\frac{1}{s-1} - \frac{1}{13}\frac{s}{s^2 + 25} + \frac{25}{13}\frac{1}{5}\frac{5}{s^2 + 25}\right\}$$

$$\underline{y(t) = \frac{1}{13}e^t - \frac{1}{13}\cos 4t + \frac{5}{13}\sin 5t}$$

Exercise

Solve using the Laplace transform: $y' - y = 1 + te^t$, $y(0) = 0$

Solution

$$\mathcal{L}\{y' - y\} = \mathcal{L}\{1 + te^t\}$$

$$sY(s) - y(0) - Y(s) = \frac{1}{s} + \frac{1}{(s-1)^2} \quad y(0) = 0$$

$$(s-1)Y(s) = \frac{s^2 - s + 1}{s(s-1)^2}$$

$$Y(s) = \frac{s^2 - s + 1}{s(s-1)^3} = \frac{A}{s} + \frac{B}{s-1} + \frac{C}{(s-1)^2} + \frac{D}{(s-1)^3}$$

$$s^2 - s + 1 = As^3 - 3As^2 + 3As - A + Bs^3 - 2Bs^2 + Bs + Cs^2 - Cs + Ds$$

$$s^3 \quad A + B = 0 \quad \underline{B = 1}$$

$$s^2 \quad -3A - 2B + C = 1 \quad \underline{C = 0}$$

$$s \quad 3A + B - C + D = -1 \quad \underline{D = 1}$$

$$s^0 \quad A = -1$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{-\frac{1}{s} + \frac{1}{s-1} + \frac{1}{(s-1)^3}\right\}$$

$$\underline{y(t) = -1 + e^t + \frac{1}{2}t^2 e^t}$$

Exercise

Solve using the Laplace transform: $y' + 3y = e^{2t}$, $y(0) = -1$

Solution

$$\mathcal{L}(y' + 3y) = \mathcal{L}(e^{2t})$$

$$\mathcal{L}(y') + 3\mathcal{L}(y) = \mathcal{L}(e^{2t})$$

$$sY(s) - y(0) + 3Y(s) = \frac{1}{s-2}$$

$$(s+3)Y(s) + 1 = \frac{1}{s-2}$$

$$(s+3)Y(s) = \frac{1}{s-2} - 1$$

$$Y(s) = \frac{1}{(s-2)(s+3)} - \frac{1}{s+3}$$

$$\frac{1}{(s-2)(s+3)} = \frac{A}{s-2} + \frac{B}{s+3}$$

$$1 = (A+B)s + 3A - 2B$$

$$\begin{cases} A+B=0 \\ 3A-2B=1 \end{cases} \Rightarrow A = \frac{1}{5} \quad B = -\frac{1}{5}$$

$$Y(s) = \frac{1}{5} \frac{1}{s-2} - \frac{1}{5} \frac{1}{s+3} - \frac{1}{s+3}$$

$$y(t) = \frac{1}{5} \mathcal{L}^{-1} \left\{ \frac{1}{s-2} \right\} - \frac{6}{5} \mathcal{L}^{-1} \left\{ \frac{1}{s+3} \right\}$$

$$= \frac{1}{5} e^{2t} - \frac{6}{5} e^{-3t}$$

Exercise

Solve using the Laplace transform: $y' + 4y = \cos t$, $y(0) = 0$

Solution

$$\mathcal{L}(y' + 4y) = \mathcal{L}(\cos t)$$

$$sY(s) - y(0) + 4Y(s) = \frac{s}{s^2 + 1} \quad y(0) = 0$$

$$(s+4)Y(s) = \frac{s}{s^2 + 1}$$

$$Y(s) = \frac{s}{(s+4)(s^2 + 1)}$$

$$\frac{s}{(s+4)(s^2+1)} = \frac{A}{s+4} + \frac{Bs+C}{s^2+1}$$

$$s = As^2 + A + Bs^2 + 4Bs + Cs + 4C$$

$$s = (A+B)s^2 + (4B+C)s + A+4C$$

$$\begin{cases} A+B=0 \\ 4B+C=1 \Rightarrow A=-\frac{4}{17} \quad B=\frac{4}{17} \quad C=\frac{1}{17} \\ A+4C=0 \end{cases}$$

$$Y(s) = -\frac{4}{17} \frac{1}{s+4} + \frac{4}{17} \frac{s}{s^2+1} + \frac{1}{17} \frac{1}{s^2+1}$$

$$\begin{aligned} y(t) &= -\frac{4}{17} \mathcal{L}^{-1} \left\{ \frac{1}{s+4} \right\} + \frac{4}{17} \mathcal{L}^{-1} \left\{ \frac{s}{s^2+1} \right\} + \frac{1}{17} \mathcal{L}^{-1} \left\{ \frac{1}{s^2+1} \right\} \\ &= \underline{-\frac{4}{17} e^{-4t} + \frac{4}{17} e^{-9t} \cos t + \frac{1}{17} \sin t} \end{aligned}$$

Exercise

Solve using the Laplace transform: $y' + 4y = e^{-4t}$, $y(0) = 2$

Solution

$$\mathcal{L}\{y' + 4y\} = \mathcal{L}\{e^{-4t}\}$$

$$sY(s) - y(0) + 4Y(s) = \frac{1}{s+4} \quad y(0) = 2$$

$$(s+4)Y(s) = \frac{1}{s+4} + 2$$

$$Y(s) = \frac{2s+9}{(s+4)^2} = \frac{A}{s+4} + \frac{B}{(s+4)^2}$$

$$2s+9 = As + 4A + B$$

$$\begin{matrix} s \\ s^0 \end{matrix} \quad \underline{A=2}$$

$$\begin{matrix} s^0 \\ s^0 \end{matrix} \quad 4A+B=9 \quad \underline{B=1}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1} \left\{ \frac{2}{s+4} + \frac{1}{(s+4)^2} \right\}$$

$$\underline{y(t) = 2e^{-4t} + te^{-4t}}$$

Exercise

Solve using the Laplace transform: $y' - 4y = t^2 e^{-2t}$, $y(0) = 1$

Solution

$$\mathcal{L}(y' - 4y) = \mathcal{L}(t^2 e^{-2t})$$

$$sY(s) - y(0) - 4Y(s) = \frac{2!}{(s+2)^3} \quad y(0) = 1$$

$$(s-4)Y(s) - 1 = \frac{2}{(s+2)^3}$$

$$(s-4)Y(s) = 1 + \frac{2}{(s+2)^3}$$

$$Y(s) = \frac{1}{s-4} + \frac{2}{(s-4)(s+2)^3}$$

$$\frac{2}{(s-4)(s+2)^3} = \frac{A}{s-4} + \frac{B}{s+2} + \frac{C}{(s+2)^2} + \frac{D}{(s+2)^3}$$

$$2 = A(s^3 + 6s^2 + 12s + 8A) + B(s^2 + 4s + 4)(s-4) + C(s-4)(s+2) + D(s-4)$$

$$2 = As^3 + 6As^2 + 12As + 8A + Bs^3 + 4Bs^2 + 4Bs - 4Bs^2 - 16Bs - 16B + Cs^2 - 2Cs - 8C + Ds - 4D$$

$$2 = (A+B)s^3 + (6A+C)s^2 + (12A-12B-2C+D)s + 8A-16B-8C-4D$$

$$\begin{cases} A+B=0 \\ 6A+C=0 \\ 12A-12B-2C+D=0 \\ 8A-16B-8C-4D=2 \end{cases} \Rightarrow \begin{matrix} A = \frac{1}{108} & B = -\frac{1}{108} \\ C = -\frac{1}{18} & D = -\frac{1}{3} \end{matrix}$$

$$Y(s) = \frac{1}{s-4} + \frac{1}{108} \frac{1}{s-4} - \frac{1}{108} \frac{1}{s+2} - \frac{1}{18} \frac{1}{(s+2)^2} - \frac{1}{3} \frac{1}{(s+2)^3}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{109}{108} \frac{1}{s-4} - \frac{1}{108} \frac{1}{s+2} - \frac{1}{18} \frac{1}{(s+2)^2} - \frac{1}{3} \frac{1}{(s+2)^3}\right\}$$

$$y(t) = \frac{109}{108} e^{4t} - \frac{1}{108} e^{-2t} - \frac{1}{18} t e^{-2t} - \frac{1}{6} t^2 e^{-2t}$$

Exercise

Solve using the Laplace transform: $y' + 9y = e^{-t}$, $y(0) = 0$

Solution

$$\mathcal{L}(y' + 9y) = \mathcal{L}(e^{-t})$$

$$Y(s) = \mathcal{L}(y)(s)$$

$$\mathcal{L}(y') + 9\mathcal{L}(y) = \mathcal{L}(e^{-t})$$

$$sY(s) - y(0) + 9Y(s) = \frac{1}{s+1}$$

$$(s+9)Y(s) = \frac{1}{s+1}$$

$$Y(s) = \frac{1}{(s+1)(s+9)}$$

$$\frac{1}{(s+1)(s+9)} = \frac{A}{s+1} + \frac{B}{s+9}$$

$$1 = (A+B)s + 9A + B$$

$$\begin{cases} A+B=0 \\ 9A+B=1 \end{cases} \Rightarrow A = \frac{1}{8} \quad B = -\frac{1}{8}$$

$$Y(s) = \frac{1}{8} \frac{1}{s+1} - \frac{1}{8} \frac{1}{s+9}$$

$$y(t) = \frac{1}{8} \mathcal{L}^{-1} \left\{ \frac{1}{s+1} \right\} - \frac{1}{8} \mathcal{L}^{-1} \left\{ \frac{1}{s+9} \right\}$$

$$= \frac{1}{8} e^{-t} - \frac{1}{8} e^{-9t}$$

Exercise

Solve using the Laplace transform: $y' + 16y = \sin 3t$, $y(0) = 1$

Solution

$$\mathcal{L}(y' + 16y) = \mathcal{L}(\sin 3t)$$

$$sY(s) - y(0) + 16Y(s) = \frac{3}{s^2 + 9} \quad y(0) = 1$$

$$(s+16)Y(s) - 1 = \frac{3}{s^2 + 9}$$

$$(s+16)Y(s) = \frac{3}{s^2 + 9} + 1$$

$$Y(s) = \frac{1}{s+16} + \frac{3}{(s+16)(s^2 + 9)}$$

$$\frac{3}{(s+16)(s^2 + 9)} = \frac{A}{s+16} + \frac{Bs+C}{s^2 + 9}$$

$$s = As^2 + 9A + Bs^2 + 16Bs + Cs + 16C$$

$$s = (A+B)s^2 + (16B+C)s + 9A+16C$$

$$\begin{cases} A+B=0 \\ 16B+C=1 \\ 9A+16C=0 \end{cases} \Rightarrow A = \frac{3}{265} \quad B = -\frac{3}{265} \quad C = \frac{48}{265}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{1}{s+16} + \frac{3}{265}\frac{1}{s+16} - \frac{3}{265}\frac{s}{s^2+9} + \frac{48}{265}\frac{1}{s^2+9}\right\}$$

$$y(t) = \frac{268}{265}e^{-16t} - \frac{3}{265}\cos 3t + \frac{16}{265}\sin 3t$$

Exercise

Solve using the Laplace transform: $y'' - y = e^{2t}$; $y(0) = 0$, $y'(0) = 1$

Solution

$$\mathcal{L}\{y'' - y\} = \mathcal{L}\{e^{2t}\}$$

$$s^2Y(s) - sy(0) - y'(0) - Y(s) = \frac{1}{s-2} \quad y(0) = 0, \quad y'(0) = 1$$

$$(s^2 - 1)Y(s) - 1 = \frac{1}{s-2}$$

$$(s^2 - 1)Y(s) = \frac{1}{s-2} + 1$$

$$(s-1)(s+1)Y(s) = \frac{s-1}{s-2}$$

$$Y(s) = \frac{1}{(s+1)(s-2)} = \frac{A}{(s+1)} + \frac{B}{(s-2)}$$

$$(A+B)s + B - 2A = 1 \quad \begin{cases} A+B=0 \\ -2A+B=1 \end{cases} \Rightarrow A = -\frac{1}{3}; B = \frac{1}{3}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \frac{1}{3}\mathcal{L}^{-1}\left\{-\frac{1}{(s+1)} + \frac{1}{(s-2)}\right\}$$

$$y(t) = \frac{1}{3}(e^{2t} - e^{-t})$$

Exercise

Solve using the Laplace transform: $y'' - y = 2t$; $y(0) = 0$, $y'(0) = -1$

Solution

$$\mathcal{L}(y'' - y) = \mathcal{L}(2t)$$

$$s^2Y(s) - sy(0) - y'(0) - Y(s) = 2\frac{1}{s^2} \quad y(0) = 0 \quad y'(0) = -1$$

$$(s^2 - 1)Y(s) + 1 = \frac{2}{s^2}$$

$$Y(s) = \frac{2}{s^2(s-1)(s+1)} - \frac{1}{(s-1)(s+1)}$$

$$\frac{2}{s^2(s-1)(s+1)} = \frac{A}{s^2} + \frac{B}{s-1} + \frac{C}{s+1}$$

$$2 = As^2 - A + Bs^3 + Bs^2 + Cs^3 - Cs^2$$

$$2 = (B+C)s^3 + (A+B-C)s^2 - A$$

$$\begin{cases} B+C=0 \\ A+B-C=0 \\ -A=2 \end{cases} \Rightarrow A=-2 \quad B=1 \quad C=-1$$

$$\frac{1}{(s-1)(s+1)} = \frac{D}{s-1} + \frac{E}{s+1}$$

$$\begin{cases} D+E=0 \\ D-E=1 \end{cases} \Rightarrow D=\frac{1}{2} \quad E=-\frac{1}{2}$$

$$Y(s) = -\frac{2}{s^2} + \frac{1}{s-1} - \frac{1}{s+1} - \frac{1}{2} \frac{1}{s-1} + \frac{1}{2} \frac{1}{s+1}$$

$$= -\frac{2}{s^2} + \frac{1}{2} \frac{1}{s-1} - \frac{1}{2} \frac{1}{s+1}$$

$$y(t) = 2\mathcal{L}^{-1}\left\{\frac{1}{s^2}\right\} + \frac{1}{2}\mathcal{L}^{-1}\left\{\frac{1}{s-1}\right\} - \frac{1}{2}\mathcal{L}^{-1}\left\{\frac{1}{s+1}\right\}$$

$$\underline{y(t) = -2t + \frac{1}{2}e^t - \frac{1}{2}e^{-t}}$$

Exercise

Solve using the Laplace transform: $y'' - y = t - 2$; $y(2) = 3$, $y'(2) = 0$

Solution

$$\text{Let: } w(t) = y(t+2) \leftrightarrow y(t) = w(t-2)$$

$$\mathcal{L}\{y'' - y\} = \mathcal{L}\{t+2\}$$

$$\mathcal{L}\{w'' - w\} = \mathcal{L}\{t\}$$

$$s^2W(s) - sw(0) - w'(0) - W(s) = \frac{1}{s}$$

$$y(2) = w(2-2) = w(0) = 3, \quad y'(2) = w'(0) = 0$$

$$(s^2 - 1)W(s) = \frac{1}{s} + 3s$$

$$W(s) = \frac{1+3s^2}{s(s^2-1)} = \frac{A}{s} + \frac{B}{s-1} + \frac{C}{s+1}$$

$$1+3s^2 = As^2 - A + Bs^2 + Bs + Cs^2 - Cs$$

$$s^2 \quad A + C = 3 \quad \underline{C = 2}$$

$$s^1 \quad B - C = 0 \quad \underline{B = 2}$$

$$s^0 \quad \underline{A = -1}$$

$$\mathcal{L}^{-1}\{W(s)\} = \mathcal{L}^{-1}\left\{-\frac{1}{s} + \frac{2}{s-1} + \frac{2}{s+1}\right\}$$

$$w(t) = -t + 2e^t + 2e^{-t}$$

$$\begin{aligned} y(t) &= w(t-2) = -(t-2) + 2e^{t-2} + 2e^{-(t-2)} \\ &= \underline{2-t + 2e^{t-2} + 2e^{-t+2}} \end{aligned}$$

Exercise

Solve using the Laplace transform: $y'' + y = t$; $y(\pi) = y'(\pi) = 0$

Solution

$$\text{Let: } w(t) = y(t + \pi) \leftrightarrow y(t) = w(t - \pi)$$

$$y'' + y = t \rightarrow w'' + w = t + \pi$$

$$\mathcal{L}\{w'' + w\} = \mathcal{L}\{t + \pi\}$$

$$s^2 W(s) - sw(0) - w'(0) + W(s) = \frac{1}{s^2} + \frac{\pi}{s}$$

$$y(\pi) = w(\pi - \pi) = w(0) = 0, \quad y'(\pi) = w'(0) = 0$$

$$(s^2 + 1)W(s) = \frac{1 + \pi s}{s^2}$$

$$W(s) = \frac{1 + \pi s}{s^2(s^2 + 1)} = \frac{A}{s} + \frac{B}{s^2} + \frac{Cs + D}{s^2 + 1}$$

$$1 + \pi s = As^3 + As^2 + Bs^2 + B + Cs^3 + Ds^2$$

$$s^3 \quad A + C = 0 \quad \underline{C = -\pi}$$

$$s^2 \quad B + D = 0 \quad \underline{D = -1}$$

$$s^1 \quad \underline{A = \pi}$$

$$s^0 \quad \underline{B = 1}$$

$$\mathcal{L}^{-1}\{W(s)\} = \mathcal{L}^{-1}\left\{\frac{\pi}{s} + \frac{1}{s^2} - \frac{\pi s}{s^2 + 1} - \frac{1}{s^2 + 1}\right\}$$

$$w(t) = \pi + t - \pi \cos t - \sin t$$

$$\begin{aligned} y(t) &= w(t - \pi) = \pi + (t - \pi) - \pi \cos(t - \pi) - \sin(t - \pi) \\ &= t - \pi(\cos t \cos \pi - \sin t \sin \pi) - (\cos t \sin \pi - \cos \pi \sin t) \\ &= \underline{t + \pi \cos t + \sin t} \end{aligned}$$

Exercise

Solve using the Laplace transform: $y'' - 2y' + 5y = -8e^{\pi-t}$; $y(\pi) = 2, y'(\pi) = 12$

Solution

$$\text{Let: } w(t) = y(t + \pi) \leftrightarrow y(t) = w(t - \pi)$$

$$y'' - 2y' + 5y = -8e^{\pi-t} \rightarrow w'' - 2w' + 5w = -8e^{-t}$$

$$\mathcal{L}\{w'' - 2w' + 5w\} = \mathcal{L}\{-8e^{-t}\}$$

$$s^2W(s) - sw(0) - w'(0) - 2sW(s) + 2w(0) + 5W(s) = -\frac{8}{s+1} \quad y(\pi) = w(0) = 2, \quad y'(\pi) = w'(0) = 12$$

$$(s^2 - 2s + 5)W(s) = -\frac{8}{s+1} + 2s + 12 - 4$$

$$(s^2 - 2s + 5)W(s) = \frac{2s^2 + 10s}{s+1}$$

$$W(s) = \frac{2s^2 + 10s}{(s+1)((s-1)^2 + 4)} = \frac{A}{s+1} + \frac{B(s-1) + C}{(s-1)^2 + 4}$$

$$2s^2 + 10s = As^2 - 2As + 5A + Bs^2 - B + Cs + C$$

$$s^2 \quad A + B = 2$$

$$s^1 \quad -2A + C = 10$$

$$s^0 \quad 5A - B + C = 0$$

$$\Delta = \begin{vmatrix} 1 & 1 & 0 \\ -2 & 0 & 1 \\ 5 & -1 & 1 \end{vmatrix} = 8 \quad \Delta_A = \begin{vmatrix} 2 & 1 & 0 \\ 10 & 0 & 1 \\ 0 & -1 & 1 \end{vmatrix} = -8$$

$$\underline{A = -1 \quad B = 3 \quad C = 8}$$

$$\mathcal{L}^{-1}\{W(s)\} = \mathcal{L}^{-1}\left\{-\frac{1}{s+1} + \frac{3(s-1)}{(s-1)^2 + 2^2} + \frac{4(2)}{(s-1)^2 + 2^2}\right\}$$

$$w(t) = -e^{-t} + 3e^t \cos 2t + 4e^t \sin 2t$$

$$y(t) = w(t - \pi) = -e^{-(t-\pi)} + 3e^{t-\pi} \cos 2(t-\pi) + 4e^{t-\pi} \sin 2(t-\pi)$$

$$\underline{= -e^{-t+\pi} + 3e^{t-\pi} \cos 2t + 4e^{t-\pi} \sin 2t}$$

Exercise

Solve using the Laplace transform: $y'' + y = t^2 + 2$; $y(0) = 1, y'(0) = -1$

Solution

$$\mathcal{L}\{y'' + y\} = \mathcal{L}\{t^2 + 2\}$$

$$s^2 Y(s) - sy(0) - y'(0) + Y(s) = \frac{2}{s^3} + \frac{2}{s}$$

$$y(0) = 1; \quad y'(0) = -1$$

$$s^2 Y(s) - s + 1 + Y(s) = \frac{2 + 2s^2}{s^3}$$

$$(s^2 + 1)Y(s) = \frac{2 + 2s^2}{s^3} + s - 1$$

$$Y(s) = \frac{2 + 2s^2 + s^4 - s^3}{s^3(s^2 + 1)} = \frac{A}{s} + \frac{B}{s^2} + \frac{C}{s^3} + \frac{Ds + E}{s^2 + 1}$$

$$s^4 - s^3 + 2s^2 + 2 = As^4 + As^2 + Bs^3 + Bs + Cs^2 + C + Ds^4 + Es^3$$

$$s^4 \quad A + D = 1 \quad \underline{D = 1}$$

$$s^3 \quad B + E = -1 \quad \underline{E = -1}$$

$$s^2 \quad A + C = 2 \quad \underline{A = 0}$$

$$s \quad \underline{B = 0}$$

$$s^0 \quad \underline{C = 2}$$

$$Y(s) = \frac{2}{s^3} + \frac{s}{s^2 + 1} - \frac{1}{s^2 + 1}$$

$$\underline{y(t) = t^2 + \cos t - \sin t}$$

Exercise

Solve using the Laplace transform: $y'' + y = \sqrt{2} \sin \sqrt{2}t$; $y(0) = 10$, $y'(0) = 0$

Solution

$$\mathcal{L}\{y'' + y\} = \mathcal{L}\{\sqrt{2} \sin \sqrt{2}t\}$$

$$s^2 Y(s) - sy(0) - y'(0) + Y(s) = \sqrt{2} \frac{\sqrt{2}}{s^2 + 2}$$

$$y(0) = 10; \quad y'(0) = 0$$

$$(s^2 + 1)Y(s) - 10s = \frac{2}{s^2 + 2}$$

$$(s^2 + 1)Y(s) = \frac{2}{s^2 + 2} + 10s$$

$$(s^2 + 1)Y(s) = \frac{10s^3 + 20s + 2}{s^2 + 2}$$

$$Y(s) = \frac{10s^3 + 20s + 2}{(s^2 + 1)(s^2 + 2)} = \frac{As + B}{s^2 + 1} + \frac{Cs + D}{s^2 + 2}$$

$$As^3 + 2As + Bs^2 + 2B + Cs^3 + Cs + Ds^2 + D = 10s^3 + 20s + 2$$

$$\begin{array}{rcl} s^3 & A + C = 10 & \\ s^2 & B + D = 0 & \\ s^1 & 2A + C = 20 & \\ s^0 & 2B + D = 2 & \end{array} \rightarrow \begin{cases} A + C = 10 \\ 2A + C = 20 \end{cases} \rightarrow \underline{A = 10, C = 0} \\ \rightarrow \begin{cases} B + D = 0 \\ 2B + D = 2 \end{cases} \rightarrow \underline{B = 2, D = -2}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{10s}{s^2+1} + \frac{2}{s^2+1} - \sqrt{2}\frac{\sqrt{2}}{s^2+2}\right\}$$

$$\underline{y(t) = 10\cos t + 2\sin t - \sqrt{2}\sin\sqrt{2}t}$$

Exercise

Solve using the Laplace transform: $y'' + y = -2\cos 2t$; $y(0) = 1$, $y'(0) = -1$

Solution

$$\mathcal{L}(y'' + y)(s) = \mathcal{L}(-2\cos 2t)(s)$$

$$s^2Y(s) - sy(0) - y'(0) + Y(s) = -2\frac{s}{s^2+4} \quad y(0) = 1 \quad y'(0) = -1$$

$$s^2Y(s) - s + 1 + Y(s) = -2\frac{s}{s^2+4}$$

$$(s^2+1)Y(s) = \frac{-2s}{s^2+4} + s - 1$$

$$Y(s) = \frac{-2s}{(s^2+4)(s^2+1)} + \frac{s}{s^2+1} - \frac{1}{s^2+1}$$

$$= \frac{As}{s^2+4} + \frac{Bs}{s^2+1} + \frac{s}{s^2+1} - \frac{1}{s^2+1}$$

$$\frac{-2s}{(s^2+4)(s^2+1)} = \frac{As}{s^2+4} + \frac{Bs}{s^2+1}$$

$$\Rightarrow -2s = As(s^2+1) + Bs(s^2+4)$$

$$-2s = As^3 + As + Bs^3 + 4Bs$$

$$-2s = (A+B)s^3 + (A+4B)s$$

$$\begin{cases} A+B=0 \\ A+4B=-2 \end{cases} \Rightarrow \boxed{A = \frac{2}{3}} \quad \boxed{B = -\frac{2}{3}}$$

$$Y(s) = \frac{2}{3}\frac{s}{s^2+4} - \frac{2}{3}\frac{s}{s^2+1} + \frac{s}{s^2+1} - \frac{1}{s^2+1}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{2}{3}\frac{s}{s^2+4} + \frac{1}{3}\frac{s}{s^2+1} - \frac{1}{s^2+1}\right\}$$

$$\underline{y(t) = \frac{2}{3}\cos 2t + \frac{1}{3}\cos t - \sin t}$$

Exercise

Solve using the Laplace transform: $y'' - y' = e^t \cos t$; $y(0) = 0, \quad y'(0) = 0$

Solution

$$\mathcal{L}\{y'' - y'\} = \mathcal{L}\{e^t \cos t\}$$

$$s^2 Y(s) - sy(0) - y'(0) + sY(s) - y(0) = \frac{s-1}{(s-1)^2 + 1} \quad y(0) = 0; \quad y'(0) = 0$$

$$(s^2 + s)Y(s) = \frac{s-1}{s^2 - 2s + 2}$$

$$Y(s) = \frac{s-1}{(s^2 + s)(s^2 - 2s + 2)} = \frac{A}{s} + \frac{B}{s+1} + \frac{Cs+D}{s^2 - 2s + 2}$$

$$As^3 - As^2 + 2A + Bs^3 - 2Bs^2 + 2Bs + Cs^3 + Cs^2 +Ds^2 +Ds = s-1$$

$$\begin{array}{lcl} s^3 & A+B+C+D=0 \\ s^2 & -A-2B+C+D=0 \\ s^1 & 2B+D=1 \\ s^0 & 2A=-1 \rightarrow A=-\frac{1}{2} \end{array} \rightarrow \left\{ \begin{array}{l} B+C+D=\frac{1}{2} \\ -2B+C+D=-\frac{1}{2} \\ 2B+D=1 \end{array} \Rightarrow \underline{B=\frac{1}{3} \quad C=-\frac{1}{6} \quad D=\frac{1}{3}} \right.$$

$$Y(s) = -\frac{1}{2}\frac{1}{s} + \frac{1}{3}\frac{1}{s+1} - \frac{1}{6}\frac{s-2}{(s-1)^2 + 1}$$

$$= -\frac{1}{2}\frac{1}{s} + \frac{1}{3}\frac{1}{s+1} - \frac{1}{6}\frac{s-1-1}{(s-1)^2 + 1}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{-\frac{1}{2}\frac{1}{s} + \frac{1}{3}\frac{1}{s+1} - \frac{1}{6}\frac{s-1}{(s-1)^2 + 1} + \frac{1}{6}\frac{1}{(s-1)^2 + 1}\right\}$$

$$\underline{y(t) = -\frac{1}{2}t + \frac{1}{3}e^{-t} - \frac{1}{6}e^t \cos t + \frac{1}{6}e^t \sin t}$$

Exercise

Solve using the Laplace transform: $y'' + y' - y = t^3$; $y(0) = 1$, $y'(0) = 0$

Solution

$$\mathcal{L}(y'' - y' - y) = \mathcal{L}(t^3)$$

$$s^2 Y(s) - sy(0) - y'(0) + sY(s) - y(0) - Y(s) = \frac{6}{s^4} \quad y(0) = 1 \quad y'(0) = 0$$

$$(s^2 + s - 1)Y(s) - s - 1 = \frac{6}{s^4}$$

$$(s^2 + s - 1)Y(s) = \frac{6}{s^4} + s + 1$$

$$Y(s) = \frac{s^5 + s^4 + 6}{s^4(s^2 + s - 1)} = \frac{A}{s} + \frac{B}{s^2} + \frac{C}{s^3} + \frac{D}{s^4} + \frac{E}{s + \frac{1}{2} - \frac{\sqrt{5}}{2}} + \frac{F}{s + \frac{1}{2} + \frac{\sqrt{5}}{2}}$$

$$s^5 \quad A + E + F = 1 \quad E + F = 19$$

$$s^4 \quad A + B + \left(\frac{1+\sqrt{5}}{2}\right)E + \left(\frac{1-\sqrt{5}}{2}\right)F = 1 \quad \left(\frac{1+\sqrt{5}}{2}\right)E + \left(\frac{1-\sqrt{5}}{2}\right)F = 31$$

$$s^3 \quad -A + B + C = 0 \quad A = -18$$

$$s^2 \quad -B + C + D = 0 \quad B = -12$$

$$s^1 \quad -C + D = 0 \quad C = -6$$

$$s^0 \quad -D = 6 \quad D = -6$$

$$F = \frac{19}{2} - \frac{43\sqrt{5}}{10} \quad E = -\frac{19}{2} + \frac{43\sqrt{5}}{10}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{-\frac{18}{s} - \frac{12}{s^2} - \frac{6}{s^3} - \frac{6}{s^4} + \frac{-\frac{19}{2} + \frac{43\sqrt{5}}{10}}{s + \frac{1}{2} - \frac{\sqrt{5}}{2}} + \frac{\frac{19}{2} - \frac{43\sqrt{5}}{10}}{s + \frac{1}{2} + \frac{\sqrt{5}}{2}}\right\}$$

$$y(t) = -18 - 12t - 3t^2 - t^3 + \left(-\frac{19}{2} + \frac{43\sqrt{5}}{10}\right)e^{\left(\frac{-1+\sqrt{5}}{2}\right)t} + \left(\frac{19}{2} - \frac{43\sqrt{5}}{10}\right)e^{\left(\frac{1+\sqrt{5}}{2}\right)t}$$

Exercise

Solve using the Laplace transform: $y'' - y' - 2y = 4t^2$, $y(0) = 1$, $y'(0) = 4$

Solution

$$\mathcal{L}\{y'' - y' - 2y\}(s) = \mathcal{L}\{4t^2\}(s)$$

$$\mathcal{L}\{t^n\}(s) = \frac{n!}{s^{n+1}}$$

$$s^2 Y(s) - sy(0) - y'(0) - sY(s) + y(0) - 2Y(s) = \frac{8}{s^3} \quad y(0) = 1, \quad y'(0) = 4$$

$$(s^2 - s - 2)Y(s) - s - 4 + 1 = \frac{8}{s^3}$$

$$(s+1)(s-2)Y(s) = \frac{8}{s^3} + s + 3$$

$$Y(s) = \frac{s^4 + 3s^3 + 8}{s^3(s+1)(s-2)} = \frac{A}{s} + \frac{B}{s^2} + \frac{C}{s^3} + \frac{D}{s+1} + \frac{E}{s-2}$$

$$\begin{aligned} s^4 + 3s^3 + 8 &= As^2(s^2 - s - 2) + Bs(s^2 - s - 2) + Cs^2 - Cs - 2C + Ds^3(s-2) + Es^3(s+1) \\ &= As^4 - As^3 - 2As^2 + Bs^3 - Bs^2 - 2Bs + Cs^2 - Cs - 2C + Ds^4 - 2Ds^3 + Es^4 + Es^3 \end{aligned}$$

$$\begin{cases} s^4 & A + D + E = 1 & D + E = 4 \\ s^3 & -A + B - 2D + E = 3 & -2D + E = -2 \\ s^2 & -2A - B + C = 0 & \rightarrow \underline{A = -3} \\ s & -2B - C = 0 & \rightarrow \underline{B = 2} \\ s^0 & -2C = 8 & \rightarrow \underline{C = -4} \end{cases} \Rightarrow \underline{D = 2, E = 2}$$

$$\mathcal{L}^{-1}\{Y(s)\}(t) = \mathcal{L}^{-1}\left\{-\frac{3}{s} + \frac{2}{s^2} - \frac{4}{s^3} + \frac{2}{s+1} + \frac{2}{s-2}\right\}(t)$$

$$\underline{y(t) = -3 + 2t - 2t^2 + 2e^{-t} + 2e^{2t}}$$

Exercise

Solve using the Laplace transform: $y'' - y' - 2y = e^{2t}$; $y(0) = -1$, $y'(0) = 0$

Solution

$$\mathcal{L}(y'' - y' - 2y) = \mathcal{L}(e^{2t})$$

$$s^2Y(s) - sy(0) - y'(0) - (sY(s) - y(0)) - 2Y(s) = \frac{1}{s-2} \quad y(0) = -1 \quad y'(0) = 0$$

$$s^2Y(s) + s - sY(s) - 1 - 2Y(s) = \frac{1}{s-2}$$

$$(s^2 - s - 2)Y(s) = \frac{1}{s-2} - s + 1$$

$$(s+1)(s-2)(Y(s)) = \frac{1}{s-2} - s + 1$$

$$\begin{aligned} Y(s) &= \frac{1}{(s+1)(s-2)^2} - \frac{s-1}{(s+1)(s-2)} \\ &= \frac{1 - (s-1)(s-2)}{(s+1)(s-2)^2} \end{aligned}$$

$$Y(s) = \frac{-s^2 + 3s - 1}{(s+1)(s-2)^2} = \frac{A}{s+1} + \frac{B}{s-2} + \frac{C}{(s-2)^2}$$

$$-s^2 + 3s - 1 = As^2 - 4As + 4A + Bs^2 - Bs - 2B + Cs + C$$

$$-s^2 + 3s - 1 = (A+B)s^2 + (-4A-B+C)s + 4A-2B+C$$

$$\begin{cases} A+B=-1 \\ -4A-B+C=3 \\ 4A-2B+C=-1 \end{cases} \Rightarrow A=-\frac{5}{9} \quad B=-\frac{4}{9} \quad C=\frac{1}{3}$$

$$Y(s) = -\frac{5}{9} \frac{1}{s+1} - \frac{4}{9} \frac{1}{s-2} + \frac{1}{3} \frac{1}{(s-2)^2}$$

$$y(t) = -\frac{5}{9} \mathcal{L}^{-1} \left\{ \frac{1}{s+1} \right\} - \frac{4}{9} \mathcal{L}^{-1} \left\{ \frac{1}{s-2} \right\} + \frac{1}{3} \mathcal{L}^{-1} \left\{ \frac{1}{(s-2)^2} \right\}$$

$$\underline{y(t) = -\frac{5}{9}e^{-t} - \frac{4}{9}e^{2t} + \frac{1}{3}te^{2t} \quad |}$$

Exercise

Solve using the Laplace transform: $y'' - y' - 2y = 0$, $y(0) = -2$, $y'(0) = 5$

Solution

$$\mathcal{L}(y'' - y' - 2y) = 0$$

$$s^2 Y(s) - sy(0) - y'(0) - sY(s) + y(0) - 2Y(s) = 0 \quad y(0) = -2 \quad y'(0) = 5$$

$$(s^2 - s - 2)Y(s) = 7 - 2s$$

$$Y(s) = \frac{7-2s}{s^2-s-2} = \frac{A}{s+1} + \frac{B}{s-2}$$

$$\begin{array}{l} s \quad A+B=-2 \\ s^0 \quad -2A+B=7 \end{array} \rightarrow \underline{A=-3, B=1}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{ \frac{-3}{s+1} + \frac{1}{s-2} \right\}$$

$$\underline{y(t) = e^{2t} - 3e^{-t} \quad |}$$

Exercise

Solve using the Laplace transform: $y'' - y' - 2y = -8\cos t - 2\sin t$; $y\left(\frac{\pi}{2}\right) = 1$, $y'\left(\frac{\pi}{2}\right) = 0$

Solution

$$\text{Let: } w(t) = y\left(t + \frac{\pi}{2}\right) \leftrightarrow y(t) = w\left(t - \frac{\pi}{2}\right)$$

$$y'' - y' - 2y = -8\cos t - 2\sin t \rightarrow w'' - w' - 2w = -8\cos\left(t + \frac{\pi}{2}\right) - 2\sin\left(t + \frac{\pi}{2}\right)$$

$$w'' - w' - 2w = -8\left(\cos t \cos \frac{\pi}{2} - \sin t \sin \frac{\pi}{2}\right) - 2\left(\sin t \cos \frac{\pi}{2} + \cos t \sin \frac{\pi}{2}\right)$$

$$\mathcal{L}\{w'' - w' - 2w\} = \mathcal{L}\{8\sin t - 2\cos t\}$$

$$s^2W(s) - sw(0) - w'(0) - sW(s) + w(0) - 2W(s) = \frac{8}{s^2 + 1} - \frac{2s}{s^2 + 1}$$

$$y\left(\frac{\pi}{2}\right) = w(0) = 1, \quad y'\left(\frac{\pi}{2}\right) = w'(0) = 0$$

$$(s^2 - s - 2)W(s) = \frac{8 - 2s}{s^2 + 1} + s - 1$$

$$W(s) = \frac{s^3 - s^2 - s + 7}{(s+1)(s-2)(s^2+1)} = \frac{A}{s+1} + \frac{B}{s-2} + \frac{Cs+D}{s^2+1}$$

$$s^3 - s^2 - s + 7 = As^3 - 2As^2 + As - 2A + Bs^3 + Bs^2 + Bs + B + Cs^3 - Cs^2 - 2Cs + Ds^2 - Ds - 2D$$

$$\begin{cases} s^3 & A + B + C = 1 \\ s^2 & -2A + B - C + D = -1 \\ s^1 & A + B - 2C - D = -1 \\ s^0 & -2A + B - 2D = 7 \end{cases} \rightarrow A = -1, B = \frac{3}{5}, C = \frac{7}{5}, D = -\frac{11}{5}$$

$$\mathcal{L}^{-1}\{W(s)\} = \mathcal{L}^{-1}\left\{-\frac{13}{3}\frac{1}{s+1} + \frac{74}{15}\frac{1}{s-2} + \frac{7}{5}\frac{s}{s^2+1} - \frac{11}{5}\frac{1}{s^2+1}\right\}$$

$$w(t) = -e^{-t} + \frac{3}{5}e^{2t} + \frac{7}{5}\cos t - \frac{11}{5}\sin t$$

$$y(t) = w\left(t - \frac{\pi}{2}\right) = -e^{-t+\frac{\pi}{2}} + \frac{3}{5}e^{2\left(t-\frac{\pi}{2}\right)} + \frac{7}{5}\cos\left(t - \frac{\pi}{2}\right) - \frac{11}{5}\sin\left(t - \frac{\pi}{2}\right)$$

$$= -e^{-t+\frac{\pi}{2}} + \frac{3}{5}e^{2t-\pi} + \frac{7}{5}\left(\cos t \cos \frac{\pi}{2} + \sin t \sin \frac{\pi}{2}\right) - \frac{11}{5}\left(\sin t \cos \frac{\pi}{2} - \cos t \sin \frac{\pi}{2}\right)$$

$$= \frac{3}{5}e^{2t-\pi} - e^{-t+\frac{\pi}{2}} + \frac{7}{5}\sin t + \frac{11}{5}\cos t$$

Exercise

Solve using the Laplace transform: $x'' - x' - 6x = 0$; $x(0) = 2$, $x'(0) = -1$

Solution

$$\mathcal{L}\{x'' - x' - 6x\} = 0$$

$$s^2 X(s) - sx(0) - x'(0) - sX(s) + x(0) - 6X(s) = 0 \quad x(0) = 2, \quad x'(0) = -1$$

$$(s^2 - s - 6)X(s) - 2s + 1 + 2 = 0$$

$$(s^2 - s - 6)X(s) = 2s - 3$$

$$X(s) = \frac{2s-3}{s^2-s-6} = \frac{A}{s-3} + \frac{B}{s+2}$$

$$As + 2A + Bs - 3B = 2s - 3$$

$$\begin{cases} A + B = 2 \\ 2A - 3B = -3 \end{cases} \quad \begin{vmatrix} 1 & 1 \\ 2 & -3 \end{vmatrix} = -5 \quad \begin{vmatrix} 2 & 1 \\ -3 & -3 \end{vmatrix} = -3 \quad \rightarrow A = \frac{3}{5}, \quad B = \frac{7}{5}$$

$$\mathcal{L}^{-1}\{X(s)\} = \mathcal{L}^{-1}\left\{\frac{3}{5} \frac{1}{s-3} + \frac{7}{5} \frac{1}{s+2}\right\}$$

$$\underline{x(t) = \frac{3}{5}e^{3t} + \frac{7}{5}e^{-2t}}$$

Exercise

Solve using the Laplace transform: $y'' + 2y' + y = 0, \quad y(0) = 1, \quad y'(0) = 1$

Solution

$$\mathcal{L}\{y'' + 2y' + y\} = 0$$

$$s^2 Y(s) - sy(0) - y'(0) + 2sY(s) - 2y(0) + Y(s) = 0 \quad y(0) = 1 \quad y'(0) = 1$$

$$(s^2 + 2s + 1)Y(s) - s - 1 - 2 = 0$$

$$Y(s) = \frac{s+3}{(s+1)^2} = \frac{A}{s+1} + \frac{B}{(s+1)^2}$$

$$s + 3 = As + A + B$$

$$s \quad \underline{A = 1}$$

$$s^0 \quad A + B = 3 \quad \underline{B = 2}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{1}{s+1} + \frac{2}{(s+1)^2}\right\}$$

$$\underline{y(t) = e^{-t} + 2te^{-t}}$$

Exercise

Solve using the Laplace transform: $y'' + 2y' + y = t$, $y(0) = -3$, $y(1) = -1$

Solution

$$\mathcal{L}\{y'' + 2y' + y\}(s) = \mathcal{L}\{t\}(s)$$

$$s^2 Y(s) - sy(0) - y'(0) + 2sY(s) - 2y(0) + Y(s) = \frac{1}{s^2} \quad y(0) = -3$$

$$(s^2 + 2s + 1)Y(s) + 3s - y'(0) + 6 = \frac{1}{s^2}$$

$$(s+1)^2 Y(s) = \frac{1}{s^2} - 3s - 6 + y'(0)$$

$$Y(s) = \frac{-3s^3 + (y'(0) - 6)s^2 + 1}{s^2(s+1)^2} \quad \text{Let } k = y'(0) - 6$$

$$= \frac{-3s^3 + ks^2 + 1}{s^2(s+1)^2} = \frac{A}{s} + \frac{B}{s^2} + \frac{C}{s+1} + \frac{D}{(s+1)^2}$$

$$-3s^3 + ks^2 + 1 = As^3 + 2As^2 + As + Bs^2 + 2Bs + B + Cs^3 + Cs^2 + Ds^2$$

$$\begin{cases} s^3 & A + C = -3 & \rightarrow \underline{C = -1} \\ s^2 & 2A + B + C + D = k & \rightarrow \underline{D = k + 4} \\ s & A + 2B = 0 & \rightarrow \underline{A = -2} \\ s^0 & B = 1 \end{cases}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{-\frac{2}{s} + \frac{1}{s^2} - \frac{1}{s+1} + \frac{k+4}{(s+1)^2}\right\}$$

$$y(t) = -2 + t - e^{-t} + (k+4)te^{-t} \quad y(1) = -1$$

$$-1 = -1 - e^{-1} + (k+4)e^{-1}$$

$$(k+4)e^{-1} = e^{-1}$$

$$k+4 = 1 \rightarrow \underline{k = -3}$$

$$\underline{y(t) = -2 + t - e^{-t} + te^{-t}}$$

Exercise

Solve using the Laplace transform: $y'' - 2y' - y = e^{2t} - e^t$; $y(0) = 1$, $y'(0) = 3$

Solution

$$\mathcal{L}\{y'' - 2y' - y\} = \mathcal{L}\{e^{2t} - e^t\}$$

$$s^2 Y(s) - sy(0) - y'(0) - 2sY(s) + 2y(0) - Y(s) = \frac{1}{s-2} - \frac{1}{s-1} \quad y(0)=1 \quad y'(0)=3$$

$$(s^2 - 2s - 1)Y(s) = \frac{1}{(s-2)(s-1)} + s + 1$$

$$Y(s) = \frac{s^3 - 2s^2 - s + 3}{(s-2)(s-1)(s-1-\sqrt{2})(s-1+\sqrt{2})} = \frac{A}{s-2} + \frac{B}{s-1} + \frac{C}{s-1-\sqrt{2}} + \frac{D}{s-1+\sqrt{2}}$$

$$s^3 - 2s^2 - s + 3 = A(s-1)(s^2 - 2s - 1) + B(s-2)(s^2 - 2s - 1) + C(s-1+\sqrt{2})(s^2 - 3s + 2) + D(s-1-\sqrt{2})(s^2 - 3s + 2)$$

$$s^3 \quad A + B + C + D = 1$$

$$s^2 \quad -3A - 4B + (-4 + \sqrt{2})C + (-4 - \sqrt{2})D = -2$$

$$s^1 \quad A + 3B + (5 - 3\sqrt{2})C + (5 + 3\sqrt{2})D = -1$$

$$s^0 \quad A + 2B + (-2 + 2\sqrt{2})C - 2(1 + \sqrt{2})D = 3$$

$$\underline{A = -1 \quad B = \frac{1}{2} \quad C = \frac{3}{4}(1 + \sqrt{2}) \quad D = \frac{3}{4}(1 - \sqrt{2})}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{-1}{s-2} + \frac{1}{2} \frac{1}{s-1} + \frac{3}{4}(1 + \sqrt{2}) \frac{1}{s-1-\sqrt{2}} + \frac{3}{4}(1 - \sqrt{2}) \frac{1}{s-1+\sqrt{2}}\right\}$$

$$\underline{y(t) = -e^{2t} + \frac{1}{2}e^t + \left(\frac{3}{4} + \frac{3\sqrt{2}}{4}\right)e^{(1+\sqrt{2})t} + \left(\frac{3}{4} - \frac{3\sqrt{2}}{4}\right)e^{(1-\sqrt{2})t}}$$

Exercise

Solve using the Laplace transform: $y'' - 2y' + y = 6t - 2$; $y(-1) = 3$, $y'(-1) = 7$

Solution

$$\text{Let: } w(t) = y(t-1) \leftrightarrow y(t) = w(t+1)$$

$$w'' - 2w' + w = 6(t-1) - 2 = 6t - 8$$

$$\mathcal{L}\{w'' - 2w' + w\} = \mathcal{L}\{6t - 8\}$$

$$s^2 W(s) - sw(0) - w'(0) - 2sW(s) + 2w(0) + W(s) = \frac{6}{s^2} - \frac{8}{s} \quad y(-1) = w(0) = 3, \quad y'(-1) = w'(0) = 7$$

$$(s^2 - 2s + 1)W(s) = \frac{6-8s}{s^2} + 3s + 1$$

$$W(s) = \frac{3s^3 + s^2 - 8s + 6}{s^2(s-1)^2} = \frac{A}{s} + \frac{B}{s^2} + \frac{C}{s-1} + \frac{D}{(s-1)^2}$$

$$3s^3 + s^2 - 8s + 6 = As(s^2 - 2s + 1) + B(s^2 - 2s + 1) + Cs^2(s - 1) + Ds^2$$

$$s^3 \quad A + C = 3 \quad \underline{C = -1}$$

$$s^2 \quad -2A + B - C + D = 1 \quad \underline{D = 2}$$

$$s^1 \quad A - 2B = -8 \quad \underline{A = 4}$$

$$s^0 \quad \underline{B = 6}$$

$$\mathcal{L}^{-1}\{W(s)\} = \mathcal{L}^{-1}\left\{\frac{4}{s} + \frac{6}{s^2} - \frac{1}{s-1} + \frac{2}{(s-1)^2}\right\}$$

$$\underline{w(t) = 4 + 6t - e^t + 2te^t}$$

$$y(t) = w(t+1) = 4 + 6(t+1) - e^{t+1} + 2(t+1)e^{t+1}$$

$$\underline{= 6t + 10 + 2te^{t+1} + e^{t+1}}$$

Exercise

Solve using the Laplace transform: $y'' - 2y' + y = \cos t - \sin t$; $y(0) = 1, \quad y'(0) = 3$

Solution

$$\mathcal{L}\{y'' - 2y' + y\} = \mathcal{L}\{\cos t - \sin t\}$$

$$s^2Y(s) - sy(0) - y'(0) - 2sY(s) + 2y(0) + Y(s) = \frac{s}{s^2+1} - \frac{1}{s^2+1} \quad y(0) = 1, \quad y'(0) = 3$$

$$(s^2 - 2s + 1)Y(s) = \frac{s-1}{s^2+1} + s + 1$$

$$Y(s) = \frac{s^3 + s^2 + 2s}{(s^2+1)(s-1)^2} = \frac{As+B}{s^2+1} + \frac{C}{s-1} + \frac{D}{(s-1)^2}$$

$$s^3 + s^2 + 2s = (As+B)(s^2 - 2s + 1) + C(s^2 + 1) + D(s^2 + 1)$$

$$s^3 \quad A + C = 1$$

$$s^2 \quad -2A + B - C + D = 1$$

$$s^1 \quad A - 2B + C = 2$$

$$s^0 \quad B - C + D = 0$$

$$\underline{A = -\frac{1}{2} \quad B = -\frac{1}{2} \quad C = \frac{3}{2} \quad D = 2}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{-\frac{1}{2}\frac{s}{s^2+1} - \frac{1}{2}\frac{1}{s^2+1} + \frac{3}{2}\frac{1}{s-1} + \frac{2}{(s-1)^2}\right\}$$

$$\underline{y(t) = -\frac{1}{2}\cos t - \frac{1}{2}\sin t + \frac{3}{2}e^t + 2te^t}$$

Exercise

Solve using the Laplace transform: $y'' - 2y' + 5y = 0$, $y(0) = 2$, $y'(0) = 4$

Solution

$$\mathcal{L}\{y'' - 2y' + 5y\} = 0$$

$$s^2 Y(s) - sy(0) - y'(0) - 2sY(s) + 2y(0) + 5Y(s) = 0 \quad y(0) = 2 \quad y'(0) = 4$$

$$(s^2 - 2s + 5)Y(s) = 2s$$

$$Y(s) = \frac{2s}{(s-1)^2 + 4}$$

$$= \frac{2(s-1) + 2}{(s-1)^2 + 4}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{2(s-1)}{(s-1)^2 + 4} + \frac{2}{(s-1)^2 + 4}\right\}$$

$$\underline{y(t) = 2e^t \cos 2t + e^t \sin 2t}$$

Exercise

Solve using the Laplace transform: $y'' - 2y' + 5y = 1 + t$, $y(0) = 0$, $y'(0) = 0$

Solution

$$\mathcal{L}\{y'' - 2y' + 5y\} = \mathcal{L}\{1 + t\}$$

$$s^2 Y(s) - sy(0) - y'(0) - 2sY(s) + 2y(0) + 5Y(s) = \frac{1}{s} + \frac{1}{s^2} \quad y(0) = 0 \quad y'(0) = 0$$

$$(s^2 - 2s + 5)Y(s) = \frac{s+1}{s^2}$$

$$Y(s) = \frac{s+1}{s^2((s-1)^2 + 4)} = \frac{A}{s} + \frac{B}{s^2} + \frac{C(s-1) + D}{(s-1)^2 + 4}$$

$$s+1 = As^3 - 2As^2 + 5As + Bs^2 - 2Bs + 5B + Cs^3 - Cs^2 + Ds^2$$

$$s^3 \quad A + C = 0 \quad \underline{C = -\frac{7}{25}}$$

$$s^2 \quad -2A + B - C + D = 0 \quad \underline{D = \frac{2}{25}}$$

$$s^1 \quad 5A - 2B = 1 \quad \underline{A = \frac{7}{25}}$$

$$s^0 \quad 5B = 1 \quad \underline{B = \frac{1}{5}}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{7}{25}\frac{1}{s} + \frac{1}{5}\frac{1}{s^2} - \frac{7}{25}\frac{s-1}{(s-1)^2 + 2^2} + \frac{1}{25}\frac{2}{(s-1)^2 + 2^2}\right\}$$

$$\underline{y(t) = \frac{7}{25} + \frac{1}{5}t - \frac{7}{25}e^t \cos 2t + \frac{1}{25}e^t \sin 2t}$$

Exercise

Solve using the Laplace transform: $y'' + 3y' = -3t$; $y(0) = -1$, $y'(0) = 1$

Solution

$$\mathcal{L}(y'' + 3y') = \mathcal{L}(-3t)$$

$$s^2Y(s) - sy(0) - y'(0) + 3(sY(s) - y(0)) = -3\frac{1}{s^2} \quad y(0) = -1 \quad y'(0) = 1$$

$$s^2Y(s) + s - 1 + 3sY(s) + 3 = -\frac{3}{s^2}$$

$$(s^2 + 3s)Y(s) = -\frac{3}{s^2} - s - 2$$

$$Y(s) = -\frac{3}{s^3(s+3)} - \frac{s+2}{s(s+3)}$$

$$\frac{3}{s^3(s+3)} = \frac{A}{s} + \frac{B}{s^2} + \frac{C}{s^3} + \frac{D}{s+3}$$

$$3 = As^2(s+3) + Bs(s+3) + C(s+3) + Ds^3$$

$$3 = (A+D)s^3 + (3A+B)s^2 + (3B+C)s + 3C$$

$$\begin{cases} A+D=0 \\ 3A+B=0 \\ 3B+C=0 \\ 3C=3 \end{cases} \Rightarrow \begin{matrix} A=\frac{1}{9} & B=-\frac{1}{3} \\ C=1 & D=-\frac{1}{9} \end{matrix}$$

$$\frac{s+2}{s(s+3)} = \frac{E}{s} + \frac{F}{s+3}$$

$$\begin{cases} E+F=1 \\ 3E=2 \end{cases} \Rightarrow E=\frac{2}{3} \quad F=\frac{1}{3}$$

$$Y(s) = -\left(\frac{1}{9}\frac{1}{s} - \frac{1}{3}\frac{1}{s^2} + \frac{1}{s^3} - \frac{1}{9}\frac{1}{s+3}\right) - \left(\frac{2}{3}\frac{1}{s} + \frac{1}{3}\frac{1}{s+3}\right)$$

$$= -\frac{1}{9}\frac{1}{s} + \frac{1}{3}\frac{1}{s^2} - \frac{1}{s^3} + \frac{1}{9}\frac{1}{s+3} - \frac{2}{3}\frac{1}{s} - \frac{1}{3}\frac{1}{s+3}$$

$$= -\frac{7}{9}\frac{1}{s} + \frac{1}{3}\frac{1}{s^2} - \frac{1}{s^3} - \frac{2}{9}\frac{1}{s+3}$$

$$y(t) = -\frac{7}{9} \mathcal{L}^{-1} \left\{ \frac{1}{s} \right\} + \frac{1}{3} \mathcal{L}^{-1} \left\{ \frac{1}{s^2} \right\} - \mathcal{L}^{-1} \left\{ \frac{1}{2!} \frac{2!}{s^3} \right\} - \frac{2}{9} \mathcal{L}^{-1} \left\{ \frac{1}{s+3} \right\}$$

$$\underline{y(t) = -\frac{7}{9} + \frac{1}{3}t - \frac{1}{2}t^2 - \frac{2}{9}e^{-3t}}$$

Exercise

Solve using the Laplace transform: $y'' + 3y = t^3$; $y(0) = 0$, $y'(0) = 0$

Solution

$$\mathcal{L}\{y'' + 3y'\} = \mathcal{L}\{t^3\}$$

$$s^2 Y(s) - sy(0) - y'(0) + 3Y(s) = \frac{6}{s^4}$$

$$y(0) = 0 \quad y'(0) = 0$$

$$(s^2 + 3)Y(s) = \frac{6}{s^4}$$

$$Y(s) = \frac{6}{s^4(s^2 + 3)} = \frac{A}{s} + \frac{B}{s^2} + \frac{C}{s^3} + \frac{D}{s^4} + \frac{Es + F}{s^2 + 3}$$

$$6 = As^5 + 3As^3 + Bs^4 + 3Bs^2 + Cs^3 + 3Cs + Ds^2 + 3D + Es^5 + Fs^4$$

$$s^5 \quad A + E = 0 \quad \underline{E = 0}$$

$$s^4 \quad B + F = 0 \quad \underline{F = -\frac{2}{3}}$$

$$s^3 \quad 3A + C = 0 \quad \underline{A = 0}$$

$$s^2 \quad 3B + D = 0 \quad \underline{B = -\frac{2}{3}}$$

$$s^1 \quad 3C = 0 \quad \underline{C = 0}$$

$$s^0 \quad 3D = 6 \quad \underline{D = 2}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1} \left\{ -\frac{2}{3} \frac{1}{s^2} + \frac{2}{s^4} + \frac{2}{3} \frac{1}{s^2 + (\sqrt{3})^2} \right\}$$

$$y(t) = -\frac{2}{3}t + \frac{1}{3!}t^3 + \frac{2}{3} \frac{1}{\sqrt{3}} \sin(\sqrt{3}t)$$

$$\underline{= -\frac{2}{3}t + \frac{1}{3}t^3 + \frac{2\sqrt{3}}{9} \sin(\sqrt{3}t)}$$

Exercise

Solve using the Laplace transform: $y'' - 3y' + 2y = e^{-t}$, $y(1) = 0$, $y'(1) = 0$

Solution

$$\text{Let } v = t - 1 \rightarrow t = v + 1$$

$$x(v) = y(t) = y(v+1) \quad y(1) = x(0) = 0 \quad y'(1) = x'(0) = 0$$

$$y''(t) - 3y'(t) + 2y(t) = e^{-t}$$

$$\rightarrow y''(v+1) - 3y'(v+1) + 2y(v+1) = e^{-(v+1)}$$

$$x''(v) - 3x'(v) + 2x(v) = e^{-(v+1)}$$

$$\mathcal{L}\{x'' - 3x' + 2x\} = \mathcal{L}\{e^{-1}e^{-v}\}$$

$$s^2 X(s) - sx(0) - x'(0) - 3sX(s) + 3x(0) + 2X(s) = \frac{e^{-1}}{s+1}$$

$$(s^2 - 3s + 2)X(s) = \frac{e^{-1}}{s+1}$$

$$X(s) = e^{-1} \frac{1}{(s+1)(s-1)(s-2)}$$

$$\frac{1}{(s+1)(s-1)(s-2)} = \frac{A}{s+1} + \frac{B}{s-1} + \frac{C}{s-2}$$

$$1 = As^2 - 3As + 2A + Bs^2 - Bs - 2B + Cs^2 - C$$

$$\begin{cases} s^2 & A + B + C = 0 \\ s & -3A - B = 0 \\ s^0 & 2A - 2B - C = 1 \end{cases}$$

$$\Delta = \begin{vmatrix} 1 & 1 & 1 \\ -3 & -1 & 0 \\ 2 & -2 & -1 \end{vmatrix} = 6 \quad \Delta_A = \begin{vmatrix} 0 & 1 & 1 \\ 0 & -1 & 0 \\ 1 & -2 & -1 \end{vmatrix} = 1 \quad \Delta_B = \begin{vmatrix} 1 & 0 & 1 \\ -3 & 0 & 0 \\ 2 & 1 & -1 \end{vmatrix} = -3$$

$$\underline{A = \frac{1}{6}, \quad B = -\frac{1}{2}, \quad C = \frac{1}{3}}$$

$$\mathcal{L}^{-1}\{X(s)\} = \mathcal{L}^{-1}\left\{e^{-1}\left(\frac{1}{6} \frac{1}{s+1} - \frac{1}{2} \frac{1}{s-1} + \frac{1}{3} \frac{1}{s-2}\right)\right\}$$

$$x(v) = e^{-1}\left(\frac{1}{6}e^{-v} - \frac{1}{2}e^v + \frac{1}{3}e^{2v}\right) \quad v = t - 1$$

$$y(t) = e^{-1}\left(\frac{1}{6}e^{-t+1} - \frac{1}{2}e^{t-1} + \frac{1}{3}e^{2(t-1)}\right)$$

$$\underline{= \frac{1}{6}e^{-t} - \frac{1}{2}e^{t-2} + \frac{1}{3}e^{2t-3}}$$

Exercise

Solve using the Laplace transform: $y'' - 3y' + 2y = \cos t$; $y(0) = 0$, $y'(0) = -1$

Solution

$$\mathcal{L}\{y'' - 3y' + 2y\} = \mathcal{L}\{\cos t\}$$

$$s^2 Y(s) - sy(0) - y'(0) - 3sY(s) + 3y(0) + 2Y(s) = \frac{s}{s^2 + 1} \quad y(0) = 0 \quad y'(0) = -1$$

$$(s^2 - 3s + 2)Y(s) = \frac{s}{s^2 + 1} - 1$$

$$Y(s) = \frac{-s^2 + s - 1}{(s-1)(s-2)(s^2 + 1)} = \frac{A}{s-1} + \frac{B}{s-2} + \frac{Cs + D}{s^2 + 1}$$

$$\begin{cases} s^3 & A + B + C = 0 \\ s^2 & -2A - B - 3C + D = -1 \\ s^1 & A + B + 2C - 3D = 1 \\ s^0 & -2A - B + 2D = -1 \end{cases} \rightarrow \underline{A = \frac{1}{2}, B = -\frac{3}{5}, C = \frac{1}{10}, D = -\frac{3}{10}}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{1}{2} \frac{1}{s-1} - \frac{3}{5} \frac{1}{s-2} + \frac{1}{10} \frac{s}{s^2 + 1} - \frac{3}{10} \frac{1}{s^2 + 1}\right\}$$

$$\underline{y(t) = \frac{1}{2}e^t - \frac{3}{5}e^{2t} + \frac{1}{10}\cos t - \frac{3}{10}\sin t}$$

Exercise

Solve using the Laplace transform: $y'' - 4y = e^{-t}$; $y(0) = -1$, $y'(0) = 0$

Solution

$$\mathcal{L}(y'' - 4y) = \mathcal{L}(e^{-t})$$

$$s^2 Y(s) - sy(0) - y'(0) - 4Y(s) = \frac{1}{s+1} \quad y(0) = -1 \quad y'(0) = 0$$

$$(s^2 - 4)Y(s) + s = \frac{1}{s+1}$$

$$Y(s) = \frac{1}{(s+1)(s^2 - 4)} - \frac{s}{s^2 - 4}$$

$$\frac{1}{(s+1)(s^2 - 4)} = \frac{A}{s+1} + \frac{B}{s-2} + \frac{C}{s+2}$$

$$1 = As^2 - 4A + Bs^2 + 3Bs + 2B + Cs^2 - Cs - 2C$$

$$1 = (A + B + C)s^2 + (3B - C)s + 2B - 4A - 2C$$

$$\begin{cases} A + B + C = 0 \\ 3B - C = 0 \\ -4A + 2B - 2C = 1 \end{cases} \Rightarrow A = -\frac{1}{3} \quad B = \frac{1}{12} \quad C = \frac{1}{4}$$

$$\frac{s}{s^2-4} = \frac{D}{s-2} + \frac{E}{s+2}$$

$$s = (D + E)s + 2D - 2E$$

$$\begin{cases} D + E = 1 \\ 2D - 2E = 0 \end{cases} \Rightarrow D = \frac{1}{2} \quad E = \frac{1}{2}$$

$$Y(s) = -\frac{1}{3} \frac{1}{s+1} + \frac{1}{12} \frac{1}{s-2} + \frac{1}{4} \frac{1}{s+2} - \frac{1}{2} \frac{1}{s-2} - \frac{1}{2} \frac{1}{s+2}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{-\frac{1}{3} \frac{1}{s+1} - \frac{5}{12} \frac{1}{s-2} - \frac{1}{4} \frac{1}{s+2}\right\}$$

$$\underline{y(t) = -\frac{1}{3}e^{-t} - \frac{5}{12}e^{2t} - \frac{1}{4}e^{-2t}}$$

Exercise

Solve using the Laplace transform: $y'' - 4y' = 6e^{3t} - 3e^{-t}$, $y(0) = 1$ $y'(0) = -1$

Solution

$$\mathcal{L}\{y'' - 4y'\} = \mathcal{L}\{6e^{3t} - 3e^{-t}\}$$

$$s^2Y(s) - sy(0) - y'(0) - 4sY(s) + 4y(0) = 0$$

$$y(0) = 1; \quad y'(0) = -1$$

$$(s^2 - 4s)Y(s) - s + 1 + 4 = 0$$

$$Y(s) = \frac{s-5}{s(s-4)} = \frac{A}{s} + \frac{B}{s-4}$$

$$As - 4A + Bs = s - 5$$

$$\begin{cases} A + B = 1 \\ -4A = -5 \end{cases} \rightarrow \underline{A = \frac{5}{4}, B = -\frac{1}{4}}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{5}{4} \frac{1}{s} - \frac{1}{4} \frac{1}{s-4}\right\}$$

$$\underline{y(t) = \frac{5}{4}t - \frac{1}{4}e^{4t}}$$

Exercise

Solve using the Laplace transform: $y'' - 4y' + 4y = t^3e^{2t}$; $y(0) = 0$, $y'(0) = 0$

Solution

$$\mathcal{L}\{y'' - 4y' + 4y\} = \mathcal{L}\{t^3 e^{2t}\}$$

$$s^2 Y(s) - sy(0) - y'(0) - 4sY(s) + 4y(0) + 4Y(s) = \frac{6}{(s-2)^4} \quad y(0)=0; \quad y'(0)=0$$

$$(s^2 - 4s + 4)Y(s) = \frac{6}{(s-2)^4}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{6}{(s-2)^4}\right\}$$

$$\underline{y(t) = \frac{1}{20}t^5 e^{2t}}$$

Exercise

Solve using the Laplace transform: $y'' - 4y' + 4y = t^3, \quad y(0)=1, \quad y'(0)=0$

Solution

$$\mathcal{L}\{y'' - 4y' + 4y\} = \mathcal{L}\{t^3\}$$

$$s^2 Y(s) - sy(0) - y'(0) - 4sY(s) + 4y(0) + 4Y(s) = \frac{6}{s^4} \quad y(0)=1; \quad y'(0)=0$$

$$(s^2 - 4s + 4)Y(s) = \frac{6}{s^4} + s - 4$$

$$Y(s) = \frac{s^5 - 4s^4 + 6}{s^4(s-2)^2} = \frac{A}{s} + \frac{B}{s^2} + \frac{C}{s^3} + \frac{D}{s^4} + \frac{E}{s-2} + \frac{F}{(s-2)^2}$$

$$s^5 \quad A + E = 1 \quad E = -\frac{1}{4}$$

$$s^4 \quad -4A + B - 2E + F = -4 \quad F = -\frac{13}{8}$$

$$s^3 \quad 4A - 4B + C = 0 \quad A = \frac{3}{4}$$

$$s^2 \quad 4B - 4C + D = 0 \quad B = \frac{9}{8}$$

$$s^1 \quad 4C - 4D = 0 \quad C = \frac{3}{2}$$

$$s^0 \quad 4D = 6 \quad D = \frac{3}{2}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{3}{4}\frac{1}{s} + \frac{9}{8}\frac{1}{s^2} + \frac{3}{2}\frac{1}{s^3} + \frac{3}{2}\frac{1}{s^4} + \frac{1}{4}\frac{1}{s-2} - \frac{13}{8}\frac{1}{(s-2)^2}\right\}$$

$$\underline{y(t) = \frac{3}{4} + \frac{9}{8}t + \frac{3}{2}t^2 + \frac{3}{2}t^3 + \frac{1}{4}e^{2t} - \frac{13}{8}te^{2t}}$$

Exercise

Solve using the Laplace transform: $x'' + 4x' + 4x = t^2$; $x(0) = x'(0) = 0$

Solution

$$\mathcal{L}\{x'' + 4x' + 4x\} = \mathcal{L}\{t^2\}$$

$$s^2 X(s) - sx(0) - x'(0) + 4sX(s) - 4x(0) + 4X(s) = \frac{2}{s^3} \quad x(0) = x'(0) = 0$$

$$(s^2 + 4s + 4)X(s) = \frac{2}{s^3}$$

$$X(s) = \frac{2}{s^3(s+2)^2} = \frac{A}{s^3} + \frac{B}{s^2} + \frac{C}{s} + \frac{D}{s+2} + \frac{E}{(s+2)^2}$$

$$As^2 + 4As + 4A + Bs^3 + 4Bs^2 + 4Bs + Cs^4 + 4Cs^3 + 4Cs^2 + Ds^4 + 2Ds^3 + Es^3 = 3$$

$$s^4 \quad C + D = 0 \quad \rightarrow D = -\frac{9}{16}$$

$$s^3 \quad B + 4C + 2D + E = 0 \quad \rightarrow E = -\frac{3}{8}$$

$$s^2 \quad A + 4B + 4C = 0 \quad \rightarrow C = \frac{9}{16}$$

$$s^1 \quad 4A + 4B = 0 \quad \rightarrow B = -\frac{3}{4}$$

$$s^0 \quad 4A = 3 \rightarrow A = \frac{3}{4}$$

$$\mathcal{L}^{-1}\{X(s)\} = \mathcal{L}^{-1}\left\{\frac{3}{4}\frac{1}{s^3} - \frac{3}{4}\frac{1}{s^2} + \frac{9}{16}\frac{1}{s} - \frac{9}{16}\frac{1}{s+2} - \frac{3}{8}\frac{1}{(s+2)^2}\right\}$$

$$x(t) = \frac{3}{8}t^2 - \frac{3}{4}t + \frac{9}{16} - \frac{9}{16}e^{-2t} - \frac{3}{8}te^{-2t}$$

Exercise

Solve using the Laplace transform: $y'' + 4y = 4t^2 - 4t + 10$; $y(0) = 0$, $y'(0) = 3$

Solution

$$\mathcal{L}\{y'' + 4y\} = \mathcal{L}\{4t^2 - 4t + 10\}$$

$$s^2 Y(s) - sy(0) - y'(0) + 4Y(s) = \frac{8}{s^3} - \frac{4}{s^2} + \frac{10}{s} \quad y(0) = 0 \quad y'(0) = 3$$

$$(s^2 + 4)Y(s) = \frac{8 - 4s + 10s^2}{s^3} + 3$$

$$Y(s) = \frac{3s^3 + 10s^2 - 4s + 8}{s^3(s^2 + 4)} = \frac{A}{s} + \frac{B}{s^2} + \frac{C}{s^3} + \frac{Ds + E}{s^2 + 4}$$

$$s^4 \quad A + D = 0 \quad \underline{D = -2}$$

$$s^3 \quad B + E = 3 \quad \underline{E = 4}$$

$$s^2 \quad 4A + C = 10 \quad \underline{A = 2}$$

$$s^1 \quad 4B = -4 \quad \underline{B = -1}$$

$$s^0 \quad 4C = 8 \quad \underline{C = 2}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{2}{s} - \frac{1}{s^2} + \frac{2}{s^3} - \frac{2s}{s^2 + 2^2} + \frac{4}{s^2 + 2^2}\right\}$$

$$\underline{y(t) = 2 - t + t^2 - 2\cos 2t + 2\sin 2t}$$

Exercise

Solve using the Laplace transform: $y'' - 4y = 4t - 8e^{-2t}$; $y(0) = 0$, $y'(0) = 5$

Solution

$$\mathcal{L}\{y'' - 4y\} = \mathcal{L}\{4t - 8e^{-2t}\}$$

$$s^2Y(s) - sy(0) - y'(0) - 4Y(s) = \frac{4}{s^2} - \frac{8}{s+2} \quad y(0) = 0 \quad y'(0) = 5$$

$$(s^2 - 4)Y(s) = \frac{-8s^2 + 4s + 8}{s^2(s+2)} + 5$$

$$Y(s) = \frac{5s^3 + 2s^2 + 4s + 8}{s^2(s+2)^2(s-2)} = \frac{A}{s} + \frac{B}{s^2} + \frac{C}{s+2} + \frac{D}{(s+2)^2} + \frac{E}{s-2}$$

$$5s^3 + 2s^2 + 4s + 8 = A(s^2 + 2s)(s^2 - 4) + B(s+2)(s^2 - 4) + C(s^4 - 4s^2) + D(s^3 - 2s^2) + Es^2(s^2 + 4s + 4)$$

$$\begin{cases} s^4 & A + C + E = 0 & C + E = 0 \\ s^3 & 2A + B + D + 4E = 5 & D + 4E = 6 \\ s^2 & -4A + 2B - 4C - 2D + 4E = 2 & -4C - 2D + 4E = 4 \\ s^1 & -8A - 4B = 4 & \underline{A = 0} \\ s^0 & -8B = 8 & \underline{B = -1} \end{cases}$$

$$\underline{C = -1 \quad D = 2 \quad E = 1}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{-\frac{1}{s^2} - \frac{1}{s+2} + \frac{2}{(s+2)^2} + \frac{1}{s-2}\right\}$$

$$\underline{y(t) = -t - e^{-2t} + 2te^{-2t} + e^{2t}}$$

Exercise

Solve using the Laplace transform: $y'' + 4y' = \cos(t-3) + 4t$, $y(3) = 0$, $y'(3) = 7$

Solution

$$y''(t) + 4y'(t) = \cos(t-3) + 4t$$

$$\text{Let } v = t - 3 \rightarrow t = v + 3$$

$$x(v) = y(t) = y(v+3) \quad y(3) = x(0) = 0 \quad y'(3) = x'(0) = 7$$

$$x''(v) + 4x'(v) = \cos v + 4v + 12$$

$$\mathcal{L}\{x'' + 4x'\} = \mathcal{L}\{\cos v + 4v + 12\}$$

$$s^2 X(s) - sx(0) - x'(0) + 4sX(s) - 4x(0) = \frac{s}{s^2+1} + \frac{4}{s^2} + \frac{12}{s}$$

$$(s^2 + 4s)X(s) = \frac{s}{s^2+1} + \frac{4+12s}{s^2} + 7$$

$$s(s+4)X(s) = \frac{s}{s^2+1} + \frac{7s^2+12s+4}{s^2}$$

$$X(s) = \frac{1}{(s+4)(s^2+1)} + \frac{7s^2+12s+4}{s^3(s+4)}$$

$$\frac{1}{(s+4)(s^2+1)} = \frac{A_1}{s+4} + \frac{A_2 s + A_3}{s^2+1}$$

$$1 = A_1 s^2 + A_1 + A_2 s^2 + 4A_2 s + A_3 s + 4A_3$$

$$\begin{cases} s^2 & A_1 + A_2 = 0 \quad \rightarrow A_1 = -A_2 & \underline{A_1 = \frac{1}{17}} \\ s & 4A_2 + A_3 = 0 \quad \rightarrow A_3 = -4A_2 & \underline{A_3 = \frac{4}{17}} \\ s^0 & A_1 + 4A_3 = 1 \Rightarrow -A_2 - 16A_2 = 1 \rightarrow A_2 = -\frac{1}{17} \end{cases}$$

$$\frac{1}{(s+4)(s^2+1)} = \frac{1}{17} \frac{1}{s+4} + \frac{1}{17} \frac{-s+4}{s^2+1}$$

$$\frac{7s^2+12s+4}{s^3(s+4)} = \frac{B_1}{s} + \frac{B_2}{s^2} + \frac{B_3}{s^3} + \frac{B_4}{s+4}$$

$$7s^2 + 12s + 4 = B_1 s^3 + 4B_1 s^2 + B_2 s^2 + 4B_2 s + B_3 s + 4B_3 + B_4 s^3$$

$$\begin{cases} s^3 & B_1 + B_4 = 0 & \rightarrow B_4 = -\frac{17}{16} \\ s^2 & 4B_1 + B_2 = 7 & \rightarrow B_1 = \frac{17}{16} \\ s & 4B_2 + B_3 = 12 & \rightarrow B_2 = \frac{11}{4} \\ s^0 & 4B_3 = 4 & \rightarrow B_3 = 1 \end{cases}$$

$$\frac{7s^2 + 12s + 4}{s^3(s+4)} = \frac{17}{16} \frac{1}{s} + \frac{11}{4} \frac{1}{s^2} + \frac{1}{s^3} - \frac{17}{16} \frac{1}{s+4}$$

$$\begin{aligned} X(s) &= \frac{1}{17} \frac{1}{s+4} + \frac{1}{17} \frac{-s+4}{s^2+1} + \frac{17}{16} \frac{1}{s} + \frac{11}{4} \frac{1}{s^2} + \frac{1}{s^3} - \frac{17}{16} \frac{1}{s+4} \\ &= \frac{17}{16} \frac{1}{s} + \frac{11}{4} \frac{1}{s^2} + \frac{1}{s^3} - \frac{273}{272} \frac{1}{s+4} - \frac{1}{17} \frac{s}{s^2+1} + \frac{4}{17} \frac{1}{s^2+1} \end{aligned}$$

$$\mathcal{L}^{-1}\{X(s)\} = \mathcal{L}^{-1}\left\{\frac{17}{16} \frac{1}{s} + \frac{11}{4} \frac{1}{s^2} + \frac{1}{s^3} - \frac{273}{272} \frac{1}{s+4} - \frac{1}{17} \frac{s}{s^2+1} + \frac{4}{17} \frac{1}{s^2+1}\right\}$$

$$x(v) = \frac{17}{16} + \frac{11}{4}v + \frac{1}{2}v^2 - \frac{273}{272}e^{-4v} - \frac{1}{17}\cos v + \frac{4}{17}\sin v \quad v = t - 3$$

$$\begin{aligned} y(t) &= \frac{17}{16} + \frac{11}{4}(t-3) + \frac{1}{2}(t-3)^2 - \frac{273}{272}e^{-4(t-3)} - \frac{1}{17}\cos(t-3) + \frac{4}{17}\sin(t-3) \\ &= \frac{17}{16} + \frac{11}{4}t - \frac{33}{4} + \frac{1}{2}t^2 - 3t + \frac{9}{2} - \frac{273}{272}e^{-4(t-3)} + \frac{1}{17}(4\sin(t-3) - \cos(t-3)) \\ &= \frac{43}{16} - \frac{1}{4}t + \frac{1}{2}t^2 - \frac{273}{272}e^{-4(t-3)} + \frac{1}{17}(4\sin(t-3) - \cos(t-3)) \end{aligned}$$

Exercise

Solve using the Laplace transform: $y'' + 4y' + 8y = \sin t$, $y(0) = 1$, $y'(0) = 0$

Solution

$$\mathcal{L}\{y'' + 4y' + 8y\}(s) = \mathcal{L}\{\sin t\}(s)$$

$$s^2Y(s) - sy(0) - y'(0) + 4sY(s) - 4y(0) + 8Y(s) = \frac{1}{s^2+1} \quad y(0) = 1, \quad y'(0) = 0$$

$$(s^2 + 4s + 8)Y(s) - s - 4 = \frac{1}{s^2+1}$$

$$(s^2 + 4s + 8)Y(s) = \frac{1}{s^2+1} + s + 4$$

$$Y(s) = \frac{s^3 + 4s^2 + s + 5}{(s^2+1)(s^2+4s+8)} = \frac{As+B}{s^2+1} + \frac{Cs+D}{s^2+4s+8}$$

$$s^3 + 4s^2 + s + 5 = As^3 + 4As^2 + 8As + Bs^2 + 4Bs + 8B + Cs^3 + Cs + Ds^2 + D$$

$$\begin{cases} s^3 & A + C = 1 \\ s^2 & 4A + B + D = 4 \\ s & 8A + 4B + C = 1 \\ s^0 & 8B + D = 5 \end{cases} \Rightarrow \begin{matrix} A = -\frac{4}{65} & B = \frac{7}{65} \\ C = \frac{69}{65} & D = \frac{269}{65} \end{matrix}$$

$$Y(s) = \frac{1}{65} \left(-4 \frac{s}{s^2+1} + \frac{7}{s^2+1} + \frac{69(s+2) - 138 + 269}{(s+2)^2+4} \right)$$

$$= \frac{1}{65} \left(-4 \frac{s}{s^2+1} + \frac{7}{s^2+1} + \frac{69s}{(s+2)^2+4} + \frac{2}{2} \frac{131}{(s+2)^2+4} \right)$$

$$\mathcal{L}^{-1}\{Y(s)\} = \frac{1}{65} \mathcal{L}^{-1} \left\{ -4 \frac{s}{s^2+1} + \frac{7}{s^2+1} + \frac{69s}{(s+2)^2+4} + \frac{131}{2} \frac{2}{(s+2)^2+4} \right\}$$

$$y(t) = -\frac{4}{65} \cos t + \frac{7}{65} \sin t + \frac{69}{65} e^{-2t} \cos 2t + \frac{131}{130} e^{-2t} \sin 2t$$

Exercise

Solve using the Laplace transform: $y'' + 5y' - y = e^t - 1$; $y(0)=1$, $y'(0)=1$

Solution

$$\mathcal{L}\{y'' + 5y' - y\} = \mathcal{L}\{e^t - 1\}$$

$$s^2 Y(s) - sy(0) - y'(0) + 5sY(s) - 5y(0) - Y(s) = \frac{1}{s-1} - \frac{1}{s} \quad y(0)=1 \quad y'(0)=1$$

$$(s^2 + 5s - 1)Y(s) = \frac{1}{s(s-1)} + s + 6$$

$$Y(s) = \frac{s^3 + 5s^2 - 6s + 1}{s(s-1)\left(s + \frac{5}{2} - \frac{\sqrt{29}}{2}\right)\left(s + \frac{5}{2} + \frac{\sqrt{29}}{2}\right)}$$

$$= \frac{A}{s} + \frac{B}{s-1} + \frac{C}{s + \frac{5}{2} - \frac{\sqrt{29}}{2}} + \frac{D}{s + \frac{5}{2} + \frac{\sqrt{29}}{2}}$$

$$\begin{cases} s^3 & A + C + D = 1 \\ s^2 & 4A + B + \left(\frac{3}{2} + \frac{\sqrt{29}}{2}\right)C + \left(\frac{3}{2} - \frac{\sqrt{29}}{2}\right)D = 5 \\ s^1 & -6A + 5B - \left(\frac{5}{2} + \frac{\sqrt{29}}{2}\right)C - \left(\frac{5}{2} - \frac{\sqrt{29}}{2}\right)D = -6 \\ s^0 & A - B = 1 \end{cases}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{1}{s} + \frac{1}{5} \frac{1}{s-1} + \left(-\frac{1}{10} + \frac{3}{10\sqrt{29}}\right) \frac{1}{s + \frac{5}{2} - \frac{\sqrt{29}}{2}} - \left(\frac{1}{10} + \frac{3}{10\sqrt{29}}\right) \frac{1}{s + \frac{5}{2} + \frac{\sqrt{29}}{2}}\right\}$$

$$\underline{y(t) = 1 + \frac{1}{5}e^t + \left(-\frac{1}{10} + \frac{3\sqrt{29}}{290}\right)e^{\frac{-5+\sqrt{29}}{2}t} - \left(\frac{1}{10} + \frac{3\sqrt{29}}{290}\right)e^{\frac{-5-\sqrt{29}}{2}t}}$$

Exercise

Solve using the Laplace transform: $y'' + 5y' - 6y = 21e^{t-1}$ $y(1) = -1$, $y'(1) = 9$

Solution

Let: $w(t) = y(t+1) \leftrightarrow y(t) = w(t-1)$

$$\mathcal{L}\{w'' + 5w' - 6w\} = \mathcal{L}\{21e^t\}$$

$$s^2W(s) - sw(0) - w'(0) + 5sW(s) - 5w(0) - 6W(s) = 21\frac{1}{s-1} \quad y(1) = w(0) = -1, \quad y'(1) = w'(0) = 9$$

$$(s^2 + 5s - 6)W(s) = \frac{21}{s-1} - s + 4$$

$$W(s) = \frac{-s^2 + 5s + 17}{(s+6)(s-1)^2} = \frac{A}{s+6} + \frac{B}{s-1} + \frac{C}{(s-1)^2}$$

$$s^2 \quad A + B = -1$$

$$s^1 \quad -2A + 5B + C = 5$$

$$s^0 \quad A - 6B + 6C = 17$$

$$\Delta = \begin{vmatrix} 1 & 1 & 0 \\ -2 & 5 & 1 \\ 1 & -6 & 6 \end{vmatrix} = 49 \quad \Delta_A = \begin{vmatrix} -1 & 1 & 0 \\ 5 & 5 & 1 \\ 17 & -6 & 6 \end{vmatrix} = -49 \quad \Delta_B = \begin{vmatrix} 1 & -1 & 0 \\ -2 & 5 & 1 \\ 1 & 17 & 6 \end{vmatrix} = 0 \quad \Delta_C = \begin{vmatrix} 1 & 1 & -1 \\ -2 & 5 & 5 \\ 1 & -6 & 17 \end{vmatrix} = 147$$

$$\underline{A = -1, \quad B = 0, \quad C = 3}$$

$$\mathcal{L}^{-1}\{W(s)\} = \mathcal{L}^{-1}\left\{\frac{-1}{s+6} + \frac{3}{(s-1)^2}\right\}$$

$$w(t) = -e^{-6t} + 3te^t$$

$$\underline{y(t) = w(t-1) = -e^{-6(t-1)} + 3(t-1)e^{(t-1)}}$$

Exercise

Solve using the Laplace transform: $y'' + 5y' + 4y = 0$; $y(0) = 1$, $y'(0) = 0$

Solution

$$\mathcal{L}\{y'' + 5y' + 4y\} = 0$$

$$s^2 Y(s) - sy(0) - y'(0) + 5sY(s) - 5y(0) + 4Y(s) = 0 \quad y(0) = 1; \quad y'(0) = 0$$

$$(s^2 + 5s + 4)Y(s) - s - 5 = 0$$

$$Y(s) = \frac{s+5}{s^2 + 5s + 4} = \frac{A}{s+1} + \frac{B}{s+4}$$

$$As + 4A + Bs + B = s + 5$$

$$\begin{cases} A + B = 1 \\ 4A + B = 5 \end{cases} \rightarrow \underline{A = \frac{4}{3}; B = -\frac{1}{3}}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{4}{3} \frac{1}{s+1} - \frac{1}{3} \frac{1}{s+4}\right\}$$

$$\underline{y(t) = \frac{4}{3}e^{-t} - \frac{1}{3}e^{-4t}}$$

Exercise

Solve using the Laplace transform: $y'' + 6y = t^2 - 1$; $y(0) = 0$, $y'(0) = -1$

Solution

$$\mathcal{L}\{y'' + 6y\} = \mathcal{L}\{t^2 - 1\}$$

$$s^2 Y(s) - sy(0) - y'(0) + 6Y(s) = \frac{2}{s^3} - \frac{1}{s} \quad y(0) = 0 \quad y'(0) = -1$$

$$(s^2 + 6)Y(s) = \frac{2-s^2}{s^3} - 1$$

$$Y(s) = \frac{2-s^2-s^3}{s^3(s^2+6)} = \frac{A}{s} + \frac{B}{s^2} + \frac{C}{s^3} + \frac{Ds+E}{s^2+6}$$

$$s^4 \quad A + D = 0 \quad \underline{D = \frac{2}{9}}$$

$$s^3 \quad B + E = -1 \quad \underline{E = -1}$$

$$s^2 \quad 6A + C = -1 \quad \underline{A = -\frac{2}{9}}$$

$$s \quad \underline{B = 0}$$

$$s^0 \quad 6C = 2 \quad \underline{C = \frac{1}{3}}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{-\frac{2}{9}\frac{1}{s} + \frac{1}{3}\frac{1}{s^3} + \frac{1}{9}\frac{s}{s^2+6} - \frac{1}{s^2+6}\right\}$$

$$\underline{y(t) = -\frac{2}{9} + \frac{1}{6}t^2 + \frac{1}{9}\cos\sqrt{6}t - \frac{1}{\sqrt{6}}\sin\sqrt{6}t}$$

Exercise

Solve using the Laplace transform: $y'' - 6y' + 9y = t$; $y(0) = 0$, $y'(0) = 1$

Solution

$$\mathcal{L}\{y'' - 6y' + 9y\} = \mathcal{L}\{t\}$$

$$s^2Y(s) - sy(0) - y'(0) - 6sY(s) + 6y(0) + 9Y(s) = \frac{1}{s^2} \quad y(0) = 0 \quad y'(0) = 1$$

$$(s^2 - 6s + 9)Y(s) = \frac{1}{s^2} + 1$$

$$Y(s) = \frac{s^2 + 1}{s^2(s-3)^2} = \frac{A}{s} + \frac{B}{s^2} + \frac{C}{s-3} + \frac{D}{(s-3)^2}$$

$$\begin{cases} s^3 & A + C = 0 & C = -\frac{2}{27} \\ s^2 & -6A + B - 3C + D = 1 & D = \frac{10}{9} \\ s & 9A - 6B = 0 & A = \frac{2}{27} \\ s^0 & 9B = 1 & B = \frac{1}{9} \end{cases}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{2}{27}\frac{1}{s} + \frac{1}{9}\frac{1}{s^2} - \frac{2}{27}\frac{1}{s-3} + \frac{10}{9}\frac{1}{(s-3)^2}\right\}$$

$$\underline{y(t) = \frac{2}{27} + \frac{1}{9}t - \frac{2}{27}e^{3t} + \frac{10}{9}te^{3t}}$$

Exercise

Solve using the Laplace transform: $y'' - 6y' + 15y = 2\sin 3t$, $y(0) = -1$, $y'(0) = -4$

Solution

$$\mathcal{L}\{y'' - 6y' + 15y\}(s) = \mathcal{L}\{2\sin 3t\}(s)$$

$$s^2Y(s) - sy(0) - y'(0) - 6sY(s) + 6y(0) + 15Y(s) = \frac{6}{s^2 + 9} \quad y(0) = -1, \quad y'(0) = -4$$

$$(s^2 - 6s + 15)Y(s) + s + 4 - 6 = \frac{6}{s^2 + 9}$$

$$(s^2 - 6s + 15)Y(s) = \frac{6}{s^2 + 9} - s + 2$$

$$Y(s) = \frac{-s^3 + 2s^2 - 9s + 24}{(s^2 + 9)(s^2 - 6s + 15)} = \frac{As + B}{s^2 + 9} + \frac{Cs + D}{s^2 - 6s + 15}$$

$$-s^3 + 2s^2 - 9s + 24 = As^3 - 6As^2 + 15As + Bs^2 - 6Bs + 15B + Cs^3 + 9Cs + Ds^2 + 9D$$

$$\begin{cases} s^3 & A + C = -1 \\ s^2 & -6A + B + D = 2 \\ s & 15A - 6B + 9C = -9 \\ s^0 & 15B + 9D = 24 \end{cases} \rightarrow \begin{cases} A = \frac{1}{10} & B = \frac{1}{10} \\ C = -\frac{11}{10} & D = \frac{5}{2} \end{cases}$$

$$Y(s) = \frac{1}{10} \frac{s}{s^2 + 9} + \frac{1}{10} \frac{1}{s^2 + 9} + \frac{1}{10} \frac{-11(s-3) - 33 + 25}{(s-3)^2 - 9 + 15}$$

$$= \frac{1}{10} \frac{s}{s^2 + 9} + \frac{1}{10} \frac{1}{s^2 + 9} \frac{3}{3} - \frac{11}{10} \frac{s-3}{(s-3)^2 + 6} - \frac{1}{10} \frac{8}{(s-3)^2 + 6} \frac{\sqrt{6}}{\sqrt{6}}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{1}{10} \frac{s}{s^2 + 9} + \frac{1}{30} \frac{3}{s^2 + 9} - \frac{11}{10} \frac{s-3}{(s-3)^2 + 6} - \frac{8}{10\sqrt{6}} \frac{\sqrt{6}}{(s-3)^2 + 6}\right\}$$

$$y(t) = \frac{1}{10} \cos 3t + \frac{1}{30} \sin 3t - \frac{11}{10} e^{3t} \cos \sqrt{6}t - \frac{8}{10\sqrt{6}} e^{3t} \sin \sqrt{6}t$$

Exercise

Solve using the Laplace transform: $y'' - 6y' + 13y = 0$; $y(0) = 0$, $y'(0) = -3$

Solution

$$\mathcal{L}\{y'' - 6y' + 13y\} = 0$$

$$s^2 Y(s) - sy(0) - y'(0) - 6sY(s) + 6y(0) + 13Y(s) = 0 \quad y(0) = 0 \quad y'(0) = -3$$

$$(s^2 - 6s + 13)Y(s) = -3$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{-3}{(s-3)^2 + 4}\right\}$$

$$y(t) = -\frac{3}{2} e^{3t} \sin 2t$$

Exercise

Solve using the Laplace transform: $y'' + 6y' + 9y = 0$, $y(0) = -1$, $y'(0) = 6$

Solution

$$\mathcal{L}\{y'' + 6y' + 9y\} = 0$$

$$s^2Y(s) - sy(0) - y'(0) + 6Y(s) - 6y(0) + 9Y(s) = 0 \quad y(0) = -1 \quad y'(0) = 6$$

$$(s^2 + 6s + 9)Y(s) = -s$$

$$Y(s) = -\frac{s}{(s+3)^2} = \frac{A}{s+3} + \frac{B}{(s+3)^2}$$

$$\begin{cases} s^1 & A = -1 \\ s^0 & 3A + B = 0 \quad B = 3 \end{cases}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{-1}{s+3} + \frac{3}{(s+3)^2}\right\}$$

$$\underline{y(t) = -e^{-3t} + 3te^{-3t}}$$

Exercise

Solve using the Laplace transform: $y'' + 6y' + 5y = 12e^t$, $y(0) = -1$, $y'(0) = 7$

Solution

$$\mathcal{L}\{y'' + 6y' + 5y\} = \mathcal{L}\{12e^t\}$$

$$s^2Y(s) - sy(0) - y'(0) + 6Y(s) - 6y(0) + 5Y(s) = \frac{12}{s-1} \quad y(0) = -1 \quad y'(0) = 7$$

$$(s^2 + 6s + 5)Y(s) = \frac{12}{s-1} - s + 1$$

$$Y(s) = \frac{-s^2 + 2s + 11}{(s+1)(s+5)(s-1)} = \frac{A}{s+1} + \frac{B}{s+5} + \frac{C}{s-1}$$

$$\begin{cases} s^2 & A + B + C = -1 \\ s^1 & 4A + 6C = 2 \\ s^0 & -5A - B + 5C = 11 \end{cases}$$

$$\Delta = \begin{vmatrix} 1 & 1 & 1 \\ 4 & 0 & 6 \\ -5 & -1 & 5 \end{vmatrix} = -48 \quad \Delta_A = \begin{vmatrix} -1 & 1 & 1 \\ 2 & 0 & 6 \\ 11 & -1 & 5 \end{vmatrix} = 48 \quad \Delta_B = \begin{vmatrix} 1 & -1 & 1 \\ 4 & 2 & 6 \\ -5 & 11 & 5 \end{vmatrix} = 48$$

$$\underline{A = -1 \quad B = -1 \quad C = 1}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{-1}{s+1} - \frac{1}{s+5} + \frac{1}{s-1}\right\}$$

$$\underline{y(t) = -e^{-t} - e^{-5t} + e^t}$$

Exercise

Solve using the Laplace transform: $y'' - 7y' + 10y = 9\cos t + 7\sin t$; $y(0) = 5$, $y'(0) = -4$

Solution

$$\mathcal{L}\{y'' - 7y' + 10y\} = \mathcal{L}\{9\cos t + 7\sin t\}$$

$$s^2Y(s) - sy(0) - y'(0) - 7sY(s) + 7y(0) + 10Y(s) = \frac{9s}{s^2+1} + \frac{7}{s^2+1} \quad y(0) = 5, \quad y'(0) = -4$$

$$(s^2 - 7s + 10)Y(s) = \frac{9s+7}{s^2+1} + 5s - 39$$

$$Y(s) = \frac{5s^3 - 39s^2 + 14s - 32}{(s-2)(s-5)(s^2+1)} = \frac{A}{s-2} + \frac{B}{s-5} + \frac{Cs+D}{s^2+1}$$

$$\begin{cases} s^3 & A+B+C=5 \\ s^2 & -5A-2B-7C+D=-39 \\ s & A+B+10C-7D=14 \\ s^0 & -5A-2B+10D=-32 \end{cases} \rightarrow \underline{A=8, B=-4, C=1, D=0}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{8}{s-2} - \frac{4}{s-5} + \frac{s}{s^2+1}\right\}$$

$$\underline{y(t) = 8e^{2t} - 4e^{5t} + \cos t}$$

Exercise

Solve using the Laplace transform: $y'' + 8y' + 25y = 0$, $y(\pi) = 0$, $y'(\pi) = 6$

Solution

$$\text{Let: } w(t) = y(t + \pi) \leftrightarrow y(t) = w(t - \pi)$$

$$y''(t) + 8y'(t) + 25y(t) = 0$$

$$\mathcal{L}\{w'' + 8w' + 25w\} = 0$$

$$s^2W(s) - sw(0) - w'(0) + 8sW(s) - 8w(0) + 25W(s) = 0 \quad y(\pi) = w(0) = 0, \quad y'(\pi) = w'(0) = 6$$

$$(s^2 + 8s + 25)W(s) - 6 = 0$$

$$W(s) = \frac{6}{(s+4)^2 - 16 + 25}$$

$$\mathcal{L}^{-1}\{W(s)\} = \mathcal{L}^{-1}\left\{\frac{2(3)}{(s+4)^2 + 9}\right\}$$

$$w(t) = 2e^{-4t} \sin 3t$$

$$y(t) = w(t - \pi)$$

$$y(t) = 2e^{-4(t-\pi)} \sin 3(t - \pi)$$

$$\sin(3t - 3\pi) = \sin 3t \cos 3\pi - \cos 3t \sin 3\pi = \sin 3t(-1) - 0 = -\sin 3t$$

$$= -2e^{-4(t-\pi)} \sin 3t$$

Exercise

Solve using the Laplace transform: $y'' + 9y = 2 \sin 2t$; $y(0) = 0$, $y'(0) = -1$

Solution

$$\mathcal{L}(y'' + 9y) = \mathcal{L}(2 \sin 2t)$$

$$s^2 Y(s) - sy(0) - y'(0) + 9Y(s) = 2 \frac{2}{s^2 + 2^2} \quad y(0) = 0 \quad y'(0) = -1$$

$$(s^2 + 9)Y(s) + 1 = \frac{4}{s^2 + 4}$$

$$Y(s) = \frac{4}{(s^2 + 9)(s^2 + 4)} - \frac{1}{s^2 + 9}$$

$$\frac{4}{(s^2 + 9)(s^2 + 4)} = \frac{A}{s^2 + 9} + \frac{B}{s^2 + 4}$$

$$4 = (A + B)s^2 + 4A + 9B$$

$$\begin{cases} A + B = 0 \\ 4A + 9B = 4 \end{cases} \Rightarrow A = -\frac{4}{5} \quad B = \frac{4}{5}$$

$$Y(s) = -\frac{4}{5} \frac{1}{s^2 + 9} + \frac{4}{5} \frac{1}{s^2 + 4} - \frac{1}{s^2 + 9}$$

$$= \frac{4}{5} \frac{1}{s^2 + 4} - \frac{9}{5} \frac{1}{s^2 + 9}$$

$$= \frac{4}{5} \frac{1}{2} \frac{2}{s^2 + 4} - \frac{3}{5} \frac{3}{s^2 + 9}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{2}{5} \frac{2}{s^2 + 4} - \frac{3}{5} \frac{3}{s^2 + 9}\right\}$$

$$y(t) = \frac{2}{5} \sin 2t - \frac{3}{5} \sin 3t$$

Exercise

Solve using the Laplace transform: $y'' + 9y = 3\sin 2t$; $y(0) = 0$, $y'(0) = -1$

Solution

$$\mathcal{L}(y'' + 9y)(s) = \mathcal{L}(3\sin 2t)(s)$$

$$s^2 Y(s) - sy(0) - y'(0) + 9Y(s) = 3 \frac{1}{s^2 + 4} \quad y(0) = 0 \quad y'(0) = -1$$

$$s^2 Y(s) + 1 + Y(s) = \frac{3}{s^2 + 4}$$

$$(s^2 + 1)Y(s) = \frac{3}{s^2 + 4} - 1$$

$$(s^2 + 1)Y(s) = \frac{-s^2 - 1}{s^2 + 4}$$

$$Y(s) = \frac{-s^2 - 1}{(s^2 + 4)(s^2 + 1)} = \frac{A}{s^2 + 4} + \frac{B}{s^2 + 1}$$

$$As^2 + A + Bs^2 + 4B = -s^2 - 1$$

$$\begin{cases} s^2 \\ s^0 \end{cases} \begin{cases} A + B = -1 \\ A + 4B = 1 \end{cases} \Rightarrow \begin{cases} A = \frac{2}{3} \\ B = -\frac{5}{3} \end{cases}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{2}{3} \frac{1}{s^2 + 4} - \frac{5}{3} \frac{1}{s^2 + 1}\right\}$$

$$\underline{y(t) = \frac{2}{3} \sin 2t - \frac{5}{3} \sin t}$$

Exercise

Solve using the Laplace transform: $y'' + 16y = 2\sin 4t$; $y(0) = -\frac{1}{2}$, $y'(0) = 0$

Solution

$$\mathcal{L}\{y'' + 16y\}(s) = \mathcal{L}\{2\sin 4t\}(s)$$

$$s^2 Y(s) - sy(0) - y'(0) + 16Y(s) = \frac{8}{s^2 + 16}$$

$$(s^2 + 16)Y(s) + \frac{s}{2} = \frac{8}{s^2 + 16}$$

$$(s^2 + 16)Y(s) = \frac{8}{s^2 + 16} - \frac{s}{2}$$

$$Y(s) = \frac{8}{(s^2 + 16)^2} - \frac{1}{2} \frac{s}{s^2 + 16}$$

$$\mathcal{L}^{-1} \frac{2a^3}{(s^2 + a^2)^2} = \sin(at) - at \cos(at)$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{8}{128} \frac{128}{(s^2 + 4^2)^2} - \frac{1}{2} \frac{s}{s^2 + 16}\right\}$$

$$\underline{y(t) = \frac{1}{16}(\sin 4t - 4t \cos 4t) - \frac{1}{2} \cos 4t}$$

Exercise

Solve using the Laplace transform: $y'' - 10y' + 9y = 5t$; $y(0) = -1$, $y'(0) = 2$

Solution

$$\mathcal{L}\{y'' - 10y' + 9y\}(s) = \mathcal{L}\{5t\}(s)$$

$$s^2 Y(s) - sy(0) - y'(0) - 10sY(s) + 10y(0) + 9Y(s) = \frac{5}{s^2}$$

$$(s^2 - 10s + 9)Y(s) = \frac{5}{s^2} - s + 12$$

$$Y(s) = \frac{-s^3 + 12s^2 + 5}{s^2(s-9)(s-1)} = \frac{A}{s} + \frac{B}{s^2} + \frac{C}{s-9} + \frac{D}{s-1}$$

$$-s^3 + 12s^2 + 5 = As^3 - 10As^2 + 9As + Bs^2 - 10Bs + 9B + Cs^3 - Cs^2 + Ds^3 - 9Ds^2$$

$$\begin{cases} s^3 & A + C + D = -1 \\ s^2 & -10A + B - C - 9D = -12 \\ s^1 & 9A - 10B = 0 \rightarrow A = \frac{50}{81} \\ s^0 & 9B = 5 \rightarrow B = \frac{5}{9} \end{cases}$$

$$\begin{cases} C + D = -\frac{131}{81} \\ C - 9D = -\frac{517}{81} \end{cases} \rightarrow \begin{cases} C = \frac{31}{81} \\ D = -2 \end{cases}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{50}{81} \frac{1}{s} + \frac{5}{9} \frac{1}{s^2} + \frac{31}{81} \frac{1}{s-9} - \frac{2}{s-1}\right\}$$

$$\underline{y(t) = \frac{50}{81} + \frac{5}{9}t + \frac{31}{81}e^{9t} - 2e^t}$$

Exercise

Solve using the Laplace transform: $2y'' + 3y' - 2y = te^{-2t}$, $y(0) = 0$, $y'(0) = -2$

Solution

$$\mathcal{L}\{2y'' + 3y' - 2y\}(s) = \mathcal{L}\{te^{-2t}\}(s)$$

$$2s^2Y(s) - 2sy(0) - 2y'(0) + 3sY(s) - 3y(0) - 2Y(s) = \frac{1}{(s+2)^2}$$

$$(2s^2 + 3s - 2)Y(s) + 4 = \frac{1}{(s+2)^2}$$

$$(2s-1)(s+2)Y(s) = \frac{1}{(s+2)^2} - 4$$

$$Y(s) = \frac{-4s^2 - 16s - 15}{(2s-1)(s+2)^3} = \frac{A}{2s-1} + \frac{B}{s+2} + \frac{C}{(s+2)^2} + \frac{D}{(s+2)^3}$$

$$-4s^2 - 16s - 15 = As^3 + 6As^2 + 12As + 8A + (2Bs - B)(s^2 + 4s + 4) + 2Cs^2 + 2Cs - 2C + 2Ds - D$$

$$\begin{cases} s^3 & A + 2B = 0 \\ s^2 & 6A + 7B + 2C = -4 \\ s^1 & 12A + 4B + 3C + 2D = -16 \\ s^0 & 8A - 4B - 2C - D = -15 \end{cases}$$

$$\underline{A = -\frac{192}{125} \quad B = \frac{96}{125} \quad C = -\frac{2}{25} \quad D = -\frac{1}{5}}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{-\frac{192}{125} \frac{1}{2\left(s - \frac{1}{2}\right)} + \frac{96}{125} \frac{1}{s+2} - \frac{2}{25} \frac{1}{(s+2)^2} - \frac{1}{5} \frac{1}{(s+2)^3}\right\}$$

$$\underline{y(t) = -\frac{96}{125}e^{t/2} + \frac{96}{125}e^{-2t} - \frac{2}{25}te^{-2t} - \frac{1}{5}t^2e^{-2t}}$$

Exercise

Solve using the Laplace transform: $2y'' + 20y' + 51y = 0$, $y(0) = 2$, $y'(0) = 0$

Solution

$$\mathcal{L}\{2y'' + 20y' + 51y\} = 0$$

$$2s^2Y(s) - 2sy(0) - 2y'(0) + 20sY(s) - 20y(0) + 51Y(s) = 0 \quad y(0) = 2 \quad y'(0) = 0$$

$$(2s^2 + 20s + 51)Y(s) = 4s + 40$$

$$\begin{aligned}
Y(s) &= \frac{4s+40}{2\left(s^2+10s+\frac{51}{2}\right)} \\
&= \frac{2s+20}{(s+5)^2+\frac{1}{2}} \\
&= \frac{2s}{(s+5)^2+\left(\frac{1}{\sqrt{2}}\right)^2} + \frac{20}{\sqrt{2}} \frac{\frac{1}{\sqrt{2}}}{(s+5)^2+\left(\frac{1}{\sqrt{2}}\right)^2} \\
\mathcal{L}^{-1}\{Y(s)\} &= \mathcal{L}^{-1}\left\{\frac{2s}{(s+5)^2+\left(\frac{\sqrt{2}}{2}\right)^2} + 10\sqrt{2} \frac{\frac{1}{\sqrt{2}}}{(s+5)^2+\left(\frac{\sqrt{2}}{2}\right)^2}\right\} \\
y(t) &= \underline{2e^{-5t} \cos \frac{\sqrt{2}t}{2} + 10\sqrt{2}e^{-5t} \sin \frac{\sqrt{2}t}{2}}
\end{aligned}$$

Exercise

Solve using the Laplace transform: $y''' + y' = e^t$, $y(0) = y'(0) = y''(0) = 0$

Solution

$$\begin{aligned}
\mathcal{L}\{y''' + y'\} &= \mathcal{L}\{e^t\} \\
s^3Y(s) - s^2y(0) - sy'(0) - y''(0) + sY(s) - y(0) &= \frac{1}{s-1} \\
(s^3 + s)Y(s) &= \frac{1}{s-1} \\
Y(s) &= \frac{1}{s(s-1)(s^2+1)} = \frac{A}{s} + \frac{B}{s-1} + \frac{Cs+D}{s^2+1} \\
1 &= As^3 - As^2 + As - A + Bs^3 + Bs + Cs^3 - Cs^2 +Ds^2 -Ds \\
&\Rightarrow \begin{cases} s^3 & A+B+C=0 & B+C=1 \\ s^2 & -A-C+D=0 & -C+D=-1 \\ s & A+B-D=0 & B-D=1 \\ s^0 & -A=1 & \rightarrow \underline{A=-1} \end{cases} \Rightarrow B=\frac{1}{2} \quad C=\frac{1}{2} \quad D=-\frac{1}{2} \\
\mathcal{L}^{-1}\{Y(s)\} &= \mathcal{L}^{-1}\left\{-\frac{1}{s} + \frac{1}{2} \frac{B}{s-1} + \frac{1}{2} \frac{s}{s^2+1} - \frac{1}{2} \frac{1}{s^2+1}\right\} \\
y(t) &= \underline{-1 + \frac{1}{2}e^t + \frac{1}{2}\cos t - \frac{1}{2}\sin t}
\end{aligned}$$

Exercise

Solve using the Laplace transform: $2y^{(3)} + 3y'' - 3y' - 2y = e^{-t}$; $y(0) = 0$, $y'(0) = 0$, $y''(0) = 1$

Solution

$$\mathcal{L}\{2y^{(3)} + 3y'' - 3y' - 2y\} = \mathcal{L}\{e^{-t}\}$$

$$2s^3Y(s) - 2s^2y(0) - 2sy'(0) - 2y''(0) + 3s^2Y(s) - 3sy(0) - 3y'(0) - 3sY(s) + 3y(0) - 2Y(s) = \frac{1}{s+1}$$

$$(2s^3 + 3s^2 - 3s - 2)Y(s) - 2 = \frac{1}{s+1} \quad y(0) = y'(0) = 0, \quad y''(0) = 1$$

$$(s-1)(2s+1)(s+2)Y(s) = \frac{1}{s+1} + 2$$

$$Y(s) = \frac{2s+3}{(s-1)(2s+1)(s+2)(s+1)} = \frac{A}{s-1} + \frac{B}{2s+1} + \frac{C}{s+2} + \frac{D}{s+1}$$

$$A(2s+1)(s^2+3s+2) + B(s+2)(s^2-1) + C(2s+1)(s^2-1) + D(2s+1)(s^2+s-2) = 2s+3$$

$$\begin{cases} s^3 & 2A + B + 2C + 2D = 0 \\ s^2 & 7A + 2B + C + 3D = 0 \\ s^1 & 7A - B - 2C - 3D = 2 \\ s^0 & 2A - 2B - C - 2D = 3 \end{cases} \Rightarrow A = \frac{5}{18} \quad B = -\frac{16}{9} \quad C = \frac{1}{9} \quad D = \frac{1}{2}$$

$$Y(s) = \frac{5}{18} \frac{1}{s-1} - \frac{16}{9} \frac{1}{2s+1} + \frac{1}{9} \frac{1}{s+2} + \frac{1}{2} \frac{1}{s+1}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{5}{18} \frac{1}{s-1} - \frac{8}{9} \frac{1}{s+\frac{1}{2}} + \frac{1}{9} \frac{1}{s+2} + \frac{1}{2} \frac{1}{s+1}\right\}$$

$$y(t) = \frac{5}{18}e^t - \frac{8}{9}e^{-t/2} + \frac{1}{9}e^{-2t} + \frac{1}{2}e^{-t}$$

Exercise

Solve using the Laplace transform: $y^{(3)} + 2y'' - y' - 2y = \sin 3t$; $y(0) = 0$, $y'(0) = 0$, $y''(0) = 1$

Solution

$$\mathcal{L}\{y^{(3)} + 2y'' - y' - 2y\} = \mathcal{L}\{\sin 3t\}$$

$$s^3Y(s) - s^2y(0) - sy'(0) - y''(0) + 2s^2Y(s) - 2sy(0) - 2y'(0) - sY(s) + y(0) - 2Y(s) = \frac{3}{s^2+9}$$

$$(s^3 + 2s^2 - s - 2)Y(s) - 1 = \frac{3}{s^2+9} \quad y(0) = y'(0) = 0, \quad y''(0) = 1$$

$$(s-1)(s+1)(s+2)Y(s) = \frac{3}{s^2+9} + 1$$

$$Y(s) = \frac{s^2 + 12}{(s-1)(s+1)(s+2)(s^2+9)} = \frac{A}{s-1} + \frac{B}{s+1} + \frac{C}{s+2} + \frac{Ds+E}{s^2+9}$$

$$A(s^2+3s+2)(s^2+9) + B(s^2+s-2)(s^2+9) + C(s^2-1)(s^2+9) + (Ds+E)(s^3+2s^2-s-2)$$

$$\begin{cases} s^4 & A+B+C+D=0 \\ s^3 & 3A+B+2D+E=0 \\ s^2 & 11A+7B+8C-D+2E=1 \\ s^1 & 27A+9B-2D-E=0 \\ s^0 & 18A-18B-9C-2E=12 \end{cases} \rightarrow \begin{matrix} A = \frac{13}{60} & B = -\frac{13}{20} & C = \frac{16}{39} \\ D = \frac{3}{130} & E = -\frac{3}{65} \end{matrix}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{13}{60}\frac{1}{s-1} - \frac{13}{20}\frac{1}{s+1} + \frac{16}{39}\frac{1}{s+2} + \frac{3}{130}\frac{s}{s^2+9} - \frac{1}{65}\frac{3}{s^2+9}\right\}$$

$$y(t) = \frac{13}{60}e^t - \frac{13}{20}e^{-t} + \frac{16}{39}e^{-2t} + \frac{3}{130}\cos 3t - \frac{1}{65}\sin 3t$$

Exercise

Solve using the Laplace transform: $y^{(3)} - y'' + y' - y = 0$; $y(0)=1$, $y'(0)=1$, $y''(0)=3$

Solution

$$s^3Y(s) - s^2y(0) - sy'(0) - y''(0) - s^2Y(s) + sy(0) + y'(0) + sY(s) - y(0) - Y(s) = 0$$

$$(s^3 - s^2 + s - 1)Y(s) - s^2 - s - 3 + s = 0 \quad y(0)=1 \quad y'(0)=1 \quad y''(0)=3$$

$$s^3 - s^2 + s - 1 = s^2(s-1) + (s-1)$$

$$Y(s) = \frac{s^2+3}{(s-1)(s^2+1)} = \frac{A}{s-1} + \frac{Bs+C}{s^2+1}$$

$$\begin{cases} s^2 & A+B=1 \\ s & -B+C=0 \\ s^0 & A-C=3 \end{cases} \quad \begin{matrix} A+C=1 \\ A-C=3 \end{matrix} \quad \underline{A=2, C=-1, B=-1}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{2}{s-1} - \frac{s}{s^2+1} - \frac{1}{s^2+1}\right\}$$

$$y(t) = 2e^t - \cos t - \sin t$$

Exercise

Solve using the Laplace transform: $y^{(3)} + 4y'' + y' - 6y = -12$; $y(0)=1, \quad y'(0)=4, \quad y''(0)=-2$

Solution

$$s^3 Y(s) - s^2 y(0) - sy'(0) - y''(0) + 4s^2 Y(s) - 4sy(0) - 4y'(0) + sY(s) - y(0) - 6Y(s) = -\frac{12}{s}$$

$$(s^3 + 4s^2 + s - 6)Y(s) = s^2 - 8s - 15 - \frac{12}{s}$$

$$s^3 + 4s^2 + s - 6 = (s-1)(s^2 + 5s + 6)$$

$$\begin{array}{c|cccc} 1 & 1 & 4 & 1 & -6 \\ & & 1 & 5 & 6 \\ \hline & 1 & 5 & 6 & 0 \end{array}$$

$$Y(s) = \frac{s^3 - 8s^2 - 15s - 12}{s(s-1)(s+2)(s+3)} = \frac{A}{s} + \frac{B}{s-1} + \frac{C}{s+2} + \frac{D}{s+3}$$

$$\begin{cases} s^3 & A + B + C + D = 1 \\ s^2 & 4A + 5B + 2C + D = -8 \\ s & A + 6B - 3C - 2D = -15 \\ s^0 & -6A = -12 \end{cases} \quad \underline{A=2, \quad B=1, \quad C=-3, \quad D=1}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{2}{s} + \frac{1}{s-1} - \frac{3}{s+2} + \frac{1}{s+3}\right\}$$

$$\underline{y(t) = 2 + e^t - 3e^{-2t} + e^{-3t}}$$

Exercise

Solve using the Laplace transform: $y^{(3)} + 3y'' + 3y' + y = 0$; $y(0)=-4, \quad y'(0)=4, \quad y''(0)=-2$

Solution

$$\mathcal{L}\{y^{(3)} + 3y'' + 3y' + y\}(s) = 0$$

$$s^3 Y(s) - s^2 y(0) - sy'(0) - y''(0) + 3s^2 Y(s) - 3sy(0) - 3y'(0) + 3sY(s) - 3y(0) + Y(s) = 0$$

$$(s^3 + 3s^2 + 3s + 1)Y(s) = -4s^2 - 8s - 2$$

$$Y(s) = \frac{-4s^2 - 8s - 2}{(s+1)^3} = \frac{A}{s+1} + \frac{B}{(s+1)^2} + \frac{C}{(s+1)^3}$$

$$\begin{cases} s^2 & \underline{A=-4} \\ s & 2A + B = -8 \quad \underline{B=0} \\ s^0 & A + B + C = -2 \quad \underline{C=2} \end{cases}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{-4}{s+1} + \frac{2}{(s+1)^3}\right\}$$

$$\underline{y(t) = -4e^{-t} + t^2 e^{-t}}$$

Exercise

Solve using the Laplace transform: $y^{(3)} - 3y'' + 3y' - y = t^2 e^t$, $y(0) = 1$, $y'(0) = 2$, $y''(0) = 3$

Solution

$$\mathcal{L}\{y^{(3)} - 3y'' + 3y' - y\}(s) = \mathcal{L}\{t^2 e^t\}(s) \qquad \mathcal{L}\{t^n e^{-at}\}(s) = \frac{n!}{(s+a)^{n+1}}$$

$$s^3 Y(s) - s^2 y(0) - sy'(0) - y''(0) - 3s^2 Y(s) + 3sy(0) + 3y'(0) + 3sY(s) - 3y(0) - Y(s) = \frac{2}{(s-1)^3}$$

$$(s^3 - 3s^2 + 3s - 1)Y(s) - s^2 - 2s - 3 + 3s + 6 - 3 = \frac{2}{(s-1)^3}$$

$$(s-1)^3 Y(s) = \frac{2}{(s-1)^3} + s^2 - s$$

$$Y(s) = \frac{2 + (s^2 - s)(s^3 - 3s^2 + 3s - 1)}{(s-1)^6}$$

$$= \frac{s^5 - 4s^4 + 6s^3 - 4s^2 + s + 2}{(s-1)^6} = \frac{A}{s-1} + \frac{B}{(s-1)^2} + \frac{C}{(s-1)^3} + \frac{D}{(s-1)^4} + \frac{E}{(s-1)^5} + \frac{F}{(s-1)^6}$$

$$s^5 - 4s^4 + 6s^3 - 4s^2 + s + 2 = A(s-1)^5 + B(s-1)^4 + C(s-1)^3 + D(s-1)^2 + E(s-1) + F$$

$$(s-1)^5 = s^5 - 5s^4 + 10s^3 - 10s^2 + 5s - 1 \qquad (s-1)^4 = s^4 - 4s^3 + 6s^2 - 4s + 1$$

$$\begin{cases} s^4 & \underline{A=1} \\ s^4 & -5A + B = -4 \quad \rightarrow \underline{B=1} \\ s^3 & 10A - 4B + C = 6 \quad \rightarrow \underline{C=0} \\ s^2 & -10A + 6B - 3C + D = -4 \quad \rightarrow \underline{D=0} \\ s^1 & 5A - 4B + 3C - 2D + E = 1 \quad \rightarrow \underline{E=0} \\ s^0 & -A + B - C + D - E + F = 2 \quad \rightarrow \underline{F=2} \end{cases}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{1}{s-1} + \frac{1}{(s-1)^2} + \frac{2}{(s-1)^6}\right\}$$

$$\mathcal{L}^{-1}\left\{\frac{1}{(s+a)^{n+1}}\right\} = \frac{1}{n!} t^n e^{-at}$$

$$\underline{y(t) = e^t + te^t + \frac{2}{5!} t^5 e^t = e^t \left(1 + t + \frac{1}{60} t^5\right)}$$

Exercise

Solve using the Laplace transform: $y^{(3)} + y'' + 3y' - 5y = 16e^{-t}$; $y(0) = 0$, $y'(0) = 2$, $y''(0) = -4$

Solution

$$\mathcal{L}\{y^{(3)} + y'' + 3y' - 5y\}(s) = \mathcal{L}\{16e^{-t}\}(s)$$

$$s^3Y(s) - s^2y(0) - sy'(0) - y''(0) + s^2Y(s) - sy(0) - y'(0) + 3sY(s) - 3y(0) - 5Y(s) = \frac{16}{s+1}$$

$$(s^3 + s^2 + 3s - 5)Y(s) = \frac{16}{s+1} + 2s - 2$$

$$s^3 + s^2 + 3s - 5 = (s-1)(s^2 + 2s + 5) \quad \begin{array}{c|ccc} 1 & 1 & 1 & 3 & -5 \\ & & 1 & 2 & 5 \\ \hline & 1 & 2 & 5 & 0 \end{array}$$

$$Y(s) = \frac{2s^2 + 14}{(s+1)(s-1)(s^2 + 2s + 5)} = \frac{A}{s+1} + \frac{B}{s-1} + \frac{C(s+1) + D}{(s+1)^2 + 4}$$

$$\begin{cases} s^3 & A + B + C = 0 \\ s^2 & A + 3B + C + D = 2 \\ s & 3A + 7B - C = 0 \\ s^0 & -5A + 5B - C - D = 14 \end{cases} \quad \underline{A = -2, \quad B = 1, \quad C = 1, \quad D = 0}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{-2}{s+1} + \frac{1}{s-1} + \frac{(s+1)}{(s+1)^2 + 4}\right\}$$

$$\underline{y(t) = -2e^{-t} + e^t + e^{-t} \cos 2t}$$

Exercise

Solve using the Laplace transform: $y''' + 4y'' + 5y' + 2y = 10\cos t$, $y(0) = y'(0) = 0$, $y''(0) = 3$

Solution

$$\mathcal{L}\{y''' + 4y'' + 5y' + 2y\} = \mathcal{L}\{10\cos t\}$$

$$s^3Y(s) - s^2y(0) - sy'(0) - y''(0) + 4s^2Y(s) - 4sy(0) - 4y'(0) + 5sY(s) - 5y(0) + 2Y(s) = \frac{10s}{s^2 + 1}$$

$$(s^3 + 4s^2 + 5s + 2)Y(s) = \frac{10s}{s^2 + 1} + 3$$

$$\begin{array}{c|ccc} -1 & 1 & 4 & 5 & 2 \\ & & -1 & -3 & -2 \\ \hline & 1 & 3 & 2 & 0 \end{array} \rightarrow s^2 + 3s + 2 = 0 \quad \underline{s = -1, -1, -2}$$

$$\begin{aligned}
Y(s) &= \frac{3s^2 + 10s + 3}{(s+2)(s^2+1)(s+1)^2} = \frac{A}{s+2} + \frac{Bs+C}{s^2+1} + \frac{D}{s+1} + \frac{E}{(s+1)^2} \\
3s^2 + 10s + 3 &= A(s^2+1)(s^2+2s+1) + (Bs+C)(s+2)(s^2+2s+1) \\
&\quad + D(s+1)(s+2)(s^2+1) + E(s^2+1)(s+2) \\
&= A(s^2+1)(s^2+2s+1) + (Bs^2+2Bs+Cs+2C)(s^2+2s+1) \\
&\quad + D(s^2+3s+2)(s^2+1) + Es^3+2Es^2+Es+2E \\
&= As^4+2As^3+2As^2+2As+A+B s^4+4Bs^3+5Bs^2+2Bs+Cs^3+4Cs^2 \\
&\quad + 5Cs+2C+Ds^4+3Ds^3+3Ds^2+3Ds+2D+Es^2+3Es+2E \\
\begin{cases} s^4 & A+B+D=0 \\ s^3 & 2A+4B+C+3D+E=0 \\ s^2 & 2A+5B+4C+3D+2E=3 \\ s^1 & 2A+2B+5C+3D+E=10 \\ s^0 & A+2C+2D+2E=3 \end{cases} &\rightarrow \begin{matrix} A=-1 & B=-1 & C=2 \\ D=2 & E=-2 \end{matrix}
\end{aligned}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{-1}{s+2} - \frac{s}{s^2+1} + \frac{2}{s^2+1} + \frac{2}{s+1} - \frac{2}{(s+1)^2}\right\}$$

$$\underline{y(t) = -e^{-2t} - \cos t + 2 \sin t + 2e^{-t} - 2te^{-t}}$$

Exercise

Solve using the Laplace transform: $y^{(4)} + 2y'' + y = 4te^t$; $y(0) = y'(0) = y''(0) = y^{(3)}(0) = 0$

Solution

$$\mathcal{L}\{y^{(4)} + 2y'' + y\} = \mathcal{L}\{4te^t\}$$

$$s^4 Y(s) - s^3 y(0) - s^2 y'(0) - s y''(0) - y'''(0) + 2s^2 Y(s) - 2s y(0) - 2y'(0) + Y(s) = \frac{4}{(s-1)^2}$$

$$(s^4 + 2s^2 + 1)Y(s) = \frac{4}{(s-1)^2} \quad y(0) = y'(0) = y''(0) = y^{(3)}(0) = 0$$

$$(s^2 + 1)^2 Y(s) = \frac{4}{(s-1)^2}$$

$$Y(s) = \frac{4}{(s-1)^2(s^2+1)^2} = \frac{A}{s-1} + \frac{B}{(s-1)^2} + \frac{Cs+D}{s^2+1} + \frac{Es+F}{(s^2+1)^2}$$

$$A(s-1)(s^4+2s^2+1) + B(s^4+2s^2+1) + (Cs+D)(s^2-2s+1)(s^2+1) + (Es+F)(s^2-2s+1) = 4$$

$$\begin{cases} s^5 & A+C=0 \\ s^4 & -A-2C+D=0 \\ s^3 & 2A+C-2D+E=0 \\ s^2 & -2A+2B-2C+2D-2E+F=0 \\ s^1 & A+C-2D+E-2F=0 \\ s^0 & -A+B+D+F=4 \end{cases} \rightarrow \begin{matrix} A = -\frac{16}{17} & B = \frac{28}{17} & C = \frac{16}{17} \\ D = \frac{16}{17} & E = \frac{48}{17} & F = \frac{8}{17} \end{matrix}$$

$$Y(s) = -\frac{16}{17} \frac{1}{s-1} + \frac{28}{17} \frac{1}{(s-1)^2} + \frac{16}{17} \frac{s}{s^2+1} + \frac{16}{17} \frac{1}{s^2+1} + \frac{48}{17} \frac{s}{(s^2+1)^2} + \frac{8}{17} \frac{1}{(s^2+1)^2}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{-\frac{16}{17} \frac{1}{s-1} + \frac{28}{17} \frac{1}{(s-1)^2} + \frac{16}{17} \frac{s}{s^2+1} + \frac{16}{17} \frac{1}{s^2+1} + \frac{24}{17} \frac{2s}{(s^2+1)^2} + \frac{4}{17} \frac{2}{(s^2+1)^2}\right\}$$

$$\underline{y(t) = -\frac{16}{17}e^t + \frac{28}{17}te^t + \frac{16}{17}\cos t + \frac{16}{17}\sin t + \frac{27}{17}t\sin t + \frac{4}{17}\sin t - t\cos t}$$

Exercise

Solve using the Laplace transform: $y^{(4)} - y = 0$; $y(0)=1$, $y'(0)=0$, $y''(0)=0$, $y^{(3)}(0)=0$

Solution

$$\mathcal{L}\{y^{(4)} - y\}(s) = 0$$

$$s^4Y(s) - s^3y(0) - s^2y'(0) - sy''(0) - y'''(0) - Y(s) = 0$$

$$(s^4 - 1)Y(s) = s^3$$

$$Y(s) = \frac{s^3}{(s-1)(s+1)(s^2+1)} = \frac{A}{s-1} + \frac{B}{s+1} + \frac{Cs+D}{s^2+1}$$

$$s^3 = As^3 + As^2 + As + A + Bs^3 - Bs^2 + Bs - B + Cs^3 - Cs + Ds^2 - D$$

$$\begin{cases} s^3 & A+B+C=1 \\ s^2 & A-B+D=0 \\ s & A+B-C=0 \\ s^0 & A-B-D=0 \end{cases} \rightarrow \begin{matrix} A=\frac{1}{4} & B=\frac{1}{4} \\ C=\frac{1}{2} & D=0 \end{matrix}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{1}{4}\frac{1}{s-1} + \frac{1}{4}\frac{1}{s+1} + \frac{1}{2}\frac{s}{s^2+1}\right\}$$

$$\underline{y(t) = \frac{1}{4}e^t + \frac{1}{4}e^{-t} + \frac{1}{2}\cos t}$$

Exercise

Solve using the Laplace transform: $y^{(4)} - 4y = 0$; $y(0)=1, y'(0)=0, y''(0)=-2, y^{(3)}(0)=0$

Solution

$$\mathcal{L}\{y^{(4)} - 4y\} = 0$$

$$s^4 Y(s) - s^3 y(0) - s^2 y'(0) - s y''(0) - y'''(0) - 4Y(s) = 0 \quad y(0)=1, y'(0)=0, y''(0)=-2, y^{(3)}(0)=0$$

$$(s^4 - 4)Y(s) - s^3 + 2s = 0$$

$$Y(s) = \frac{s(s^2 - 2)}{(s^2 - 2)(s^2 + 2)}$$

$$= \frac{s}{s^2 + 2}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{s}{s^2 + 2}\right\}$$

$$\underline{y(t) = \cos \sqrt{2}t}$$

Exercise

Solve using the Laplace transform:

$$y^{(4)} - 4y^{(3)} + 6y'' - 4y' + y = 0; \quad y(0)=0, y'(0)=1, y''(0)=0, y^{(3)}(0)=1$$

Solution

$$\mathcal{L}\{y^{(4)} - 4y^{(3)} + 6y'' - 4y' + y\} = 0$$

$$s^4 Y(s) - s^3 y(0) - s^2 y'(0) - sy''(0) - y'''(0) - 4s^3 Y(s) + 4s^2 y(0) + 4sy'(0) + 4y''(0) + 6s^2 Y(s) - 6sy(0) - 6y'(0) - 4sY(s) + 4y(0) + Y(s) = 0$$

$$(s^4 - 4s^3 + 6s^2 - 4s + 1)Y(s) - s^2 - 1 + 4s - 6 = 0$$

$$(s+1)^4 Y(s) = s^2 - 4s + 7$$

$$Y(s) = \frac{s^2 - 4s + 7}{(s+1)^4} = \frac{A}{s+1} + \frac{B}{(s+1)^2} + \frac{C}{(s+1)^3} + \frac{D}{(s+1)^4}$$

$$As^3 + 3As^2 + 3As + A + Bs^2 + 2Bs + B + Cs + C + D = s^2 - 4s + 7$$

$$\begin{cases} s^3 & A = 0 \\ s^2 & 3A + B = 1 \\ s^1 & 3A + 2B + C = -4 \\ s^0 & A + B + C + D = 7 \end{cases} \Rightarrow B = 1 \quad C = -6 \quad D = 13$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{1}{(s+1)^2} - \frac{6}{(s+1)^3} + \frac{13}{(s+1)^4}\right\}$$

$$y(t) = te^t - 3t^2 e^t + \frac{13}{6}t^3 e^t$$

$$= \left(t - 3t^2 + \frac{13}{6}t^3\right)e^t$$

Exercise

Given: $y'' - 4y' + 3y = 0$, $y(0) = 1$ $y'(0) = -1$

a) Show that the general solution is: $y(t) = C_1 e^{3t} + C_2 e^t$ and find C_1 and C_2

b) Use Laplace transform to solve the system

Solution

$$a) \lambda^2 - 4\lambda + 3 = 0 \Rightarrow \boxed{\lambda = 3, 1}$$

That implies to the general solution: $y = C_1 e^{3t} + C_2 e^t$

$$1 = C_1 e^{3(0)} + C_2 e^{(0)}$$

$$\underline{1 = C_1 + C_2}$$

$$y' = 3C_1 e^{3t} + C_2 e^t$$

$$-1 = 3C_1 e^{3(0)} + C_2 e^{(0)}$$

$$\underline{-1 = 3C_1 + C_2}$$

$$\begin{cases} C_1 + C_2 = 1 \\ 3C_1 + C_2 = -1 \end{cases} \Rightarrow \boxed{C_1 = -1} \quad \boxed{C_2 = 2}$$

Therefore; the general solution is: $\boxed{y = -e^{3t} + 2e^t}$

b) $\mathcal{L}(y'' - 4y' + 3y)(s) = 0$

$$s^2 Y(s) - sy(0) - y'(0) - 4(sY(s) - y(0)) + 3Y(s) = 0$$

$$s^2 Y(s) - s + 1 - 4(sY(s) - 1) + 3Y(s) = 0$$

$$s^2 Y(s) - s + 1 - 4sY(s) + 4 + 3Y(s) = 0$$

$$(s^2 - 4s + 3)Y(s) = s - 5$$

$$Y(s) = \frac{s-5}{s^2 - 4s + 3}$$

$$= \frac{s-5}{(s-1)(s-3)}$$

$$= \frac{A}{s-1} + \frac{B}{s-3} = \frac{(A+B)s - 3A - B}{(s-1)(s-3)}$$

$$\begin{cases} A+B=1 \\ -3A-B=-5 \end{cases} \Rightarrow \boxed{A=2} \quad \boxed{B=-1}$$

$$= \frac{2}{s-1} - \frac{1}{s-3}$$

That implies: $\boxed{y(t) = 2e^t - e^{3t}}$

Exercise

Solve the initial value problem $x'' + 4x = \sin 3t$; $x(0) = x'(0) = 0$.

Such problem arises in the motion of a mass-and-spring system with external force as shown below.

Solution


$$\mathcal{L}\{x'' + 4x\} = \mathcal{L}\{\sin 3t\}$$

$$s^2 X(s) - sx(0) - x'(0) + 4X(s) = \frac{3}{s^2 + 9}$$

$$(s^2 + 4)X(s) = \frac{3}{s^2 + 9}$$

$$X(s) = \frac{3}{(s^2 + 4)(s^2 + 9)} = \frac{As + B}{s^2 + 4} + \frac{Cs + D}{s^2 + 9}$$

$$As^3 + 9AS + Bs^2 + 9B + Cs^3 + 4Cs + Ds^2 + 4D = 3$$



$$\begin{cases} s^3 & A + C = 0 \\ s^2 & B + D = 0 \\ s^1 & 9A + 4C = 0 \\ s^0 & 9B + 4D = 3 \end{cases} \rightarrow A = C = 0 \quad 5B = 3 \Rightarrow B = \frac{3}{5} \quad D = -\frac{3}{5}$$

$$X(s) = \frac{3}{5} \frac{1}{s^2 + 4} - \frac{3}{5} \frac{1}{s^2 + 9}$$

$$\mathcal{L}^{-1}\{X(s)\} = \mathcal{L}^{-1}\left\{\frac{3}{10} \frac{2}{s^2 + 4} - \frac{1}{5} \frac{3}{s^2 + 9}\right\}$$

$$\underline{x(t) = \frac{3}{10} \sin 2t - \frac{1}{5} \sin 3t}$$

Exercise

Solve the system
$$\begin{cases} 2x'' = -6x + 2y \\ y'' = 2x - 2y + 40 \sin 3t \end{cases}$$

Subject to the initial conditions $x(0) = x'(0) = y(0) = y'(0) = 0$

Thus the force $f(t) = 40 \sin 3t$ is applied to the second mass as shown below, beginning at time $t = 0$ when the system is at rest in its equilibrium position.

Solution

$$\begin{cases} \mathcal{L}\{2x''\} = \mathcal{L}\{-6x + 2y\} \\ \mathcal{L}\{y''\} = \mathcal{L}\{2x - 2y + 40 \sin 3t\} \end{cases}$$


$$\begin{cases} 2s^2 X(s) - 2sx(0) - 2x'(0) = -6X(s) + 2Y(s) \\ s^2 Y(s) - sy(0) - y'(0) = 2X(s) - 2Y(s) + \frac{120}{s^2 + 9} \end{cases}$$

Given: $x(0) = x'(0) = y(0) = y'(0) = 0$

$$\begin{cases} 2s^2 X(s) = -6X(s) + 2Y(s) \\ s^2 Y(s) = 2X(s) - 2Y(s) + \frac{120}{s^2 + 9} \end{cases}$$

$$\begin{cases} (s^2 + 3)X(s) - Y(s) = 0 & (1) \\ -2X(s) + (s^2 + 2)Y(s) = \frac{120}{s^2 + 9} & (2) \end{cases}$$

$$\begin{vmatrix} s^2 + 3 & -1 \\ -2 & s^2 + 2 \end{vmatrix} = s^4 + 5s^2 + 4 = (s^2 + 1)(s^2 + 4)$$



$$\begin{vmatrix} 0 & -1 \\ \frac{120}{s^2+9} & s^2+2 \end{vmatrix} = \frac{120}{s^2+9} \rightarrow X(s) = \frac{120}{(s^2+1)(s^2+4)(s^2+9)}$$

$$\begin{vmatrix} s^2+3 & 0 \\ -2 & \frac{120}{s^2+9} \end{vmatrix} = 120 \frac{s^2+3}{s^2+9} \rightarrow Y(s) = \frac{120(s^2+3)}{(s^2+1)(s^2+4)(s^2+9)}$$

$$X(s) = \frac{120}{(s^2+1)(s^2+4)(s^2+9)} = \frac{As+B}{s^2+1} + \frac{Cs+D}{s^2+4} + \frac{Es+F}{s^2+9}$$

$$(As+B)(s^2+4)(s^2+9) + (Cs+D)(s^2+1)(s^2+9) + (Es+F)(s^2+1)(s^2+4) = 120$$

$$(As+B)(s^4+13s^2+36) + (Cs+D)(s^4+10s^2+9) + (Es+F)(s^4+5s^2+4) = 120$$

$$\begin{cases} s^5 & A+C+E=0 \\ s^4 & B+D+F=0 \\ s^3 & 13A+10C+5E=0 \\ s^2 & 13B+10D+5F=0 \\ s & 36A+9C+4E=0 \\ & 36B+9D+4F=120 \end{cases} \Rightarrow \begin{cases} A+C+E=0 \\ 13A+10C+5E=0 \rightarrow A+C+E=0 \\ 36A+9C+4E=0 \\ B+D+F=0 \\ 13B+10D+5F=0 \rightarrow B=5; D=-8; F=3 \\ 36B+9D+4F=120 \end{cases}$$

$$Y(s) = \frac{120(s^2+3)}{(s^2+1)(s^2+4)(s^2+9)} = \frac{As+B}{s^2+1} + \frac{Cs+D}{s^2+4} + \frac{Es+F}{s^2+9}$$

$$(As+B)(s^2+4)(s^2+9) + (Cs+D)(s^2+1)(s^2+9) + (Es+F)(s^2+1)(s^2+4) = 120s^2 + 360$$

$$\begin{cases} s^5 & A+C+E=0 \\ s^4 & B+D+F=0 \\ s^3 & 13A+10C+5E=0 \\ s^2 & 13B+10D+5F=120 \\ s & 36A+9C+4E=0 \\ & 36B+9D+4F=360 \end{cases} \Rightarrow \begin{cases} A+C+E=0 \\ 13A+10C+5E=0 \rightarrow A+C+E=0 \\ 36A+9C+4E=0 \\ B+D+F=0 \\ 13B+10D+5F=120 \rightarrow B=10; D=8; F=-18 \\ 36B+9D+4F=360 \end{cases}$$

$$X(s) = \frac{5}{s^2+1} - \frac{8}{s^2+4} + \frac{3}{s^2+9}$$

$$\mathcal{L}^{-1}\{X(s)\} = \mathcal{L}^{-1}\left\{5\frac{1}{s^2+1} - 4\frac{2}{s^2+4} + \frac{3}{s^2+9}\right\}$$

$$\underline{x(t) = 5\sin t - 4\sin 2t + \sin 3t}$$


$$Y(s) = \frac{10}{s^2+1} + \frac{8}{s^2+4} - \frac{18}{s^2+9}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{10\frac{1}{s^2+1} + 4\frac{2}{s^2+4} - 6\frac{3}{s^2+9}\right\}$$

$$\underline{y(t) = 10\sin t + 4\sin 2t - 6\sin 3t}$$

Exercise

Consider a mass-spring system with $m = \frac{1}{2}$, $k = 17$, and $c = 3$.



Let $x(t)$ be the displacement of the mass m from its equilibrium position. If the mass is set in motion with $x(0) = 3$ and $x'(0) = 1$, find $x(t)$ for the resulting damped free oscillations.

Solution

$$\frac{1}{2}x'' + 3x' + 17x = 0 \quad mx'' + cx' + kx = 0$$

$$x'' + 6x' + 34x = 0 \quad x(0) = 3; \quad x'(0) = 1$$

$$\mathcal{L}\{x'' + 6x' + 34x\} = 0$$

$$s^2X(s) - sx(0) - x'(0) + 6sX(s) - 6x(0) + 34X(s) = 0$$


$$s^2X(s) - 3s - 1 + 6sX(s) - 18 + 34X(s) = 0$$

$$(s^2 + 6s + 34)X(s) = 3s + 19$$

$$\begin{aligned} X(s) &= \frac{3s + 19}{s^2 + 6s + 34} \\ &= \frac{3s + 19}{(s + 3)^2 + 25} \\ &= \frac{3s + 9 + 10}{(s + 3)^2 + 25} \\ &= \frac{3(s + 3)}{(s + 3)^2 + 25} + \frac{10}{(s + 3)^2 + 25} \end{aligned}$$

$$\mathcal{L}^{-1}\{X(s)\} = \mathcal{L}^{-1}\left\{3 \cdot \frac{(s + 3)}{(s + 3)^2 + 25} + 5 \cdot \frac{2}{(s + 3)^2 + 25}\right\}$$

$$\underline{x(t) = (3\cos 5t + 2\sin 5t)e^{-3t}}$$



Exercise

A 4-lb weight stretches a spring 2 feet. The weight is released from rest 18 inches above the equilibrium position, and the resulting motion takes place in a medium offering a damping force numerically equal to $\frac{7}{8}$ times the instantaneous velocity. Use the Laplace transform to find the equation of motion $x(t)$.

Solution

$$m = \frac{4}{32} = \frac{1}{8} \quad (w = mg)$$

$$k = \frac{4}{2} = 2 \quad (xk = mg)$$

$$c = \frac{7}{8}$$

$$\frac{1}{8}x'' + \frac{7}{8}x' + 2x = 0 \quad mx'' + cx' + kx = f(t)$$

$$x'' + 7x' + 16x = 0; \quad x(0) = -\frac{18}{12} = -\frac{3}{2}, \quad x'(0) = 0$$

$$\mathcal{L}\{x'' + 7x' + 16x\} = 0$$

$$s^2X(s) - sx(0) - x'(0) + 7sX(s) - 7x(0) + 16X(s) = 0$$

$$(s^2 + 7s + 16)X(s) = -\frac{3}{2}s - \frac{21}{2}$$

$$X(s) = -\frac{3}{2} \frac{s+7}{\left(s+\frac{7}{2}\right)^2 + \frac{15}{4}}$$

$$\frac{s+7}{\left(s+\frac{7}{2}\right)^2 + \frac{15}{4}} = \frac{A\left(s+\frac{7}{2}\right)}{\left(s+\frac{7}{2}\right)^2 + \frac{15}{4}} + \frac{B}{\left(s+\frac{7}{2}\right)^2 + \frac{15}{4}}$$

$$\begin{cases} \textcolor{red}{s} & \underline{A=1} \\ \textcolor{red}{s^0} & \frac{7}{2}A + B = 7 \quad \underline{B = \frac{7}{2}} \end{cases}$$

$$\{X(s)\} = -\frac{3}{2} \mathcal{L}^{-1} \left\{ \frac{s+\frac{7}{2}}{\left(s+\frac{7}{2}\right)^2 + \left(\frac{\sqrt{15}}{2}\right)^2} \right\} - \frac{3}{2} \frac{7}{2} \frac{\textcolor{red}{2}}{\sqrt{15}} \mathcal{L}^{-1} \left\{ \frac{\frac{\sqrt{15}}{2}}{\left(s+\frac{7}{2}\right)^2 + \left(\frac{\sqrt{15}}{2}\right)^2} \right\}$$

$$\underline{x(t) = -\frac{3}{2}e^{7t/2} \cos \frac{\sqrt{15}}{2}t - \frac{7\sqrt{15}}{10}e^{7t/2} \sin \frac{\sqrt{15}}{2}t}$$

Exercise

Consider a mass-spring-dashpot system with $m = \frac{1}{2}$, $k = 17$, $c = 3$, and $f(t) = 15 \sin 2t$ with initial conditions $x(0) = x'(0) = 0$. Let $x(t)$ be the displacement of the mass m from its equilibrium position. Find the resulting transient motion and steady periodic motion of the mass..

Solution

$$\frac{1}{2}x'' + 3x' + 17x = 15 \sin 2t \quad mx'' + cx' + kx = 0$$

$$x'' + 6x' + 34x = 30 \sin 2t \quad x(0) = x'(0) = 0$$

$$\mathcal{L}\{x'' + 6x' + 34x\} = \mathcal{L}\{30 \sin 2t\}$$

$$s^2 X(s) - sx(0) - x'(0) + 6sX(s) - 6x(0) + 34X(s) = \frac{60}{s^2 + 4}$$

$$(s^2 + 6s + 34)X(s) = \frac{60}{s^2 + 4}$$

$$X(s) = \frac{60}{(s^2 + 4)((s + 3)^2 + 25)} = \frac{As + B}{s^2 + 4} + \frac{Cs + D}{(s + 3)^2 + 25}$$

$$As^3 + 6As^2 + 34sA + Bs^2 + 6sB + 34B + Cs^3 + 4Cs + Ds^2 + 4D = 60$$

$$\begin{cases} s^3 & A + C = 0 \\ s^2 & 6A + B + D = 0 \\ s^1 & 34A + 6B + 4C = 0 \\ s^0 & 34B + 4D = 60 \end{cases}$$

$$\begin{cases} C = -A & 6A - \frac{15}{2}B = -15 \\ & 30A + 6B = 0 \\ D = 15 - \frac{17}{2}B \end{cases} \rightarrow \begin{cases} -30A + \frac{75}{2}B = 75 \\ 30A + 6B = 0 \end{cases} \quad \underline{B = \frac{150}{87} = \frac{50}{29}}$$

$$A = -\frac{10}{29}; \quad B = \frac{50}{29}; \quad C = \frac{10}{29}; \quad D = \frac{10}{29}$$

$$X(s) = \frac{10}{29} \left(\frac{-s+5}{s^2+4} + \frac{s+1}{(s+3)^2+25} \right)$$

$$= \frac{10}{29} \left(\frac{5}{s^2+4} - \frac{s}{s^2+4} + \frac{s+3}{(s+3)^2+25} - \frac{2}{(s+3)^2+25} \right)$$

$$\mathcal{L}^{-1}\{X(s)\} = \frac{10}{29} \mathcal{L}^{-1} \left(\frac{5}{2} \cdot \frac{2}{s^2+4} - \frac{s}{s^2+4} + \frac{s+3}{(s+3)^2+25} - \frac{2}{5} \cdot \frac{5}{(s+3)^2+25} \right)$$

$$x(t) = \frac{10}{29} \left(\frac{5}{2} \sin 2t - \cos 2t + e^{-3t} \left(\cos 5t - \frac{2}{5} \sin 5t \right) \right)$$

$$= \frac{5}{29}(5\sin 2t - 2\cos 2t) + \frac{2}{29}e^{-3t}(5\cos 5t - 2\sin 5t) \Big|$$

Exercise

A 8-kg mass is attached to a spring hanging vertically and come to rest at equilibrium. The damping constant is given by 3 N-sec/m and the spring constant is 40 N/m. If the mass is driven by an external force equal to $f(t) = 2\sin 2t \cos 2t$ N. Find the solution.

Solution

Given: $m = 8$ $k = 40$ $c = 3$

$$8y'' + 3y' + 40y = 2\sin 2t \cos 2t \\ = \sin 4t$$

$$my'' + cy' + ky = F(t)$$

$$8y'' + 3y' + 40y = \sin 4t ; \quad y(0) = 0, \quad y'(0) = 0$$

$$\mathcal{L}\{8y'' + 3y' + 40y\} = \mathcal{L}\{\sin 4t\}$$

$$8s^2Y(s) - 8sy(0) - 8y'(0) + 3sY(s) - 3y(0) + 40Y(s) = \frac{4}{s^2 + 16}$$

$$(8s^2 + 3s + 40)Y(s) = \frac{4}{s^2 + 16}$$

$$Y(s) = \frac{4}{8\left(s^2 + \frac{3}{8}s + 5\right)(s^2 + 16)}$$

$$= \frac{\frac{1}{2}}{\left(\left(s + \frac{3}{16}\right)^2 + \frac{1271}{256}\right)(s^2 + 16)} = \frac{A\left(s + \frac{3}{16}\right) + B}{\left(s + \frac{3}{16}\right)^2 + \frac{1271}{256}} + \frac{Cs + D}{s^2 + 16}$$

$$s^3 \quad A + C = 0$$

$$s^2 \quad \frac{3}{16}A + B + \frac{3}{8}C + D = 0$$

$$s \quad 16A + 5C + \frac{3}{8}D = 0$$

$$s^0 \quad 3A + 16B + 5D = \frac{1}{2}$$

$$A = \frac{3}{1972} \quad B = \frac{1417}{31552} \quad C = -\frac{3}{1972} \quad D = -\frac{22}{493} \Big|$$

$$\mathcal{L}^{-1}\{Y(s)\} = \left\{ \frac{3}{1972} \frac{s + \frac{3}{16}}{\left(s + \frac{3}{16}\right)^2 + \frac{1271}{256}} + \frac{1417}{31552} \frac{1}{\left(s + \frac{3}{16}\right)^2 + \frac{1271}{256}} - \frac{3}{1972} \frac{s}{s^2 + 16} - \frac{22}{493} \frac{1}{s^2 + 16} \right\}$$

$$y(t) = \frac{3}{1972} e^{-3t/16} \cos \frac{\sqrt{1271}}{16} t + \frac{1417}{31552} \frac{16}{\sqrt{1271}} e^{-3t/16} \sin \frac{\sqrt{1271}}{16} t - \frac{3}{1972} \cos 4t - \frac{22}{493} \frac{1}{4} \sin 4t$$

$$= \frac{3}{1972} e^{-3t/16} \cos \frac{\sqrt{1271}}{16} t + \frac{1417}{1972\sqrt{1271}} e^{-3t/16} \sin \frac{\sqrt{1271}}{16} t - \frac{3}{1972} \cos 4t - \frac{11}{986} \sin 4t \Big|$$

Exercise

A 2-kg mass is attached to a spring hanging vertically and come to rest at equilibrium. The damping constant is given by $c = 8 \text{ kg/sec}$ and the spring constant is $k = 80 \text{ N/m}$. At time $t = 0$, the resulting spring-mass system is disturbed from its rest state by the force $F(t) = 20e^{-t} \text{ N}$. (t in seconds). Find the equation of motion.

Solution

$$2y'' + 8y' + 80y = 20e^{-t} ; \quad y(0) = 0, \quad y'(0) = 0 \qquad my'' + cy' + ky = F(t)$$

$$\mathcal{L}^{-1}\{2y'' + 8y' + 80y\} = \mathcal{L}^{-1}\{20e^{-t}\}$$

$$2s^2Y(s) - 2sy(0) - 2y'(0) + 8sY(s) - 8y(0) + 80Y(s) = \frac{20}{s+1}$$

$$2(s^2 + 4s + 40)Y(s) = \frac{20}{s+1}$$

$$Y(s) = \frac{10}{(s+1)((s+2)^2 + 36)} = \frac{A}{s+1} + \frac{B(s+2) + C}{(s+2)^2 + 36}$$

$$\begin{cases} s^2 & A + B = 0 \\ s & 4A + 3B + C = 0 \\ s^0 & 40A + 2B + C = 10 \end{cases}$$

$$\Delta = \begin{vmatrix} 1 & 1 & 0 \\ 4 & 3 & 1 \\ 40 & 2 & 1 \end{vmatrix} = 37 \quad \Delta_A = \begin{vmatrix} 0 & 1 & 0 \\ 0 & 3 & 1 \\ 10 & 2 & 1 \end{vmatrix} = 10$$

$$A = \frac{10}{37} \quad B = -\frac{10}{37} \quad C = -\frac{10}{37}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{10}{37} \frac{1}{s+1} - \frac{10}{37} \frac{(s+2)}{(s+2)^2 + 6^2} - \frac{10}{37} \frac{1}{6} \frac{6}{(s+2)^2 + 6^2}\right\}$$

$$y(t) = \frac{10}{37} e^{-t} - \left(\frac{10}{37} \cos 6t + \frac{5}{111} \sin 6t\right) e^{-2t}$$

Exercise

A 10-kg mass is attached to a spring having a spring constant of 140 N/m . The mass is started in motion initially from the equilibrium position with an initial velocity 1 m/sec in the upward direction and with an applied external force $F(t) = 5 \sin t$. If the force due to air resistance is $-90y' \text{ N}$. Find the equation motion of the mass.

Solution

$$10y'' + 90y' + 140y = 5\sin t$$

$$y'' + 9y' + 14y = \frac{1}{2}\sin t ; \quad y(0) = 0, \quad y'(0) = -1$$

$$s^2Y(s) - sy(0) - y'(0) + 9sY(s) - 9y(0) + 14Y(s) = \frac{1}{2} \frac{1}{s^2 + 1}$$

$$(s^2 + 9s + 14)Y(s) = \frac{1}{2} \frac{1}{s^2 + 1} - 1$$

$$Y(s) = \frac{-s^2 - \frac{1}{2}}{(s+2)(s+7)(s^2+1)} = \frac{A}{s+2} + \frac{B}{s+7} + \frac{Cs+D}{s^2+1}$$

$$\begin{cases} s^3 & A+B+C=0 \\ s^2 & 7A+2B+9C+D=-1 \\ s & A+B+14C+9D=0 \\ s^0 & 7A+2B+14D=-\frac{1}{2} \end{cases} \quad \underline{A = -\frac{9}{50} \quad B = \frac{99}{500} \quad C = -\frac{9}{500} \quad D = \frac{13}{500}}$$

$$\{Y(s)\} = \left\{ -\frac{9}{50} \frac{1}{s+2} + \frac{99}{500} \frac{1}{s+7} - \frac{9}{500} \frac{s}{s^2+1} + \frac{13}{500} \frac{1}{s^2+1} \right\}$$

$$y(t) = -\frac{9}{50}e^{-2t} + \frac{99}{500}e^{-7t} - \frac{9}{500}\cos t + \frac{13}{500}\sin t$$

$$= \frac{1}{500} \left(99e^{-7t} - 90e^{-2t} + 13\sin t - 9\cos t \right)$$

Exercise

A 128-*lb* weight is attached to a spring having a spring constant of 64 *lb/ft*. The weight is started in motion initially by displacing it 6 *in* above the equilibrium position with no initial velocity and with an applied external force $F(t) = 8\sin 4t$. Assume no air resistance. Find the equation motion of the mass.

Solution

$$m = \frac{128}{32} = 4$$

$$4y'' + 64y = 8\sin 4t$$

$$y'' + 16y = 2\sin 4t ; \quad y(0) = -\frac{6}{12} = -\frac{1}{2}, \quad y'(0) = 0$$

$$s^2Y(s) - sy(0) - y'(0) + 16Y(s) = \frac{8}{s^2 + 16}$$

$$(s^2 + 16)Y(s) = \frac{8}{s^2 + 16} - \frac{1}{2}s$$

$$Y(s) = \frac{8}{(s^2 + 16)^2} - \frac{1}{2} \frac{s}{s^2 + 16}$$

$$\frac{8}{(s^2+16)^2} = \frac{As+B}{s^2+16} + \frac{C(s^2-16)}{(s^2+16)^2} + \frac{Ds}{(s^2+16)^2}$$

$$\begin{array}{lcl} s^3 & A=0 \\ s^2 & B+C=0 \\ s & 16A+D=0 \\ s^0 & 16B-16C=8 \end{array} \quad \underline{A=0 \quad B=\frac{1}{4} \quad C=-\frac{1}{4} \quad D=0}$$

$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{1}{4}\frac{1}{s^2+16} - \frac{1}{4}\frac{s^2-16}{(s^2+16)^2} - \frac{1}{2}\frac{s}{s^2+16}\right\}$$

$$\underline{y(t) = \frac{1}{16}\sin 4t - \frac{1}{4}t \cos 4t - \frac{1}{2}\cos 4t}$$

Exercise

Find the motion of a damped mass-and-spring system with $m=1$, $c=2$, and $k=26$ under the influence of an external force $F(t)=82\cos 4t$ with $x(0)=6$ and $x'(0)=0$.

Solution

Given: $m=1$, $c=2$, $k=26$, and $F(t)=82\cos 4t$ $x(0)=6$; $x'(0)=0$

$$x'' + 2x' + 26x = 82\cos 4t \quad mx'' + cx' + kx = F(t)$$

$$\mathcal{L}\{x'' + 2x' + 26x\} = \mathcal{L}\{82\cos 4t\}$$

$$s^2X(s) - sx(0) - x'(0) + 2sX(s) - 2x(0) + 26X(s) = \frac{82s}{s^2+16}$$

$$(s^2 + 2s + 26)X(s) = \frac{82s}{s^2+16} + 6s + 12$$

$$X(s) = \frac{6s^3 + 12s^2 + 178s + 192}{(s^2+16)((s+1)^2+25)} = \frac{As+B}{s^2+16} + \frac{C(s+1)+D}{(s+1)^2+25}$$

$$\begin{cases} s^3 & A+C=6 \\ s^2 & 2A+B+C+D=12 \\ s & 26A+2B+16C=178 \\ s^0 & 26B+16C+16D=192 \end{cases}$$

$$\underline{A=5 \quad B=16 \quad C=1 \quad D=-15}$$

$$\mathcal{L}^{-1}\{X(s)\} = \mathcal{L}^{-1}\left\{\frac{5s}{s^2+16} + \frac{16}{s^2+16} + \frac{s+1}{(s+1)^2+25} - \frac{15}{(s+1)^2+5^2}\right\}$$

$$\underline{x(t) = 5\cos 4t + 4\sin 4t + e^{-t}(\cos 5t - 3\sin 5t)}$$

Exercise

A spring with a mass of 2-kg has natural length 0.5 m. A force of 25.6 N is required to maintain it stretched to a length of 0.7 m. The spring is immersed in a fluid with damping constant $c = 40$. If the spring is started from the equilibrium position and is given a push to start it with initial velocity 0.6 m/s. Find the position of the mass at any time t .

Solution

$$k = \frac{25.6}{0.7-0.5} = 128 \quad k(x_2 - x_1) = F$$

$$2x'' + 40x + 128 = 0; \quad x(0) = 0, \quad x'(0) = 0.6$$

$$\mathcal{L}\{x'' + 20x + 64\} = 0$$

$$s^2 X(s) - sx(0) - x'(0) + 20sX(s) - 20x(0) + 64X(s) = 0$$

$$(s^2 + 20s + 64)X(s) = \frac{6}{10}$$

$$X(s) = \frac{3}{5} \frac{1}{(s+16)(s+4)} = \frac{3}{5} \left(\frac{A}{s+16} + \frac{B}{s+4} \right)$$

$$\begin{array}{l} s^1 \quad A + B = 0 \\ s^0 \quad 4A + 16B = 1 \end{array} \rightarrow \underline{A = -\frac{1}{12}, B = \frac{1}{12}}$$

$$\mathcal{L}^{-1}\{X(s)\} = \frac{3}{5} \mathcal{L}^{-1}\left\{-\frac{1}{12} \frac{1}{s+16} + \frac{1}{12} \frac{1}{s+4}\right\}$$

$$\begin{aligned} x(t) &= \frac{3}{5} \left(-\frac{1}{12} e^{-16t} + \frac{1}{12} e^{-t} \right) \\ &= \underline{\frac{1}{20} e^{-4t} - \frac{1}{20} e^{-16t}} \end{aligned}$$

Exercise

A spring with a mass of 3-kg is held stretched 0.6 m beyond its natural length by a force of 20 N. If the spring begins at its equilibrium and with initial velocity 1.2 m/s. Find the position of the mass.

Solution

$$k = \frac{20}{0.6} = \frac{100}{3} \quad kx = F$$

$$3x'' + \frac{100}{3}x = 0; \quad x(0) = 0, \quad x'(0) = 1.2 = \frac{6}{5} \quad mx'' + cx' + kx = F(t)$$

$$\mathcal{L}\{9x'' + 100x\} = 0$$

$$9s^2 X(s) - 9sx(0) - 9x'(0) + 100X(s) = 0$$

$$(9s^2 + 100)X(s) = \frac{36}{5}$$

$$\mathcal{L}^{-1}\{X(s)\} = \frac{6}{5} \mathcal{L}^{-1}\left\{\frac{1}{s^2 + \frac{100}{9}}\right\}$$

$$x(t) = \frac{6}{5} \frac{3}{10} \sin \frac{10}{3} t$$

$$= \frac{9}{25} \sin \frac{10}{3} t$$

Exercise

A spring with a mass of 2-*kg* is held stretched 0.5 *m*, has damping constant 14, and a force of 6 *N*. If the spring is stretched 1 *m* beyond at its equilibrium and with no initial velocity. Find the position of the mass at any time *t*.

Solution

$$k = \frac{6}{.5} = 12 \quad kx = F$$

$$2x'' + 14x' + 12x = 0 ; \quad x(0) = 1, \quad x'(0) = 0 \quad mx'' + cx' + kx = F(t)$$

$$\mathcal{L}\{x'' + 7x' + 6x\} = 0$$

$$s^2 X(s) - sx(0) - x'(0) + 7sX(s) - 7x(0) + 6X(s) = 0$$

$$(s^2 + 7s + 6)X(s) = s + 7$$

$$X(s) = \frac{s+7}{(s+1)(s+6)} = \frac{A}{s+1} + \frac{B}{s+6}$$

$$s + 7 = As + 6A + Bs + B$$

$$\begin{cases} s & A + B = 1 \\ s^0 & 6A + B = 7 \end{cases} \rightarrow \underline{A = \frac{6}{5}, B = -\frac{1}{5}}$$

$$\mathcal{L}^{-1}\{X(s)\} = \mathcal{L}^{-1}\left\{\frac{6}{5} \frac{1}{s+1} - \frac{1}{5} \frac{1}{s+6}\right\}$$

$$x(t) = \frac{6}{5} e^{-t} - \frac{1}{5} e^{-6t}$$

Exercise

When a uniform beam is supported by an elastic foundation, the differential equation for its deflection $y(x)$ is

$$EI \frac{d^4 y}{dx^4} + ky = w(x)$$

Where k is the modulus of the foundation and $-ky$ is the restoring force of the foundation that acts in the direction opposite to that of the load $w(x)$. For algebraic convenience, suppose that the differential equation is written as

$$\frac{d^4 y}{dx^4} + 4a^4 y = \frac{w(x)}{EI}$$

Where $a = \left(\frac{k}{4EI}\right)^{1/4}$. Assume $L = \pi$ and $a = 1$. Find the deflection $y(x)$ of a beam that is supported on an elastic foundation when

- The beam is simply supported at both ends and a constant load w_0 is uniformly distributed along its length,
- The beam is embedded at both ends and $w(x)$ is concentrated load w_0 applied at $x = \frac{\pi}{2}$

Solution

$$a) \quad y(0) = y''(0) = 0 \quad \& \quad y(\pi) = y''(\pi) = 0$$

$$\text{Let: } y'(0) = c_1 \quad y'(0) = c_2$$

$$\mathcal{L}\left\{\frac{d^4 y}{dx^4} + 4y\right\} = \mathcal{L}\left\{\frac{w(x)}{EI}\right\}$$

$$s^4 Y(s) - s^3 y(0) - s^2 y'(0) - sy''(0) - y'''(0) + 4Y(s) = \frac{w_0}{EI} \frac{1}{s}$$

$$(s^4 + 4)Y(s) = \frac{w_0}{EI} \frac{1}{s} + c_1 s^2 + c_2$$

$$Y(s) = \frac{w_0}{EI} \frac{1}{s(s^4 + 4)} + \frac{c_1 s^2}{s^4 + 4} + \frac{c_2}{s^4 + 4}$$

$$\frac{1}{s(s^4 + 4)} = \frac{A}{s} + \frac{B(s-1) + C}{(s-1)^2 + 1} + \frac{D(s+1) + E}{(s+1)^2 + 1}$$

$$s^4 \quad A + B + D = 0$$


$$s^3 \quad B + C - D + E = 0$$

$$s^2 \quad 2C - 2E = 0$$

$$s \quad -2B + 2C + 2D + 2E = 0$$

$$s^0 \quad 4A = 1$$

$$\left. A = \frac{1}{4}, B = -\frac{1}{8}, C = 0, D = -\frac{1}{8}, E = 0 \right|$$



$$\mathcal{L}^{-1}\{Y(s)\} = \mathcal{L}^{-1}\left\{\frac{w_0}{EI}\left(\frac{1}{4}\frac{1}{s} - \frac{1}{8}\frac{s-1}{(s-1)^2+1} - \frac{1}{8}\frac{s+1}{(s+1)^2+1}\right) + \frac{c_1}{2}\frac{2s^2}{s^4+4} + \frac{c_2}{4}\frac{4}{s^4+4}\right\}$$

$$y(x) = \frac{w_0}{EI}\left(\frac{1}{4} - \frac{1}{8}e^x \cos x - \frac{1}{8}e^{-x} \cos x\right) + \frac{c_1}{2}(\sin x \cosh x + \cos x \sinh x) \\ + \frac{c_2}{4}(\sin x \cosh x - \cos x \sinh x)$$

$$y(x) = \frac{w_0}{4EI}(1 - \cos x \cosh x) + \frac{c_1}{2}(\sin x \cosh x + \cos x \sinh x) + \frac{c_2}{4}(\sin x \cosh x - \cos x \sinh x)$$

$$y(\pi) = \frac{w_0}{4EI}(1 + \cosh \pi) - \frac{1}{2}c_1 \sinh \pi + \frac{1}{4}c_2 \sinh \pi = 0$$

$$\underline{2c_1 \sinh \pi - c_2 \sinh \pi = \frac{w_0}{EI}(1 + \cosh \pi)}$$

$$y' = \frac{w_0}{4EI}(\sin x \cosh x - \cos x \sinh x) + c_1 \cos x \cosh x + \frac{1}{2}c_2 \sin x \sinh x$$

$$y'' = \frac{w_0}{2EI} \sin x \sinh x + c_1(-\sin x \cosh x + \cos x \sinh x) + \frac{1}{2}c_2(\cos x \sinh x + \sin x \cosh x)$$

$$y''(\pi) = -c_1 \sinh \pi - \frac{1}{2}c_2 \sinh \pi = 0$$

$$\underline{c_1 = -\frac{1}{2}c_2}$$

$$2c_1 \sinh \pi + 2c_1 \sinh \pi = \frac{w_0}{EI}(1 + \cosh \pi)$$

$$\underline{c_1 = \frac{w_0}{4EI}(1 + \cosh \pi) \operatorname{csch} \pi \quad c_2 = -\frac{w_0}{2EI}(1 + \cosh \pi) \operatorname{csch} \pi}$$

$$y(x) = \frac{w_0}{4EI}(1 - \cos x \cosh x) + \frac{w_0}{8EI}(1 + \cosh \pi) \operatorname{csch} \pi (\sin x \cosh x + \cos x \sinh x) \\ - \frac{w_0}{4EI}(1 + \cosh \pi) \operatorname{csch} \pi (\sin x \cosh x - \cos x \sinh x)$$

$$b) \quad \mathcal{L}\left\{\frac{d^4 y}{dx^4} + 4y\right\} = \mathcal{L}\left\{\delta\left(t - \frac{\pi}{2}\right)\right\}$$

$$s^4 Y(s) - s^3 y(0) - s^2 y'(0) - s y''(0) - y'''(0) + 4Y(s) = \frac{w_0}{EI} e^{-\pi s/2}$$

$$Y(s) = \frac{w_0}{4EI} \frac{4}{s^4+4} e^{-\pi s/2} + \frac{c_1}{2} \frac{2s^2}{s^4+4} + \frac{c_2}{4} \frac{4}{s^4+4}$$

$$y(x) = \frac{w_0}{4EI} \left(\sin\left(x - \frac{\pi}{2}\right) \cosh\left(x - \frac{\pi}{2}\right) - \cos\left(x - \frac{\pi}{2}\right) \sinh\left(x - \frac{\pi}{2}\right) \right) u\left(x - \frac{\pi}{2}\right) \\ + \frac{c_1}{2} \sin x \sinh x + \frac{c_2}{4} (\sin x \cosh x - \cos x \sinh x)$$

$$y(\pi) = \frac{w_0}{4EI} \cosh \frac{\pi}{2} + \frac{c_2}{4} \sinh \pi = 0 \rightarrow c_2 = -\frac{w_0}{EI} \frac{\cosh \frac{\pi}{2}}{\sinh \pi}$$


$$c_1 = \frac{w_0}{EI} \frac{\sinh \frac{\pi}{2}}{\sinh \pi}$$

$$y(x) = \frac{w_0}{4EI} \left(\sin \left(x - \frac{\pi}{2} \right) \cosh \left(x - \frac{\pi}{2} \right) - \cos \left(x - \frac{\pi}{2} \right) \sinh \left(x - \frac{\pi}{2} \right) \right) u \left(x - \frac{\pi}{2} \right) \\ + \frac{w_0}{2EI} \frac{\sinh \frac{\pi}{2}}{\sinh \pi} \sin x \sinh x - \frac{w_0}{4EI} \frac{\cosh \frac{\pi}{2}}{\sinh \pi} (\sin x \cosh x - \cos x \sinh x)$$

Exercise

Suppose two identical pendulums are coupled by means of a spring with constant k . when the displacement angles $\theta_1(t)$ and $\theta_2(t)$ are small, the system of linear differential equations describing the motion is

$$\begin{cases} \theta_1'' + \frac{g}{l} \theta_1 = -\frac{k}{m} (\theta_1 - \theta_2) \\ \theta_2'' + \frac{g}{l} \theta_2 = \frac{k}{m} (\theta_1 - \theta_2) \end{cases}$$



a) Use Laplace transform to solve the system when

$$\theta_1'(0) = 0 \quad \theta_1(0) = \theta_0 \quad \theta_2'(0) = 0 \quad \theta_2(0) = \psi_0$$

Where θ_0 and ψ_0 constants. Let $\omega^2 = \frac{g}{l}$, $K = \frac{k}{m}$

b) Use the solution in part (a) to discuss the motion of the coupled pendulums in the special case when the initial conditions are $\theta_1'(0) = 0$, $\theta_1(0) = \theta_0$, $\theta_2'(0) = \theta_0$, $\theta_2(0) = 0$

c) Use the solution in part (a) to discuss the motion of the coupled pendulums in the special case when the initial conditions are $\theta_1'(0) = 0$, $\theta_1(0) = \theta_0$, $\theta_2'(0) = -\theta_0$, $\theta_2(0) = 0$

Solution

$$a) \begin{cases} \theta_1'' + \omega^2 \theta_1 = -K \theta_1 + K \theta_2 \\ \theta_2'' + \omega^2 \theta_2 = K \theta_1 - K \theta_2 \end{cases}$$

$$\begin{cases} s^2 \theta_1(s) - s \theta_1(0) - \theta_1'(0) + \omega^2 \theta_1(s) + K \theta_1(s) = +K \theta_2(s) \\ s^2 \theta_2(s) - s \theta_2(0) - \theta_2'(0) + \omega^2 \theta_2(s) + K \theta_2(s) = K \theta_1(s) \end{cases}$$

$$\begin{cases} (s^2 + \omega^2 + K)\theta_1(s) - K\theta_2(s) = s\theta_0 \\ -K\theta_1(s) + (s^2 + \omega^2 + K)\theta_2(s) = s\psi_0 \end{cases}$$

$$\Delta = \begin{vmatrix} s^2 + \omega^2 + K & -K \\ -K & s^2 + \omega^2 + K \end{vmatrix} = (s^2 + \omega^2 + K)^2 - K^2 = (s^2 + \omega^2)(s^2 + \omega^2 + 2K)$$

$$\Delta_1 = \begin{vmatrix} s\theta_0 & -K \\ s\psi_0 & s^2 + \omega^2 + K \end{vmatrix} = s^3\theta_0 + (\omega^2\theta_0 + K\theta_0 + K\psi_0)s$$

$$\Delta_2 = \begin{vmatrix} s^2 + \omega^2 + K & s\theta_0 \\ -K & s\psi_0 \end{vmatrix} = s^3\psi_0 + (\omega^2\psi_0 + K\psi_0 + K\theta_0)s$$

$$\theta_1(s) = \frac{s^3\theta_0 + (\omega^2\theta_0 + K\theta_0 + K\psi_0)s}{(s^2 + \omega^2)(s^2 + \omega^2 + 2K)} = \frac{As + B}{s^2 + \omega^2} + \frac{Cs + D}{s^2 + (\omega^2 + 2K)}$$

$$\begin{cases} s^3 & A + C = \theta_0 \\ s^2 & B + D = 0 \\ s & (\omega^2 + 2K)A + \omega^2 C = \omega^2\theta_0 + K\theta_0 + K\psi_0 \\ s^0 & (\omega^2 + 2K)B + \omega^2 D = 0 \end{cases}$$

$$(\omega^2 + 2K)A + \omega^2(\theta_0 - A) = \omega^2\theta_0 + K\theta_0 + K\psi_0$$

$$2KA = K\theta_0 + K\psi_0 \rightarrow \underline{A = \frac{1}{2}(\theta_0 + \psi_0)}$$

$$C = \theta_0 - A \rightarrow \underline{C = \frac{1}{2}(\theta_0 - \psi_0)}$$

$$\underline{B = D = 0}$$

$$\{\theta_1(s)\} = \frac{1}{2}(\theta_0 + \psi_0)\mathcal{L}^{-1}\left\{\frac{s}{s^2 + \omega^2}\right\} + \frac{1}{2}(\theta_0 - \psi_0)\mathcal{L}^{-1}\left\{\frac{s}{s^2 + (\sqrt{\omega^2 + 2K})^2}\right\}$$

$$\underline{\theta_1(t) = \frac{1}{2}(\theta_0 + \psi_0)\cos\omega t + \frac{1}{2}(\theta_0 - \psi_0)\cos\sqrt{\omega^2 + 2K}t}$$

$$\theta_2(s) = \frac{s^3\psi_0 + (\omega^2\psi_0 + K\psi_0 + K\theta_0)s}{(s^2 + \omega^2)(s^2 + \omega^2 + 2K)} = \frac{as + b}{s^2 + \omega^2} + \frac{cs + d}{s^2 + (\omega^2 + 2K)}$$

$$\begin{cases} s^3 & a + c = \psi_0 \\ s^2 & b + d = 0 \\ s & (\omega^2 + 2K)a + \omega^2 c = \omega^2 \psi_0 + K\psi_0 + K\theta_0 \\ s^0 & (\omega^2 + 2K)b + \omega^2 d = 0 \end{cases}$$

$$(\omega^2 + 2K)a + \omega^2(\psi_0 - a) = \omega^2 \psi_0 + K\psi_0 + K\theta_0$$

$$2Ka = K\theta_0 + K\psi_0 \rightarrow a = \frac{1}{2}(\theta_0 + \psi_0)$$

$$C = \psi_0 - A \rightarrow C = -\frac{1}{2}(\theta_0 - \psi_0)$$

$$b = d = 0$$

$$\theta_2(t) = \frac{1}{2}(\theta_0 + \psi_0)\cos\omega t - \frac{1}{2}(\theta_0 - \psi_0)\cos\sqrt{\omega^2 + 2K}t$$

$$b) \quad \theta'_1(0) = 0, \quad \theta_1(0) = \theta_0, \quad \theta'_2(0) = \theta_0, \quad \theta_2(0) = 0$$

$$\Rightarrow \psi_0 = \theta_0$$

$$\theta_1(t) = \theta_0 \cos\omega t \quad \& \quad \theta_2(t) = \theta_0 \cos\omega t$$

\therefore This means that both pendulums swing in the same direction (free) and the spring exerts no influence on the motion.

$$c) \quad \theta'_1(0) = 0, \quad \theta_1(0) = \theta_0, \quad \theta'_2(0) = -\theta_0, \quad \theta_2(0) = 0$$


$$\Rightarrow \psi_0 = -\theta_0$$

$$\theta_1(t) = \theta_0 \cos\sqrt{\omega^2 + 2K}t \quad \& \quad \theta_2(t) = -\theta_0 \cos\sqrt{\omega^2 + 2K}t$$

\therefore This means that both pendulums swing in the opposite directions, stretching and compressing the spring. The amplitude of both displacements is $|\theta_0|$. Which the spring is stretched to its maximum.

Exercise

A projectile, such as the canon ball, has weight $w = mg$ and initial velocity v_0 that is tangent to its path of motion.



If air resistance and all other forces except its weight are ignored, that motion of the projectile is describe by the system of linear differential equations:

$$\begin{cases} m \frac{d^2 x}{dt^2} = 0 \\ m \frac{d^2 y}{dt^2} = -mg \end{cases}$$

- a) Use Laplace transform to solve the system when

$$x(0) = 0 \quad x'(0) = v_0 \cos \theta \quad y(0) = 0 \quad y'(0) = v_0 \sin \theta$$

Where $v_0 = |v|$ is constant and θ is the constant angle of elevation.

The solutions $x(t)$ and $y(t)$ are parametric equations of the trajectory of the projectile.

- b) Use $x(t)$ in part (a) to eliminate the parameter t in $y(t)$. Use the resulting equation for y to show that the horizontal range R of the projectile is given by

$$R = \frac{v_0^2}{g} \sin 2\theta$$

- c) From the formula in part (b), we see that R is a maximum when $\sin 2\theta = 1$ or when $\theta = \frac{\pi}{4}$. Show that the same range – less than the maximum – can be obtained by firing the gun at either of two complementary angles θ and $\frac{\pi}{2} - \theta$. The only difference is that the smaller angle results in a low trajectory whereas the larger angle gives a high trajectory.

- d) Suppose $g = 32 \text{ ft/s}^2$, $\theta = 30^\circ$, and $v_0 = 300 \text{ ft/s}$. Use part (b) to find the horizontal range of the projectile.

- e) Find the time when the projectile hits the ground.

- f) Use the parametric equations $x(t)$ and $y(t)$ in part (a) along with the numerical data in part (d) to plot the ballistic curve of the projectile.

- g) Repeat with $\theta = 52^\circ$ and $v_0 = 300 \text{ ft/s}$.

- h) Superimpose both curves part (f) & (g) on the same coordinate system.

Solution

$$a) \quad \begin{cases} \frac{d^2x}{dt^2} = 0 \\ \frac{d^2y}{dt^2} = -g \end{cases} \rightarrow \begin{cases} s^2X(s) - sx(0) - x'(0) = 0 \\ s^2Y(s) - sy(0) - y'(0) = -\frac{g}{s} \end{cases}$$

$$x(0) = 0 \quad x'(0) = v_0 \cos \theta \quad y(0) = 0 \quad y'(0) = v_0 \sin \theta$$

$$X(s) = v_0 \cos \theta \frac{1}{s^2}$$

$$\underline{x(t) = (v_0 \cos \theta)t}$$

$$Y(s) = v_0 \sin \theta \frac{1}{s^2} - \frac{g}{s^3}$$

$$\underline{y(t) = (v_0 \sin \theta)t - \frac{1}{2}gt^2}$$

$$b) \quad t = \frac{x}{v_0 \cos \theta}$$

$$\begin{aligned} y(x) &= (v_0 \sin \theta)t - \frac{1}{2}gt^2 \\ &= (v_0 \sin \theta) \frac{x}{v_0 \cos \theta} - \frac{1}{2}g \frac{x^2}{v_0^2 \cos^2 \theta} \\ &= \frac{1}{2v_0^2 \cos^2 \theta} (2v_0^2 \cos \theta \sin \theta x - gx^2) \\ &= \frac{1}{2v_0^2 \cos^2 \theta} (v_0^2 \sin 2\theta - gx)x = 0 \end{aligned}$$

At $x = y = 0$, the projectile hits the ground.

$$v_0^2 \sin 2\theta - gx = 0$$

$$\underline{x = R(\theta) = \frac{1}{g} v_0^2 \sin 2\theta}$$

$$\begin{aligned} c) \quad R\left(\frac{\pi}{2} - \theta\right) &= \frac{1}{g} v_0^2 \sin(\pi - 2\theta) \\ &= \frac{1}{g} v_0^2 \sin 2\theta \\ &= R(\theta) \end{aligned}$$

$$\sin(\pi - \alpha) = \sin \pi \cos \alpha - \cos \pi \sin \alpha$$

$$d) \quad \text{Given: } g = 32 \text{ ft/s}^2, \theta = 30^\circ, \text{ and } v_0 = 300 \text{ ft/s}$$


$$R(30^\circ) = \frac{1}{32} (300)^2 \sin 60^\circ \approx \underline{2,436 \text{ ft}}$$

$$e) \quad x = (v_0 \cos \theta)t = 2,436$$

$$t = \frac{2,436}{300 \cos 30^\circ} \approx \underline{9.38 \text{ sec}}$$

$$f) \quad y(x) = \frac{1}{2v_0^2 \cos^2 \theta} \left(\left(v_0^2 \sin 2\theta \right) x - gx^2 \right)$$

$$= 0.57735x - 0.000237x^2$$




g) **Given:** $g = 32 \text{ ft/s}^2$, $\theta = 52^\circ$, and $v_0 = 300 \text{ ft/s}$

$$R(30^\circ) = \frac{1}{32} (300)^2 \sin 104^\circ \approx 2729 \text{ ft}$$

$$h) \quad y(x) = \frac{1}{2v_0^2 \cos^2 \theta} \left(\left(v_0^2 \sin 2\theta \right) x - gx^2 \right)$$

$$= 1.2799x - 0.000469x^2$$



Exercise

Derive the system of differential equations describing the straight-line vertical motion of the coupled springs. Use the Laplace transform to solve the system when

$$k_1 = 1, k_2 = 1, k_3 = 1, m_1 = 1, m_2 = 1 \quad \text{and} \quad x_1(0) = 0, \quad x_1'(0) = -1, \quad x_2(0) = 0, \quad x_2'(0) = 1$$

Solution

$$x_1'' + 2x_1 - x_2 = 0$$

$$x_2'' + 2x_2 - x_1 = 0$$

$$m_1 x_1'' = -k_1 x_1 + k_2 (x_2 - x_1)$$

$$m_2 x_2'' = -k_2 (x_2 - x_1) - k_3 x_2$$

$$\begin{cases} s^2 X_1(s) - s x_1(0) - x_1'(0) + 2X_1(s) - X_2(s) = 0 \\ s^2 X_2(s) - s x_2(0) - x_2'(0) + 2X_2(s) - X_1(s) = 0 \end{cases}$$

$$\begin{cases} (s^2 + 2)X_1(s) - X_2(s) = -1 \\ -X_1(s) + (s^2 + 2)X_2(s) = 1 \end{cases}$$

$$\Delta = \begin{vmatrix} s^2 + 2 & -1 \\ -1 & s^2 + 2 \end{vmatrix} = (s^2 + 2)^2 - 1$$

$$\Delta_1 = \begin{vmatrix} -1 & -1 \\ 1 & s^2 + 2 \end{vmatrix} = -s^2 - 1 \quad \Delta_2 = \begin{vmatrix} s^2 + 2 & -1 \\ -1 & 1 \end{vmatrix} = s^2 + 1$$

$$\begin{aligned} X_1(s) &= \frac{-(s^2 + 1)}{s^4 + 4s^2 + 3} \\ &= -\frac{s^2 + 1}{(s^2 + 1)(s^2 + 3)} \end{aligned}$$

$$\mathcal{L}^{-1}\{X_1(s)\} = \mathcal{L}^{-1}\left\{-\frac{1}{s^2 + 3}\right\}$$

$$\underline{x_1(t) = -\sin\sqrt{3}t}$$

$$X_2(s) = \frac{s^2 + 1}{(s^2 + 1)(s^2 + 3)} = \frac{1}{s^2 + 3}$$

$$\mathcal{L}^{-1}\{X_2(s)\} = \mathcal{L}^{-1}\left\{-\frac{1}{s^2 + 3}\right\}$$

$$\underline{x_2(t) = -\sin\sqrt{3}t}$$

