

Section 3.6 – Alternating Series, Absolute and Conditional Convergence

A series in which the terms are alternately positive and negative is an alternating series.

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \dots + \frac{(-1)^{n+1}}{n} + \dots$$
$$-2 + 1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{8} + \dots + \frac{(-1)^n 4}{2^n} + \dots$$

Theorem – The Alternating Series Test (Leibniz's Test)

The series $\sum_{n=1}^{\infty} (-1)^{n+1} u_n = u_1 - u_2 + u_3 - u_4 + \dots$

Converges if all three of the following conditions are satisfied:

1. The u_n 's are all positive.
2. The positive u_n 's are (eventually) non-increasing: $u_n \geq u_{n+1}$ for all $n \geq N$, for some integer N .
3. $u_n \rightarrow 0$

Example

The alternating harmonic series $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n} = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$

Solution

1. $\frac{1}{n} > 0$
2. $n < n+1 \rightarrow \frac{1}{n} > \frac{1}{n+1}$
3. $\frac{1}{n} \rightarrow 0$

Therefore, the series converges.

Theorem – The Alternating Series Estimation Theorem

IF the alternating series $\sum_{n=1}^{\infty} (-1)^{n+1} u_n$ satisfies the three conditions, then for $n \geq N$

$$s_n = u_1 - u_2 + u_3 - \cdots + (-1)^{n+1} u_n$$

Approximates the sum L of the series with an error whose absolute value is less than u_{n+1} , the absolute value of the first unused term. Furthermore, the sum L lies between any two successive partial sums s_n and s_{n+1} and the remainder, $L - s_n$, has the same sign as the first unused term.

Absolute and Conditional Convergence

Definition

A series $\sum a_n$ **converges absolutely** (is **absolutely convergent**) if the corresponding series of absolute values, $\sum |a_n|$, converges.

Definition

A series converges but does not converge absolutely **converges conditionally**.

Theorem

If $\sum_{n=1}^{\infty} |a_n|$ converges, then $\sum_{n=1}^{\infty} a_n$ converges.

Example

For $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n^2} = 1 - \frac{1}{4} + \frac{1}{9} - \frac{1}{16} + \cdots$ the corresponding series of absolute values is the convergent series

$$\sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n^2} = 1 + \frac{1}{4} + \frac{1}{9} + \frac{1}{16} + \cdots$$

The original series converges because it converges absolutely.

Example

For $\sum_{n=1}^{\infty} \frac{\sin n}{n^2} = \frac{\sin 1}{1} + \frac{\sin 2}{4} + \frac{\sin 3}{9} + \dots$, which contains both positive and negative terms, the corresponding series of absolute values is

$$\sum_{n=1}^{\infty} \left| \frac{\sin n}{n^2} \right| = \frac{|\sin 1|}{1} + \frac{|\sin 2|}{4} + \frac{|\sin 3|}{9} + \dots$$

Which converges by comparison with $\sum_{n=1}^{\infty} \frac{1}{n^2}$ because $|\sin n| \leq 1$ for every n .

The original series converges absolutely; therefore, it converges.

Rearranging Series

Theorem

If $\sum_{n=1}^{\infty} a_n$ converges absolutely, and $b_1, b_2, \dots, b_n, \dots$ is any arrangement of the sequence $\{a_n\}$, then

$$\sum b_n \text{ converges absolutely and } \sum_{n=1}^{\infty} b_n = \sum_{n=1}^{\infty} a_n$$

Exercises Section 3.6 – Alternating Series, Absolute and Conditional Convergence

Determine if the alternating series converges or diverges

1. $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{\sqrt{n}}$

9. $\sum_{n=1}^{\infty} \frac{(-1)^n}{3^n}$

17. $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{n+1}{\ln(n+1)}$

2. $\sum_{n=2}^{\infty} (-1)^n \frac{4}{(\ln n)^2}$

10. $\sum_{n=1}^{\infty} \frac{(-1)^n}{e^n}$

18. $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{\ln(n+1)}{n+1}$

3. $\sum_{n=1}^{\infty} (-1)^n \frac{n}{n^2+1}$

11. $\sum_{n=1}^{\infty} (-1)^n \frac{5n-1}{4n+1}$

19. $\sum_{n=1}^{\infty} \sin \frac{(2n-1)\pi}{2}$

4. $\sum_{n=1}^{\infty} (-1)^n \frac{n^2+5}{n^2+4}$

12. $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{n}{n^2+5}$

20. $\sum_{n=1}^{\infty} \frac{1}{n} \cos n\pi$

5. $\sum_{n=1}^{\infty} (-1)^{n+1} \left(\frac{n}{10}\right)^n$

13. $\sum_{n=1}^{\infty} (-1)^n \frac{n}{\ln(n+1)}$

21. $\sum_{n=0}^{\infty} \frac{(-1)^n}{n!}$

6. $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{\sqrt{n}+1}{n+1}$

14. $\sum_{n=1}^{\infty} \frac{(-1)^n}{\ln(n+1)}$

22. $\sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!}$

7. $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n+1}$

15. $\sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n}}$

23. $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{\sqrt{n}}{n+2}$

8. $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{n}{3n+2}$

16. $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{n^2}{n^2+4}$

24. $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{\sqrt{n}}{\sqrt[3]{n}}$

Determine if the series converge absolutely or conditionally, or diverges

25. $\sum_{n=1}^{\infty} (-1)^{n+1} (0.1)^n$

28. $\sum_{n=1}^{\infty} (-1)^n \frac{\sin n}{n^2}$

31. $\sum_{n=1}^{\infty} \frac{n \cos(n\pi)}{2^n}$

26. $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{(0.1)^n}{n}$

29. $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{3+n}{5+n}$

32. $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{2n-1}$

27. $\sum_{n=1}^{\infty} \frac{(-1)^n}{1+\sqrt{n}}$

30. $\sum_{n=2}^{\infty} (-1)^{n+1} \frac{1}{n \ln n}$

33. $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^2+2n+1}$

$$\begin{array}{lll}
34. \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{\sqrt{n}} & 42. \sum_{n=10}^{\infty} \frac{\sin\left(n + \frac{1}{2}\right)\pi}{\ln \ln n} & 51. \sum_{n=2}^{\infty} \frac{(-1)^{n+1}}{n \ln n} \\
35. \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2 + \ln n} & 43. \sum_{n=1}^{\infty} \frac{(-1)^n}{2^n} & 52. \sum_{n=0}^{\infty} (-1)^n e^{-n^2} \\
36. \sum_{n=0}^{\infty} \frac{(-1)^n (n^2 - 1)}{n^2 + 1} & 44. \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^2} & 53. \sum_{n=2}^{\infty} (-1)^n \frac{n}{n^3 - 5} \\
37. \sum_{n=1}^{\infty} \frac{\cos(n\pi)}{(n+1)\ln(n+1)} & 45. \sum_{n=1}^{\infty} \frac{(-1)^n}{n!} & 54. \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^{4/3}} \\
38. \sum_{n=1}^{\infty} \frac{(-2)^n}{n!} & 46. \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n+3} & 55. \sum_{n=0}^{\infty} \frac{(-1)^{n+1}}{\sqrt{n+4}} \\
39. \sum_{n=1}^{\infty} \frac{(-1)^n}{n\pi^n} & 47. \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{\sqrt{n}} & 56. \sum_{n=0}^{\infty} \frac{\cos n\pi}{n+1} \\
40. \sum_{n=1}^{\infty} \frac{100\cos(n\pi)}{2n+3} & 48. \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n\sqrt{n}} & 57. \sum_{n=1}^{\infty} (-1)^{n+1} \arctan n \\
41. \sum_{n=1}^{\infty} \frac{n!}{(-100)^n} & 49. \sum_{n=1}^{\infty} (-1)^{n+1} \frac{n^2}{(n+1)^2} & 58. \sum_{n=1}^{\infty} \frac{\cos n\pi}{n^2} \\
& 50. \sum_{n=1}^{\infty} (-1)^{n+1} \frac{2n+3}{n+10} & 59. \sum_{n=1}^{\infty} \frac{\sin\left[(n-1)\frac{\pi}{2}\right]}{n}
\end{array}$$

For what values of x does the series converge absolutely? Converge conditionally? Diverge?

$$\begin{array}{lll}
60. \sum_{n=1}^{\infty} \frac{(x-5)^n}{n 2^n} & 63. \sum_{n=0}^{\infty} (-1)^n \frac{(x-1)^n}{2n+3} & 66. \sum_{n=1}^{\infty} \frac{(4x+1)^n}{n^3} \\
61. \sum_{n=1}^{\infty} \frac{(x-2)^n}{n^2 2^{2n}} & 64. \sum_{n=1}^{\infty} \frac{1}{2n-1} \left(\frac{3x+2}{-5}\right)^n & 67. \sum_{n=1}^{\infty} \frac{(2x+3)^n}{n^{1/3} 4^n} \\
62. \sum_{n=1}^{\infty} (n+1)^2 \left(\frac{x}{x+2}\right)^n & 65. \sum_{n=2}^{\infty} \frac{x^n}{2^n \ln n} & 68. \sum_{n=1}^{\infty} \frac{1}{n} \left(1 + \frac{1}{x}\right)^n
\end{array}$$

Use any method to determine if the series converges or diverges.

$$69. \sum_{n=1}^{\infty} (-1)^n \frac{\sqrt{n}}{n+1}$$

$$70. \sum_{n=2}^{\infty} \frac{(-1)^n}{n \ln n}$$

$$71. \sum_{n=1}^{\infty} (-1)^{n+1} \frac{5}{n}$$

$$72. \sum_{n=1}^{\infty} (-1)^{n+1} \frac{3^{n-1}}{n!}$$

$$73. \sum_{n=1}^{\infty} (-1)^{n+1} \frac{3^n}{n2^n}$$

$$74. \sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n}$$

$$75. \sum_{n=1}^{\infty} \frac{n}{(-2)^{n-1}}$$

$$76. \sum_{n=1}^{\infty} \frac{10}{n^{3/2}}$$

$$77. \sum_{n=1}^{\infty} \frac{2}{n^2 + 5}$$

$$78. \sum_{n=1}^{\infty} \frac{3^n}{n^2}$$

$$79. \sum_{n=1}^{\infty} \frac{1}{2^n + 1}$$

$$80. \sum_{n=1}^{\infty} 5\left(\frac{7}{8}\right)^n$$

$$81. \sum_{n=1}^{\infty} \frac{3n^2}{2n^2 + 1}$$

$$82. \sum_{n=1}^{\infty} 100e^{-n/2}$$

$$83. \sum_{n=1}^{\infty} \frac{(-1)^n}{n+4}$$

$$84. \sum_{n=1}^{\infty} (-1)^{n+1} \frac{4}{3n^2 - 1}$$

$$85. \sum_{n=2}^{\infty} \frac{\ln n}{n}$$

$$86. \sum_{k=1}^{\infty} \frac{2}{k^{3/2}}$$

$$87. \sum_{k=1}^{\infty} k^{-2/3}$$

$$88. \sum_{k=1}^{\infty} \frac{2k^2 + 1}{\sqrt{k^3 + 2}}$$

$$89. \sum_{k=1}^{\infty} \frac{2^k}{e^k}$$

$$90. \sum_{k=1}^{\infty} \left(\frac{k}{k+3}\right)^{2k}$$

$$91. \sum_{k=1}^{\infty} \frac{2^k k!}{k^k}$$

$$92. \sum_{k=1}^{\infty} \frac{1}{\sqrt{k} \sqrt{k+1}}$$

$$93. \sum_{k=1}^{\infty} \frac{3}{2 + e^k}$$

$$94. \sum_{k=1}^{\infty} k \sin \frac{1}{k}$$

$$95. \sum_{k=1}^{\infty} \frac{\sqrt[k]{k}}{k^3}$$

$$96. \sum_{k=1}^{\infty} \frac{1}{1 + \ln k}$$

$$97. \sum_{k=1}^{\infty} k^5 e^{-k}$$

$$98. \sum_{k=4}^{\infty} \frac{1}{k^2 - 10}$$

$$99. \sum_{k=1}^{\infty} \frac{\ln k^2}{k^2}$$

$$100. \sum_{k=1}^{\infty} k e^{-k}$$

$$101. \sum_{k=0}^{\infty} \frac{2 \cdot 4^k}{(2k+1)!}$$

$$102. \sum_{k=0}^{\infty} \frac{9^k}{(2k)!}$$

$$103. \sum_{k=1}^{\infty} \frac{\coth k}{k}$$

$$104. \sum_{k=1}^{\infty} \frac{1}{\sinh k}$$

$$105. \sum_{k=1}^{\infty} \tanh k$$

$$106. \sum_{k=0}^{\infty} \operatorname{sech} k$$

$$107. \sum_{k=2}^{\infty} \frac{(-1)^k}{k^2 - 1}$$

$$108. \sum_{k=1}^{\infty} (-1)^{k+1} \frac{k^2 + 4}{2k^2 + 1}$$

$$109. \sum_{k=1}^{\infty} (-1)^k k e^{-k}$$

$$110. \sum_{k=1}^{\infty} \frac{(-1)^k}{\sqrt{k^2 + 1}}$$

$$111. \sum_{k=1}^{\infty} (-1)^{k+1} \frac{10^k}{k!}$$

$$112. \sum_{k=1}^{\infty} \frac{(-2)^{k+1}}{k^2}$$

$$113. \sum_{k=0}^{\infty} \frac{(-1)^k}{e^k + e^{-k}}$$

$$114. \sum_{k=2}^{\infty} \frac{(-1)^k}{k \ln k}$$

$$115. \sum_{k=2}^{\infty} (-1)^k \frac{\ln k}{k^2}$$

$$116. \sum_{k=2}^{\infty} \frac{(-1)^k}{k \ln^2 k}$$

$$117. \sum_{k=2}^{\infty} \frac{(-1)^k}{\ln k}$$

$$118. \sum_{k=2}^{\infty} 3e^{-k}$$

$$119. \sum_{k=1}^{\infty} \frac{2^k}{e^k - 1}$$

$$120. \sum_{k=1}^{\infty} (-1)^{k+1} \frac{e^k}{(k+1)!}$$

$$121. \sum_{k=1}^{\infty} \frac{k}{(k^2 + 1)^3}$$

$$122. \sum_{k=2}^{\infty} \frac{k^e}{k^\pi}$$

$$123. \sum_{k=3}^{\infty} \frac{1}{(k-2)^4}$$

$$124. \sum_{k=1}^{\infty} \left(\frac{2k}{k+1} \right)^k$$

$$125. \sum_{k=1}^{\infty} \frac{k^2}{2^k}$$

$$126. \sum_{k=1}^{\infty} \frac{k^2 - 1}{k^3 + 4}$$

$$127. \sum_{k=1}^{\infty} \frac{1}{k^2} \sin \frac{\pi}{2k}$$

128. Use a Riemann sum argument to show that $\ln n! \geq \int_1^n \ln t \, dt = n \ln n - n + 1$

Then for what values of x does the series $\sum_{n=1}^{\infty} \frac{n! x^n}{n^n}$ converge absolutely? Converge conditionally?

Diverge? (Use the ratio test first)

129. Let S_n be the n th partial sum of $\sum_{k=1}^{\infty} a_k = 8$. Find the $\lim_{k \rightarrow \infty} a_k$ and $\lim_{n \rightarrow \infty} S_n$

- 130.** It can be proved that if a series converges absolutely, then its terms may be summed in any order without changing the value of the series. However, if a series converges conditionally, then the value of the series depends on the order of summation. For example, the (conditionally convergent) alternating harmonic series has the value

$$1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots = \ln 2$$

Show that by rearranging the terms (so the sign pattern is $++-$),

$$1 + \frac{1}{3} - \frac{1}{2} + \frac{1}{5} + \frac{1}{7} - \frac{1}{4} + \dots = \frac{3}{2} \ln 2$$

- 131.** A crew of workers is constructing a tunnel through a mountain. Understandably, the rate of construction decreases because rocks and earth must be removed a greater distance as the tunnel gets longer. Suppose that each week the crew digs 0.95 of the distance it dug the previous week. In the first week, the crew constructed 100 m of tunnel.
- How far does the crew dig in 10 weeks? 20 weeks? N weeks?
 - What is the longest tunnel the crew can build at this rate?
 - The time required to dig 100 m increases by 10% each week, starting with 1 week to dig the first 100 m. Can the crew complete a 1.5 km tunnel in 10 weeks? Explain.

- 132.** Consider the alternating series

$$\sum_{k=1}^{\infty} (-1)^{k+1} a_k \quad \text{where} \quad a_k = \begin{cases} \frac{4}{k+1} & \text{if } k \text{ is odd} \\ \frac{2}{k} & \text{if } k \text{ is even} \end{cases}$$


- Write out the first ten terms of the series, group them in pairs, and show that the even partial sums of the series form the (divergent) harmonic series.
 - Show that $\lim_{k \rightarrow \infty} a_k = 0$
 - Explain why the series diverges even though the terms of the series approach zero.
- 133.** The concentration in the blood resulting from a single dose of a drug normally decreases with time as the drug is eliminated from the body. Doses may therefore need to be repeated periodically to keep the concentration from dropping below some particular level. One model for the effect of repeated doses gives the residual concentration just before the $(n+1)$ st dose as

$$R_n = C_0 e^{-kt_0} + C_0 e^{-2kt_0} + \dots + C_0 e^{-nkt_0}$$

Where C_0 = the change in concentration achievable by a single dose (mg / mL),


k = the elimination constant (h^{-1}), and

t_0 = time between doses (h).



- a) Write R_n in closed form as a single fraction, and find $R = \lim_{n \rightarrow \infty} R_n$
- b) Calculate R_1 and R_{10} for $C_0 = 1 \text{ mg/mL}$, $k = 0.1 \text{ h}^{-1}$, and $t_0 = 10 \text{ h}$. How good as estimate of R is R_{10}
- c) If $k = 0.01 \text{ h}^{-1}$ and $t_0 = 10 \text{ h}$, find the smallest n such that $R_n > \frac{1}{2}R$.

134. If a drug is known to be ineffective below a concentration C_L and harmful above some higher concentration C_H , one needs to find values of C_0 and t_0 that will produce a concentration that is safe (not above C_H) but effective (not below C_L).



We want to find values for C_0 and t_0 for which

$$R = C_L \quad \text{and} \quad C_0 + R = C_H$$

Thus, $C_0 = C_H - C_L$. The resulting equation simplifies to

$$t_0 = \frac{1}{k} \ln \frac{C_H}{C_L}$$

To reach an effective level rapidly, one might administer a “loading” dose that would produce a concentration of $C_H \text{ mg/mL}$. This could be followed every t_0 hours by a dose that raises the concentration by $C_0 = C_H - C_L \text{ mg/mL}$.

- a) Verify the preceding equation for t_0 .
- b) If $k = 0.05 \text{ h}^{-1}$ and the highest safe concentration is e times the lowest effective concentration, find the length of time between doses that will assure safe and effective concentrations.
- c) Given $C_H = 2 \text{ mg / mL}$, $C_L = 0.5 \text{ mg / mL}$, and $k = 0.02 \text{ h}^{-1}$, determine a scheme for administering the drug.
- d) Suppose that $k = 0.2 \text{ h}^{-1}$ and the smallest effective concentration is 0.03 mg/mL . A single dose that produces a concentration of 0.1 mg/mL is administered. About how long will the drug remain effective?