

Solution **Section 3.5 – Least Squares Analysis**

Exercise

Find the equation of the line that best fits the given points in the least-squares sense and find the error.

$$\{(0, 2), (1, 2), (2, 0)\}$$

Solution

$$\{(0, 2), (1, 2), (2, 0)\}$$

Let $y = mx + b$ be the equation of the line that best fits the given points. Then

$$\begin{bmatrix} 0 & 1 \\ 1 & 1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} m \\ b \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \\ 0 \end{bmatrix}$$

$$\text{where } A = \begin{bmatrix} 0 & 1 \\ 1 & 1 \\ 2 & 1 \end{bmatrix} \quad \vec{x} = \begin{bmatrix} m \\ b \end{bmatrix} \quad \vec{y} = \begin{bmatrix} 2 \\ 2 \\ 0 \end{bmatrix}$$

The normal equation formula: $A^T A \vec{x} = A^T \vec{y}$

$$\begin{bmatrix} 0 & 1 & 2 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} m \\ b \end{bmatrix} = \begin{bmatrix} 0 & 1 & 2 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 2 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} 5 & 3 \\ 3 & 3 \end{bmatrix} \begin{bmatrix} m \\ b \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$$

$$m = \frac{\begin{vmatrix} 2 & 3 \\ 4 & 3 \end{vmatrix}}{\begin{vmatrix} 5 & 3 \\ 3 & 3 \end{vmatrix}}$$


$$= \frac{-6}{6}$$

$$= -1$$


$$b = \frac{\begin{vmatrix} 5 & 2 \\ 3 & 4 \end{vmatrix}}{6}$$

$$= \frac{7}{3}$$

$$\text{Thus, } y = -x + \frac{7}{3}$$



$$\begin{aligned}
 A\vec{x} &= \begin{pmatrix} 0 & 1 \\ 1 & 1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} -1 \\ \frac{7}{3} \end{pmatrix} \\
 &= \begin{pmatrix} \frac{7}{3} \\ \frac{4}{3} \\ \frac{1}{3} \end{pmatrix} \\
 \vec{y} - A\vec{x} &= \begin{pmatrix} 2 \\ 2 \\ 0 \end{pmatrix} - \begin{pmatrix} \frac{7}{3} \\ \frac{4}{3} \\ \frac{1}{3} \end{pmatrix} \\
 &= \begin{pmatrix} \frac{1}{3} \\ \frac{2}{3} \\ -\frac{1}{3} \end{pmatrix}
 \end{aligned}$$



Error:

$$\begin{aligned}
 \|\vec{y} - A\vec{x}\| &= \sqrt{\frac{1}{9} + \frac{4}{9} + \frac{1}{9}} \\
 &= \frac{\sqrt{6}}{3} \\
 &\approx 0.8164966
 \end{aligned}$$

The **second order** equation:

$$y = -x^2 + x + 2$$

Error = 0.00000

The **third order** equation:

$$y = -0.1250x^3 - 0.1250x^2 + 0.25x + 2$$

Error = 2.01556

Exercise

Find the equation of the line that best fits the given points in the least-squares sense and find the error.

$$\{(1, 5), (2, 4), (3, 1), (4, 1), (5, -1)\}$$

Solution

$$\{(1, 5), (2, 4), (3, 1), (4, 1), (5, -1)\}$$

Let $y = mx + b$ be the equation of the line that best fits the given points. Then

$$\begin{bmatrix} 1 & 1 \\ 2 & 1 \\ 3 & 1 \\ 4 & 1 \\ 5 & 1 \end{bmatrix} \begin{bmatrix} m \\ b \end{bmatrix} = \begin{bmatrix} 5 \\ 4 \\ 1 \\ 1 \\ -1 \end{bmatrix} \quad \text{where } A = \begin{bmatrix} 1 & 1 \\ 2 & 1 \\ 3 & 1 \\ 4 & 1 \\ 5 & 1 \end{bmatrix} \quad \vec{x} = \begin{bmatrix} m \\ b \end{bmatrix} \quad \vec{y} = \begin{bmatrix} 5 \\ 4 \\ 1 \\ 1 \\ -1 \end{bmatrix}$$

The normal equation: $A^T A \vec{x} = A^T \vec{y}$


$$\begin{bmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 2 & 1 \\ 3 & 1 \\ 4 & 1 \\ 5 & 1 \end{bmatrix} \begin{bmatrix} m \\ b \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 1 & 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 5 \\ 4 \\ 1 \\ 1 \\ -1 \end{bmatrix}$$

$$\begin{bmatrix} 55 & 15 \\ 15 & 5 \end{bmatrix} \begin{bmatrix} m \\ b \end{bmatrix} = \begin{bmatrix} 15 \\ 10 \end{bmatrix}$$

$$m = \frac{\begin{vmatrix} 15 & 15 \\ 10 & 5 \end{vmatrix}}{\begin{vmatrix} 55 & 15 \\ 15 & 5 \end{vmatrix}} = \frac{-75}{50} = -\frac{3}{2}$$

$$b = \frac{\begin{vmatrix} 55 & 15 \\ 15 & 10 \end{vmatrix}}{50} = \frac{325}{50} = \frac{13}{2}$$

$$\text{Thus, } \underline{y = -\frac{3}{2}x + \frac{13}{2}} \quad \text{or} \quad \underline{y = -1.5x + 6.5}$$




$$A\vec{x} = \begin{pmatrix} 1 & 1 \\ 2 & 1 \\ 3 & 1 \\ 4 & 1 \\ 5 & 1 \end{pmatrix} \begin{pmatrix} -\frac{3}{2} \\ \frac{13}{2} \end{pmatrix}$$

$$= \begin{pmatrix} 5 \\ \frac{7}{2} \\ 2 \\ \frac{1}{2} \\ -1 \end{pmatrix}$$

$$\vec{y} - A\vec{x} = \begin{pmatrix} 5 \\ 4 \\ 1 \\ 1 \\ -1 \end{pmatrix} - \begin{pmatrix} 5 \\ \frac{7}{2} \\ 2 \\ \frac{1}{2} \\ -1 \end{pmatrix}$$

$$= \begin{pmatrix} 0 \\ \frac{1}{2} \\ -1 \\ \frac{1}{2} \\ 0 \end{pmatrix}$$



Error:

$$\begin{aligned} \|\vec{y} - A\vec{x}\| &= \sqrt{\frac{1}{4} + 1 + \frac{1}{4}} \\ &= \frac{\sqrt{6}}{2} \\ &\approx 1.224745 \end{aligned}$$

The **second order** equation:

$$y = 0.07143x^2 - 1.92857x + 7$$

Error = 1.19523

The **third order** equation:

$$y = 0.0x^3 + 0.07143x^2 - 1.92857x + 7$$

Error = 1.19523

Exercise

Find the equation of the line that best fits the given points in the least-squares sense and find the error.

$$\{(0, 1), (1, 3), (2, 4), (3, 4)\}$$

Solution

Let $y = mx + b$ be the equation of the line that best fits the given points. Then

$$\begin{pmatrix} 0 & 1 \\ 1 & 1 \\ 2 & 1 \\ 3 & 1 \end{pmatrix} \begin{pmatrix} m \\ b \end{pmatrix} = \begin{pmatrix} 1 \\ 3 \\ 4 \\ 4 \end{pmatrix}$$

$$\text{where } A = \begin{pmatrix} 0 & 1 \\ 1 & 1 \\ 2 & 1 \\ 3 & 1 \end{pmatrix} \quad \vec{x} = \begin{pmatrix} m \\ b \end{pmatrix} \quad \vec{y} = \begin{pmatrix} 1 \\ 3 \\ 4 \\ 4 \end{pmatrix}$$

The normal equation: $A^T A \vec{x} = A^T \vec{y}$

$$\begin{pmatrix} 0 & 1 & 2 & 3 \\ 1 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 1 \\ 2 & 1 \\ 3 & 1 \end{pmatrix} \begin{pmatrix} m \\ b \end{pmatrix} = \begin{pmatrix} 0 & 1 & 2 & 3 \\ 1 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 3 \\ 4 \\ 4 \end{pmatrix}$$


$$\begin{pmatrix} 14 & 6 \\ 6 & 4 \end{pmatrix} \begin{pmatrix} m \\ b \end{pmatrix} = \begin{pmatrix} 23 \\ 12 \end{pmatrix}$$

$$\begin{aligned} \begin{pmatrix} m \\ b \end{pmatrix} &= \frac{1}{20} \begin{pmatrix} 4 & -6 \\ -6 & 14 \end{pmatrix} \begin{pmatrix} 23 \\ 12 \end{pmatrix} & X = A^{-1}B \\ &= \frac{1}{20} \begin{pmatrix} 20 \\ 30 \end{pmatrix} \\ &= \begin{pmatrix} 1 \\ \frac{3}{2} \end{pmatrix} \end{aligned}$$

We have: $m = 1$ and $b = \frac{3}{2}$.

Thus, $y = x + \frac{3}{2}$

$$A\vec{x} = \begin{pmatrix} 0 & 1 \\ 1 & 1 \\ 2 & 1 \\ 3 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ \frac{3}{2} \end{pmatrix}$$



$$= \begin{pmatrix} \frac{3}{2} \\ \frac{5}{2} \\ \frac{7}{2} \\ \frac{9}{2} \end{pmatrix}$$

$$\vec{y} - A\vec{x} = \begin{pmatrix} 1 \\ 3 \\ 4 \\ 4 \end{pmatrix} - \begin{pmatrix} \frac{3}{2} \\ \frac{5}{2} \\ \frac{7}{2} \\ \frac{9}{2} \end{pmatrix}$$

$$= \begin{pmatrix} -\frac{1}{2} \\ \frac{1}{2} \\ \frac{1}{2} \\ -\frac{1}{2} \end{pmatrix}$$

Error:

$$\|\vec{y} - A\vec{x}\| = \sqrt{4\left(\frac{1}{4}\right)}$$

$$= 1$$

The **second order** equation:


$$y = -0.50x^2 + 2.5x + 1.0$$

Error = 0.0000

The **third order** equation:

$$y = 0.0x^3 - 0.5x^2 + 2.5x + 1$$

Error = 0.00000



Exercise

Find the equation of the line that best fits the given points in the least-squares sense and find the error.

$$\{(-2, 0), (-1, 0), (0, 1), (1, 3), (2, 5)\}$$

Solution

Let $y = mx + b$ be the equation of the line that best fits the given points. Then

$$\begin{pmatrix} -2 & 1 \\ -1 & 1 \\ 0 & 1 \\ 1 & 1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} m \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 3 \\ 5 \end{pmatrix}$$

$$\text{where } A = \begin{pmatrix} -2 & 1 \\ -1 & 1 \\ 0 & 1 \\ 1 & 1 \\ 2 & 1 \end{pmatrix} \quad \vec{x} = \begin{pmatrix} m \\ b \end{pmatrix} \quad \vec{y} = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 3 \\ 5 \end{pmatrix}$$

The normal equation: $A^T A \vec{x} = A^T \vec{y}$


$$\begin{pmatrix} -2 & -1 & 0 & 1 & 2 \\ 1 & 1 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} -2 & 1 \\ -1 & 1 \\ 0 & 1 \\ 1 & 1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} m \\ b \end{pmatrix} = \begin{pmatrix} -2 & -1 & 0 & 1 & 2 \\ 1 & 1 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 0 \\ 0 \\ 1 \\ 3 \\ 5 \end{pmatrix}$$

$$\begin{pmatrix} 10 & 0 \\ 0 & 5 \end{pmatrix} \begin{pmatrix} m \\ b \end{pmatrix} = \begin{pmatrix} 13 \\ 9 \end{pmatrix}$$

$$\begin{pmatrix} m \\ b \end{pmatrix} = \frac{1}{50} \begin{pmatrix} 5 & 0 \\ 0 & 10 \end{pmatrix} \begin{pmatrix} 13 \\ 9 \end{pmatrix} = \begin{pmatrix} \frac{13}{10} \\ \frac{9}{5} \end{pmatrix}$$


We have: $m = 1.3$ and $b = 1.8$

$$\text{Thus, } \underline{y = \frac{13}{10}x + \frac{9}{5}}$$



$$A\vec{x} = \begin{pmatrix} -2 & 1 \\ -1 & 1 \\ 0 & 1 \\ 1 & 1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} \frac{13}{10} \\ \frac{9}{5} \end{pmatrix} = \begin{pmatrix} -\frac{4}{5} \\ \frac{1}{2} \\ \frac{9}{5} \\ \frac{31}{10} \\ \frac{22}{5} \end{pmatrix}$$

$$\vec{y} - A\vec{x} = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 3 \\ 5 \end{pmatrix} - \begin{pmatrix} -\frac{4}{5} \\ \frac{1}{2} \\ \frac{9}{5} \\ \frac{31}{10} \\ \frac{22}{5} \end{pmatrix} = \begin{pmatrix} \frac{4}{5} \\ \frac{1}{2} \\ -\frac{4}{5} \\ -\frac{1}{10} \\ \frac{3}{5} \end{pmatrix}$$



Error: $\|\vec{y} - A\vec{x}\| = \sqrt{\frac{16}{25} + \frac{1}{4} + \frac{16}{25} + \frac{1}{100} + \frac{9}{25}}$

$$= \sqrt{\frac{41}{25} + \frac{26}{100}}$$

$$= \frac{\sqrt{190}}{10}$$

$$\approx 1.37840$$

The *second order* equation:

$$y = 0.35714x^2 + 1.30x + 1.08571$$

Error = 0.33806

The *third order* equation:

$$y = -0.08333x^3 + 0.35714x^2 + 1.58333x + 1.08571$$

Error = 0.11952

Exercise

Find the equation of the line that best fits the given points in the least-squares sense and find the error.

$$\{(2, 3), (3, 2), (5, 1), (6, 0)\}$$

Solution

Let $y = mx + b$ be the equation of the line that best fits the given points. Then

$$\begin{pmatrix} 2 & 1 \\ 3 & 1 \\ 5 & 1 \\ 6 & 1 \end{pmatrix} \begin{pmatrix} m \\ b \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ 1 \\ 0 \end{pmatrix}$$

$$\text{where } A = \begin{pmatrix} 2 & 1 \\ 3 & 1 \\ 5 & 1 \\ 6 & 1 \end{pmatrix} \quad \vec{x} = \begin{pmatrix} m \\ b \end{pmatrix} \quad \vec{y} = \begin{pmatrix} 3 \\ 2 \\ 1 \\ 0 \end{pmatrix}$$

The normal equation: $A^T A \vec{x} = A^T \vec{y}$

$$\begin{pmatrix} 2 & 3 & 5 & 6 \\ 1 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 2 & 1 \\ 3 & 1 \\ 5 & 1 \\ 6 & 1 \end{pmatrix} \begin{pmatrix} m \\ b \end{pmatrix} = \begin{pmatrix} 2 & 3 & 5 & 6 \\ 1 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 3 \\ 2 \\ 1 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 74 & 16 \\ 16 & 4 \end{pmatrix} \begin{pmatrix} m \\ b \end{pmatrix} = \begin{pmatrix} 17 \\ 6 \end{pmatrix}$$

$$\Delta = \begin{vmatrix} 74 & 16 \\ 16 & 4 \end{vmatrix} = 40 \quad \Delta_m = \begin{vmatrix} 17 & 16 \\ 6 & 4 \end{vmatrix} = -28 \quad \Delta_b = \begin{vmatrix} 74 & 17 \\ 16 & 6 \end{vmatrix} = 172$$

$$m = -\frac{28}{40} = -\frac{7}{10}$$

$$b = \frac{172}{40} = \frac{43}{10}$$


$$\text{Thus, } y = -\frac{7}{10}x + \frac{43}{10}$$

$$A\vec{x} = \begin{pmatrix} 2 & 1 \\ 3 & 1 \\ 5 & 1 \\ 6 & 1 \end{pmatrix} \begin{pmatrix} -\frac{7}{10} \\ \frac{43}{10} \end{pmatrix}$$

$$= \begin{pmatrix} \frac{29}{10} \\ \frac{22}{10} \\ \frac{8}{10} \\ \frac{1}{10} \end{pmatrix}$$

$$\vec{y} - A\vec{x} = \begin{pmatrix} 3 \\ 2 \\ 1 \\ 0 \end{pmatrix} - \begin{pmatrix} \frac{29}{10} \\ \frac{22}{10} \\ \frac{8}{10} \\ \frac{1}{10} \end{pmatrix}$$

$$= \begin{pmatrix} \frac{1}{10} \\ -\frac{1}{5} \\ \frac{1}{5} \\ -\frac{1}{10} \end{pmatrix}$$



Error:

$$\begin{aligned} \|\vec{y} - A\vec{x}\| &= \sqrt{2\left(\frac{1}{100} + \frac{1}{25}\right)} \\ &= \frac{\sqrt{10}}{10} \\ &= 0.31623 \end{aligned}$$

The **second order** equation:

$$y = 0.0x^2 - 0.7x + 4.3$$

Error = 0.31623

The **third order** equation:

$$y = -0.08333x^3 + x^2 - 4.41667x + 8.5$$

Error = 0.00000

Exercise

Find the equation of the line that best fits the given points in the least-squares sense and find the error.

$$\{(-1, 0), (0, 1), (1, 2), (2, 4)\}$$

Solution

Let $y = mx + b$ be the equation of the line that best fits the given points. Then

$$\begin{pmatrix} -1 & 1 \\ 0 & 1 \\ 1 & 1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} m \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \\ 2 \\ 4 \end{pmatrix}$$

$$\text{where } A = \begin{pmatrix} -1 & 1 \\ 0 & 1 \\ 1 & 1 \\ 2 & 1 \end{pmatrix} \quad \vec{x} = \begin{pmatrix} m \\ b \end{pmatrix} \quad \vec{y} = \begin{pmatrix} 0 \\ 1 \\ 2 \\ 4 \end{pmatrix}$$

The normal equation: $A^T A \vec{x} = A^T \vec{y}$

$$\begin{pmatrix} -1 & 0 & 1 & 2 \\ 0 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} -1 & 1 \\ 0 & 1 \\ 1 & 1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} m \\ b \end{pmatrix} = \begin{pmatrix} -1 & 0 & 1 & 2 \\ 0 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \\ 2 \\ 4 \end{pmatrix}$$

$$\begin{pmatrix} 6 & 2 \\ 2 & 4 \end{pmatrix} \begin{pmatrix} m \\ b \end{pmatrix} = \begin{pmatrix} 10 \\ 7 \end{pmatrix}$$

$$\Delta = \begin{vmatrix} 6 & 2 \\ 2 & 4 \end{vmatrix} = 20 \quad \Delta_m = \begin{vmatrix} 10 & 2 \\ 7 & 4 \end{vmatrix} = 26 \quad \Delta_b = \begin{vmatrix} 6 & 10 \\ 2 & 7 \end{vmatrix} = 22$$

$$m = \frac{26}{20} = \frac{13}{10}$$

$$b = \frac{22}{20} = \frac{11}{10}$$


$$\text{Thus, } y = \frac{13}{10}x + \frac{11}{10}$$

$$A\vec{x} = \begin{pmatrix} -1 & 1 \\ 0 & 1 \\ 1 & 1 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} \frac{13}{10} \\ \frac{11}{10} \end{pmatrix}$$

$$= \begin{pmatrix} -\frac{1}{5} \\ \frac{11}{10} \\ \frac{12}{5} \\ \frac{37}{10} \end{pmatrix}$$

$$\vec{y} - A\vec{x} = \begin{pmatrix} 0 \\ 1 \\ 2 \\ 4 \end{pmatrix} - \begin{pmatrix} -\frac{1}{5} \\ \frac{11}{10} \\ \frac{12}{5} \\ \frac{37}{10} \end{pmatrix}$$

$$= \begin{pmatrix} \frac{1}{5} \\ -\frac{1}{10} \\ -\frac{2}{5} \\ \frac{3}{10} \end{pmatrix}$$



Error:

$$\begin{aligned} \|\vec{y} - A\vec{x}\| &= \sqrt{\frac{1}{25} + \frac{1}{100} + \frac{4}{25} + \frac{9}{100}} \\ &= \sqrt{\frac{4+1+16+9}{100}} \\ &= \frac{\sqrt{30}}{10} \\ &= 0.54772 \end{aligned}$$

The **second order** equation:

$$y = 0.25x^2 + 1.05x + 0.85$$

Error = 0.22361

The **third order** equation:

$$y = 0.16667x^3 + 0.82222x + 1$$

Error = 0.00000

Exercise

Find the equation of the line that best fits the given points in the least-squares sense and find the error.

$$\{(1, 0), (2, 1), (4, 2), (5, 3)\}$$

Solution

Let $y = mx + b$ be the equation of the line that best fits the given points. Then

$$\begin{pmatrix} 1 & 1 \\ 2 & 1 \\ 4 & 1 \\ 5 & 1 \end{pmatrix} \begin{pmatrix} m \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \\ 2 \\ 3 \end{pmatrix}$$

$$\text{where } A = \begin{pmatrix} 1 & 1 \\ 2 & 1 \\ 4 & 1 \\ 5 & 1 \end{pmatrix} \quad \vec{x} = \begin{pmatrix} m \\ b \end{pmatrix} \quad \vec{y} = \begin{pmatrix} 0 \\ 1 \\ 2 \\ 3 \end{pmatrix}$$

The normal equation: $A^T A \vec{x} = A^T \vec{y}$

$$\begin{pmatrix} 1 & 2 & 4 & 5 \\ 1 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 2 & 1 \\ 4 & 1 \\ 5 & 1 \end{pmatrix} \begin{pmatrix} m \\ b \end{pmatrix} = \begin{pmatrix} 1 & 2 & 4 & 5 \\ 1 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \\ 2 \\ 3 \end{pmatrix}$$

$$\begin{pmatrix} 46 & 12 \\ 12 & 4 \end{pmatrix} \begin{pmatrix} m \\ b \end{pmatrix} = \begin{pmatrix} 25 \\ 6 \end{pmatrix}$$

$$\Delta = \begin{vmatrix} 46 & 12 \\ 12 & 4 \end{vmatrix} = 40 \quad \Delta_m = \begin{vmatrix} 25 & 12 \\ 6 & 4 \end{vmatrix} = 28 \quad \Delta_b = \begin{vmatrix} 46 & 25 \\ 12 & 6 \end{vmatrix} = -24$$

$$m = \frac{28}{40} = \frac{7}{10}$$

$$b = -\frac{24}{40} = -\frac{3}{5}$$


$$\text{Thus, } y = \frac{7}{10}x - \frac{3}{5}$$

$$A\vec{x} = \begin{pmatrix} 1 & 1 \\ 2 & 1 \\ 4 & 1 \\ 5 & 1 \end{pmatrix} \begin{pmatrix} \frac{7}{10} \\ -\frac{3}{5} \end{pmatrix}$$

$$= \begin{pmatrix} \frac{1}{10} \\ \frac{4}{5} \\ \frac{11}{5} \\ \frac{29}{10} \end{pmatrix}$$

$$\vec{y} - A\vec{x} = \begin{pmatrix} 0 \\ 1 \\ 2 \\ 3 \end{pmatrix} - \begin{pmatrix} \frac{1}{10} \\ \frac{4}{5} \\ \frac{11}{5} \\ \frac{29}{10} \end{pmatrix}$$

$$= \begin{pmatrix} -\frac{1}{10} \\ \frac{1}{5} \\ -\frac{1}{5} \\ \frac{1}{10} \end{pmatrix}$$



Error:

$$\begin{aligned} \|\vec{y} - A\vec{x}\| &= \sqrt{2\left(\frac{1}{100} + \frac{1}{25}\right)} \\ &= \frac{\sqrt{10}}{10} \\ &= 0.31623 \end{aligned}$$

The **second order** equation:

$$y = 0.0x^2 + 0.7x - .6$$

$$\mathbf{Error} = 0.31623$$

The **third order** equation:

$$y = 0.08333x^3 - 0.75x^2 + 2.66667x - 2$$

$$\mathbf{Error} = 0.00000$$

Exercise

Find the orthogonal projection of the vector \vec{u} on the subspace of \mathbb{R}^4 spanned by the vectors

$$\vec{u} = (-3, -3, 8, 9); \quad \vec{v}_1 = (3, 1, 0, 1), \quad \vec{v}_2 = (1, 2, 1, 1), \quad \vec{v}_3 = (-1, 0, 2, -1)$$

Solution

$$\text{Let } A = \begin{pmatrix} 3 & 1 & -1 \\ 1 & 2 & 0 \\ 0 & 1 & 2 \\ 1 & 1 & -1 \end{pmatrix}$$

$$A^T A = \begin{pmatrix} 3 & 1 & 0 & 1 \\ 1 & 2 & 1 & 1 \\ -1 & 0 & 2 & -1 \end{pmatrix} \begin{pmatrix} 3 & 1 & -1 \\ 1 & 2 & 0 \\ 0 & 1 & 2 \\ 1 & 1 & -1 \end{pmatrix}$$

$$= \begin{pmatrix} 11 & 6 & -4 \\ 6 & 7 & 0 \\ -4 & 0 & 6 \end{pmatrix}$$

$$A^T \vec{u} = \begin{pmatrix} 3 & 1 & 0 & 1 \\ 1 & 2 & 1 & 1 \\ -1 & 0 & 2 & -1 \end{pmatrix} \begin{pmatrix} -3 \\ -3 \\ 8 \\ 9 \end{pmatrix}$$

$$= \begin{pmatrix} -3 \\ 8 \\ 10 \end{pmatrix}$$

The normal solution is $A^T A \vec{x} = A^T \vec{u}$

$$\begin{pmatrix} 11 & 6 & -4 \\ 6 & 7 & 0 \\ -4 & 0 & 6 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -3 \\ 8 \\ 10 \end{pmatrix}$$

$$\Delta = \begin{vmatrix} 11 & 6 & -4 \\ 6 & 7 & 0 \\ -4 & 0 & 6 \end{vmatrix} = 134$$

$$\Delta_1 = \begin{vmatrix} -3 & 6 & -4 \\ 8 & 7 & 0 \\ 10 & 0 & 6 \end{vmatrix} = -134$$

$$\Delta_2 = \begin{vmatrix} 11 & -3 & -4 \\ 6 & 8 & 0 \\ -4 & 10 & 6 \end{vmatrix} = 268$$

$$\underline{x_1 = \frac{-134}{134} = -1} \quad \underline{x_2 = \frac{268}{134} = 2} \quad \underline{x_3 = \frac{134}{134} = 1}$$

$$\vec{x} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -1 \\ 2 \\ 1 \end{pmatrix}$$

So $\text{proj}_W \vec{u} = A\vec{x}$

$$= \begin{pmatrix} 3 & 1 & -1 \\ 1 & 2 & 0 \\ 0 & 1 & 2 \\ 1 & 1 & -1 \end{pmatrix} \begin{pmatrix} -1 \\ 2 \\ 1 \end{pmatrix}$$

$$= \begin{pmatrix} -2 \\ 3 \\ 4 \\ 0 \end{pmatrix}$$

$$\underline{\text{proj}_W \vec{u} = (-2, 3, 4, 0)}$$

Exercise

Find the orthogonal projection of the vector \vec{u} on the subspace of \mathbb{R}^4 spanned by the vectors

$$\vec{u} = (6, 3, 9, 6); \quad \vec{v}_1 = (2, 1, 1, 1), \quad \vec{v}_2 = (1, 0, 1, 1), \quad \vec{v}_3 = (-2, -1, 0, -1)$$

Solution

$$\text{Let } A = \begin{pmatrix} 2 & 1 & -2 \\ 1 & 0 & -1 \\ 1 & 1 & 0 \\ 1 & 1 & -1 \end{pmatrix}$$

$$A^T A = \begin{pmatrix} 2 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ -2 & -1 & 0 & -1 \end{pmatrix} \begin{pmatrix} 2 & 1 & -2 \\ 1 & 0 & -1 \\ 1 & 1 & 0 \\ 1 & 1 & -1 \end{pmatrix}$$

$$= \begin{pmatrix} 7 & 4 & -6 \\ 4 & 3 & -3 \\ -6 & -3 & 6 \end{pmatrix}$$

$$A^T \vec{u} = \begin{pmatrix} 2 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ -2 & -1 & 0 & -1 \end{pmatrix} \begin{pmatrix} 6 \\ 3 \\ 9 \\ 6 \end{pmatrix}$$

$$= \begin{pmatrix} 30 \\ 21 \\ -21 \end{pmatrix}$$

The normal solution is $A^T A\vec{x} = A^T \vec{u}$

$$\begin{pmatrix} 7 & 4 & -6 \\ 4 & 3 & -3 \\ -6 & -3 & 6 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 30 \\ 21 \\ -21 \end{pmatrix}$$

$$\Delta = \begin{vmatrix} 7 & 4 & -6 \\ 4 & 3 & -3 \\ -6 & -3 & 6 \end{vmatrix} = 3$$

$$\Delta_1 = \begin{vmatrix} 30 & 4 & -6 \\ 21 & 3 & -3 \\ -21 & -3 & 6 \end{vmatrix} = 18$$

$$\Delta_2 = \begin{vmatrix} 7 & 30 & -6 \\ 4 & 21 & -3 \\ -6 & -21 & 6 \end{vmatrix} = 9$$

$$\underline{x_1 = \frac{18}{3} = 6 \quad x_2 = \frac{9}{3} = 3 \quad x_3 = 4}$$

$$\vec{x} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 6 \\ 3 \\ 4 \end{pmatrix}$$

$$\text{So } \text{proj}_W \vec{u} = A\vec{x}$$

$$= \begin{pmatrix} 2 & 1 & -2 \\ 1 & 0 & -1 \\ 1 & 1 & 0 \\ 1 & 1 & -1 \end{pmatrix} \begin{pmatrix} 6 \\ 3 \\ 4 \end{pmatrix}$$

$$= \begin{pmatrix} 7 \\ 2 \\ 9 \\ 5 \end{pmatrix}$$

$$\underline{\text{proj}_W \vec{v} = (7, 2, 9, 5)}$$

Exercise

Find the orthogonal projection of the vector \vec{u} on the subspace of \mathbb{R}^4 spanned by the vectors

$$\vec{u} = (-2, 0, 2, 4); \quad v_1 = (1, 1, 3, 0), \quad \vec{v}_2 = (-2, -1, -2, 1), \quad \vec{v}_3 = (-3, -1, 1, 3)$$

Solution

$$\text{Let } A = \begin{pmatrix} 1 & -2 & -3 \\ 1 & -1 & -1 \\ 3 & -2 & 1 \\ 0 & 1 & 3 \end{pmatrix}$$

$$A^T A = \begin{pmatrix} 1 & 1 & 3 & 0 \\ -2 & -1 & -2 & 1 \\ -3 & -1 & 1 & 3 \end{pmatrix} \begin{pmatrix} 1 & -2 & -3 \\ 1 & -1 & -1 \\ 3 & -2 & 1 \\ 0 & 1 & 3 \end{pmatrix}$$

$$= \begin{pmatrix} 11 & -9 & -1 \\ -9 & 10 & 8 \\ -1 & 8 & 20 \end{pmatrix}$$

$$A^T \vec{u} = \begin{pmatrix} 1 & 1 & 3 & 0 \\ -2 & -1 & -2 & 1 \\ -3 & -1 & 1 & 3 \end{pmatrix} \begin{pmatrix} -2 \\ 0 \\ 2 \\ 4 \end{pmatrix}$$

$$= \begin{pmatrix} 4 \\ 4 \\ 20 \end{pmatrix}$$

The normal solution is $A^T A \vec{x} = A^T \vec{u}$

$$\begin{pmatrix} 11 & -9 & -1 \\ -9 & 10 & 8 \\ -1 & 8 & 20 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 4 \\ 4 \\ 20 \end{pmatrix}$$

$$\Delta = \begin{vmatrix} 11 & -9 & -1 \\ -9 & 10 & 8 \\ -1 & 8 & 20 \end{vmatrix} = 10 \quad \Delta_1 = \begin{vmatrix} 4 & -9 & -1 \\ 4 & 10 & 8 \\ 20 & 8 & 20 \end{vmatrix} = -8 \quad \Delta_2 = \begin{vmatrix} 11 & 4 & -1 \\ -9 & 4 & 8 \\ -1 & 20 & 20 \end{vmatrix} = -16$$

$$\underline{x_1 = \frac{-8}{10} = -\frac{4}{5}} \quad \underline{x_2 = \frac{-16}{10} = -\frac{8}{5}} \quad \underline{x_3 = \frac{8}{5}}$$

$$\vec{x} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} -\frac{4}{5} \\ -\frac{8}{5} \\ \frac{8}{5} \end{pmatrix}$$

$$proj_W \vec{u} = A \vec{x}$$

$$= \begin{pmatrix} 1 & -2 & -3 \\ 1 & -1 & -1 \\ 3 & -2 & 1 \\ 0 & 1 & 3 \end{pmatrix} \begin{pmatrix} -\frac{4}{5} \\ -\frac{8}{5} \\ \frac{8}{5} \end{pmatrix}$$

$$= \begin{pmatrix} -\frac{12}{5} \\ -\frac{4}{5} \\ \frac{12}{5} \\ \frac{16}{5} \end{pmatrix}$$

$$\underline{\text{proj}_W \vec{u} = \left(-\frac{12}{5}, -\frac{4}{5}, \frac{12}{5}, \frac{16}{5} \right)}$$

Exercise

Find the standard matrix for the orthogonal projection P of \mathbb{R}^2 on the line passes through the origin and makes an angle θ with the positive x -axis.

Solution


Since the line l is 2-dimensional, then we can take $\vec{v} = (\cos \theta, \sin \theta)$ as a basis for this subspace

$$A = \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}$$

$$[P] = A^T A$$

$$= [\cos \theta \quad \sin \theta] \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}$$

$$= \begin{bmatrix} \cos^2 \theta & \cos \theta \sin \theta \\ \cos \theta \sin \theta & \sin^2 \theta \end{bmatrix}$$



Exercise

Hooke's law in physics states that the length x of a uniform spring is a linear function of the force y applied to it. If we express the relationship as $y = mx + b$, then the coefficient m is called the spring constant.

Suppose a particular unstretched spring has a measured length of 6.1 inches.(i.e., $x = 6.1$ when $y = 0$).

Forces of 2 pounds, 4 pounds, and 6 pounds are then applied to the spring, and the corresponding lengths are found to be 7.6 inches, 8.7 inches, and 10.4 inches. Find the spring constant.

Solution

$$M = \begin{pmatrix} 6.1 & 1 \\ 7.6 & 1 \\ 8.7 & 1 \\ 10.4 & 1 \end{pmatrix}, \quad \vec{y} = \begin{pmatrix} 0 \\ 2 \\ 4 \\ 6 \end{pmatrix}$$

The normal equation: $A^T A \vec{x} = A^T \vec{y}$

$$\begin{pmatrix} 6.1 & 7.6 & 8.7 & 10.4 \\ 1 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 6.1 & 1 \\ 7.6 & 1 \\ 8.7 & 1 \\ 10.4 & 1 \end{pmatrix} \begin{pmatrix} m \\ b \end{pmatrix} = \begin{pmatrix} 6.1 & 7.6 & 8.7 & 10.4 \\ 1 & 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 0 \\ 2 \\ 4 \\ 6 \end{pmatrix}$$


$$\begin{pmatrix} 278.82 & 32.8 \\ 32.8 & 4 \end{pmatrix} \begin{pmatrix} m \\ b \end{pmatrix} = \begin{pmatrix} 112.4 \\ 12 \end{pmatrix}$$

$$\begin{pmatrix} m \\ b \end{pmatrix} = \frac{1}{39.44} \begin{pmatrix} 4 & -32.8 \\ -32.8 & 278.2 \end{pmatrix} \begin{pmatrix} 112.4 \\ 12 \end{pmatrix}$$

$$= \frac{1}{39.44} \begin{pmatrix} 56 \\ -348.32 \end{pmatrix}$$

$$= \begin{pmatrix} 1.4 \\ -8.8 \end{pmatrix}$$

Thus, the estimated value of the spring constant is ≈ 1.4 pounds



Exercise

Prove:

If A has a linearly independent column vectors, and if \vec{b} is orthogonal to the column space of A , then the least squares solution of $A\vec{x} = \vec{b}$ is $\vec{x} = \vec{0}$.

Solution

If A has linearly independent column vectors, then $A^T A$ is invertible and the least squares solution of $A\vec{x} = \vec{b}$ is the solution of $A^T A \vec{x} = A^T \vec{b}$, but since \vec{b} is orthogonal to the column space of A .

$A^T \vec{b} = 0$, so \vec{x} is a solution of $A^T A \vec{x} = 0$.

Thus $\vec{x} = \vec{0}$ since $A^T A$ is invertible.

Exercise

Let A be an $m \times n$ matrix with linearly independent row vectors. Find a standard matrix for the orthogonal projection of \mathbb{R}^n onto the row space of A .

Solution

A^T will have linearly independent column vectors, and the column space A^T is the row space of A .

Thus, the standard matrix for the orthogonal projection of \mathbb{R}^n onto the row space of A is

$$\begin{aligned}
 [P] &= A^T \left[\begin{pmatrix} A^T \end{pmatrix}^T A^T \right]^{-1} \begin{pmatrix} A^T \end{pmatrix}^T \\
 &= A^T \left(A A^T \right)^{-1} A
 \end{aligned}$$

Exercise

Let W be the line with parametric equations $x = 2t, \quad y = -t, \quad z = 4t$

- Find a basis for W .
- Find the standard matrix for the orthogonal projection on W .
- Use the matrix in part (b) to find the orthogonal projection of a point $P_0(x_0, y_0, z_0)$ on W .
- Find the distance between the point $P_0(2, 1, -3)$ and the line W .

Solution

a) $W = \text{span}\{(2, -1, 4)\}$

So that the vector $(2, -1, 4)$ forms a basis for W (linear independence)

b) Let $A = \begin{pmatrix} 2 \\ -1 \\ 4 \end{pmatrix}$

$$[P] = A(A^T A)^{-1} A^T$$

$$= \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix} \left(\begin{bmatrix} 2 & -1 & 4 \end{bmatrix} \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix} \right)^{-1} \begin{bmatrix} 2 & -1 & 4 \end{bmatrix}$$

$$= \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix} [21]^{-1} \begin{bmatrix} 2 & -1 & 4 \end{bmatrix}$$

$$= \frac{1}{21} \begin{bmatrix} 4 & -2 & 8 \\ -2 & 1 & -4 \\ 8 & -4 & 16 \end{bmatrix}$$

$$= \begin{bmatrix} \frac{4}{21} & -\frac{2}{21} & \frac{8}{21} \\ -\frac{2}{21} & 1 & -\frac{4}{21} \\ \frac{8}{21} & -\frac{4}{21} & \frac{16}{21} \end{bmatrix}$$

$$c) \begin{bmatrix} \frac{4}{21} & -\frac{2}{21} & \frac{8}{21} \\ -\frac{2}{21} & 1 & -\frac{4}{21} \\ \frac{8}{21} & -\frac{4}{21} & \frac{16}{21} \end{bmatrix} \begin{bmatrix} x_0 \\ y_0 \\ z_0 \end{bmatrix} = \begin{bmatrix} \frac{4}{21}x_0 - \frac{2}{21}y_0 + \frac{8}{21}z_0 \\ -\frac{2}{21}x_0 + y_0 - \frac{4}{21}z_0 \\ \frac{8}{21}x_0 - \frac{4}{21}y_0 + \frac{16}{21}z_0 \end{bmatrix}$$

$$d) \begin{bmatrix} \frac{4}{21} & -\frac{2}{21} & \frac{8}{21} \\ -\frac{2}{21} & 1 & -\frac{4}{21} \\ \frac{8}{21} & -\frac{4}{21} & \frac{16}{21} \end{bmatrix} \begin{bmatrix} 2 \\ 1 \\ -3 \end{bmatrix} = \begin{bmatrix} -\frac{6}{7} \\ \frac{3}{7} \\ -\frac{12}{7} \end{bmatrix}$$

The distance between P_0 and W equals to the distance between P_0 and its projection on W .

The distance between $(2, 1, -3)$ and $\left(-\frac{6}{7}, \frac{3}{7}, -\frac{12}{7}\right)$ is

$$\begin{aligned} d &= \sqrt{\left(2 + \frac{6}{7}\right)^2 + \left(1 - \frac{3}{7}\right)^2 + \left(-3 + \frac{12}{7}\right)^2} \\ &= \sqrt{\frac{400}{49} + \frac{16}{49} + \frac{81}{49}} \\ &= \frac{\sqrt{497}}{7} \end{aligned}$$

Exercise

In R^3 , consider the line l given by the equations $x = t, \quad y = t, \quad z = t$

And the line m given by the equations $x = s, \quad y = 2s - 1, \quad z = 1$


Let P be the point on l , and let Q be a point on m . Find the values of t and s that minimize the distance between the lines by minimizing the squared distance $\|P - Q\|^2$

Solution

When $t = 1 \Rightarrow$ Let $P = (1, 1, 1)$ is on line l

When $s = 1 \Rightarrow$ Let $Q = (1, 1, 1)$ is on line m

$$\|P - Q\| = \sqrt{(1-1)^2 + (1-1)^2 + (1-1)^2} = 0 \geq 0$$



Thus, these are the values $P = (1, 1, 1)$ and $Q = (1, 1, 1)$ are the values for $s = t = 1$ that minimize the distance between the lines.

Exercise

Determine whether the statement is true or false,

- If A is an $m \times n$ matrix, then $A^T A$ is a square matrix.
- If $A^T A$ is invertible, then A is invertible.
- If A is invertible, then $A^T A$ is invertible.
- If $A\vec{x} = \vec{b}$ is a consistent linear system, then $A^T A\vec{x} = A^T \vec{b}$ is also consistent.
- If $A\vec{x} = \vec{b}$ is an inconsistent linear system, then $A^T A\vec{x} = A^T \vec{b}$ is also inconsistent.
- Every linear system has a least squares solution.
- Every linear system has a unique least squares solution.
- If A is an $m \times n$ matrix with linearly independent columns and \vec{b} is in R^m , then $A\vec{x} = \vec{b}$ has a unique least squares solution.

Solution

- True;** $A^T A$ is an $n \times n$ matrix
- False;** only square matrix has inverses, but $A^T A$ can be invertible when A is not square matrix.
- True;** if A is invertible, so is A^T , so the product $A^T A$ is also invertible
- True**
- False;** the system $A^T A\vec{x} = A^T \vec{b}$ may be consistent
- True**
- False;** the least squares solution may involve a parameter
- True;** if A has linearly independent column vectors; then $A^T A$ is invertible, so $A^T A\vec{x} = A^T \vec{b}$ has a unique solution

Exercise

A certain experiment produces the data $\{(1, 1.8), (2, 2.7), (3, 3.4), (4, 3.8), (5, 3.9)\}$.

Find the function that it will fit these data in the form of $y = \beta_1 x + \beta_2 x^2$

Solution

Given: the equation $y = \beta_1 x + \beta_2 x^2$ that best fits the given points. Then

$$\begin{pmatrix} 1 & 1 \\ 2 & 4 \\ 3 & 9 \\ 4 & 16 \\ 5 & 25 \end{pmatrix} \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix} = \begin{pmatrix} 1.8 \\ 2.7 \\ 3.4 \\ 3.8 \\ 3.9 \end{pmatrix}$$

$$\text{where } A = \begin{pmatrix} 1 & 1 \\ 2 & 4 \\ 3 & 9 \\ 4 & 16 \\ 5 & 25 \end{pmatrix} \quad \vec{x} = \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix} \quad \vec{y} = \begin{pmatrix} 1.8 \\ 2.7 \\ 3.4 \\ 3.8 \\ 3.9 \end{pmatrix}$$

The normal equation formula: $A^T A \vec{x} = A^T \vec{y}$

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 4 & 9 & 16 & 25 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 2 & 4 \\ 3 & 9 \\ 4 & 16 \\ 5 & 25 \end{pmatrix} \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 1 & 4 & 9 & 16 & 25 \end{pmatrix} \begin{pmatrix} 1.8 \\ 2.7 \\ 3.4 \\ 3.8 \\ 3.9 \end{pmatrix}$$


$$\begin{pmatrix} 55 & 225 \\ 225 & 979 \end{pmatrix} \begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix} = \begin{pmatrix} 52.1 \\ 201.5 \end{pmatrix}$$

$$\Delta = \begin{vmatrix} 55 & 225 \\ 225 & 979 \end{vmatrix} = 3,220 \quad \Delta\beta_1 = \begin{vmatrix} 52.1 & 225 \\ 201.5 & 979 \end{vmatrix} = 5,668.4 \quad \Delta\beta_2 = \begin{vmatrix} 55 & 52.1 \\ 225 & 201.5 \end{vmatrix} = -640$$

$$\beta_1 = \frac{5,668.4}{3,220}$$
$$\approx 1.76$$

$$\beta_2 = -\frac{640}{3,220}$$
$$\approx -0.199$$

$$y = 1.76x - .2x^2$$



Exercise

According to Kepler's first law, a comet should have an ellipse, parabolic, or hyperbolic orbit (with gravitational attractions from the planets ignored). In suitable polar coordinates, the position (r, ν) of a comet satisfies an equation of the form

$$r = \beta + e(r \cdot \cos \nu)$$

Where β is a constant and e is the eccentricity of the orbit, with $0 \leq e < 1$ for an ellipse, $e = 1$ for a parabolic, and $e > 1$ for a hyperbola.

Suppose observations of a newly discovered comet provide the data below.

ν	.88	1.10	1.42	1.77	2.14
r	3.00	2.30	1.65	1.25	1.01

Determine the type of orbit, and predict where the orbit will be when $\nu = 4.6$ (*radians*)?

Solution

Given: the equation in the form $r = \beta + e(r \cdot \cos \nu)$

$$3 = \beta + e(3 \cdot \cos(.88)) = \beta + 1.911e$$

$$2.3 = \beta + e(2.3 \cos(1.1)) = \beta + 1.043e$$

$$1.65 = \beta + e(1.65 \cos(1.42)) = \beta + .248e$$

$$1.25 = \beta + e(1.25 \cos(1.77)) = \beta - .247e$$

$$1.01 = \beta + e(1.01 \cos(2.14)) = \beta - .544e$$

$$\begin{pmatrix} 1 & 1.911 \\ 1 & 1.043 \\ 1 & .248 \\ 1 & -.247 \\ 1 & -.544 \end{pmatrix} \begin{pmatrix} \beta \\ e \end{pmatrix} = \begin{pmatrix} 3 \\ 2.3 \\ 1.65 \\ 1.25 \\ 1.01 \end{pmatrix}$$

$$\text{where } A = \begin{pmatrix} 1 & 1.911 \\ 1 & 1.043 \\ 1 & .248 \\ 1 & -.247 \\ 1 & -.544 \end{pmatrix} \quad \vec{v} = \begin{pmatrix} \beta \\ e \end{pmatrix} \quad \vec{r} = \begin{pmatrix} 3 \\ 2.3 \\ 1.65 \\ 1.25 \\ 1.01 \end{pmatrix}$$

The normal equation formula: $A^T A \vec{v} = A^T \vec{r}$

$$\begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ 1.911 & 1.043 & .248 & -.247 & -.544 \end{pmatrix} \begin{pmatrix} 1 & 1.911 \\ 1 & 1.043 \\ 1 & .248 \\ 1 & -.247 \\ 1 & -.544 \end{pmatrix} \begin{pmatrix} \beta \\ e \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ 1.911 & 1.043 & .248 & -.247 & -.544 \end{pmatrix} \begin{pmatrix} 3 \\ 2.3 \\ 1.65 \\ 1.25 \\ 1.01 \end{pmatrix}$$

$$\begin{pmatrix} 5 & 2.411 \\ 2.411 & 5.158 \end{pmatrix} \begin{pmatrix} \beta \\ e \end{pmatrix} = \begin{pmatrix} 9.21 \\ 7.683 \end{pmatrix}$$

$$\Delta = \begin{vmatrix} 5 & 2.411 \\ 2.411 & 5.158 \end{vmatrix} = 19.98 \quad \Delta_{\beta} = \begin{vmatrix} 9.21 & 2.411 \\ 7.683 & 5.158 \end{vmatrix} = 28.98 \quad \Delta_e = \begin{vmatrix} 5 & 9.21 \\ 2.411 & 7.683 \end{vmatrix} = 16.21$$

$$\beta = \frac{28.98}{19.98}$$

$$\approx 1.45$$



$$e = \frac{16.21}{19.98}$$

$$\approx 0.811 < 1$$

Therefore, the orbit is an *ellipse* type since $e \approx 0.811 < 1$

Since $r = \beta + e(r \cdot \cos v)$

$$\text{Then, } r(v) = \frac{1.45}{1 - 0.811 \cdot \cos v}$$



$$r(4.6) = \frac{1.45}{1 - 0.811 \cdot \cos 4.6}$$

$$\approx 1.329$$

Exercise

To measure the takeoff performance of an airplane, the horizontal position of the plane was measured every second, from $t = 0$ to $t = 12$

The position (in *feet*) were:

0, 8.8, 29.9, 62.0, 104.7, 159.1, 222.0, 294.5, 380.4, 471.1, 571.7, 686.8, and 809.2

- Find the least square cubic curve $y = \beta_0 + \beta_1 t + \beta_2 t^2 + \beta_3 t^3$ for these data.
- Estimate the velocity of the plane when $t = 4.5$ *sec*, using the result from part (a).

Solution

Given: the equation is in form $y = \beta_0 + \beta_1 t + \beta_2 t^2 + \beta_3 t^3$

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 \\ 1 & 2 & 4 & 8 \\ 1 & 3 & 9 & 27 \\ 1 & 4 & 16 & 64 \\ 1 & 5 & 25 & 125 \\ 1 & 6 & 36 & 216 \\ 1 & 7 & 49 & 343 \\ 1 & 8 & 64 & 512 \\ 1 & 9 & 81 & 729 \\ 1 & 10 & 100 & 1000 \\ 1 & 11 & 121 & 1331 \\ 1 & 12 & 144 & 1728 \end{pmatrix} \begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 8.8 \\ 29.9 \\ 62 \\ 104.7 \\ 159.1 \\ 222.0 \\ 294.5 \\ 380.4 \\ 471.1 \\ 571.7 \\ 686.8 \\ 809.2 \end{pmatrix}$$

$\mathbf{A} \quad \vec{t} = \vec{y}$

The normal equation formula: $A^T A \vec{t} = A^T \vec{y}$

$$\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & \\ 0 & 1 & 4 & 9 & 16 & 25 & 36 & 49 & 64 & 81 & 100 & 121 & 144 & \\ 0 & 1 & 8 & 27 & 64 & 125 & 216 & 343 & 512 & 729 & 1000 & 1331 & 1728 & \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 \\ 1 & 2 & 4 & 8 \\ 1 & 3 & 9 & 27 \\ 1 & 4 & 16 & 64 \\ 1 & 5 & 25 & 125 \\ 1 & 6 & 36 & 216 \\ 1 & 7 & 49 & 343 \\ 1 & 8 & 64 & 512 \\ 1 & 9 & 81 & 729 \\ 1 & 10 & 100 & 1000 \\ 1 & 11 & 121 & 1331 \\ 1 & 12 & 144 & 1728 \end{pmatrix} \begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix}$$

$$= \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 & 11 & 12 & \\ 0 & 1 & 4 & 9 & 16 & 25 & 36 & 49 & 64 & 81 & 100 & 121 & 144 & \\ 0 & 1 & 8 & 27 & 64 & 125 & 216 & 343 & 512 & 729 & 1000 & 1331 & 1728 & \end{pmatrix} \begin{pmatrix} 0 \\ 8.8 \\ 29.9 \\ 62 \\ 104.7 \\ 159.1 \\ 222.0 \\ 294.5 \\ 380.4 \\ 471.1 \\ 571.7 \\ 686.8 \\ 809.2 \end{pmatrix}$$

$$\begin{pmatrix} 13 & 78 & 650 & 6,084 \\ 78 & 650 & 6,084 & 60,710 \\ 650 & 6,084 & 60,710 & 630,708 \\ 6,084 & 60,710 & 630,708 & 6,735,950 \end{pmatrix} \begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix} = \begin{pmatrix} 3,800.2 \\ 35,127.7 \\ 348,063.9 \\ 3,599,800.9 \end{pmatrix}$$

$$\Delta = \begin{vmatrix} 13 & 78 & 650 & 6,084 \\ 78 & 650 & 6,084 & 60,710 \\ 650 & 6,084 & 60,710 & 630,708 \\ 6,084 & 60,710 & 630,708 & 6,735,950 \end{vmatrix} = 97,538,785,344$$

$$\Delta_0 = \begin{vmatrix} 3800.2 & 78 & 650 & 6,084 \\ 35,127.7 & 650 & 6,084 & 60,710 \\ 348,063.9 & 6,084 & 60,710 & 630,708 \\ 3,599,800.9 & 60,710 & 630,708 & 6,735,950 \end{vmatrix} = -83,470,691,303.8916$$

Or I use my program to find the values

```
rref = (Matrix([
[1, 0, 0, 0, -0.855769230765803],
[0, 1, 0, 0, 4.70248501498163],
[0, 0, 1, 0, 5.55536963037029],
[0, 0, 0, 1, -0.0273601398601744]]))
```

$$\beta_0 \approx -0.855769$$

$$\beta_1 \approx 4.702485$$

$$\beta_2 \approx 5.55537$$

$$\beta_3 \approx -0.02736$$

$$y(t) = -0.855769 + 4.702485t + 5.55537t^2 - 0.02736t^3$$

$$\text{Error} = 3.9734$$

