

## ***Solution***      **Section 4.2 – Series Solutions near Ordinary Points**

### ***Exercise***

Find a power series solution.  $y' = 3y$

### **Solution**

The equation  $y' = 3y$  is separable with solution

$$\frac{dy}{dx} = 3y \Rightarrow \frac{dy}{y} = 3dx \rightarrow \underline{y = Ce^{3x}}$$

$$\underline{\ln(y) = 3x + C}$$

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y' - 3y = 0$$

$$\sum_{n=0}^{\infty} (n+1) a_{n+1} x^n - 3 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} [(n+1) a_{n+1} - 3a_n] x^n = 0$$

$$(n+1) a_{n+1} - 3a_n = 0 \Rightarrow a_{n+1} = \frac{3a_n}{n+1}; \quad n \geq 0$$

With  $y(0) = 3a_0$

$$a_1 = 3a_0$$

$$a_2 = \frac{3}{2} a_1 = \frac{3 \cdot 3}{2} a_0$$

$$a_3 = \frac{3}{3} a_2 = \frac{3 \cdot 3 \cdot 3}{2 \cdot 3} a_0$$

$$a_4 = \frac{3}{4} a_3 = \frac{3 \cdot 3 \cdot 3 \cdot 3}{2 \cdot 3 \cdot 4} a_0$$

$$\vdots \quad \quad \vdots$$

$$\underline{a_n = \frac{3^n}{n!} a_0}$$

$$\begin{aligned}
 y(x) &= \sum_{n=0}^{\infty} \frac{3^n}{n!} a_0 x^n \\
 &= a_0 \sum_{n=0}^{\infty} \frac{(3x)^n}{n!} \\
 &= a_0 e^{3x} \quad \checkmark
 \end{aligned}$$

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

### Exercise

Find a power series solution.  $y' = 4y$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y' = 4y$$

$$\sum_{n=1}^{\infty} n a_n x^{n-1} = 4 \sum_{n=0}^{\infty} a_n x^n$$

$$\sum_{n=0}^{\infty} (n+1) a_{n+1} x^n = \sum_{n=0}^{\infty} 4 a_n x^n$$

$$(n+1) a_{n+1} x^n = 4 a_n x^n$$

$$(n+1) a_{n+1} = 4 a_n$$

$$a_{n+1} = \frac{4}{n+1} a_n$$

$$n=0 \rightarrow a_1 = 4a_0$$

$$n=1 \rightarrow a_2 = \frac{4}{2} a_1 = \frac{4^2}{2!} a_0$$

$$n=2 \rightarrow a_3 = \frac{4}{3} a_2 = \frac{4^3}{3!} a_0$$

$$n=3 \rightarrow a_4 = \frac{4}{4} a_3 = \frac{4^4}{4!} a_0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\left| a_n = \frac{4^n}{n!} a_0 \right|$$

$$\begin{aligned} y(x) &= \sum_{n=0}^{\infty} \frac{4^n}{n!} a_0 x^n \\ &= a_0 \left( 1 + 4x + \frac{4^2}{2!} x^2 + \frac{4^3}{3!} x^3 + \dots \right) \\ &= a_0 e^{4x} \end{aligned}$$

### Exercise

Find a power series solution.  $y' = x^2 y$

### Solution

$$\frac{dy}{dx} = x^2 y$$

$$\int \frac{dy}{y} = \int x^2 dx$$

$$\ln y = \frac{1}{3} x^3 + C_1$$

$$y = e^{\frac{1}{3} x^3 + C_1}$$

$$y = C e^{\frac{1}{3} x^3}$$

$$y(0) = C(1) = a_0 \rightarrow C = a_0$$

$$\left| y = a_0 e^{x^3/3} \right|$$

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y' - x^2 y = 0$$

$$\sum_{n=1}^{\infty} n a_n x^{n-1} - x^2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=1}^{\infty} n a_n x^{n-1} - \sum_{n=0}^{\infty} a_n x^{n+2} = 0$$

$$\sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{k=-2}^{\infty} (k+3) a_{k+3} x^{k+2}$$

$$= \sum_{n=-2}^{\infty} (n+3) a_{n+3} x^{n+2}$$

$$\sum_{n=-2}^{\infty} (n+3) a_{n+3} x^{n+2} - \sum_{n=0}^{\infty} a_n x^{n+2} = 0$$

$$a_1 + 2a_2 x + \sum_{n=-2}^{\infty} (n+3) a_{n+3} x^{n+2} - \sum_{n=0}^{\infty} a_n x^{n+2} = 0$$

$$a_1 + 2a_2 x + \sum_{n=-2}^{\infty} [(n+3) a_{n+3} - a_n] x^{n+2} = 0$$

If we set  $a_1 = a_2 = 0$ , then

$$(n+3) a_{n+3} - a_n = 0 \Rightarrow a_{n+3} = \frac{a_n}{n+3}, \quad n \geq 0$$

$$a_3 = \frac{1}{3} a_0$$

$$a_4 = \frac{1}{4} a_1 = 0$$

$$a_5 = \frac{1}{5} a_2 = 0$$

$$a_6 = \frac{1}{6} a_3 = \frac{1}{3 \cdot 6} a_0$$

$$a_7 = \frac{1}{7} a_4 = 0$$

$$a_9 = \frac{1}{9} a_6 = \frac{1}{3 \cdot 6 \cdot 9} a_0 = \frac{1}{3^3 (1 \cdot 2 \cdot 3)} a_0$$

$$a_{12} = \frac{1}{12} a_9 = \frac{1}{3 \cdot 6 \cdot 9 \cdot 12} a_0 = \frac{1}{3^4 (1 \cdot 2 \cdot 3 \cdot 4)} a_0$$

$$\left| a_{3n} = \frac{1}{3^n \cdot n!} a_0 \right|$$

$$y(x) = \sum_{n=0}^{\infty} \frac{1}{3^n \cdot n!} a_0 x^{3n}$$

$$= a_0 \sum_{n=0}^{\infty} \frac{1}{n!} \left( \frac{x^3}{3} \right)^n$$

$$\left| = a_0 e^{x^3/3} \right| \quad \checkmark$$

$$P = \lim_{n \rightarrow \infty} \left| \frac{3^k k!}{1} \right| = \infty$$

### Exercise

Find a power series solution.  $y' + 2xy = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n \Rightarrow y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y' + 2xy = 0$$

$$\sum_{n=1}^{\infty} n a_n x^{n-1} + 2x \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+1) a_{n+1} x^n + \sum_{n=0}^{\infty} 2a_n x^{n+1} = 0$$

$$a_1 + \sum_{n=0}^{\infty} (n+2) a_{n+2} x^{n+1} + \sum_{n=0}^{\infty} 2a_n x^{n+1} = 0$$

$$a_1 + \sum_{n=0}^{\infty} [(n+2) a_{n+2} + 2a_n] x^{n+1} = 0$$

$$\left\{ \begin{array}{l} \underline{a_1 = 0} \\ (n+2) a_{n+2} + 2a_n = 0 \rightarrow \underline{a_{n+2} = -\frac{2}{n+2} a_n} \end{array} \right|$$

$$n=0 \rightarrow a_2 = -a_1 \qquad n=1 \rightarrow a_3 = -\frac{2}{3} a_1 = 0$$

$$n=2 \rightarrow a_4 = -\frac{1}{2} a_2 = \frac{1}{2} a_0 \qquad n=3 \rightarrow a_5 = -\frac{2}{7} a_3 = 0$$

$$n=4 \rightarrow a_6 = -\frac{1}{3} a_4 = -\frac{1}{2 \cdot 3} a_0 \qquad \vdots \quad \vdots \quad \vdots \quad \vdots$$

$$n=6 \rightarrow a_8 = -\frac{1}{4} a_6 = \frac{1}{4!} a_0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\underline{a_{2k} = \frac{(-1)^k}{k!} a_0}$$

$$y(x) = \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} a_0 x^{2k}$$

$$= a_0 \left( 1 - x^2 + \frac{1}{2!} x^4 - \frac{1}{3!} x^6 + \dots \right)$$

$$\underline{= a_0 e^{-x^2}}$$

$$P = \lim_{n \rightarrow \infty} \left| \frac{n+2}{-2} \right| = \underline{\underline{\infty}}$$

### Exercise

Find a power series solution.  $(x-2)y' + y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n \Rightarrow y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$(x-2) \sum_{n=1}^{\infty} n a_n x^{n-1} + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$x \sum_{n=1}^{\infty} n a_n x^{n-1} - 2 \sum_{n=1}^{\infty} n a_n x^{n-1} + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=1}^{\infty} n a_n x^n - 2 \sum_{n=1}^{\infty} n a_n x^{n-1} + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} n a_n x^n - 2 \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} [n a_n - 2(n+1) a_{n+1} + a_n] x^n = 0$$

$$n a_n - 2(n+1) a_{n+1} + a_n = 0$$

$$2(n+1) a_{n+1} = (n+1) a_n$$

$$\underline{a_{n+1} = \frac{1}{2} a_n}$$

$$n=0 \rightarrow a_1 = \frac{1}{2} a_0$$

$$n=1 \rightarrow a_2 = \frac{1}{2} a_1 = \frac{1}{2^2} a_0$$

$$n=2 \rightarrow a_3 = \frac{1}{2} a_2 = \frac{1}{2^3} a_0$$

$$n=3 \rightarrow a_4 = \frac{1}{2} a_3 = \frac{1}{2^4} a_0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\underline{a_n = \frac{1}{2^n} a_0}$$

$$y(x) = \sum_{n=0}^{\infty} \frac{1}{2^n} a_0 x^n$$

$$\begin{aligned}
&= a_0 \left( 1 + \frac{1}{2}x + \frac{1}{2^2}x^2 + \frac{1}{2^3}x^3 + \dots \right) \\
&= a_0 \left( 1 + \frac{x}{2} + \left(\frac{x}{2}\right)^2 + \left(\frac{x}{2}\right)^3 + \dots \right) \\
&= a_0 \frac{1}{1 - \frac{x}{2}} \\
&= \frac{2a_0}{2-x}
\end{aligned}$$

### Exercise

Find a power series solution.  $(2x-1)y' + 2y = 0$

### Solution

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} a_n x^n \Rightarrow y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} \\
(2x-1) \sum_{n=1}^{\infty} n a_n x^{n-1} + 2 \sum_{n=0}^{\infty} a_n x^n &= 0 \\
0 + \sum_{n=1}^{\infty} 2n a_n x^n - \sum_{n=1}^{\infty} n a_n x^{n-1} + \sum_{n=0}^{\infty} 2a_n x^n &= 0 \\
\sum_{n=0}^{\infty} 2n a_n x^n - \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n + \sum_{n=0}^{\infty} 2a_n x^n &= 0 \\
\sum_{n=0}^{\infty} [2n a_n - (n+1) a_{n+1} + 2a_n] x^n &= 0 \\
2n a_n - (n+1) a_{n+1} + 2a_n &= 0 \\
(n+1) a_{n+1} &= 2(n+1) a_n \\
a_{n+1} &= 2a_n
\end{aligned}$$

$$n=0 \rightarrow a_1 = 2a_0$$

$$n=1 \rightarrow a_2 = 2a_1 = 2^2 a_0$$

$$n=2 \rightarrow a_3 = 2a_2 = 2^3 a_0$$

$$n=3 \rightarrow a_4 = 2a_3 = 2^4 a_0$$

$$\begin{array}{cccc} \vdots & \vdots & \vdots & \vdots \\ \hline a_n = 2^n a_0 \end{array}$$

$$\begin{aligned} y(x) &= \sum_{n=0}^{\infty} 2^n a_0 x^n \\ &= a_0 (1 + 2x + 2^2 x^2 + 2^3 x^3 + \dots) \\ &= a_0 (1 + 2x + (2x)^2 + (2x)^3 + \dots) \\ &= \frac{a_0}{1-2x} \end{aligned}$$

### Exercise

Find a power series solution.  $2(x-1)y' = 3y$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n \Rightarrow y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$(2x-2) \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} 3 a_n x^n$$

$$2x \sum_{n=1}^{\infty} n a_n x^{n-1} - \sum_{n=1}^{\infty} 2 n a_n x^{n-1} = \sum_{n=0}^{\infty} 3 a_n x^n$$

$$0 + \sum_{n=1}^{\infty} 2 n a_n x^n - \sum_{n=1}^{\infty} 2 n a_n x^{n-1} = \sum_{n=0}^{\infty} 3 a_n x^n$$

$$\sum_{n=0}^{\infty} 2 n a_n x^n - \sum_{n=0}^{\infty} 2(n+1) a_{n+1} x^n = \sum_{n=0}^{\infty} 3 a_n x^n$$

$$\sum_{n=0}^{\infty} [2 n a_n - 2(n+1) a_{n+1}] x^n = \sum_{n=0}^{\infty} 3 a_n x^n$$

$$2 n a_n - 2(n+1) a_{n+1} = 3 a_n$$

$$-2(n+1) a_{n+1} = (3-2n) a_n$$

$$\underline{a_{n+1} = \frac{2n-3}{2n+2} a_n}$$

$$\rho = \lim_{n \rightarrow \infty} \frac{2n-3}{2n+2} = 1$$



$$n=0 \rightarrow a_1 = -\frac{3}{2}a_0$$

$$n=1 \rightarrow a_2 = -\frac{1}{4}a_1 = \frac{3}{8}a_0$$

$$n=2 \rightarrow a_3 = \frac{1}{6}a_2 = \frac{1}{16}a_0$$

$$n=3 \rightarrow a_4 = \frac{3}{8}a_3 = \frac{3}{128}a_0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$y(x) = a_0 \left( 1 - \frac{3}{2}x + \frac{3}{8}x^2 + \frac{1}{16}x^3 + \frac{3}{128}x^4 + \dots \right)$$

$$\frac{y'}{y} = \frac{3}{2} \frac{1}{x-1}$$

$$\int \frac{dy}{y} = \frac{3}{2} \int \frac{1}{x-1} dx$$

$$\ln y = \frac{3}{2} \ln(x-1) + \ln C$$

$$\ln y = \ln C (x-1)^{3/2}$$

$$y(x) = C(x-1)^{3/2}$$

### Exercise

Find a power series solution.  $(1+x)y' - y = 0$

### Solution

$$(1+x) \frac{dy}{dx} = y$$

$$\frac{dy}{y} = \frac{dx}{1+x}$$

$$\int \frac{dy}{y} = \int \frac{dx}{1+x}$$

$$\ln(y) = \ln(x+1) + C \Rightarrow y = C(x+1)$$

$$\text{With } y(0) = C(0+1) = a_0 \rightarrow C = a_0$$

$$\Rightarrow y = a_0(x+1)$$

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$(1+x)y' - y = 0$$

$$(1+x) \sum_{n=0}^{\infty} (n+1)a_{n+1}x^n - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+1)a_{n+1}x^n + \sum_{n=0}^{\infty} (n+1)a_{n+1}x^{n+1} - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+1)a_{n+1}x^n + \sum_{n=1}^{\infty} na_n x^n - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+1)a_{n+1}x^n + \sum_{n=0}^{\infty} na_n x^n - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left( (n+1)a_{n+1}x^n + na_n x^n - a_n x^n \right) = 0$$

$$(n+1)a_{n+1} + (n-1)a_n = 0 \rightarrow a_{n+1} = \frac{1-n}{n+1}a_n; \quad n \geq 0$$

$$a_1 = a_0 \quad a_2 = 0a_1 = 0 \quad a_3 = \frac{-1}{3}a_2 = 0$$

$$a_n = 0 \quad \text{for } n \geq 2$$

$$y(x) = a_0 + a_1 x$$

$$= a_0 + a_0 x$$

$$= a_0 (1+x) \quad \checkmark$$

### Exercise

Find a power series solution.  $(2-x)y' + 2y = 0$

### Solution

$$(2-x) \frac{dy}{dx} + 2y = 0$$

$$(2-x) \frac{dy}{dx} = -2y$$

$$\int \frac{dy}{y} = \int \frac{2d(2-x)}{2-x}$$

$$\ln y = 2 \ln(2-x) + C_1$$

$$\ln y = \ln(2-x)^2 + C_1$$

$$\ln y = \ln C(2-x)^2$$

$$y = C(2-x)^2$$

$$y(\mathbf{0}) = C(2-\mathbf{0})^2 = a_0 \rightarrow \underline{C = \frac{1}{4}a_0}$$

$$\underline{y = \frac{1}{4}a_0(2-x)^2}$$

$$y(x) = \sum_{n=0}^{\infty} a_n x^n \quad y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$(2-x)y' + 2y = 0$$

$$(2-x) \sum_{n=1}^{\infty} n a_n x^{n-1} + 2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$2 \sum_{n=1}^{\infty} n a_n x^{n-1} - \sum_{n=1}^{\infty} n a_n x^n + 2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=1}^{\infty} n a_n x^n = 0 + \sum_{n=1}^{\infty} n a_n x^n = \sum_{n=0}^{\infty} n a_n x^n$$

$$2 \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n - \sum_{n=0}^{\infty} n a_n x^n + 2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} [2(n+1)a_{n+1} - n a_n + 2a_n] x^n = 0$$

$$2(n+1)a_{n+1} - (n-2)a_n = 0$$

$$2(n+1)a_{n+1} = (n-2)a_n$$

$$a_{n+1} = \frac{n-2}{2(n+1)} a_n, \quad n \geq 0$$

$$a_1 = \frac{-2}{2} a_0 = -a_0$$

$$a_2 = \frac{-1}{4} a_1 = \frac{1}{4} a_0$$

$$a_3 = \frac{0}{6} a_0 = 0$$

$$\vdots$$

$$a_n = 0$$

$$y(x) = a_0 - a_0 x + \frac{1}{4} a_0 x^2$$

$$= a_0 \left( 1 - x + \frac{1}{4} x^2 \right)$$

$$= \frac{1}{4} a_0 (4 - 4x + x^2)$$

$$\underline{= \frac{1}{4} a_0 (2-x)^2} \quad \checkmark$$

### Exercise

Find a power series solution.  $(x-4)y' + y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n \quad y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$(x-4)y' + y = 0$$

$$(x-4) \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+1) a_{n+1} x^{n+1} - \sum_{n=0}^{\infty} 4(n+1) a_{n+1} x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=1}^{\infty} n a_n x^n + \sum_{n=0}^{\infty} [a_n - 4(n+1) a_{n+1}] x^n = 0$$

$$\sum_{n=0}^{\infty} n a_n x^n + a_0 - 4a_1 + \sum_{n=1}^{\infty} [a_n - 4(n+1) a_{n+1}] x^n = 0$$

$$a_0 - 4a_1 + \sum_{n=1}^{\infty} [(n+1) a_n - 4(n+1) a_{n+1}] x^n = 0$$

$$a_0 - 4a_1 = 0 \quad \rightarrow \quad \underline{a_1 = \frac{1}{4} a_0}$$

$$(n+1) a_n - 4(n+1) a_{n+1} = 0 \quad \rightarrow \quad \underline{a_{n+1} = \frac{1}{4} a_n}$$

$$a_2 = \frac{1}{4} a_1 = \frac{1}{4^2} a_0$$

$$a_3 = \frac{1}{4} a_2 = \frac{1}{4^3} a_0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\underline{a_n = \frac{1}{4^n} a_0}$$

$$y(x) = \sum_{n=0}^{\infty} \frac{1}{4^n} a_0 x^n$$

$$= a_0 \sum_{n=0}^{\infty} \left(\frac{x}{4}\right)^n$$

$$\begin{aligned}
&= a_0 \left( \frac{1}{1 - \frac{x}{4}} \right) \\
&= a_0 \left( \frac{4}{4 - x} \right) \\
&= \frac{-4a_0}{x - 4} \quad \checkmark
\end{aligned}$$

### Exercise

Find a power series solution.  $x^2 y' = y - x - 1$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n \quad y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$x^2 y' = y - x - 1$$

$$x^2 \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n = -x - 1 + \sum_{n=0}^{\infty} a_n x^n$$

$$\sum_{n=0}^{\infty} (n+1) a_{n+1} x^{n+2} = -x - 1 + \sum_{n=0}^{\infty} a_n x^n$$

$$\sum_{n=2}^{\infty} (n-1) a_{n-1} x^n = -x - 1 + \sum_{n=0}^{\infty} a_n x^n$$

$$\sum_{n=2}^{\infty} (n-1) a_{n-1} x^n = -x - 1 + a_0 + a_1 x + \sum_{n=2}^{\infty} a_n x^n$$

$$-x - 1 + a_0 + a_1 x = 0$$

$$a_0 + a_1 x = 1 + x \Rightarrow a_0 = 1; a_1 = 1$$

$$(n-1) a_{n-1} = a_n$$

$$a_2 = a_1 = 1$$

$$a_3 = 2a_2 = 2$$

$$a_4 = 3a_3 = 1 \cdot 2 \cdot 3$$

$$\vdots \quad \vdots$$

$$a_n = (n-1)!$$

$$y(x) = 1 + x + x^2 + 2!x^3 + 3!x^4 + \dots$$

### Exercise

Find a power series solution.  $(x-3)y' + 2y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n \quad y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$(x-3)y' + 2y = 0$$

$$(x-3) \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n + 2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+1) a_{n+1} x^{n+1} - \sum_{n=0}^{\infty} 3(n+1) a_{n+1} x^n + \sum_{n=0}^{\infty} 2a_n x^n = 0$$

$$\sum_{n=1}^{\infty} n a_n x^n - \sum_{n=0}^{\infty} 3(n+1) a_{n+1} x^n + \sum_{n=0}^{\infty} 2a_n x^n = 0$$

$$\sum_{n=0}^{\infty} n a_n x^n - \sum_{n=0}^{\infty} 3(n+1) a_{n+1} x^n + \sum_{n=0}^{\infty} 2a_n x^n = 0$$

$$\sum_{n=0}^{\infty} [n a_n - 3(n+1) a_{n+1} + 2a_n] x^n = 0$$

$$-3(n+1) a_{n+1} + (n+2) a_n = 0 \rightarrow \underline{a_{n+1} = \frac{n+2}{3(n+1)} a_n} \quad n = 0, 1, 2, \dots$$

$$a_1 = \frac{2}{3} a_0 \quad a_2 = \frac{3}{3 \cdot 2} a_1 = \frac{3}{3^2} a_0$$

$$a_3 = \frac{4}{3 \cdot 3} a_2 = \frac{4}{3^3} a_0 \quad a_4 = \frac{5}{3 \cdot 4} a_3 = \frac{5}{3^4} a_0$$

$$\vdots$$

$$a_n = \frac{n+1}{3^n} a_0 \quad n \geq 1$$

$$y(x) = \left( 1 + \frac{2}{3}x + \frac{3}{3^2}x^2 + \frac{4}{3^3}x^3 + \dots \right) = a_0 \sum_{n=0}^{\infty} \frac{n+1}{3^n} x^n$$

$$\rho = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right| = \lim_{n \rightarrow \infty} \frac{3n+3}{n+2} = 3$$

$$\underline{y(x) = \frac{1}{(3-x)^2}}$$

### Exercise

Find a power series solution.  $xy' + y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n \Rightarrow y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$xy' + y = 0$$

$$x \sum_{n=1}^{\infty} n a_n x^{n-1} + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=1}^{\infty} n a_n x^{n+1} + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} n a_n x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+1) a_n x^n = 0$$

$$(n+1) a_n = 0 \rightarrow \underline{a_n = 0}$$

$$\underline{y(x) \equiv 0}$$

$\therefore$  The equation has no non-trivial power series.

### Exercise

Find a power series solution.  $x^3 y' - 2y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n \Rightarrow y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$x^3 y' - 2y = 0$$

$$x^3 \sum_{n=1}^{\infty} n a_n x^{n-1} - 2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=1}^{\infty} n a_n x^{n+2} - 2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=3}^{\infty} (n-2) a_{n-2} x^n - 2(a_0 + a_1 x + a_2 x^2) - \sum_{n=3}^{\infty} 2 a_n x^n = 0$$

$$\sum_{n=3}^{\infty} [(n-2) a_{n-2} - 2 a_n] x^n - 2(a_0 + a_1 x + a_2 x^2) = 0$$

$$\begin{cases} a_0 = a_1 = a_2 = 0 \\ (n-2) a_{n-2} = 2 a_n \rightarrow k a_k = 2 a_{k+2} \quad (k = n-2) \end{cases}$$

$$\underline{a_{k+2} = \frac{k}{2} a_k}$$

$$a_3 = \frac{1}{2} a_1 = 0$$

$$a_4 = a_2 = 0$$

$$a_n \equiv 0$$

$$\underline{y(x) \equiv 0}$$

$\therefore$  The equation has no non-trivial power series.

### ***Exercise***

Find a power series solution.  $y'' = 4y$

### **Solution**

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' = 4y$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n = 4 \sum_{n=0}^{\infty} a_n x^n$$



$$(n+2)(n+1)a_{n+2} = 4a_n$$

$$a_{n+2} = \frac{4}{(n+2)(n+1)}a_n$$

$$n=0 \rightarrow a_2 = 2a_0 = \frac{4}{2}a_0$$

$$n=1 \rightarrow a_3 = \frac{4}{2 \cdot 3}a_1$$

$$n=2 \rightarrow a_4 = \frac{4}{3 \cdot 4}a_2 = \frac{4^2}{4!}a_0$$

$$n=3 \rightarrow a_5 = \frac{4}{4 \cdot 5}a_3 = \frac{4^2}{5!}a_1$$

$$n=4 \rightarrow a_6 = \frac{4}{5 \cdot 6}a_4 = \frac{4^3}{6!}a_0$$

$$n=5 \rightarrow a_7 = \frac{4}{6 \cdot 7}a_5 = \frac{4^3}{7!}a_1$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\left| a_{2k} = \frac{2^{2k}}{(2k)!}a_0 \right|$$

$$\left| a_{2k+1} = \frac{2^{2k}}{(2k+1)!}a_1 \right|$$

$$y(x) = a_0 \left( 1 + \frac{2^2}{2!}x^2 + \frac{2^4}{4!}x^4 + \frac{2^6}{6!}x^6 + \dots \right) + a_1 \left( x + \frac{2^3}{3!}x^3 + \frac{2^5}{5!}x^5 + \dots \right)$$

$$= a_0 \left( 1 + \frac{1}{2!}(2x)^2 + \frac{1}{4!}(2x)^4 + \frac{1}{6!}(2x)^6 + \dots \right) + a_1 \left( x + \frac{1}{3!}(2x)^3 + \frac{1}{5!}(2x)^5 + \dots \right)$$

$$\left| = a_0 \cosh 2x + a_1 \sinh 2x \right|$$

## Exercise

Find a power series solution.  $y'' = 9y$

## Solution

The equation  $y'' = 9y$  has a characteristic equation  $\lambda^2 - 9 = 0 \Rightarrow \boxed{\lambda = \pm 3}$

$\therefore$  The general solution:  $y(x) = C_1 e^{3x} + C_2 e^{-3x}$

With  $y(0) = a_0$  and  $y'(0) = a_1$

$$y(0) = C_1 e^{3(0)} + C_2 e^{-3(0)} \rightarrow C_1 + C_2 = a_0$$

$$y'(x) = 3C_1 e^{3x} - 3C_2 e^{-3x}$$

$$y'(0) = 3C_1 e^{3(0)} - 3C_2 e^{-3(0)} \rightarrow 3C_1 - 3C_2 = a_1$$

$$\begin{cases} C_1 + C_2 = a_0 \\ 3C_1 - 3C_2 = a_1 \end{cases} \rightarrow \begin{cases} 3C_1 + 3C_2 = 3a_0 \\ 3C_1 - 3C_2 = a_1 \end{cases}$$

$$6C_1 = 3a_0 + a_1 \rightarrow \left| C_1 = \frac{3a_0 + a_1}{6} \right|$$

$$C_2 = a_0 - C_1 \rightarrow \left| C_2 = a_0 - \frac{3a_0 + a_1}{6} = \frac{3a_0 - a_1}{6} \right|$$

$$y(x) = \frac{3a_0 + a_1}{6} e^{3x} + \frac{3a_0 - a_1}{6} e^{-3x}$$

$$y(x) = \sum_{n=0}^{\infty} a_n x^n \quad y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' - 9y = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} - 9 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - 9 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} [(n+2)(n+1) a_{n+2} - 9a_n] x^n = 0$$

$$(n+2)(n+1) a_{n+2} - 9a_n = 0$$

$$a_{n+2} = \frac{9}{(n+2)(n+1)} a_n, \quad n \geq 0$$

$$a_2 = \frac{9}{(2)(1)} a_0 = \frac{9}{2} a_0$$

$$a_3 = \frac{9}{(3)(2)} a_1 = \frac{9}{2 \cdot 3} a_1$$

$$a_4 = \frac{3^2}{(4)(3)} a_2 = \frac{9 \cdot 9}{2 \cdot 3 \cdot 4} a_0 = \frac{3^4}{2 \cdot 3 \cdot 4} a_0$$

$$a_5 = \frac{9}{(5)(4)} a_3 = \frac{3^4}{2 \cdot 3 \cdot 4 \cdot 5} a_1$$

$$a_6 = \frac{3^2}{(6)(5)} a_4 = \frac{3^6}{6!} a_0$$

$$a_7 = \frac{9}{(7)(6)} a_5 = \frac{3^6}{7!} a_1$$

$$a_{2n} = \frac{3^{2n}}{(2n)!} a_0$$

$$a_{2n+1} = \frac{3^{2n}}{(2n+1)!} a_1$$

$$y(x) = a_0 \left( 1 + \frac{3^2}{2!} x^2 + \frac{3^4}{4!} x^4 + \frac{3^6}{6!} x^6 + \dots \right) + a_1 \left( x + \frac{3^2}{3!} x^3 + \frac{3^4}{5!} x^5 + \frac{3^6}{7!} x^7 + \dots \right)$$

$$y(x) = \frac{3a_0 + a_1}{6} e^{3x} + \frac{3a_0 - a_1}{6} e^{-3x}$$

$$= \frac{3a_0 + a_1}{6} \left[ 1 + 3x + \frac{(3x)^2}{2!} + \frac{(3x)^3}{3!} + \dots \right] + \frac{3a_0 - a_1}{6} \left[ 1 - 3x + \frac{(-3x)^2}{2!} + \frac{(-3x)^3}{3!} + \dots \right]$$

$$\begin{aligned}
&= \frac{3a_0}{6} \left[ 1 + 3x + \frac{(3x)^2}{2!} + \frac{(3x)^3}{3!} + \frac{(3x)^4}{4!} + \dots \right] + \frac{a_1}{6} \left[ 1 + 3x + \frac{(3x)^2}{2!} + \frac{(3x)^3}{3!} + \frac{(3x)^4}{4!} + \dots \right] \\
&\quad + \frac{3a_0}{6} \left[ 1 - 3x + \frac{(3x)^2}{2!} - \frac{(3x)^3}{3!} + \dots \right] - \frac{a_1}{6} \left[ 1 - 3x + \frac{(3x)^2}{2!} - \frac{(3x)^3}{3!} + \dots \right] \\
&= \frac{1}{2} a_0 \left[ 1 + 3x + \frac{(3x)^2}{2!} + \frac{(3x)^3}{3!} + \frac{(3x)^4}{4!} + \dots + 1 - 3x + \frac{(3x)^2}{2!} - \frac{(3x)^3}{3!} + \frac{(3x)^4}{4!} + \dots \right] \\
&\quad + \frac{a_1}{6} \left[ 1 + 3x + \frac{(3x)^2}{2!} + \frac{(3x)^3}{3!} + \frac{(3x)^4}{4!} + \dots - 1 + 3x - \frac{(3x)^2}{2!} + \frac{(3x)^3}{3!} - \frac{(3x)^4}{4!} + \dots \right] \\
&= \frac{1}{2} a_0 \left[ 2 + 2 \frac{(3x)^2}{2!} + 2 \frac{(3x)^4}{4!} + \dots \right] + \frac{a_1}{6} \left[ 6x + 2 \frac{(3x)^3}{3!} + 2 \frac{(3x)^5}{5!} + \dots \right] \\
&= a_0 \left[ 1 + \frac{(3x)^2}{2!} + \frac{(3x)^4}{4!} + \dots \right] + a_1 \left[ x + \frac{3^2 x^3}{3!} + \frac{3^4 x^5}{5!} + \dots \right]
\end{aligned}$$

Which are identical.

### Exercise

Find a power series solution.  $y'' + y = 0$

### Solution

The equation  $y'' + y = 0$  has a characteristic equation  $\lambda^2 + 1 = 0 \Rightarrow \boxed{\lambda = \pm i}$

$\therefore$  The general solution:  $y(x) = C_1 \sin x + C_2 \cos x$

With  $y(0) = a_0$  and  $y'(0) = a_1$

$$y(0) = C_1 \sin(0) + C_2 \cos(0) \rightarrow C_2 = a_0$$

$$y'(x) = C_1 \cos x - C_2 \sin x$$

$$y(0) = C_1 \cos(0) - C_2 \sin(0) \rightarrow C_1 = a_1$$

$$y(x) = a_1 \sin x + a_0 \cos x$$

$$= a_0 \cos x + a_1 \sin x$$

$$y(x) = \sum_{n=0}^{\infty} a_n x^n \quad y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' + y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} [(n+2)(n+1)a_{n+2} + a_n] x^n = 0$$

$$(n+2)(n+1)a_{n+2} + a_n = 0 \rightarrow a_{n+2} = \frac{-1}{(n+2)(n+1)} a_n, \quad n \geq 0$$

$$a_2 = \frac{1}{(2)(1)} a_0 = -\frac{1}{2} a_0$$

$$a_3 = \frac{1}{(3)(2)} a_1 = -\frac{1}{2 \cdot 3} a_1$$

$$a_4 = -\frac{1}{(4)(3)} a_2 = \frac{1}{2 \cdot 3 \cdot 4} a_0$$

$$a_5 = -\frac{1}{(5)(4)} a_3 = \frac{1}{2 \cdot 3 \cdot 4 \cdot 5} a_1$$

$$a_6 = -\frac{1}{(6)(5)} a_4 = -\frac{1}{6!} a_0$$

$$a_7 = -\frac{1}{(7)(6)} a_5 = -\frac{1}{7!} a_1$$

$$a_{2n} = \frac{(-1)^n}{(2n)!} a_0$$

$$a_{2n+1} = \frac{(-1)^n}{(2n+1)!} a_1$$

$$y(x) = a_0 \left( 1 - \frac{1}{2!} x^2 + \frac{1}{4!} x^4 - \dots + \frac{(-1)^n}{(2n)!} x^{2n} + \dots \right) + a_1 \left( x - \frac{1}{3!} x^3 + \frac{1}{5!} x^5 - \dots + \frac{(-1)^n}{(2n+1)!} x^{2n+1} + \dots \right)$$

$$\underline{= a_0 \cos x + a_1 \sin x} \quad \checkmark$$

### Exercise

Find a power series solution.  $y'' - y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' - y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ (n+2)(n+1)a_{n+2} - a_n \right] x^n = 0$$

$$\left| a_{n+2} = \frac{1}{(n+2)(n+1)} a_n \right|$$

$$n=0 \rightarrow a_2 = \frac{1}{1 \cdot 2} a_0$$

$$n=1 \rightarrow a_3 = \frac{1}{2 \cdot 3} a_1$$

$$n=2 \rightarrow a_4 = \frac{1}{3 \cdot 4} a_2 = \frac{1}{4!} a_0$$

$$n=3 \rightarrow a_5 = \frac{1}{4 \cdot 5} a_3 = \frac{1}{5!} a_1$$

$$n=4 \rightarrow a_6 = \frac{1}{5 \cdot 6} a_4 = \frac{1}{6!} a_0$$

$$n=5 \rightarrow a_7 = \frac{1}{6 \cdot 7} a_5 = \frac{1}{7!} a_1$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\left| a_{2k} = \frac{1}{(2k)!} a_0 \right|$$

$$\left| a_{2k+1} = \frac{1}{(2k+1)!} a_1 \right|$$

$$y(x) = a_0 \left( 1 + \frac{1}{2!} x^2 + \frac{1}{4!} x^4 + \frac{1}{6!} x^6 + \dots \right) + a_1 \left( x + \frac{1}{3!} x^3 + \frac{1}{5!} x^5 + \dots \right)$$

$$\left| = a_0 \cosh x + a_1 \sinh x \right|$$

### Exercise

Find a power series solution.  $y'' + y = x$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' + y = x$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} + \sum_{n=0}^{\infty} a_n x^n = x$$

$$2a_2 + 6a_3 x + \sum_{n=2}^{\infty} (n+2)(n+1) a_{n+2} x^n + a_0 + a_1 x + \sum_{n=2}^{\infty} a_n x^n - x = 0$$

$$\sum_{n=2}^{\infty} \left[ (n+2)(n+1)a_{n+2} + a_n \right] x^n + a_0 + 2a_2 + (6a_3 + a_1 - 1)x = 0$$

$$\begin{cases} a_0 + 2a_2 = 0 \\ 6a_3 + a_1 - 1 = 0 \\ (n+2)(n+1)a_{n+2} + a_n = 0 \end{cases}$$

$$\begin{cases} a_2 = -\frac{1}{2}a_0 \\ a_3 = -\frac{1}{6}(a_1 - 1) \\ a_{n+2} = -\frac{1}{(n+2)(n+1)}a_n \end{cases}$$

$$n=2 \rightarrow a_4 = -\frac{1}{3 \cdot 4}a_2 = \frac{1}{4!}a_0 \quad n=3 \rightarrow a_5 = -\frac{1}{4 \cdot 5}a_3 = \frac{1}{5!}(a_1 - 1)$$

$$n=4 \rightarrow a_6 = -\frac{1}{5 \cdot 6}a_4 = -\frac{1}{6!}a_0 \quad n=5 \rightarrow a_7 = -\frac{1}{6 \cdot 7}a_5 = -\frac{1}{7!}(a_1 - 1)$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\left| a_{2k} = \frac{(-1)^k}{(2k)!}a_0 \right|$$

$$\left| a_{2k+1} = \frac{(-1)^k}{(2k+1)!}(a_1 - 1) \right|$$

$$y(x) = a_0 + a_1x + a_0 \sum_{k=2}^{\infty} \frac{(-1)^k}{(2k)!}x^{2k} + (a_1 - 1) \sum_{k=3}^{\infty} \frac{(-1)^k}{(2k+1)!}x^{2k+1}$$

$$= a_0 + a_0 \left( -\frac{1}{2!}x^2 + \frac{1}{4!}x^4 - \frac{1}{6!}x^6 + \dots \right) + (a_1 - 1) \left( -\frac{1}{3!}x^3 + \frac{1}{5!}x^5 - \dots \right) + a_1x - x + x$$

$$= a_0 \left( 1 - \frac{1}{2!}x^2 + \frac{1}{4!}x^4 - \frac{1}{6!}x^6 + \dots \right) + (a_1 - 1) \left( -\frac{1}{3!}x^3 + \frac{1}{5!}x^5 - \dots \right) + (a_1 - 1)x + x$$

$$= x + a_0 \left( 1 - \frac{1}{2!}x^2 + \frac{1}{4!}x^4 - \frac{1}{6!}x^6 + \dots \right) + (a_1 - 1) \left( x - \frac{1}{3!}x^3 + \frac{1}{5!}x^5 - \dots \right)$$

$$\left| = x + a_0 \cos x + (a_1 - 1) \sin x \right|$$

### Exercise

Find a power series solution.  $y'' - xy = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' - xy = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - x \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=0}^{\infty} a_n x^{n+1} = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=1}^{\infty} a_{n-1} x^n = 0$$

$$2a_2 + \sum_{n=1}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=1}^{\infty} a_{n-1} x^n = 0$$

$$2a_2 + \sum_{n=1}^{\infty} [(n+2)(n+1) a_{n+2} - a_{n-1}] x^n = 0$$

$$2a_2 = 0 \rightarrow \underline{a_2 = 0}$$

$$(n+2)(n+1) a_{n+2} - a_{n-1} = 0 \rightarrow \underline{a_{n+2} = \frac{a_{n-1}}{(n+1)(n+2)}}$$

$$a_3 = \frac{a_0}{2 \cdot 3} = \frac{1}{6} a_0 \quad a_4 = \frac{a_1}{3 \cdot 4} = \frac{1}{12} a_1 \quad a_5 = \frac{a_2}{4 \cdot 5} = 0$$

$$a_6 = \frac{a_3}{5 \cdot 6} = \frac{1}{180} a_0 \quad a_7 = \frac{a_4}{6 \cdot 7} = \frac{1}{504} a_1 \quad a_8 = \frac{a_5}{7 \cdot 8} = 0$$

$$\vdots \quad \vdots$$

$$\vdots \quad \vdots$$

$$\vdots \quad \vdots$$

$$\left\{ \begin{array}{l} y_1(x) = \left( 1 + \frac{1}{6} x^3 + \frac{1}{180} x^6 + \dots \right) a_0 \\ y_2(x) = \left( x + \frac{1}{12} x^4 + \frac{1}{504} x^7 + \dots \right) a_1 \end{array} \right.$$

### Exercise

Find a power series solution.  $y'' + xy = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' + xy = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + x \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + \sum_{n=0}^{\infty} a_n x^{n+1} = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + \sum_{n=1}^{\infty} a_{n-1} x^n = 0$$

$$2a_2 + \sum_{n=1}^{\infty} (n+2)(n+1) a_{n+2} x^n + \sum_{n=1}^{\infty} a_{n-1} x^n = 0$$

$$2a_2 + \sum_{n=1}^{\infty} [(n+2)(n+1) a_{n+2} + a_{n-1}] x^n = 0$$

$$\begin{cases} 2a_2 = 0 \rightarrow \underline{a_2 = 0} \\ (n+2)(n+1) a_{n+2} + a_{n-1} = 0 \end{cases}$$

$$\underline{a_{n+2} = -\frac{a_{n-1}}{(n+1)(n+2)}}$$

$$a_0$$

$$a_1$$

$$a_2 = 0$$

$$n=1 \rightarrow a_3 = -\frac{a_0}{2 \cdot 3} = -\frac{1}{6} a_0$$

$$n=2 \rightarrow a_4 = -\frac{1}{3 \cdot 4} a_1$$

$$n=3 \rightarrow a_5 = -\frac{a_2}{20} = 0$$

$$n=4 \rightarrow a_6 = -\frac{a_3}{5 \cdot 6} = -\frac{1}{180} a_0$$

$$n=5 \rightarrow a_7 = -\frac{a_4}{6 \cdot 7} = -\frac{1}{504} a_1$$

$$n=6 \rightarrow a_8 = -\frac{a_5}{56} = 0$$

$$n=7 \rightarrow a_9 = -\frac{a_6}{8 \cdot 9} = -\frac{1}{12,960} a_0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$



$$\left\{ \begin{array}{l} y_1(x) = \left( 1 - \frac{1}{6}x^3 + \frac{1}{180}x^6 - \frac{1}{12,960}x^9 + \dots \right) a_0 \\ y_2(x) = \left( x - \frac{1}{12}x^4 + \frac{1}{504}x^7 - \dots \right) a_1 \end{array} \right|$$

$$\underline{y(x) = \left( 1 - \frac{1}{6}x^3 + \frac{1}{180}x^6 - \frac{1}{12,960}x^9 + \dots \right) a_0 + \left( x - \frac{1}{12}x^4 + \frac{1}{504}x^7 - \dots \right) a_1}$$

### Exercise

Find a power series solution.  $y'' + xy' + y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' + xy' + y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + x \sum_{n=1}^{\infty} n a_n x^{n-1} + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + \sum_{n=1}^{\infty} n a_n x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + \sum_{n=0}^{\infty} n a_n x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} [(n+2)(n+1) a_{n+2} + (n+1) a_n] x^n = 0$$

$$(n+2)(n+1) a_{n+2} + (n+1) a_n = 0$$

$$\underline{a_{n+2} = -\frac{a_n}{n+2}}$$

$$a_0$$

$$n=0 \rightarrow a_2 = -\frac{1}{2} a_0$$

$$a_1$$

$$n=1 \rightarrow a_3 = -\frac{1}{3} a_1$$

$$n=2 \rightarrow a_4 = -\frac{1}{4}a_2 = \frac{1}{4 \cdot 2}a_0$$

$$n=3 \rightarrow a_5 = -\frac{a_3}{5} = \frac{1}{3 \cdot 5}a_1$$

$$n=4 \rightarrow a_6 = -\frac{a_4}{6} = -\frac{1}{6 \cdot 4 \cdot 2}a_0$$

$$n=5 \rightarrow a_7 = -\frac{a_5}{7} = -\frac{1}{7 \cdot 5 \cdot 3}a_1$$

$$n=6 \rightarrow a_8 = -\frac{a_6}{8} = \frac{1}{8 \cdot 6 \cdot 4 \cdot 2}a_0$$

$$n=7 \rightarrow a_9 = -\frac{a_7}{9} = \frac{1}{9 \cdot 7 \cdot 5 \cdot 3}a_1$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$a_{2n} = \frac{(-1)^n a_0}{(2n)(2n-2)\cdots 4 \cdot 2} = \frac{(-1)^n}{n!} a_0$$

$$a_{2n+1} = \frac{(-1)^n a_1}{(2n+1)(2n-1)\cdots 5 \cdot 3} = \frac{(-1)^n}{(2n+1)!!} a_1$$

$$y(x) = a_0 \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} x^{2n} + a_1 \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!!} x^{2n+1}$$

$$y(x) = a_0 \left( 1 - \frac{1}{2}x^2 + \frac{1}{8}x^4 - \frac{1}{48}x^6 + \cdots \right) + a_1 \left( x - \frac{1}{3}x^3 + \frac{1}{15}x^5 - \frac{1}{105}x^7 + \cdots \right)$$

## Exercise

Find a power series solution.  $y'' - xy' - y = 0$

## Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' - xy' - y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - x \sum_{n=1}^{\infty} n a_n x^{n-1} - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=1}^{\infty} n a_n x^n - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=0}^{\infty} n a_n x^n - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ (n+2)(n+1)a_{n+2} - (n+1)a_n \right] x^n = 0$$

$$(n+2)(n+1)a_{n+2} - (n+1)a_n = 0$$

$$\underline{a_{n+2} = \frac{a_n}{n+2}}$$

$a_0$

$$n=0 \rightarrow a_2 = \frac{1}{2}a_0$$

$$n=2 \rightarrow a_4 = \frac{1}{4}a_2 = \frac{1}{4 \cdot 2}a_0$$

$$n=4 \rightarrow a_6 = \frac{a_4}{6} = \frac{1}{6 \cdot 4 \cdot 2}a_0$$

$$n=6 \rightarrow a_8 = \frac{a_6}{8} = \frac{1}{8 \cdot 6 \cdot 4 \cdot 2}a_0$$

$\vdots \quad \vdots \quad \vdots \quad \vdots$

$$a_{2n} = \frac{a_0}{(2n)(2n-2)\cdots 4 \cdot 2} = \frac{1}{n! \cdot 2^n}a_0$$

$a_1$

$$n=1 \rightarrow a_3 = \frac{1}{3}a_1$$

$$n=3 \rightarrow a_5 = \frac{a_3}{5} = \frac{1}{3 \cdot 5}a_1$$

$$n=5 \rightarrow a_7 = \frac{a_5}{7} = \frac{1}{7 \cdot 5 \cdot 3}a_1$$

$$n=7 \rightarrow a_9 = \frac{a_7}{9} = \frac{1}{9 \cdot 7 \cdot 5 \cdot 3}a_1$$

$\vdots \quad \vdots \quad \vdots \quad \vdots$

$$a_{2n} = \frac{a_1}{(2n+1)(2n-1)\cdots 5 \cdot 3} = \frac{1}{(2n+1)!!}a_1$$

$$\underline{y(x) = a_0 \sum_{n=0}^{\infty} \frac{1}{n! \cdot 2^n} x^{2n} + a_1 \sum_{n=0}^{\infty} \frac{1}{(2n+1)!!} x^{2n+1}}$$

$$\underline{y(x) = a_0 \left( 1 + \frac{x^2}{2} + \frac{x^4}{2^2 \cdot 2} + \cdots + \frac{x^{2n}}{2^n \cdot n!} \right) + a_1 \left( x + \frac{x^3}{3} + \frac{x^5}{5 \cdot 3} + \cdots + \frac{2^n \cdot n!}{(2n+1)!!} x^{2n+1} \right)}$$

## Exercise

Find a power series solution.  $y'' + x^2 y = 0$

## Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' + x^2 y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n + x^2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n + \sum_{n=0}^{\infty} a_n x^{n+2} = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n + \sum_{n=2}^{\infty} a_{n-2}x^n = 0$$

$$2a_2 + 6a_3x + \sum_{n=2}^{\infty} (n+2)(n+1)a_{n+2}x^n + \sum_{n=2}^{\infty} a_{n-2}x^n = 0$$

$$2a_2 + 6a_3x + \sum_{n=2}^{\infty} [(n+2)(n+1)a_{n+2} + a_{n-2}]x^n = 0$$

$$2a_2 = 0 \rightarrow \underline{a_2 = 0} \quad 6a_3 = 0 \rightarrow \underline{a_3 = 0}$$

$$(n+2)(n+1)a_{n+2} + a_{n-2} = 0 \rightarrow a_{n+2} = \frac{-a_{n-2}}{(n+1)(n+2)}$$

$$a_4 = \frac{-a_0}{3 \cdot 4} = -\frac{1}{12}a_0$$

$$a_5 = \frac{-a_1}{4 \cdot 5} = -\frac{1}{20}a_1$$

$$a_6 = \frac{-a_2}{5 \cdot 6} = 0$$

$$a_7 = \frac{-a_3}{6 \cdot 7} = 0$$

$$a_8 = \frac{-a_4}{7 \cdot 8} = \frac{1}{672}a_0$$

$$a_9 = \frac{-a_5}{8 \cdot 9} = \frac{1}{1,440}a_1$$

$$a_{10} = \frac{-a_6}{9 \cdot 10} = 0$$

$$a_{11} = \frac{-a_7}{10 \cdot 11} = 0$$

$$\vdots$$

$$\vdots$$

$$\vdots$$

$$\vdots$$

$$\left\{ \begin{array}{l} y_1(x) = \left( 1 - \frac{1}{12}x^4 + \frac{1}{672}x^8 - \dots \right) a_0 \\ y_2(x) = \left( x - \frac{1}{20}x^5 + \frac{1}{1,440}x^9 + \dots \right) a_1 \end{array} \right.$$

## Exercise

Find a power series solution.  $y'' + k^2 x^2 y = 0$

## Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1)a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n$$

$$y'' + k^2 x^2 y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n + k^2 x^2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$2a_2 + 6a_3 x + \sum_{n=2}^{\infty} (n+2)(n+1)a_{n+2} x^n + \sum_{n=2}^{\infty} k^2 a_{n-2} x^n = 0$$

$$2a_2 + 6a_3 x + \sum_{n=2}^{\infty} \left[ (n+2)(n+1)a_{n+2} + k^2 a_{n-2} \right] x^n = 0$$

$$\begin{cases} a_2 = 0 \\ a_3 = 0 \\ (n+2)(n+1)a_{n+2} + k^2 a_{n-2} = 0 \end{cases}$$

$$\boxed{a_{n+2} = -\frac{k^2}{(n+1)(n+2)} a_{n-2} \quad (n \geq 2)}$$

$$n=2 \rightarrow a_4 = -\frac{k^2}{3 \cdot 4} a_0$$

$$n=3 \rightarrow a_5 = -\frac{k^2}{4 \cdot 5} a_1$$

$$n=6 \rightarrow a_8 = -\frac{k^2}{7 \cdot 8} a_4 = \frac{k^4}{3 \cdot 4 \cdot 7 \cdot 8} a_0$$

$$n=7 \rightarrow a_9 = -\frac{k^2}{8 \cdot 9} a_5 = \frac{k^4}{4 \cdot 5 \cdot 8 \cdot 9} a_1$$

$$n=10 \rightarrow a_{12} = -\frac{k^2}{11 \cdot 12} a_8 = \frac{k^6}{3 \cdot 4 \cdot 7 \cdot 8 \cdot 11 \cdot 12} a_0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$a_{4m} = -\frac{k^2}{(4m)(4m-1)} a_{4m-4}$$

$$a_{4m+1} = -\frac{k^2}{(4m)(4m+1)} a_{4m-3}$$

$$n=4 \rightarrow a_6 = -\frac{k^2}{5 \cdot 6} a_2 = 0$$

$$n=5 \rightarrow a_7 = -\frac{k^2}{6 \cdot 7} a_3 = 0$$

$$n=8 \rightarrow a_{10} = -\frac{k^2}{7 \cdot 8} a_6 = 0$$

$$n=9 \rightarrow a_{11} = -\frac{k^2}{10 \cdot 11} a_7 = 0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\boxed{y(x) = a_0 \left( 1 - \frac{k^2}{3 \cdot 4} x^4 + \frac{k^4}{3 \cdot 4 \cdot 7 \cdot 8} x^8 - \frac{k^6}{3 \cdot 4 \cdot 7 \cdot 8 \cdot 11 \cdot 12} x^{12} + \dots \right) + a_1 \left( x - \frac{k^2}{4 \cdot 5} x^5 + \frac{k^4}{4 \cdot 5 \cdot 8 \cdot 9} x^9 - \frac{k^6}{4 \cdot 5 \cdot 8 \cdot 9 \cdot 12 \cdot 13} x^{13} + \dots \right)}$$

### Exercise

Find a power series solution.  $y'' + 3xy' + 3y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' + 3xy' + 3y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + 3x \sum_{n=1}^{\infty} n a_n x^{n-1} + 3 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + \sum_{n=0}^{\infty} 3n a_n x^n + \sum_{n=0}^{\infty} 3a_n x^n = 0$$

$$\sum_{n=0}^{\infty} [(n+2)(n+1) a_{n+2} + (3n+3) a_n] x^n = 0$$

$$(n+2)(n+1) a_{n+2} + 3(n+1) a_n = 0$$

$$\underline{a_{n+2} = -\frac{3}{n+2} a_n}$$

$$a_0$$

$$n=2 \rightarrow a_2 = -\frac{3}{2} a_0$$

$$n=4 \rightarrow a_4 = -\frac{3}{4} a_2 = \frac{3^2}{2^2 \cdot 2} a_0$$

$$n=6 \rightarrow a_6 = -\frac{3}{6} a_4 = \frac{3^3}{2^3 \cdot 2 \cdot 3} a_0$$

$$n=8 \rightarrow a_8 = -\frac{3}{8} a_6 = \frac{3^4}{2^4 \cdot 2 \cdot 3 \cdot 4} a_0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$a_{2k} = \frac{(-3)^k}{2^k k!} a_0$$

$$a_1$$

$$n=3 \rightarrow a_3 = -\frac{3}{3} a_1$$

$$n=5 \rightarrow a_5 = -\frac{3}{5} a_3 = \frac{3^2}{3 \cdot 5} a_1$$

$$n=7 \rightarrow a_7 = -\frac{3}{7} a_5 = \frac{3^3}{3 \cdot 5 \cdot 7} a_1$$

$$n=9 \rightarrow a_9 = -\frac{3}{9} a_7 = \frac{3^3}{3 \cdot 5 \cdot 7 \cdot 9} a_1$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$a_{2k+1} = \frac{(-3)^k}{3 \cdot 5 \cdot 7 \cdots (2k+1)} a_1$$

$$y(x) = a_0 \left( 1 + \sum_{k=1}^{\infty} \frac{(-3)^k}{2^k k!} x^{2k} \right) + a_1 \left( x + \sum_{k=1}^{\infty} \frac{(-3)^k}{3 \cdot 5 \cdot 7 \cdots (2k+1)} x^{2k+1} \right)$$

$$= a_0 \left( 1 - \frac{3}{2} x^2 + \frac{9}{8} x^4 - \frac{27}{56} x^6 + \cdots \right) + a_1 \left( x - x^3 + \frac{3^2}{3 \cdot 5} x^5 - \frac{27}{3 \cdot 5 \cdot 7} x^7 + \cdots \right)$$


---

### Exercise

Find a power series solution.  $y'' - 2xy' + y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' - 2xy' + y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - 2x \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=1}^{\infty} 2n a_n x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$2a_2 + \sum_{n=1}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=1}^{\infty} 2n a_n x^n + a_0 + \sum_{n=1}^{\infty} a_n x^n = 0$$

$$2a_2 + a_0 + \sum_{n=1}^{\infty} [(n+2)(n+1) a_{n+2} - 2n a_n + a_n] x^n = 0$$

$$2a_2 + a_0 = 0 \rightarrow \underline{a_2 = -\frac{1}{2} a_0}$$

$$(n+2)(n+1) a_{n+2} - (2n-1) a_n = 0 \rightarrow \underline{a_{n+2} = \frac{(2n-1) a_n}{(n+1)(n+2)}}$$

$$a_3 = \frac{a_1}{2 \cdot 3} = \frac{1}{6} a_1$$

$$a_4 = \frac{3a_2}{3 \cdot 4} = -\frac{1}{4} \frac{1}{4} a_0 = -\frac{1}{8} a_0$$

$$\begin{aligned}
 a_5 &= \frac{5a_3}{4 \cdot 5} = \frac{1}{2 \cdot 3 \cdot 4} a_1 = \frac{1}{4!} a_1 & a_6 &= \frac{7a_4}{5 \cdot 6} = -\frac{7}{240} a_0 \\
 a_7 &= \frac{9a_5}{6 \cdot 7} = \frac{1}{112} a_1 & & \vdots \\
 & \vdots & & \vdots
 \end{aligned}$$

$$\left\{ \begin{aligned} y_1(x) &= \left( 1 - \frac{1}{2}x^2 - \frac{1}{8}x^4 - \frac{7}{240}x^6 - \dots \right) a_0 \\ y_2(x) &= \left( x + \frac{1}{6}x^3 + \frac{1}{24}x^5 + \frac{1}{112}x^7 + \dots \right) a_1 \end{aligned} \right.$$

### Exercise

Find a power series solution.  $y'' - xy' + 2y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' - xy' + 2y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - x \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n + 2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=1}^{\infty} n a_n x^n + \sum_{n=0}^{\infty} 2a_n x^n = 0$$

$$2a_2 + \sum_{n=1}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=1}^{\infty} n a_n x^n + 2a_0 + \sum_{n=1}^{\infty} 2a_n x^n = 0$$

$$2a_2 + 2a_0 + \sum_{n=1}^{\infty} [(n+2)(n+1) a_{n+2} - n a_n + 2a_n] x^n = 0$$

$$2a_2 + 2a_0 = 0 \rightarrow \underline{a_2 = -a_0}$$

$$(n+2)(n+1) a_{n+2} - (n-2) a_n = 0 \rightarrow a_{n+2} = \frac{(n-2) a_n}{(n+1)(n+2)}$$



$$a_3 = \frac{-a_1}{2 \cdot 3} = -\frac{1}{6}a_1$$

$$a_5 = \frac{a_3}{4 \cdot 5} = -\frac{1}{5!}a_1$$

$$a_7 = \frac{3a_5}{6 \cdot 7} = \frac{3}{7!}a_1$$

$$a_9 = \frac{5a_7}{8 \cdot 9} = \frac{3 \cdot 5}{9!}a_1$$

$$\vdots \quad \vdots$$

$$a_4 = 0$$

$$a_6 = \frac{2a_4}{5 \cdot 6} = 0$$

$$a_8 = \frac{4a_6}{5 \cdot 6} = 0$$

$$\vdots \quad \vdots$$

$$\left\{ \begin{array}{l} y_1(x) = 1 - x^2 \\ y_2(x) = \left( x - \frac{1}{6}x^3 + \frac{1}{5!}x^5 + \frac{3}{7!}x^7 + \dots \right) a_1 \end{array} \right.$$

### Exercise

Find a power series solution.  $y'' - xy' - x^2y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' - xy' - x^2y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - x \sum_{n=1}^{\infty} n a_n x^{n-1} - x^2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=1}^{\infty} n a_n x^n - \sum_{n=0}^{\infty} a_n x^{n+2} = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=0}^{\infty} n a_n x^n - \sum_{n=2}^{\infty} a_{n-2} x^n = 0$$

$$2a_2 + 6a_3x + \sum_{n=2}^{\infty} (n+2)(n+1) a_{n+2} x^n - a_1x - \sum_{n=2}^{\infty} n a_n x^n - \sum_{n=2}^{\infty} a_{n-2} x^n = 0$$

$$2a_2 + 6a_3x - a_1x + \sum_{n=2}^{\infty} [(n+2)(n+1)a_{n+2} - na_n - a_{n-2}]x^n = 0$$

$$\left\{ \begin{array}{l} 2a_2 = 0 \rightarrow \underline{a_2 = 0} \\ (6a_3 - a_1)x = 0 \rightarrow \underline{a_3 = \frac{1}{6}a_1} \\ (n+1)(n+2)a_{n+2} - na_n - a_{n-2} = 0 \\ \underline{a_{n+2} = \frac{na_n + a_{n-2}}{(n+1)(n+2)}} \end{array} \right.$$

$$\begin{array}{c} a_0 \\ a_2 = 0 \end{array}$$

$$\begin{array}{c} a_1 \\ a_3 = \frac{1}{6}a_1 \end{array}$$

$$n = 2 \rightarrow a_4 = \frac{2a_2 + a_0}{3 \cdot 4} = \frac{1}{12}a_0$$

$$n = 3 \rightarrow a_5 = \frac{3a_3 + a_1}{20} = \frac{1}{20}\left(\frac{3}{6} + 1\right)a_1 = \frac{1}{12}a_1$$

$$n = 4 \rightarrow a_6 = \frac{4a_4 + a_2}{5 \cdot 6} = \frac{1}{90}a_0$$

$$n = 5 \rightarrow a_7 = \frac{5a_5 + a_3}{6 \cdot 7} = \frac{1}{42}\left(\frac{5}{12} + \frac{1}{6}\right)a_1 = \frac{1}{72}a_1$$

$$n = 6 \rightarrow a_8 = \frac{6a_6 + a_4}{7 \cdot 8} = \frac{1}{56}\left(\frac{1}{15} + \frac{1}{12}\right)a_0 = \frac{3}{1120}a_0$$

$$\begin{array}{cccc} \vdots & \vdots & \vdots & \vdots \end{array}$$

$$\begin{array}{cccc} \vdots & \vdots & \vdots & \vdots \end{array}$$

$$\underline{y(x) = a_0 \left(1 + \frac{1}{12}x^2 + \frac{1}{90}x^4 + \frac{3}{1120}x^6 + \dots\right) + a_1 \left(x + \frac{1}{12}x^3 + \frac{1}{72}x^5 + \dots\right)}$$

## Exercise

Find a power series solution.  $y'' + x^2y' + xy = 0$

## Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} na_n x^{n-1} = \sum_{n=0}^{\infty} (n+1)a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1)a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n$$

$$y'' + x^2y' + xy = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n + x^2 \sum_{n=0}^{\infty} (n+1)a_{n+1} x^n + x \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n + \sum_{n=0}^{\infty} (n+1)a_{n+1}x^{n+2} + \sum_{n=0}^{\infty} a_n x^{n+1} = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n + \sum_{n=2}^{\infty} (n-1)a_{n-1}x^n + \sum_{n=1}^{\infty} a_{n-1}x^n = 0$$

$$2a_2 + 6a_3x + \sum_{n=2}^{\infty} (n+2)(n+1)a_{n+2}x^n + \sum_{n=2}^{\infty} (n-1)a_{n-1}x^n + a_0x + \sum_{n=2}^{\infty} a_{n-1}x^n = 0$$

$$2a_2 + (6a_3 + a_0)x + \sum_{n=2}^{\infty} [(n+2)(n+1)a_{n+2} + (n-1)a_{n-1} + a_{n-1}]x^n = 0$$

$$2a_2 + (6a_3 + a_0)x = 0 \Rightarrow \begin{cases} a_2 = 0 \\ a_3 = -\frac{1}{6}a_0 \end{cases}$$

$$(n+2)(n+1)a_{n+2} + na_{n-1} = 0 \Rightarrow a_{n+2} = -\frac{n}{(n+1)(n+2)}a_{n-1}$$

$$\begin{array}{lll} a_4 = -\frac{2}{3 \cdot 4}a_1 = -\frac{1}{6}a_1 & a_5 = -\frac{3}{4 \cdot 5}a_2 = 0 & a_6 = -\frac{4}{5 \cdot 6}a_3 = \frac{1}{45}a_0 \\ a_7 = -\frac{5}{6 \cdot 7}a_4 = \frac{5}{252}a_1 & a_8 = -\frac{6}{7 \cdot 8}a_5 = 0 & a_9 = -\frac{7}{8 \cdot 9}a_3 = -\frac{7}{3,240}a_0 \\ \vdots & \vdots & \vdots \end{array}$$

$$\left\{ \begin{array}{l} y_1(x) = \left( 1 - \frac{1}{6}x^3 + \frac{1}{45}x^6 - \frac{7}{3,240}x^9 + \dots \right) a_0 \\ y_2(x) = \left( x - \frac{1}{6}x^4 + \frac{5}{252}x^7 - \dots \right) a_1 \end{array} \right.$$

## Exercise

Find a power series solution.  $y'' + x^2y' + 2xy = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} na_n x^{n-1} = \sum_{n=0}^{\infty} (n+1)a_{n+1}x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1)a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n$$

$$y'' + x^2 y' + 2xy = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n + x^2 \sum_{n=0}^{\infty} (n+1)a_{n+1}x^n + 2x \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n + \sum_{n=0}^{\infty} (n+1)a_{n+1}x^{n+2} + \sum_{n=0}^{\infty} 2a_n x^{n+1} = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n + \sum_{n=2}^{\infty} (n-1)a_{n-1}x^n + \sum_{n=1}^{\infty} 2a_{n-1}x^n = 0$$

$$2a_2 + 6a_3x + \sum_{n=2}^{\infty} (n+2)(n+1)a_{n+2}x^n + \sum_{n=2}^{\infty} (n-1)a_{n-1}x^n + 2a_0x + \sum_{n=2}^{\infty} 2a_{n-1}x^n = 0$$

$$2a_2 + (6a_3 + 2a_0)x + \sum_{n=2}^{\infty} [(n+2)(n+1)a_{n+2} + (n-1)a_{n-1} + 2a_{n-1}]x^n = 0$$

$$2a_2 + (6a_3 + 2a_0)x = 0 \Rightarrow \begin{cases} a_2 = 0 \\ a_3 = -\frac{1}{3}a_0 \end{cases}$$

$$(n+2)(n+1)a_{n+2} + (n+1)a_{n-1} = 0 \Rightarrow a_{n+2} = -\frac{a_{n-1}}{n+2}$$

$$\boxed{a_{n+3} = -\frac{a_n}{n+3}}$$

$$a_0$$

$$a_1$$

$$a_2 = 0$$

$$a_3 = -\frac{1}{3}a_0$$

$$n=1 \rightarrow a_4 = -\frac{1}{4}a_1$$

$$n=2 \rightarrow a_5 = -\frac{a_2}{5} = 0$$

$$n=3 \rightarrow a_6 = -\frac{a_3}{6} = \frac{1}{2 \cdot 3^2}a_0$$

$$n=4 \rightarrow a_7 = -\frac{a_4}{7} = \frac{1}{7 \cdot 4}a_1$$

$$n=5 \rightarrow a_8 = -\frac{a_5}{8} = 0$$

$$n=6 \rightarrow a_9 = -\frac{a_6}{9} = -\frac{1}{3! \cdot 3^3}a_0$$

$$n=7 \rightarrow a_{10} = -\frac{a_7}{10} = -\frac{1}{10 \cdot 7 \cdot 4 \cdot 1}a_1$$

$$n=8 \rightarrow a_{11} = 0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$a_{3n} = \frac{(-1)^n}{n! \cdot 3^n}$$

$$a_{3n+1} = \frac{(-1)^n}{1 \cdot 4 \cdot 7 \cdots (3n+1)}$$

$$a_{3n+2} = 0$$

$$\boxed{y(x) = a_0 \sum_{n=0}^{\infty} \frac{(-1)^n}{n! \cdot 3^n} x^{3n} + a_1 \sum_{n=0}^{\infty} \frac{(-1)^n}{1 \cdot 4 \cdot 7 \cdots (3n+1)} x^{3n+1}}$$

$$\boxed{y(x) = a_0 \left( 1 - \frac{1}{3}x^3 + \frac{1}{18}x^6 - \frac{1}{162}x^9 + \cdots \right) + a_1 \left( x - \frac{1}{4}x^4 + \frac{1}{28}x^7 - \frac{1}{280}x^{10} + \cdots \right)}$$

### Exercise

Find a power series solution.  $y'' - x^2 y' - 3xy = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' - x^2 y' - 3xy = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - x^2 \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n - 3x \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=0}^{\infty} (n+1) a_{n+1} x^{n+2} - \sum_{n=0}^{\infty} 3a_n x^{n+1} = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=2}^{\infty} (n-1) a_{n-1} x^n - \sum_{n=1}^{\infty} 3a_{n-1} x^n = 0$$

$$2a_2 + 6a_3 x + \sum_{n=2}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=2}^{\infty} (n-1) a_{n-1} x^n - 3a_0 x - \sum_{n=2}^{\infty} 3a_{n-1} x^n = 0$$

$$2a_2 + 3(2a_3 - a_0)x + \sum_{n=2}^{\infty} [(n+2)(n+1) a_{n+2} - (n-1) a_{n-1}] x^n = 0$$

$$\begin{cases} 2a_2 = 0 & \rightarrow a_2 = 0 \\ 2a_3 - a_0 = 0 & \rightarrow a_3 = \frac{1}{2}a_0 \end{cases}$$

$$(n+2)(n+1) a_{n+2} = (n-1) a_{n-1}$$

$$a_{n+2} = \frac{a_{n-1}}{n+1} \rightarrow a_{n+3} = \frac{a_n}{n+2}$$

$$a_0$$

$$a_1$$

$$a_2 = 0$$

$$a_3 = \frac{1}{2}a_0$$

$$n=1 \rightarrow a_4 = \frac{1}{3}a_1$$

$$n=2 \rightarrow a_5 = \frac{a_2}{5} = 0$$

$$n=3 \rightarrow a_6 = \frac{a_3}{5} = \frac{1}{2 \cdot 5}a_0$$

$$n=4 \rightarrow a_7 = \frac{a_4}{6} = \frac{1}{2 \cdot 3^2}a_1$$

$$n=5 \rightarrow a_8 = 0$$

$$\begin{array}{lll}
 n=6 \rightarrow a_9 = \frac{a_6}{8} = -\frac{1}{2 \cdot 5 \cdot 8} a_0 & n=7 \rightarrow a_{10} = \frac{a_7}{9} = \frac{1}{2 \cdot 3 \cdot 3^3} a_1 & n=8 \rightarrow a_{11} = 0 \\
 \vdots & \vdots & \vdots
 \end{array}$$

$$a_{3n} = \frac{1}{2 \cdot 5 \cdot 8 \cdots (3n-1)}$$

$$a_{3n+1} = \frac{1}{n! 3^n}$$

$$a_{3n+2} = 0$$

$$y(x) = a_0 \sum_{n=0}^{\infty} \frac{x^{3n}}{2 \cdot 5 \cdot 8 \cdots (3n-1)} + a_1 \sum_{n=0}^{\infty} \frac{x^{3n+1}}{n! 3^n}$$

$$y(x) = a_0 \left( 1 + \frac{1}{2}x^3 + \frac{1}{10}x^6 + \frac{1}{80}x^9 + \cdots \right) + a_1 \left( x + \frac{1}{3}x^4 + \frac{1}{18}x^7 + \frac{1}{162}x^{10} + \cdots \right)$$

### Exercise

Find a power series solution.  $y'' + 2xy' + 2y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' + 2xy' + 2y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + 2x \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n + 2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + \sum_{n=0}^{\infty} 2(n+1) a_{n+1} x^{n+1} + \sum_{n=0}^{\infty} 2a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + \sum_{n=1}^{\infty} 2n a_n x^n + \sum_{n=0}^{\infty} 2a_n x^n = 0$$

$$2a_2 + \sum_{n=1}^{\infty} (n+2)(n+1) a_{n+2} x^n + \sum_{n=1}^{\infty} 2n a_n x^n + 2a_0 + \sum_{n=1}^{\infty} 2a_n x^n = 0$$

$$2a_2 + 2a_0 + \sum_{n=1}^{\infty} \left[ (n+2)(n+1)a_{n+2} + 2na_n + 2a_n \right] x^n = 0$$

$$2a_2 + 2a_0 = 0 \rightarrow \underline{a_2 = -a_0}$$

$$(n+2)(n+1)a_{n+2} + 2(n+1)a_n = 0 \rightarrow a_{n+2} = -\frac{2}{n+2}a_n \quad n=1, 2, \dots$$

$$a_3 = -\frac{2}{3}a_1$$

$$a_4 = -\frac{2}{4}a_2 = \frac{1}{2}a_0$$

$$a_5 = -\frac{2}{5}a_3 = \frac{4}{15}a_1$$

$$a_6 = -\frac{2}{6}a_4 = -\frac{1}{6}a_0$$

$$a_7 = -\frac{2}{7}a_5 = -\frac{8}{105}a_1$$

$$a_8 = -\frac{2}{8}a_6 = \frac{1}{24}a_0$$

$$\vdots$$

$$\vdots$$

$$\left\{ \begin{array}{l} y_1(x) = \left( 1 - x^2 + \frac{1}{2!}x^4 - \frac{1}{3!}x^6 + \frac{1}{4!}x^8 - \dots \right) a_0 \\ y_2(x) = \left( x - \frac{2}{3}x^3 + \frac{4}{15}x^5 - \frac{8}{105}x^7 + \dots \right) a_1 \end{array} \right.$$

### Exercise

Find a power series solution.  $2y'' + xy' + y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$2y'' + xy' + y = 0$$

$$2 \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + x \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} 2(n+2)(n+1) a_{n+2} x^n + \sum_{n=0}^{\infty} (n+1) a_{n+1} x^{n+1} + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} 2(n+2)(n+1) a_{n+2} x^n + \sum_{n=1}^{\infty} n a_n x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$4a_2 + \sum_{n=1}^{\infty} 2(n+2)(n+1)a_{n+2}x^n + \sum_{n=1}^{\infty} na_nx^n + a_0 + \sum_{n=1}^{\infty} a_nx^n = 0$$

$$4a_2 + a_0 + \sum_{n=1}^{\infty} [2(n+2)(n+1)a_{n+2} + (n+1)a_n]x^n = 0$$

$$4a_2 + a_0 = 0 \rightarrow a_2 = -\frac{1}{4}a_0$$

$$2(n+2)(n+1)a_{n+2} + (n+1)a_n = 0$$

$$a_{n+2} = -\frac{1}{2(n+2)}a_n$$

$a_0$

$$n=0 \rightarrow a_2 = -\frac{1}{4}a_0$$

$$n=2 \rightarrow a_4 = -\frac{1}{8}a_2 = \frac{1}{2^4 \cdot 2}a_0$$

$$n=4 \rightarrow a_6 = -\frac{1}{2 \cdot 6}a_4 = -\frac{1}{2^6 \cdot 2 \cdot 3}a_0$$

$$n=6 \rightarrow a_8 = -\frac{1}{2 \cdot 8}a_6 = \frac{1}{2^8 \cdot 4!}a_0$$

$\vdots \quad \vdots \quad \vdots \quad \vdots$

$$n \geq 1 \quad a_{2n} = \frac{(-1)^n}{2^{2n} n!}a_0$$

$a_1$

$$n=1 \rightarrow a_3 = -\frac{1}{6}a_1$$

$$n=3 \rightarrow a_5 = -\frac{1}{2 \cdot 5}a_3 = \frac{1}{2^2 \cdot 3 \cdot 5}a_1$$

$$n=5 \rightarrow a_7 = -\frac{1}{2 \cdot 7}a_5 = -\frac{1}{2^3 \cdot 3 \cdot 5 \cdot 7}a_1$$

$\vdots \quad \vdots \quad \vdots \quad \vdots$

$$n \geq 1 \quad a_{2n+1} = \frac{(-1)^n}{2^n 1 \cdot 3 \cdot 5 \cdots (2n+1)}a_1$$

$$= \frac{(-1)^n}{2^n (2n+1)!!}a_1$$

$$y(x) = a_0 \left( 1 - \frac{1}{4}x^2 + \frac{1}{32}x^4 - \dots \right) + a_1 \left( x - \frac{1}{6}x^3 + \frac{1}{60}x^5 - \dots \right)$$

$$y(x) = a_0 \sum_{n=0}^{\infty} \frac{(-1)^n}{2^{2n} n!} x^{2n} + a_1 \sum_{n=0}^{\infty} \frac{(-1)^n}{2^n (2n+1)!!} x^{2n+1}$$

## Exercise

Find a power series solution.  $3y'' + xy' - 4y = 0$

## Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$



$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$3y'' + xy' - 4y = 0$$

$$3 \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + x \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n - 4 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} 3(n+2)(n+1) a_{n+2} x^n + \sum_{n=0}^{\infty} (n+1) a_{n+1} x^{n+1} - \sum_{n=0}^{\infty} 4a_n x^n = 0$$

$$\sum_{n=0}^{\infty} 3(n+2)(n+1) a_{n+2} x^n + \sum_{n=1}^{\infty} n a_n x^n - \sum_{n=0}^{\infty} 4a_n x^n = 0$$

$$6a_2 + \sum_{n=1}^{\infty} 3(n+2)(n+1) a_{n+2} x^n + \sum_{n=1}^{\infty} n a_n x^n - 4a_0 - \sum_{n=1}^{\infty} 4a_n x^n = 0$$

$$6a_2 - 4a_0 + \sum_{n=1}^{\infty} [3(n+2)(n+1) a_{n+2} + (n-4) a_n] x^n = 0$$

$$6a_2 - 4a_0 \rightarrow \underline{a_2 = \frac{2}{3}a_0}$$

$$3(n+2)(n+1) a_{n+2} + (n-4) a_n = 0$$

$$\underline{a_{n+2} = -\frac{(n-4)}{3(n+1)(n+2)} a_n}$$

$$a_0$$

$$a_2 = \frac{2}{3} a_0$$

$$n=2 \rightarrow a_4 = \frac{2}{36} a_2 = \frac{1}{27} a_0$$

$$n=4 \rightarrow a_6 = 0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$a_1$$

$$n=1 \rightarrow a_3 = \frac{3}{2 \cdot 3 \cdot 3} a_1 = \frac{1}{2 \cdot 3} a_1$$

$$n=3 \rightarrow a_5 = \frac{1}{4 \cdot 5 \cdot 3} a_3 = \frac{1}{5! \cdot 3} a_1$$

$$n=5 \rightarrow a_7 = -\frac{1}{3 \cdot 6 \cdot 7} a_5 = -\frac{1}{7! \cdot 3^2} a_1$$

$$n=7 \rightarrow a_9 = -\frac{3}{3 \cdot 9 \cdot 8} a_7 = \frac{3}{9! \cdot 3^3} a_1$$

$$n=9 \rightarrow a_{11} = -\frac{5}{3 \cdot 11 \cdot 10} a_9 = -\frac{3 \cdot 5}{11! \cdot 3^4} a_1$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$n \geq 3 \quad a_{2n+1} = \frac{1 \cdot 3 \cdot 5 \cdots (2n-5)(-1)^n}{(2n+1)! \cdot 3^n} a_1$$

$$= \frac{(2n-5)!!(-1)^n}{(2n+1)! 3^n} a_1$$

$$y(x) = a_0 \left( 1 + \frac{2}{3}x^2 + \frac{1}{27}x^4 \right) + a_1 \left( x + \frac{1}{6}x^3 + \frac{1}{360}x^5 + \sum_{n=3}^{\infty} \frac{(2n-5)!!(-1)^n}{(2n+1)! 3^n} \right)$$


---


$$= a_0 \left( 1 + \frac{2}{3}x^2 + \frac{1}{27}x^4 \right) + a_1 \left( x + \frac{1}{6}x^3 + \frac{1}{360}x^5 - \frac{1}{45,360}x^7 + \dots \right)$$


---

### Exercise

Find a power series solution.  $5y'' - 2xy' + 10y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$5y'' - 2xy' + 10y = 0$$

$$5 \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - 2x \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n + 10 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} 5(n+2)(n+1) a_{n+2} x^n - \sum_{n=0}^{\infty} 2(n+1) a_{n+1} x^{n+1} + \sum_{n=0}^{\infty} 10a_n x^n = 0$$

$$\sum_{n=0}^{\infty} 5(n+2)(n+1) a_{n+2} x^n - \sum_{n=1}^{\infty} 2n a_n x^n + \sum_{n=0}^{\infty} 10a_n x^n = 0$$

$$10a_2 + \sum_{n=1}^{\infty} 5(n+2)(n+1) a_{n+2} x^n - \sum_{n=1}^{\infty} 2n a_n x^n + 10a_0 + \sum_{n=1}^{\infty} 10a_n x^n = 0$$

$$10a_2 + 10a_0 + \sum_{n=1}^{\infty} [5(n+2)(n+1) a_{n+2} - 2(n-5) a_n] x^n = 0$$

$$10a_2 + 10a_0 \rightarrow \underline{a_2 = -a_0}$$

$$5(n+2)(n+1)a_{n+2} - 2(n-5)a_n = 0$$

$$a_{n+2} = \frac{2(n-5)}{5(n+1)(n+2)} a_n$$

$$a_0$$

$$a_2 = -a_0$$

$$n=2 \rightarrow a_4 = -\frac{6}{60}a_2 = \frac{1}{10}a_0$$

$$n=4 \rightarrow a_6 = -\frac{2}{5 \cdot 5 \cdot 6}a_4 = -\frac{1}{750}a_0$$

$$n=6 \rightarrow a_8 = \frac{2}{5 \cdot 7 \cdot 8}a_6 = -\frac{2}{8! \cdot 5^2}a_0$$

$$n=8 \rightarrow a_{10} = \frac{2 \cdot 3}{5 \cdot 9 \cdot 10}a_8 = -\frac{2^2 \cdot 3}{10! \cdot 5^3}a_0$$

$$n=10 \rightarrow a_{12} = \frac{2 \cdot 5}{5 \cdot 11 \cdot 12}a_{10} = -\frac{2^3 \cdot 3 \cdot 5}{12! \cdot 5^4}a_0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$n \geq 4 \quad a_{2n} = -15 \cdot \frac{2^n (2n-7)!!}{5^n (2n)!} a_0$$

$$y(x) = a_0 \left( 1 - x^2 + \frac{1}{10}x^4 - \frac{1}{750}x^6 - \frac{1}{105,000}x^8 - \dots \right) + a_1 \left( x - \frac{4}{15}x^3 + \frac{4}{375}x^5 \right)$$

$$a_1$$

$$n=1 \rightarrow a_3 = -\frac{8}{30}a_1 = -\frac{4}{15}a_1$$

$$n=3 \rightarrow a_5 = -\frac{4}{100}a_3 = \frac{4}{375}a_1$$

$$n=5 \rightarrow a_7 = 0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

## Exercise

Find a power series solution.  $(x-1)y'' + y' = 0$

## Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(x-1)y'' + y' = 0$$

$$(x-1) \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n + \sum_{n=0}^{\infty} (n+1)a_{n+1}x^n = 0$$

$$x \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n - \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n + \sum_{n=0}^{\infty} (n+1)a_{n+1}x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^{n+1} - \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n + \sum_{n=0}^{\infty} (n+1)a_{n+1}x^n = 0$$

$$\sum_{\textcolor{red}{n}=1}^{\infty} n(n+1)a_{n+1}x^n - \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n + \sum_{n=0}^{\infty} (n+1)a_{n+1}x^n = 0$$

$$\sum_{n=1}^{\infty} n(n+1)a_{n+1}x^n - \textcolor{red}{2a_2} - \sum_{\textcolor{red}{n}=1}^{\infty} (n+2)(n+1)a_{n+2}x^n + \textcolor{blue}{a_1} + \sum_{\textcolor{red}{n}=1}^{\infty} (n+1)a_{n+1}x^n = 0$$

$$\textcolor{blue}{a_1} - \textcolor{blue}{2a_2} + \sum_{n=1}^{\infty} \left[ n(n+1)a_{n+1} - (n+2)(n+1)a_{n+2} + (n+1)a_{n+1} \right] x^n = 0$$

$$\textcolor{blue}{a_1} - \textcolor{blue}{2a_2} + \sum_{n=1}^{\infty} \left[ -(n+2)(n+1)a_{n+2} + (n+1)^2 a_{n+1} \right] x^n = 0$$

$$a_1 - 2a_2 = 0 \quad \rightarrow \quad \underline{\textcolor{blue}{a_2} = \frac{1}{2}a_1}$$

$$-(n+2)(n+1)a_{n+2} + (n+1)^2 a_{n+1}$$

$$\rightarrow \quad \underline{\textcolor{blue}{a_{n+2}} = \frac{n+1}{n+2}a_{n+1}} \quad n=1, 2, \dots$$

$$a_3 = \frac{2}{3}a_2 = \frac{1}{3}a_1$$

$$a_4 = \frac{3}{4}a_3 = \frac{1}{4}a_1$$

$$a_5 = \frac{4}{5}a_4 = \frac{1}{5}a_1$$

$$\vdots \vdots$$

$$\left\{ \begin{array}{l} y_1(x) = \textcolor{blue}{a_0} \\ y_2(x) = \left( x + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4 + \dots \right) \textcolor{blue}{a_1} \end{array} \right.$$

### Exercise

Find a power series solution.  $(x+2)y'' + xy' - y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(x+2)y'' + xy' - y = 0$$

$$(x+2) \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + x \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^{n+1} + \sum_{n=0}^{\infty} 2(n+2)(n+1) a_{n+2} x^n + \sum_{n=0}^{\infty} (n+1) a_{n+1} x^{n+1} - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=1}^{\infty} n(n+1) a_{n+1} x^n + \sum_{n=0}^{\infty} 2(n+2)(n+1) a_{n+2} x^n + \sum_{n=1}^{\infty} n a_n x^n - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=1}^{\infty} n(n+1) a_{n+1} x^n + 4a_2 + \sum_{n=1}^{\infty} 2(n+2)(n+1) a_{n+2} x^n + 4a_2 + \sum_{n=1}^{\infty} n a_n x^n - a_0 - \sum_{n=1}^{\infty} a_n x^n = 0$$

$$4a_2 - a_0 + \sum_{n=1}^{\infty} (n(n+1) a_{n+1} + 2(n+2)(n+1) a_{n+2} + (n-1) a_n) x^n = 0$$

$$4a_2 - a_0 = 0 \rightarrow \underline{a_2 = \frac{1}{4}a_0} \quad \underline{a_1}$$

$$n(n+1) a_{n+1} + 2(n+2)(n+1) a_{n+2} + (n-1) a_n = 0$$

$$\underline{a_{n+2} = -\frac{n-1}{2(n+2)(n+1)} a_n - \frac{n}{2(n+2)} a_{n+1}} \quad n = 1, 2, \dots$$

$$a_3 = -\frac{1}{6} a_2 = -\frac{1}{24} a_0$$

$$a_4 = -\frac{1}{24} a_2 - \frac{1}{4} a_3 = -\frac{1}{96} a_0 + \frac{1}{96} a_0 = 0$$

$$a_5 = -\frac{1}{20} a_3 - \frac{3}{10} a_4 = \frac{1}{480} a_0$$

$$a_6 = -\frac{3}{60}a_4 - \frac{1}{3}a_5 = -\frac{1}{1,440}a_0$$

$$\vdots \quad \vdots$$

$$\left\{ \begin{array}{l} y_1(x) = a_1 \\ y_2(x) = \left(1 + \frac{1}{4}x^2 - \frac{1}{24}x^3 + \frac{1}{480}x^5 - \dots\right)a_0 \end{array} \right.$$

### Exercise

Find a power series solution.  $y'' - (x+1)y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' - (x+1)y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - x \sum_{n=0}^{\infty} a_n x^n - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=0}^{\infty} a_n x^{n+1} - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} [(n+2)(n+1) a_{n+2} - a_n] x^n - \sum_{n=1}^{\infty} a_{n-1} x^n = 0$$

$$2a_2 - a_0 + \sum_{n=1}^{\infty} [(n+2)(n+1) a_{n+2} - a_n] x^n - \sum_{n=1}^{\infty} a_{n-1} x^n = 0$$

$$2a_2 - a_0 + \sum_{n=1}^{\infty} [(n+2)(n+1) a_{n+2} - a_n - a_{n-1}] x^n = 0$$

$$2a_2 - a_0 = 0 \rightarrow \underline{a_2 = \frac{1}{2}a_0}$$

$$(n+2)(n+1) a_{n+2} - a_n - a_{n-1} = 0$$

$$a_{n+2} = \frac{a_n + a_{n-1}}{(n+1)(n+2)}$$

$$a_0 \quad a_1$$

$$a_2 = \frac{1}{2}a_0$$

$$n=1 \rightarrow a_3 = \frac{1}{6}(a_1 + a_0)$$

$$n=2 \rightarrow a_4 = \frac{1}{12}(a_2 + a_1) = \frac{1}{12}\left(\frac{1}{2}a_0 + a_1\right)$$

$$n=3 \rightarrow a_5 = \frac{1}{20}(a_3 + a_2) = \frac{1}{20}\left(\frac{2}{3}a_0 + \frac{1}{6}a_1\right)$$

$$n=4 \rightarrow a_6 = \frac{1}{30}(a_4 + a_3) = \frac{1}{30}\left(\frac{1}{2}a_0 + a_1 + \frac{1}{6}a_0 + \frac{1}{6}a_1\right) = \frac{1}{30}\left(\frac{2}{3}a_0 + \frac{7}{6}a_1\right)$$

$$a_0 \neq 0 \quad a_1 = 0$$

$$a_2 = \frac{1}{2}a_0$$

$$a_3 = \frac{1}{6}a_0$$

$$a_4 = \frac{1}{24}a_0$$

$$a_5 = \frac{1}{30}a_0$$

$$a_6 = \frac{1}{45}a_0$$

$$a_0 = 0 \quad a_1 \neq 0$$

$$a_2 = 0$$

$$a_3 = \frac{1}{6}a_1$$

$$a_4 = \frac{1}{12}a_1$$

$$a_5 = \frac{1}{120}a_1$$

$$a_6 = \frac{7}{180}a_1$$

$$\left\{ \begin{array}{l} y_1(x) = \left(1 + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \frac{1}{30}x^5 + \frac{1}{45}x^6 + \dots\right)a_0 \\ y_2(x) = \left(x + \frac{1}{6}x^3 + \frac{1}{12}x^4 + \frac{1}{120}x^5 + \frac{7}{180}x^6 + \dots\right)a_1 \end{array} \right.$$

### Exercise

Find a power series solution.  $y'' - (x+1)y' - y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' - (x+1)y' - y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n - (x+1)\sum_{n=0}^{\infty} (n+1)a_{n+1}x^n - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n - \sum_{n=0}^{\infty} (n+1)a_{n+1}x^{n+1} - \sum_{n=0}^{\infty} (n+1)a_{n+1}x^n - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n - \sum_{n=1}^{\infty} na_n x^n - \sum_{n=0}^{\infty} (n+1)a_{n+1}x^n - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$2a_2 + \sum_{n=1}^{\infty} (n+2)(n+1)a_{n+2}x^n - \sum_{n=1}^{\infty} na_n x^n - a_1 - \sum_{n=1}^{\infty} (n+1)a_{n+1}x^n - a_0 - \sum_{n=1}^{\infty} a_n x^n = 0$$

$$2a_2 - a_1 - a_0 + \sum_{n=1}^{\infty} [(n+2)(n+1)a_{n+2} - na_n - (n+1)a_{n+1} - a_n]x^n = 0$$

$$2a_2 - a_1 - a_0 = 0 \rightarrow a_2 = \frac{1}{2}a_0 + \frac{1}{2}a_1$$

$$(n+2)(n+1)a_{n+2} - (n+1)a_{n+1} - (n+1)a_n = 0$$

$$a_{n+2} = \frac{1}{n+2}a_{n+1} + \frac{1}{n+2}a_n$$

$$a_3 = \frac{1}{3}a_2 + \frac{1}{3}a_1 = \frac{1}{6}a_0 + \frac{1}{2}a_1$$

$$a_4 = \frac{1}{4}a_3 + \frac{1}{4}a_2 = \frac{1}{24}a_0 + \frac{1}{8}a_1 + \frac{1}{8}a_0 + \frac{1}{8}a_1 = \frac{1}{6}a_0 + \frac{1}{4}a_1$$

$$a_5 = \frac{1}{5}a_4 + \frac{1}{5}a_3 = \frac{1}{30}a_0 + \frac{1}{20}a_1 + \frac{1}{30}a_0 + \frac{1}{10}a_1 = \frac{1}{15}a_0 + \frac{3}{20}a_1$$

$$\vdots$$

$$\begin{cases} y_1(x) = \left(1 + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{6}x^4 + \dots\right)a_0 \\ y_2(x) = \left(x + \frac{1}{2}x^2 + \frac{1}{2}x^3 + \frac{1}{4}x^4 + \dots\right)a_1 \end{cases}$$

### Exercise

Find a power series solution.  $(x^2 + 1)y'' - 6y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$



$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(x^2 + 1)y'' - 6y = 0$$

$$(x^2 + 1) \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - 6 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^{n+2} + \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - 6 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^n + \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - 6 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^n + 2a_2 + 6a_3 x + \sum_{n=2}^{\infty} (n+2)(n+1) a_{n+2} x^n - 6a_0 - 6a_1 x - 6 \sum_{n=2}^{\infty} a_n x^n = 0$$

$$2a_2 - 6a_0 + (6a_3 - 6a_1)x + \sum_{n=2}^{\infty} \left[ (n^2 - n - 6) a_n + (n+2)(n+1) a_{n+2} \right] x^n = 0$$

$$2a_2 - 6a_0 + (6a_3 - 6a_1)x = 0 \rightarrow \begin{cases} a_2 = 3a_0 \\ a_3 = a_1 \end{cases}$$

$$(n+2)(n-3) a_n + (n+2)(n+1) a_{n+2} = 0$$

$$\Rightarrow a_{n+2} = -\frac{n-3}{n+1} a_n \quad n = 2, 3, \dots$$

$$a_4 = \frac{1}{3} a_2 = a_0 \quad a_5 = 0$$

$$a_6 = -\frac{1}{5} a_4 = -\frac{1}{5} a_0 \quad a_7 = -\frac{1}{3} a_5 = 0$$

$$a_8 = -\frac{3}{7} a_6 = \frac{3}{35} a_0$$

$$\vdots$$

$$\left\{ \begin{array}{l} y_1(x) = \left( 1 + 3x^2 + x^4 - \frac{1}{5}x^6 + \dots \right) a_0 \\ y_2(x) = \left( x + x^3 \right) a_1 \end{array} \right.$$

### Exercise

Find a power series solution.  $(x^2 + 2)y'' + 3xy' - y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(x^2 + 2)y'' + 3xy' - y = 0$$

$$(x^2 + 2) \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + 3x \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^{n+2} + \sum_{n=0}^{\infty} 2(n+2)(n+1) a_{n+2} x^n + \sum_{n=0}^{\infty} 3(n+1) a_{n+1} x^{n+1} - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^n + \sum_{n=0}^{\infty} [2(n+2)(n+1) a_{n+2} - a_n] x^n + \sum_{n=1}^{\infty} 3n a_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^n + 4a_2 - a_0 + (12a_3 - a_1)x + \sum_{n=2}^{\infty} [2(n+2)(n+1) a_{n+2} - a_n] x^n$$

$$+ 3a_1 x + \sum_{n=2}^{\infty} 3n a_n x^n = 0$$

$$4a_2 - a_0 + (12a_3 + 2a_1)x + \sum_{n=2}^{\infty} [n(n-1) a_n + 2(n+2)(n+1) a_{n+2} + (3n-1) a_n] x^n = 0$$

$$4a_2 - a_0 + (12a_3 + 2a_1)x \Rightarrow \underline{a_2 = \frac{1}{4}a_0} \quad \underline{a_3 = -\frac{1}{6}a_1}$$

$$2(n+2)(n+1) a_{n+2} + (n^2 + 2n - 1) a_n = 0$$

$$\rightarrow a_{n+2} = -\frac{n^2 + 2n - 1}{2(n+2)(n+1)} a_n \quad n = 2, 3, \dots$$

$$a_4 = -\frac{7}{24} a_2 = -\frac{7}{96} a_0$$

$$a_5 = -\frac{7}{20} a_3 = \frac{7}{120} a_1$$

$$a_6 = -\frac{23}{60}a_4 = \frac{161}{5760}a_0$$

$$a_7 = -\frac{17}{42}a_5 = -\frac{17}{720}a_1$$

$\vdots$

$$\left\{ \begin{array}{l} y_1(x) = \left(1 + \frac{1}{4}x^2 - \frac{7}{96}x^4 + \frac{161}{5760}x^6 - \dots\right)a_0 \\ y_2(x) = \left(1 - \frac{1}{6}x^3 + \frac{7}{120}x^5 - \frac{17}{720}x^7 + \dots\right)a_1 \end{array} \right|$$

### Exercise

Find a power series solution.  $(x^2 - 1)y'' + xy' - y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(x^2 - 1)y'' + xy' - y = 0$$

$$(x^2 - 1) \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + x \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^{n+2} - \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + \sum_{n=0}^{\infty} (n+1) a_{n+1} x^{n+1} - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^n - \sum_{n=0}^{\infty} [(n+2)(n+1) a_{n+2} + a_n] x^n + \sum_{n=1}^{\infty} n a_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^n - (2a_2 + a_0) - (6a_3 + a_1)x - \sum_{n=2}^{\infty} [(n+2)(n+1) a_{n+2} + a_n] x^n$$

$$+ a_1 x + \sum_{n=2}^{\infty} n a_n x^n = 0$$

$$-2a_2 - a_0 - 6a_3 x + \sum_{n=2}^{\infty} [n(n-1) a_n - (n+2)(n+1) a_{n+2} + (n-1) a_n] x^n = 0$$

$$-2a_2 - a_0 - 6a_3x = 0 \quad \rightarrow \underline{a_2 = -\frac{1}{2}a_0} \quad \underline{a_3 = 0}$$

$$-(n+2)(n+1)a_{n+2} + (n+1)(n-1)a_n = 0$$

$$\rightarrow a_{n+2} = \frac{n-1}{n+2}a_n \quad n = 2, 3, \dots$$

$$a_4 = \frac{1}{4}a_2 = -\frac{1}{8}a_0 \quad a_5 = -\frac{2}{5}a_3 = 0$$

$$a_6 = \frac{1}{2}a_4 = -\frac{1}{16}a_0 \quad a_7 = \frac{4}{7}a_5 = 0$$

$$\vdots \vdots$$

$$\left\{ \begin{array}{l} y_1(x) = \left(1 - \frac{1}{2}x^2 - \frac{1}{8}x^4 - \frac{1}{16}x^6 - \dots\right)a_0 \\ y_2(x) = a_1 \end{array} \right.$$

$$\left\{ \begin{array}{l} y_1(x) = 1 - \frac{1}{2}x^2 - \frac{1}{8}x^4 - \frac{1}{16}x^6 - \dots \\ y_2(x) = x \end{array} \right.$$

### Exercise

Find a power series solution.  $(x^2 + 1)y'' + xy' - y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(x^2 + 1)y'' + xy' - y = 0$$

$$(x^2 + 1) \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + x \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$x^2 \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + x \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^{n+2} + \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n + \sum_{n=0}^{\infty} (n+1)a_{n+1}x^{n+1} - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1)a_n x^n + \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n + \sum_{n=1}^{\infty} na_n x^n - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1)a_n x^n + \sum_{n=1}^{\infty} na_n x^n + \sum_{n=0}^{\infty} [(n+2)(n+1)a_{n+2} - a_n]x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1)a_n x^n + a_1 x + \sum_{n=2}^{\infty} na_n x^n + (2a_2 - a_0) + (6a_3 - a_1)x + \sum_{n=2}^{\infty} [(n+2)(n+1)a_{n+2} - a_n]x^n = 0$$

$$2a_2 - a_0 + 6a_3 x + \sum_{n=2}^{\infty} [(n^2 - n + n - 1)a_n + (n+2)(n+1)a_{n+2}]x^n = 0$$

$$\begin{cases} 2a_2 - a_0 = 0 & \rightarrow a_2 = \frac{1}{2}a_0 \\ 6a_3 x = 0 & \rightarrow a_3 = 0 \end{cases}$$

$$(n-1)(n+1)a_n + (n+2)(n+1)a_{n+2} = 0$$

$$a_{n+2} = -\frac{n-1}{n+2}a_n$$

$a_0$	$a_1$
$n=0 \rightarrow a_2 = \frac{1}{2}a_0$	$n=1 \rightarrow a_3 = 0$
$n=2 \rightarrow a_4 = -\frac{1}{4}a_2 = -\frac{1}{2^3}a_0$	$n=3 \rightarrow a_5 = -\frac{2}{5}a_3 = 0$
$n=4 \rightarrow a_6 = -\frac{3}{6}a_4 = \frac{1 \cdot 3}{2^3 \cdot 3!}a_0$	$n=5 \rightarrow a_7 = 0$
$n=6 \rightarrow a_8 = -\frac{5}{8}a_6 = -\frac{1 \cdot 3 \cdot 5}{2^4 \cdot 4!}a_0$	$\vdots \quad \vdots \quad \vdots \quad \vdots$
$\vdots \quad \vdots \quad \vdots \quad \vdots$	

$$a_{2n} = (-1)^{n-1} \frac{1 \cdot 3 \cdot 5 \cdots (2n-3)}{2^n \cdot n!} a_0 \quad (n \geq 2)$$

$$y(x) = a_0 \left( 1 + \frac{1}{2}x^2 - \frac{1}{2^2 \cdot 2!}x^4 + \frac{1 \cdot 3}{2^3 \cdot 3!}x^6 - \frac{1 \cdot 3 \cdot 5}{2^4 \cdot 4!}x^8 + \cdots \right) + a_1 x$$

$$= a_0 \left( 1 + \frac{1}{2}x^2 + \sum_{n=2}^{\infty} (-1)^{n-1} \frac{(2n-3)!!}{2^n \cdot n!} x^{2n} \right) + a_1 x$$

### Exercise

Find a power series solution.  $(x^2 + 1)y'' - xy' + y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(x^2 + 1)y'' - xy' + y = 0$$

$$(x^2 + 1) \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - x \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$x^2 \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - x \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^{n+2} + \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=0}^{\infty} (n+1) a_{n+1} x^{n+1} + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^n + \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=1}^{\infty} n a_n x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^n - \sum_{n=1}^{\infty} n a_n x^n + \sum_{n=0}^{\infty} [(n+2)(n+1) a_{n+2} + a_n] x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^n - a_1 x - \sum_{n=2}^{\infty} n a_n x^n + (2a_2 + a_0) + (6a_3 + a_1)x + \sum_{n=2}^{\infty} [(n+2)(n+1) a_{n+2} + a_n] x^n = 0$$

$$2a_2 + a_0 + 6a_3 x + \sum_{n=2}^{\infty} [(n^2 - 2n + 1) a_n + (n+2)(n+1) a_{n+2}] x^n = 0$$

$$\begin{cases} 2a_2 + a_0 = 0 & \rightarrow a_2 = -\frac{1}{2}a_0 \\ 6a_3 = 0 & \rightarrow a_3 = 0 \end{cases}$$

$$(n-1)^2 a_n + (n+1)(n+2) a_{n+2} = 0$$

$$\boxed{a_{n+2} = -\frac{(n-1)^2}{(n+1)(n+2)}a_n}$$

$a_0$	$a_1$
$n=0 \rightarrow a_2 = -\frac{1}{2}a_0$	$n=1 \rightarrow a_3 = 0$
$n=2 \rightarrow a_4 = -\frac{1}{12}a_2 = \frac{1}{4!}a_0$	$n=3 \rightarrow a_5 = -\frac{2}{5}a_3 = 0$
$n=4 \rightarrow a_6 = -\frac{3^2}{5 \cdot 6}a_4 = -\frac{3^2}{6!}a_0$	$n=5 \rightarrow a_7 = 0$
$n=6 \rightarrow a_8 = -\frac{5^2}{7 \cdot 8}a_6 = \frac{1 \cdot 3^2 \cdot 5^2}{8!}a_0$	$\vdots \quad \vdots \quad \vdots \quad \vdots$
$\vdots \quad \vdots \quad \vdots \quad \vdots$	

$$a_{2n} = (-1)^{n-1} \frac{1 \cdot 3^2 \cdot 5^2 \cdots (2n-3)^2}{(2n)!} a_0 \quad (n \geq 3)$$

$$\boxed{y(x) = a_0 \left( 1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{9}{6!}x^6 + \frac{1 \cdot 3^2 \cdot 5^2}{8!}x^8 - \cdots \right) + a_1 x}$$

$$\boxed{= a_0 \left( 1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 + \sum_{n=3}^{\infty} (-1)^n \frac{(2n-3)^2!!}{(2n)!} x^{2n} \right) + a_1 x}$$

### Exercise

Find a power series solution.  $(1-x^2)y'' - 6xy' - 4y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(1-x^2)y'' - 6xy' - 4y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - x^2 \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} - 6x \sum_{n=1}^{\infty} n a_n x^{n-1} - 4 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n - \sum_{n=2}^{\infty} n(n-1)a_n x^n - \sum_{n=1}^{\infty} 6na_n x^n - \sum_{n=0}^{\infty} 4a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n - \sum_{n=0}^{\infty} n(n-1)a_n x^n - \sum_{n=0}^{\infty} 6na_n x^n - \sum_{n=0}^{\infty} 4a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ (n+2)(n+1)a_{n+2} - (n(n-1) + 6n + 4)a_n \right] x^n = 0$$

$$(n+2)(n+1)a_{n+2} - (n^2 + 5n + 4)a_n = 0$$

$$(n+2)(n+1)a_{n+2} = (n+4)(n+1)a_n$$

$$\underline{a_{n+2} = \frac{n+4}{n+2}a_n}$$

$$a_0$$

$$n=2 \rightarrow a_2 = 2a_0$$

$$n=4 \rightarrow a_4 = \frac{6}{4}a_2 = 3a_0$$

$$n=6 \rightarrow a_6 = \frac{8}{6}a_3 = 4a_0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$a_{2k} = (k+1)a_0$$

$$a_1$$

$$n=3 \rightarrow a_3 = \frac{5}{3}a_1$$

$$n=5 \rightarrow a_5 = \frac{7}{5}a_3 = \frac{7}{3}a_1$$

$$n=7 \rightarrow a_7 = \frac{11}{7}a_5 = \frac{11}{3}a_1$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$a_{2k+1} = \frac{2k+3}{3}a_1$$

$$y(x) = a_0 \left( 1 + 2x^2 + 3x^4 + 4x^6 + \dots \right) + a_1 \left( x + \frac{5}{3}x^3 + \frac{7}{3}x^5 + \frac{11}{3}x^7 + \dots \right)$$

$$\underline{= \frac{a_0}{(1-x^2)^2} + \frac{3x-x^3}{3(1-x^2)^2}a_1}$$

## Exercise

Find a power series solution.  $y'' + (x-1)^2 y' - 4(x-1)y = 0$

### Solution

$$\text{Let } z = x-1 \rightarrow dz = dx$$

$$y(x) = \sum_{n=0}^{\infty} a_n z^n$$

$$y'(x) = \sum_{n=1}^{\infty} na_n z^{n-1} = \sum_{n=0}^{\infty} (n+1)a_{n+1} z^n$$



$$y''(x) = \sum_{n=2}^{\infty} n(n-1)a_n z^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} z^n$$

$$y'' + z^2 y' - 4zy = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} z^n + z^2 \sum_{n=1}^{\infty} n a_n z^{n-1} - 4z \sum_{n=0}^{\infty} a_n z^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} z^n + \sum_{n=1}^{\infty} n a_n z^{n+1} - \sum_{n=0}^{\infty} 4a_n z^{n+1} = 0$$

$$2a_2 + \sum_{n=1}^{\infty} (n+2)(n+1)a_{n+2} z^n + \sum_{n=1}^{\infty} (n-4)a_n z^{n+1} = 0$$

$$2a_2 + \sum_{n=0}^{\infty} (n+2)(n+3)a_{n+3} z^{n+1} + \sum_{n=1}^{\infty} (n-4)a_n z^{n+1} = 0$$

$$2a_2 + \sum_{n=0}^{\infty} [(n+2)(n+3)a_{n+3} + (n-4)a_n] z^{n+1} = 0$$

$$\underline{a_2 = 0}$$

$$(n+2)(n+3)a_{n+3} + (n-4)a_n = 0$$

$$\underline{a_{n+3} = -\frac{n-4}{(n+2)(n+3)}a_n}$$

$$a_0$$

$$a_1$$

$$a_2 = 0$$

$$n=0 \rightarrow a_3 = \frac{4}{2 \cdot 3} a_0$$

$$n=1 \rightarrow a_4 = \frac{3}{3 \cdot 4} a_1$$

$$n=2 \rightarrow a_5 = \frac{2}{20} a_2 = 0$$

$$n=3 \rightarrow a_6 = \frac{1}{5 \cdot 6} a_3 = \frac{4}{2 \cdot 3 \cdot 5 \cdot 6} a_0$$

$$n=4 \rightarrow a_7 = 0$$

$$n=5 \rightarrow a_8 = 0$$

$$n=6 \rightarrow a_9 = -\frac{2}{8 \cdot 9} a_6 = -\frac{8}{2 \cdot 3 \cdot 5 \cdot 6 \cdot 8 \cdot 9} a_0$$

$$n=7 \rightarrow a_{10} = 0$$

$$n=8 \rightarrow a_9 = 0$$

$$\vdots$$

$$\vdots$$

$$\vdots$$

$$\vdots$$

$$\vdots$$

$$\vdots$$

$$\vdots$$

$$\vdots$$

$$y(x) = a_0 \left( 1 + \frac{4}{2 \cdot 3} z^3 + \frac{4}{2 \cdot 3 \cdot 5 \cdot 6} z^6 - \frac{8}{2 \cdot 3 \cdot 5 \cdot 6 \cdot 8 \cdot 9} z^9 + \dots \right) + a_1 \left( z + \frac{1}{4} z^4 \right)$$

$$\underline{= a_0 \left( 1 + \frac{2}{3} (x-1)^3 + \frac{1}{45} (x-1)^6 - \frac{1}{1,620} (x-1)^9 + \dots \right) + a_1 \left( x-1 + \frac{1}{4} (x-1)^4 \right)}$$

## Exercise

Find a power series solution.  $(2 - x^2)y'' - xy' + 16y = 0$

## Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(2 - x^2)y'' - xy' + 16y = 0$$

$$2 \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - x^2 \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} - x \sum_{n=1}^{\infty} n a_n x^{n-1} + 16 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} 2(n+2)(n+1) a_{n+2} x^n - \sum_{n=2}^{\infty} n(n-1) a_n x^n - \sum_{n=1}^{\infty} n a_n x^n + \sum_{n=0}^{\infty} 16 a_n x^n = 0$$

$$\sum_{n=0}^{\infty} 2(n+2)(n+1) a_{n+2} x^n - \sum_{n=0}^{\infty} n(n-1) a_n x^n - \sum_{n=0}^{\infty} n a_n x^n + \sum_{n=0}^{\infty} 16 a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ 2(n+2)(n+1) a_{n+2} - (n^2 - n + n - 16) a_n \right] x^n = 0$$

$$2(n+1)(n+2) a_{n+2} - (n^2 - 16) a_n = 0$$

$$2(n+1)(n+2) a_{n+2} = (n+4)(n-4) a_n$$

$$a_{n+2} = \frac{(n+4)(n-4)}{2(n+1)(n+2)} a_n$$

$$a_0$$

$$n=0 \rightarrow a_2 = -4a_0$$

$$n=2 \rightarrow a_4 = -\frac{1}{2}a_2 = 2a_0$$

$$n=4 \rightarrow a_6 = 0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$a_1$$

$$n=1 \rightarrow a_3 = -\frac{5}{4}a_1$$

$$n=3 \rightarrow a_5 = -\frac{7}{40}a_3 = \frac{7}{320}a_1$$

$$n=5 \rightarrow a_7 = \frac{9}{70}a_5 = \frac{9}{320}a_1$$

$$n=7 \rightarrow a_9 = \frac{33}{144}a_7 = \frac{33}{5,120}a_1$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$a_{2n+1} = \frac{(2n-5)!! (2n+3)!!}{2^n (2n+1)!} a_1$$

$$y(x) = a_0 \left( 1 - 4x^2 + 2x^4 \right) + a_1 \left( x - \frac{5}{4}x^3 + \frac{7}{32}x^5 + \frac{9}{320}x^7 + \cdots \right)$$

$$= a_0 \left( 1 - 4x^2 + 2x^4 \right) + a_1 \sum_{n=0}^{\infty} \frac{(2n-5)!! (2n+3)!!}{2^n (2n+1)!} x^{2n+1}$$

### Exercise

Find a power series solution.  $(x^2 + 1)y'' - y' + y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(x^2 + 1)y'' - y' + y = 0$$

$$x^2 \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} + \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=1}^{\infty} n a_n x^{n-1} + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^n + \sum_{n=0}^{\infty} (n+1)(n+2) a_{n+2} x^n - \sum_{n=1}^{\infty} n a_n x^{n-1} + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} n(n-1) a_n x^n + \sum_{n=0}^{\infty} (n+1)(n+2) a_{n+2} x^n - \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ (n+1)(n+2) a_{n+2} - (n+1) a_{n+1} + (n^2 - n + 1) a_n \right] x^n = 0$$

$$(n+1)(n+2) a_{n+2} - (n+1) a_{n+1} + (n^2 - n + 1) a_n = 0$$

$$a_{n+2} = \frac{(n+1)a_{n+1} - (n^2 - n + 1)a_n}{(n+1)(n+2)}$$

$$n=0 \rightarrow a_2 = \frac{1}{2}(a_1 - a_0)$$

$$n=1 \rightarrow a_3 = \frac{1}{6}(2a_2 - a_1) = \frac{1}{6}(a_1 - a_0 - a_1) = -\frac{1}{6}a_0$$

$$n=2 \rightarrow a_4 = \frac{1}{12}(3a_3 - 3a_2) = \frac{1}{4}\left(-\frac{1}{6}a_0 - \frac{1}{2}a_1 + \frac{1}{2}a_0\right) = \frac{1}{12}a_0 - \frac{1}{8}a_1$$

$$n=3 \rightarrow a_5 = \frac{1}{20}(4a_4 - 7a_3) = \frac{1}{20}\left(\frac{1}{3}a_0 - \frac{1}{2}a_1 + \frac{7}{6}a_0\right) = \frac{3}{40}a_0 - \frac{1}{40}a_1$$

$$n=4 \rightarrow a_6 = \frac{1}{30}(5a_5 - 13a_4) = \frac{1}{30}\left(\frac{3}{8}a_0 - \frac{1}{8}a_1 - \frac{13}{12}a_0 + \frac{13}{8}a_1\right) = -\frac{17}{720}a_0 + \frac{1}{20}a_1$$

$$y(x) = a_0 + a_1x + \left(\frac{1}{2}a_0 - \frac{1}{2}a_1\right)x^2 - \frac{1}{6}a_0x^3 + \left(\frac{1}{12}a_0 - \frac{1}{8}a_1\right)x^4 + \left(\frac{3}{40}a_0 - \frac{1}{40}a_1\right)x^5 \\ + \left(-\frac{17}{720}a_0 + \frac{1}{20}a_1\right)x^6 + \dots$$

$$y(x) = a_0 \left(1 + \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{12}x^4 + \frac{3}{40}x^5 - \frac{17}{720}x^6 + \dots\right) \\ + a_1 \left(x - \frac{1}{2}x^2 - \frac{1}{8}x^4 - \frac{1}{40}x^5 + \frac{1}{20}x^6 + \dots\right)$$

### Exercise

Find a power series solution.  $(x^2 + 1)y'' + 6xy' + 4y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(x^2 + 1)y'' + 6xy' + 4y = 0$$

$$x^2 \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} + \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + 6x \sum_{n=1}^{\infty} n a_n x^{n-1} + 4 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1)a_n x^n + \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n + \sum_{n=1}^{\infty} 6na_n x^n + \sum_{n=0}^{\infty} 4a_n x^n = 0$$

$$\sum_{n=0}^{\infty} n(n-1)a_n x^n + \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n + \sum_{n=0}^{\infty} 6na_n x^n + \sum_{n=0}^{\infty} 4a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ (n+2)(n+1)a_{n+2} + (n^2 + 5n + 4)a_n \right] x^n = 0$$

$$(n+1)(n+2)a_{n+2} + (n+1)(n+4)a_n = 0$$

$$\underline{a_{n+2} = -\frac{n+4}{n+2}a_n}$$

$$\begin{array}{cccc} a_0 & & & \\ n=0 \rightarrow a_2 = -2a_0 & & & \\ n=2 \rightarrow a_4 = -\frac{3}{2}a_2 = 3a_0 & & & \\ n=4 \rightarrow a_6 = -\frac{8}{6}a_4 = -4a_0 & & & \\ \vdots & \vdots & \vdots & \vdots \end{array}$$

$$\begin{array}{cccc} a_1 & & & \\ n=1 \rightarrow a_3 = -\frac{5}{3}a_1 & & & \\ n=3 \rightarrow a_5 = -\frac{7}{5}a_3 = \frac{7}{3}a_1 & & & \\ n=5 \rightarrow a_7 = -\frac{9}{7}a_5 = -\frac{9}{3}a_1 & & & \\ \vdots & \vdots & \vdots & \vdots \end{array}$$

$$a_{2n} = (-1)^n (n+1)a_0$$

$$a_{2n+1} = (-1)^n (2n+3)a_1$$

$$\underline{y(x) = a_0 \sum_{n=0}^{\infty} (-1)^n (n+1)x^{2n} + \frac{1}{3}a_1 \sum_{n=0}^{\infty} (-1)^n (2n+3)x^{2n+1}}$$

$$\underline{y(x) = a_0 \left( 1 - 2x^2 + 3x^4 - 4x^6 + \dots \right) + \frac{1}{3}a_1 \left( x - \frac{5}{3}x^3 + \frac{7}{3}x^5 - \frac{9}{3}x^7 + \dots \right)}$$

### Exercise

Find a power series solution.  $(x^2 - 1)y'' - 6xy' + 12y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} na_n x^{n-1} = \sum_{n=0}^{\infty} (n+1)a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1)a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n$$

$$(x^2 - 1)y'' - 6xy' + 12y = 0$$

$$x^2 \sum_{n=2}^{\infty} n(n-1)a_n x^{n-2} - \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n - 6x \sum_{n=1}^{\infty} na_n x^{n-1} + 12 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1)a_n x^n - \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n - \sum_{n=1}^{\infty} 6na_n x^n + \sum_{n=0}^{\infty} 12a_n x^n = 0$$

$$\sum_{n=0}^{\infty} n(n-1)a_n x^n - \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n - \sum_{n=0}^{\infty} 6na_n x^n + \sum_{n=0}^{\infty} 12a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ -(n+2)(n+1)a_{n+2} + (n^2 - 7n + 12)a_n \right] x^n = 0$$

$$-(n+1)(n+2)a_{n+2} + (n-3)(n-4)a_n = 0$$

$$a_{n+2} = \frac{(n-3)(n-4)}{(n+1)(n+2)} a_n$$

$$a_0$$

$$n=0 \rightarrow a_2 = 6a_0$$

$$n=2 \rightarrow a_4 = \frac{1}{6}a_2 = a_0$$

$$n=4 \rightarrow a_6 = 0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$y(x) = a_0(1 + 6x^2 + x^4) + a_1(x + x^3)$$

$$a_1$$

$$n=1 \rightarrow a_3 = a_1$$

$$n=3 \rightarrow a_5 = 0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

### Exercise

Find a power series solution.  $(x^2 - 1)y'' + 8xy' + 12y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} na_n x^{n-1} = \sum_{n=0}^{\infty} (n+1)a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1)a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n$$

$$(x^2 - 1)y'' + 8xy' + 12y = 0$$

$$x^2 \sum_{n=2}^{\infty} n(n-1)a_n x^{n-2} - \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n + 8x \sum_{n=1}^{\infty} na_n x^{n-1} + 12 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1)a_n x^n - \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n + \sum_{n=1}^{\infty} 8na_n x^n + \sum_{n=0}^{\infty} 12a_n x^n = 0$$

$$\sum_{n=0}^{\infty} n(n-1)a_n x^n - \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n + \sum_{n=0}^{\infty} 8na_n x^n + \sum_{n=0}^{\infty} 12a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ -(n+2)(n+1)a_{n+2} + (n^2 + 7n + 12)a_n \right] x^n = 0$$

$$-(n+1)(n+2)a_{n+2} + (n+3)(n+4)a_n = 0$$

$$a_{n+2} = \frac{(n+3)(n+4)}{(n+1)(n+2)} a_n$$

$$\begin{array}{cccc} & & a_0 & \\ n=0 \rightarrow & a_2 = 6a_0 & & \\ n=2 \rightarrow & a_4 = \frac{5}{2}a_2 = 15a_0 & & \\ n=4 \rightarrow & a_6 = \frac{5}{2}a_4 = 28a_0 & & \\ \vdots & \vdots & \vdots & \vdots \\ & a_{2n} = (n+1)(2n+1)a_0 & & \end{array}$$

$$\begin{array}{cccc} & & a_1 & \\ n=1 \rightarrow & a_3 = \frac{10}{3}a_1 & & \\ n=3 \rightarrow & a_5 = \frac{42}{20}a_3 = \frac{21}{3}a_1 & & \\ n=5 \rightarrow & a_7 = \frac{12}{7}a_5 = \frac{36}{3}a_1 & & \\ \vdots & \vdots & \vdots & \vdots \\ & a_{2n+1} = \frac{1}{3}(n+1)(2n+3)a_1 & & \end{array}$$

$$y(x) = a_0 \left( 1 + 6x^2 + 15x^4 + 28x^6 + \dots \right) + a_1 \left( x + \frac{10}{3}x^3 + 7x^5 + 12x^7 + \dots \right)$$

$$= a_0 \sum_{n=0}^{\infty} (n+1)(2n+1)x^{2n} + \frac{1}{3}a_1 \sum_{n=0}^{\infty} (n+1)(2n+3)x^{2n+1}$$

### Exercise

Find a power series solution.  $(x^2 - 1)y'' + 4xy' + 2y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(x^2 - 1)y'' + 4xy' + 2y = 0$$

$$x^2 \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} - \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + 4x \sum_{n=1}^{\infty} n a_n x^{n-1} + 2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^n - \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + \sum_{n=1}^{\infty} 4n a_n x^n + \sum_{n=0}^{\infty} 2a_n x^n = 0$$

$$\sum_{n=0}^{\infty} n(n-1) a_n x^n - \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + \sum_{n=0}^{\infty} 4n a_n x^n + \sum_{n=0}^{\infty} 2a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ -(n+2)(n+1) a_{n+2} + (n^2 + 3n + 2) a_n \right] x^n = 0$$

$$-(n+1)(n+2) a_{n+2} + (n+1)(n+2) a_n = 0$$

$$\underline{a_{n+2} = a_n}$$

$$a_0$$

$$a_1$$

$$n=0 \rightarrow a_2 = a_0$$

$$n=1 \rightarrow a_3 = a_1$$

$$n=2 \rightarrow a_4 = a_2 = a_0$$

$$n=3 \rightarrow a_5 = a_1$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$y(x) = a_0 (1 + x^2 + x^4 + x^6 + \dots) + a_1 (x + x^3 + x^5 + x^7 + \dots)$$

$$= a_0 \sum_{n=0}^{\infty} x^{2n} + a_1 \sum_{n=0}^{\infty} x^{2n+1}$$



$$= \frac{a_0 + a_1 x}{1 - x^2} \Bigg|$$

### Exercise

Find a power series solution.  $(x^2 + 1)y'' - 4xy' + 6y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(x^2 + 1)y'' - 4xy' + 6y = 0$$

$$x^2 \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} + \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - 4x \sum_{n=1}^{\infty} n a_n x^{n-1} + 6 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^n + \sum_{n=0}^{\infty} (n+1)(n+2) a_{n+2} x^n - \sum_{n=1}^{\infty} 4n a_n x^n + \sum_{n=0}^{\infty} 6a_n x^n = 0$$

$$\sum_{n=0}^{\infty} n(n-1) a_n x^n + \sum_{n=0}^{\infty} (n+1)(n+2) a_{n+2} x^n - \sum_{n=0}^{\infty} 4n a_n x^n + \sum_{n=0}^{\infty} 6a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ (n+1)(n+2) a_{n+2} + (n^2 - 5n + 6) a_n \right] x^n = 0$$

$$(n+1)(n+2) a_{n+2} + (n-2)(n-3) a_n = 0$$

$$a_{n+2} = -\frac{(n-2)(n-3)}{(n+1)(n+2)} a_n \Bigg|$$

$$a_0$$

$$n=0 \rightarrow a_2 = -3a_0$$

$$n=2 \rightarrow a_4 = 0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$y(x) = a_0 \left( 1 - 3x^2 \right) + a_1 \left( x - \frac{1}{3}x^3 \right) \Bigg|$$

$$a_1$$

$$n=1 \rightarrow a_3 = -\frac{1}{3}a_1$$

$$n=3 \rightarrow a_5 = 0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

### Exercise

Find a power series solution.  $(x^2 + 2)y'' + 4xy' + 2y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(x^2 + 2)y'' + 4xy' + 2y = 0$$

$$x^2 \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} + 2 \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + 4x \sum_{n=1}^{\infty} n a_n x^{n-1} + 2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^n + \sum_{n=0}^{\infty} 2(n+2)(n+1) a_{n+2} x^n + \sum_{n=1}^{\infty} 4n a_n x^n + \sum_{n=0}^{\infty} 2a_n x^n = 0$$

$$\sum_{n=0}^{\infty} n(n-1) a_n x^n + \sum_{n=0}^{\infty} 2(n+2)(n+1) a_{n+2} x^n + \sum_{n=0}^{\infty} 4n a_n x^n + \sum_{n=0}^{\infty} 2a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ 2(n+2)(n+1) a_{n+2} + (n^2 + 3n + 2) a_n \right] x^n = 0$$

$$2(n+1)(n+2) a_{n+2} + (n+1)(n+2) a_n = 0$$

$$\underline{a_{n+2} = -\frac{1}{2} a_n}$$

$$a_0$$

$$n=0 \rightarrow a_2 = -\frac{1}{2} a_0$$

$$n=2 \rightarrow a_4 = -\frac{1}{2} a_2 = \frac{1}{2^2} a_0$$

$$n=4 \rightarrow a_6 = -\frac{1}{2} a_4 = -\frac{1}{2^3} a_0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$a_{2n} = (-1)^n \frac{1}{2^n} a_0$$

$$a_1$$

$$n=1 \rightarrow a_3 = -\frac{1}{2} a_1$$

$$n=3 \rightarrow a_5 = -\frac{1}{2} a_3 = \frac{1}{2^2} a_1$$

$$n=5 \rightarrow a_7 = -\frac{1}{2} a_5 = -\frac{1}{2^3} a_1$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$a_{2n+1} = (-1)^n \frac{1}{2^n} a_1$$

$$y(x) = a_0 \left( 1 - \frac{1}{2}x^2 + \frac{1}{4}x^4 - \frac{1}{8}x^6 + \dots \right) + a_1 \left( x - \frac{1}{2}x^3 + \frac{1}{4}x^5 - \frac{1}{8}x^7 + \dots \right)$$


---


$$= a_0 \sum_{n=0}^{\infty} \frac{(-1)^n}{2^n} x^{2n} + a_1 \sum_{n=0}^{\infty} \frac{(-1)^n}{2^n} x^{2n+1}$$


---

### Exercise

Find a power series solution.  $(x^2 - 3)y'' + 2xy' = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(x^2 - 3)y'' + 2xy' = 0$$

$$x^2 \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} - 3 \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + 2x \sum_{n=1}^{\infty} n a_n x^{n-1} = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^n - \sum_{n=0}^{\infty} 3(n+2)(n+1) a_{n+2} x^n + \sum_{n=1}^{\infty} 2n a_n x^n = 0$$

$$\sum_{n=0}^{\infty} n(n-1) a_n x^n - \sum_{n=0}^{\infty} 3(n+1)(n+2) a_{n+2} x^n + \sum_{n=0}^{\infty} 2n a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ -3(n+1)(n+2) a_{n+2} + (n^2 + n) a_n \right] x^n = 0$$

$$-3(n+1)(n+2) a_{n+2} + n(n+1) a_n = 0$$

$$a_{n+2} = \frac{1}{3} \frac{n}{n+2} a_n$$

$$a_0$$

$$n=0 \rightarrow a_2 = 0$$

$$a_1$$

$$n=1 \rightarrow a_3 = \frac{1}{3^2} a_1$$

$$n=2 \rightarrow a_4 = \frac{2}{12}a_2 = 0$$

$$n=4 \rightarrow a_6 = 0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$n=3 \rightarrow a_5 = \frac{1}{3} \frac{3}{5} a_3 = \frac{1}{3^2 \cdot 5} a_1$$

$$n=5 \rightarrow a_7 = \frac{1}{3} \frac{5}{7} a_5 = \frac{1}{3^3 \cdot 7} a_1$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$a_{2n+1} = (-1)^n \frac{1}{(2n+1)3^n} a_1$$

$$y(x) = a_0 + a_1 \left( x + \frac{1}{9}x^3 + \frac{1}{45}x^5 + \frac{1}{189}x^7 + \dots \right)$$

$$= a_0 + a_1 \sum_{n=0}^{\infty} \frac{1}{(2n+1)3^n} x^{2n+1}$$

### Exercise

Find a power series solution.  $(x^2 + 3)y'' - 7xy' + 16y = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(x^2 + 3)y'' - 7xy' + 16y = 0$$

$$x^2 \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} + 3 \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - 7x \sum_{n=1}^{\infty} n a_n x^{n-1} + 16 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^n + \sum_{n=0}^{\infty} 3(n+1)(n+2) a_{n+2} x^n - \sum_{n=1}^{\infty} 7n a_n x^n + \sum_{n=0}^{\infty} 16 a_n x^n = 0$$

$$\sum_{n=0}^{\infty} n(n-1) a_n x^n + \sum_{n=0}^{\infty} 3(n+1)(n+2) a_{n+2} x^n - \sum_{n=0}^{\infty} 7n a_n x^n + \sum_{n=0}^{\infty} 16 a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ 3(n+1)(n+2)a_{n+2} + (n^2 - 8n + 16)a_n \right] x^n = 0$$

$$3(n+1)(n+2)a_{n+2} + (n-4)^2 a_n = 0$$

$$a_{n+2} = -\frac{(n-4)^2}{3(n+1)(n+2)} a_n$$

$a_0$

$a_1$

$$n=0 \rightarrow a_2 = -\frac{16}{6}a_0 = -\frac{8}{3}a_0$$

$$n=1 \rightarrow a_3 = -\frac{9}{18}a_1 = -\frac{1}{2}a_1$$

$$n=2 \rightarrow a_4 = -\frac{1}{9}a_2 = \frac{8}{27}a_0$$

$$n=3 \rightarrow a_5 = -\frac{1}{60}a_3 = \frac{1}{120}a_1$$

$$n=4 \rightarrow a_6 = 0$$

$$n=5 \rightarrow a_7 = -\frac{1}{126}a_5 = -\frac{1}{560 \cdot 3}a_1$$

$\vdots \quad \vdots \quad \vdots \quad \vdots$

$\vdots \quad \vdots \quad \vdots \quad \vdots$

$$y(x) = a_0 \left( 1 - \frac{8}{3}x^2 + \frac{8}{27}x^4 \right) + a_1 \left( x - \frac{1}{2}x^3 + \frac{1}{120}x^5 + \frac{1}{15,120}x^7 + \dots \right)$$

### Exercise

Find the series solution to the initial value problem  $y'' + 4y = 0$  ;  $y(0) = 0$ ,  $y'(0) = 3$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' + 4y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + 4 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ (n+2)(n+1) a_{n+2} + 4a_n \right] x^n = 0$$

$$(n+2)(n+1) a_{n+2} + 4a_n = 0$$

$$a_{n+2} = -\frac{4}{(n+1)(n+2)} a_n$$

Given:  $y(0) = 0 = a_0$ ,  $y'(0) = 3 = a_1$

$$a_0 = 0$$

$$n=0 \rightarrow a_2 = -2a_0 = 0$$

$$n=2 \rightarrow a_4 = -\frac{4}{12}a_2 = 0$$

$$n=4 \rightarrow a_6 = 0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$a_1 = 3$$

$$n=1 \rightarrow a_3 = -\frac{4}{6}a_1 = -\frac{2^2}{3!}a_1 = -2$$

$$n=3 \rightarrow a_5 = -\frac{4}{20}a_3 = -\frac{2^4}{5!}a_1 = \frac{2}{5}$$

$$n=5 \rightarrow a_7 = -\frac{4}{42}a_5 = -\frac{2^6}{7!}a_1 = -\frac{4}{105}$$

$$a_{2k+1} = \frac{(-1)^k 2^{2k}}{(2k+1)!} a_1$$

$$y(x) = 3x - 2x^3 + \frac{2}{5}x^5 - \frac{4}{105}x^7 + \dots$$

$$= 3 \left( x - \frac{2^2}{3!}x^3 + \frac{2^4}{5!}x^5 - \frac{2^6}{7!}x^7 + \dots \right)$$

$$= \frac{3}{2} \left( (2x) - \frac{1}{3!}(2x)^3 + \frac{1}{5!}(2x)^5 - \frac{1}{7!}(2x)^7 + \dots \right)$$

$$= \frac{3}{2} \sin 2x$$

## Exercise

Find the series solution to the initial value problem  $y'' + x^2 y = 0$ ;  $y(0) = 1$ ,  $y'(0) = 0$

## Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' + x^2 y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + x^2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n + \sum_{n=0}^{\infty} a_n x^{n+2} = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}x^n + \sum_{n=2}^{\infty} a_{n-2}x^n = 0$$

$$2a_2 + 6a_3x + \sum_{n=2}^{\infty} (n+2)(n+1)a_{n+2}x^n + \sum_{n=2}^{\infty} a_{n-2}x^n = 0$$

$$2a_2 + 6a_3x + \sum_{n=2}^{\infty} [(n+1)(n+2)a_{n+2} + a_{n-2}]x^n = 0$$

$$(n+1)(n+2)a_{n+2} + a_{n-2} = 0$$

$$a_{n+2} = -\frac{1}{(n+1)(n+2)}a_{n-2}$$

Given:  $y(0) = 1 = a_0$ ,  $y'(0) = 0 = a_1$

$$2a_2 + 6a_3x = 0 \rightarrow a_2 = a_3 = 0$$

$$a_0 = 1$$

$$n=2 \rightarrow a_4 = -\frac{1}{12}a_0 = -\frac{1}{12}$$

$$a_1 = a_2 = a_3 = 0$$

$$n=3 \rightarrow a_5 = -\frac{1}{20}a_1 = 0$$

$$n=4 \rightarrow a_6 = *a_2 = 0$$

$$n=5 \rightarrow a_7 = *a_3 = 0$$

$$n=6 \rightarrow a_8 = -\frac{1}{56}a_4 = \frac{1}{672}$$

$$n=10 \rightarrow a_{12} = -\frac{1}{132}a_8 = \frac{1}{88,704}$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$y(x) = 1 - \frac{1}{12}x^4 + \frac{1}{672}x^8 - \frac{1}{88,704}x^{12} + \dots$$

### Exercise

Find the series solution to the initial value problem  $y'' - 2xy' + 8y = 0$ ;  $y(0) = 3$ ,  $y'(0) = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1)a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n$$

$$y'' - 2xy' + 8y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n - 2x \sum_{n=0}^{\infty} (n+1)a_{n+1} x^n + 8 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n - \sum_{n=0}^{\infty} 2(n+1)a_{n+1} x^{n+1} + \sum_{n=0}^{\infty} 8a_n x^n = 0$$

$$\sum_{n=0}^{\infty} [(n+2)(n+1)a_{n+2} + 8a_n] x^n - \sum_{n=1}^{\infty} 2na_n x^n = 0$$

$$2a_2 + 8a_0 + \sum_{n=1}^{\infty} [(n+2)(n+1)a_{n+2} + 8a_n] x^n - \sum_{n=1}^{\infty} 2na_n x^n = 0$$

$$2a_2 + 8a_0 + \sum_{n=1}^{\infty} [(n+2)(n+1)a_{n+2} + (8-2n)a_n] x^n = 0$$

$$\text{Given: } y(0) = 3 = a_0, \quad y'(0) = 0 = a_1$$

$$2a_2 + 8a_0 = 0 \rightarrow a_2 = -4a_0 = -12$$

$$(n+2)(n+1)a_{n+2} + (8-2n)a_n = 0$$

$$\rightarrow a_{n+2} = \frac{2n-8}{(n+1)(n+2)} a_n \quad n=1, 2, \dots$$

$$a_3 = -a_1 = 0$$

$$a_4 = -\frac{1}{3}a_2 = 4$$

$$a_5 = -\frac{1}{10}a_3 = 0$$

$$a_6 = 0a_4 = 0$$

$$a_7 = \frac{1}{21}a_5 = 0$$

$$a_8 = 0a_6 = 0$$

$$y(x) = 3 - 12x^2 + 4x^4$$

### Exercise

Find the series solution to the initial value problem  $y'' + y' - 2y = 0$ ;  $y(0) = 1$ ,  $y'(0) = -2$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$



$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' + y' - 2y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n - 2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} [(n+2)(n+1) a_{n+2} + (n+1) a_{n+1} - 2a_n] x^n = 0$$

$$(n+2)(n+1) a_{n+2} + (n+1) a_{n+1} - 2a_n = 0$$

$$a_{n+2} = \frac{2a_n - (n+1)a_{n+1}}{(n+1)(n+2)}$$

$$\text{Given: } y(0) = 1 = a_0, \quad y'(0) = -2 = a_1$$

$$a_0 = 1$$

$$a_1 = -2$$

$$n=0 \rightarrow a_2 = \frac{2a_0 - a_1}{2} = 2$$

$$n=1 \rightarrow a_3 = \frac{2a_1 - 2a_2}{6} = -\frac{8}{6}$$

$$n=2 \rightarrow a_4 = \frac{2a_2 - 3a_3}{12} = \frac{2}{3}$$

$$n=3 \rightarrow a_5 = \frac{2a_3 - 4a_4}{20} = -\frac{4}{15}$$

$$n=4 \rightarrow a_6 = \frac{2a_4 - 5a_5}{30} = \frac{4}{45}$$

$$n=5 \rightarrow a_7 = \frac{2a_5 - 6a_6}{42} = \frac{8}{315}$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$y(x) = 1 - 2x + 2x^2 - \frac{4}{3}x^3 + \frac{2}{3}x^4 - \frac{4}{15}x^5 + \frac{4}{45}x^6 - \frac{8}{315}x^7 + \dots$$

$$= 1 + (-2x) + \frac{1}{2!}(-2x)^2 + \frac{1}{3!}(-2x)^3 + \frac{1}{4!}(-2x)^4 + \frac{1}{5!}(-2x)^5 + \frac{1}{6!}(-2x)^6 + \frac{1}{7!}(-2x)^7 + \dots$$

$$= e^{-2x}$$

### Exercise

Find the series solution to the initial value problem  $y'' - 2y' + y = 0$ ;  $y(0) = 0$ ,  $y'(0) = 1$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' - 2y' + y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - 2 \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} [(n+2)(n+1) a_{n+2} - 2(n+1) a_{n+1} + a_n] x^n = 0$$

$$(n+2)(n+1) a_{n+2} - 2(n+1) a_{n+1} + a_n = 0$$

$$a_{n+2} = \frac{2(n+1) a_{n+1} - a_n}{(n+1)(n+2)}$$

$$\text{Given: } y(0) = 0 = a_0, \quad y'(0) = 1 = a_1$$

$$a_0 = 0$$

$$a_1 = 1$$

$$n=0 \rightarrow a_2 = \frac{2a_1 - a_0}{2} = 1$$

$$n=1 \rightarrow a_3 = \frac{4a_2 - a_1}{6} = \frac{1}{2}$$

$$n=2 \rightarrow a_4 = \frac{6a_3 - a_2}{12} = \frac{1}{6}$$

$$n=3 \rightarrow a_5 = \frac{8a_4 - a_3}{20} = \frac{1}{20} \left( \frac{4}{3} - \frac{1}{2} \right) = \frac{1}{24}$$

$$n=4 \rightarrow a_6 = \frac{10a_5 - a_4}{30} = \frac{1}{30} \left( \frac{5}{12} - \frac{1}{6} \right) = \frac{1}{120} \quad n=5 \rightarrow a_7 = \frac{12a_6 - a_5}{42} = \frac{1}{42} \left( \frac{1}{10} - \frac{1}{24} \right) = \frac{1}{720}$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots$$

$$y(x) = x + x^2 + \frac{1}{2}x^3 + \frac{1}{6}x^4 + \frac{1}{24}x^5 + \frac{1}{120}x^6 + \frac{1}{720}x^7 + \dots$$

$$= x \left( 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \frac{1}{5!}x^5 + \frac{1}{6!}x^6 + \dots \right)$$

$$= x e^x$$

## Exercise

Find the series solution to the initial value problem  $y'' + xy' + y = 0$   $y(0) = 1$   $y'(0) = 0$

## Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' + xy' + y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + x \sum_{n=1}^{\infty} n a_n x^{n-1} + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + \sum_{n=0}^{\infty} n a_n x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ (n+2)(n+1) a_{n+2} + n a_n + a_n \right] x^n = 0$$

$$(n+2)(n+1) a_{n+2} + (n+1) a_n = 0$$

$$(n+2)(n+1) a_{n+2} = -(n+1) a_n$$

$$a_{n+2} = -\frac{1}{n+2} a_n$$

$$a_0 = y(0) = 1$$

$$a_1 = y'(0) = 0$$

$$a_2 = -\frac{1}{2} a_0 = -\frac{1}{2}$$

$$a_3 = -\frac{1}{3} a_1 = 0$$

$$a_4 = -\frac{1}{4} a_2 = \frac{1}{2 \cdot 4} = \frac{1}{2^2 \cdot 1 \cdot 2}$$

$$a_5 = -\frac{1}{5} a_3 = 0$$

$$a_6 = -\frac{1}{6} a_4 = -\frac{1}{2^3 \cdot 1 \cdot 2 \cdot 3}$$

$$a_7 = -\frac{1}{7} a_7 = 0$$

$$y(x) = \sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + a_4 x^4 + a_5 x^5 + \dots$$

$$= 1 - \frac{1}{2} x^2 + \frac{1}{2^2 2!} x^4 - \frac{1}{2^3 3!} x^5 + \dots$$

### Exercise

Find the series solution to the initial value problem  $y'' - xy' - y = 0$   $y(0) = 2$   $y'(0) = 1$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' - xy' - y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - x \sum_{n=1}^{\infty} n a_n x^{n-1} - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=0}^{\infty} n a_n x^n - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ (n+2)(n+1) a_{n+2} - n a_n - a_n \right] x^n = 0$$

$$(n+2)(n+1) a_{n+2} - (n+1) a_n = 0$$

$$(n+2)(n+1) a_{n+2} = -(n+1) a_n$$

$$\underline{a_{n+2} = \frac{1}{n+2} a_n}$$

$$a_0 = y(0) = 2$$

$$a_1 = y'(0) = 1$$

$$a_2 = \frac{1}{2} a_0 = 1$$

$$a_3 = \frac{1}{3} a_1 = \frac{1}{3}$$

$$a_4 = \frac{1}{4} a_2 = \frac{1}{4}$$

$$a_5 = \frac{1}{5} a_3 = \frac{1}{3 \cdot 5}$$

$$a_6 = \frac{1}{6} a_4 = \frac{1}{4 \cdot 6} = \frac{1}{24}$$

$$a_7 = \frac{1}{7} a_5 = \frac{1}{3 \cdot 5 \cdot 7}$$

$$\vdots \quad \vdots \quad \vdots$$

$$\vdots \quad \vdots \quad \vdots$$

$$\underline{y(x) = 2 + x + x^2 + \frac{1}{3}x^3 + \frac{1}{4}x^4 + \frac{1}{15}x^5 + \frac{1}{24}x^6 + \dots}$$

### Exercise

Find the series solution to the initial value problem  $y'' - xy' - y = 0$  ;  $y(0) = 1$   $y'(0) = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' - xy' - y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - x \sum_{n=1}^{\infty} n a_n x^{n-1} - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=0}^{\infty} n a_n x^n - \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ (n+2)(n+1) a_{n+2} - n a_n - a_n \right] x^n = 0$$

$$(n+2)(n+1) a_{n+2} - (n+1) a_n = 0$$

$$(n+2)(n+1) a_{n+2} = -(n+1) a_n$$

$$\underline{a_{n+2} = \frac{1}{n+2} a_n}$$

$$\text{Given: } a_0 = y(0) = 1$$

$$a_1 = y'(0) = 0$$

$$a_2 = \frac{1}{2} a_0 = \frac{1}{2}$$

$$a_3 = \frac{1}{3} a_1 = 0$$

$$a_4 = \frac{1}{4} a_2 = \frac{1}{2 \cdot 2^2}$$

$$a_5 = \frac{1}{5} a_3 = 0$$

$$a_6 = \frac{1}{6} a_4 = \frac{1}{2^3 \cdot 3!}$$

$$\vdots \quad \vdots \quad \vdots$$

$$\vdots \quad \vdots \quad \vdots$$

$$\underline{y(x) = 1 + \frac{1}{2}x^2 + \frac{1}{8}x^4 + \frac{1}{48}x^6 + \dots}$$

### Exercise

Find a power series solution.  $y'' + xy' - 2y = 0$  ;  $y(0) = 1$   $y'(0) = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' + xy' - 2y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + x \sum_{n=1}^{\infty} n a_n x^{n-1} - \sum_{n=0}^{\infty} 2 a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + \sum_{n=0}^{\infty} n a_n x^n - \sum_{n=0}^{\infty} 2 a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ (n+1)(n+2) a_{n+2} + (n-2) a_n \right] x^n = 0$$

$$(n+2)(n+1) a_{n+2} + (n-2) a_n = 0$$

$$a_{n+2} = -\frac{n-2}{(n+1)(n+2)} a_n$$

$$a_0 = y(0) = 1$$

$$a_1 = y'(0) = 0$$

$$n=0 \rightarrow a_2 = \frac{2}{2} a_0 = 1$$

$$n=1 \rightarrow a_3 = \frac{1}{6} a_1 = 0$$

$$n=2 \rightarrow a_4 = 0$$

$$n=3 \rightarrow a_5 = 0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$y(x) = 1 + x^2$$

### Exercise

Find the series solution to the initial value problem  $y'' + (x-1)y' + y = 0$   $y(1) = 2$   $y'(1) = 0$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$y'' + (x-1)y' + y = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} (x-1)^n + (x-1) \sum_{n=1}^{\infty} n a_n (x-1)^{n-1} + \sum_{n=0}^{\infty} a_n (x-1)^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} (x-1)^n + \sum_{n=1}^{\infty} n a_n (x-1)^n + \sum_{n=0}^{\infty} a_n (x-1)^n = 0$$

$$\sum_{n=0}^{\infty} [(n+2)(n+1) a_{n+2} + n a_n + a_n] x^{n-1} = 0$$

$$(n+2)(n+1) a_{n+2} + (n+1) a_n = 0$$

$$a_{n+2} = -\frac{1}{n+2} a_n$$

$$a_0 = y(1) = 2$$

$$a_1 = y'(1) = 0$$

$$a_2 = -\frac{1}{2} a_0 = -1$$

$$a_3 = -\frac{1}{3} a_1 = 0$$

$$a_4 = -\frac{1}{4} a_2 = \frac{1}{2 \cdot 4} a_0 = \frac{1}{4}$$

$$a_5 = -\frac{1}{5} a_3 = 0$$

$$a_6 = -\frac{1}{6} a_4 = -\frac{1}{24}$$

$$a_7 = -\frac{1}{7} a_5 = 0$$

$$y(x) = \sum_{n=0}^{\infty} a_n (x-1)^n = a_0 + a_1 (x-1) + a_2 (x-1)^2 + a_3 (x-1)^3 + a_4 (x-1)^4 + \dots$$

$$= 2 - (x-1)^2 + \frac{1}{4} (x-1)^4 - \frac{1}{24} (x-1)^6 + \dots$$

$$= \sum_{n=0}^{\infty} \frac{(-1)^n (x-1)^{2n}}{n! 2^n}$$

### Exercise

Find the series solution to the initial value problem  $(x-1)y'' - xy' + y = 0$ ;  $y(0) = -2$ ,  $y'(0) = 6$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(x-1)y'' - xy' + y = 0$$

$$(x-1) \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - x \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^{n+1} - \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=0}^{\infty} (n+1) a_{n+1} x^{n+1} + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=1}^{\infty} n(n+1) a_{n+1} x^n - \sum_{n=1}^{\infty} n a_n x^n + \sum_{n=0}^{\infty} [a_n - (n+2)(n+1) a_{n+2}] x^n = 0$$

$$\sum_{n=1}^{\infty} [n(n+1) a_{n+1} - n a_n] x^n + a_0 - 2a_2 + \sum_{n=1}^{\infty} [a_n - (n+2)(n+1) a_{n+2}] x^n = 0$$

$$a_0 - 2a_2 + \sum_{n=1}^{\infty} [n(n+1) a_{n+1} - (n-1) a_n - (n+2)(n+1) a_{n+2}] x^n = 0$$

$$\text{Given: } y(0) = -2 = a_0, \quad y'(0) = 6 = a_1$$

$$a_0 - 2a_2 = 0 \rightarrow a_2 = \frac{1}{2} a_0 = -1$$

$$n(n+1) a_{n+1} - (n-1) a_n - (n+2)(n+1) a_{n+2} = 0$$

$$\rightarrow a_{n+2} = \frac{n}{n+2} a_{n+1} - \frac{n-1}{(n+2)(n+1)} a_n$$

$$a_3 = \frac{1}{3} a_2 - 0 a_1 = -\frac{1}{3}$$

$$a_4 = \frac{1}{2} a_3 - \frac{1}{12} a_2 = -\frac{1}{6} + \frac{1}{12} = -\frac{1}{12}$$

$$a_5 = \frac{3}{5} a_4 - \frac{1}{10} a_3 = -\frac{3}{60} + \frac{1}{30} = -\frac{1}{60}$$

$$a_6 = \frac{2}{3} a_5 - \frac{1}{10} a_4 = -\frac{1}{90} + \frac{1}{120} = -\frac{1}{360}$$

$$y(x) = \left( -2 - x^2 - \frac{1}{3} x^3 - \frac{1}{12} x^4 - \dots \right) + a_1 x$$



$$\begin{aligned}
&= -2\left(1 + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \dots\right) + 6x \\
&= -2\left(1 + \textcolor{red}{x} + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{1}{24}x^4 + \dots\right) + 6x + \textcolor{red}{2x} \\
&= \underline{8x - 2e^x}
\end{aligned}$$

### Exercise

Find the series solution to the initial value problem

$$(x+1)y'' - (2-x)y' + y = 0; \quad y(0) = 2, \quad y'(0) = -1$$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$\textcolor{red}{(x+1)y'' - (2-x)y' + y = 0}$$

$$(x+1) \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - (2-x) \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^{n+1} + \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=0}^{\infty} 2(n+1) a_{n+1} x^n$$

$$+ \sum_{n=0}^{\infty} (n+1) a_{n+1} x^{n+1} + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} [(n+2)(n+1) a_{n+2} + (n+1) a_{n+1}] x^{n+1} + \sum_{n=0}^{\infty} [(n+2)(n+1) a_{n+2} - 2(n+1) a_{n+1} + a_n] x^n = 0$$

$$\sum_{\textcolor{red}{n=1}}^{\infty} [n(n+1) a_{n+1} + n a_n] x^n + \sum_{n=0}^{\infty} [(n+2)(n+1) a_{n+2} - 2(n+1) a_{n+1} + a_n] x^n = 0$$

$$\sum_{n=1}^{\infty} [n(n+1) a_{n+1} + n a_n] x^n + \textcolor{blue}{2a_2 - 2a_1 + a_0} + \sum_{\textcolor{red}{n=1}}^{\infty} [(n+2)(n+1) a_{n+2} - 2(n+1) a_{n+1} + a_n] x^n = 0$$

$$2a_2 - 2a_1 + a_0 + \sum_{n=1}^{\infty} \left[ (n+2)(n+1)a_{n+2} + (n-2)(n+1)a_{n+1} + (n+1)a_n \right] x^n = 0$$

Given:  $y(0) = 2 = a_0$ ,  $y'(0) = -1 = a_1$

$$2a_2 - 2a_1 + a_0 = 0 \rightarrow \underline{a_2} = \frac{1}{2}(2a_1 - a_0) = \underline{-2}$$

$$(n+2)(n+1)a_{n+2} + (n-2)(n+1)a_{n+1} + (n+1)a_n = 0$$

$$\rightarrow a_{n+2} = -\frac{n-2}{n+2}a_{n+1} - \frac{1}{n+2}a_n$$

$$a_3 = \frac{1}{3}a_2 - \frac{1}{3}a_1 = \frac{2}{3} + \frac{1}{3} = 1$$

$$a_4 = 0a_3 - \frac{1}{4}a_2 = \frac{1}{2}$$

$$a_5 = -\frac{1}{5}a_4 - \frac{1}{5}a_3 = -\frac{1}{10} - \frac{1}{5} = -\frac{3}{10}$$

$$a_6 = -\frac{1}{3}a_5 - \frac{1}{6}a_4 = \frac{1}{10} - \frac{1}{12} = \frac{1}{60}$$

$$\underline{y(x) = 2 - x - 2x^2 + x^3 + \frac{1}{2}x^4 - \frac{3}{10}x^5 + \dots}$$

### Exercise

Find the series solution to the initial value problem

$$(1-x)y'' + xy' - 2y = 0; \quad y(0) = 0, \quad y'(0) = 1$$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(1-x)y'' + xy' - 2y = 0$$

$$(1-x) \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + x \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n - 2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n - \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^{n+1} + \sum_{n=0}^{\infty} (n+1) a_{n+1} x^{n+1} - \sum_{n=0}^{\infty} 2a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ (n+2)(n+1) a_{n+2} - 2a_n \right] x^n - \sum_{n=0}^{\infty} \left[ (n+2)(n+1) a_{n+2} - (n+1) a_{n+1} \right] x^{n+1} = 0$$

$$\begin{aligned}
& \sum_{n=0}^{\infty} \left[ (n+2)(n+1)a_{n+2} - 2a_n \right] x^n - \sum_{n=1}^{\infty} \left[ n(n+1)a_{n+1} - na_n \right] x^n = 0 \\
& 2a_2 - 2a_0 + \sum_{n=1}^{\infty} \left[ (n+2)(n+1)a_{n+2} - 2a_n \right] x^n - \sum_{n=1}^{\infty} \left[ n(n+1)a_{n+1} - na_n \right] x^n = 0 \\
& 2a_2 - 2a_0 + \sum_{n=1}^{\infty} \left[ (n+2)(n+1)a_{n+2} - n(n+1)a_{n+1} + (n-2)a_n \right] x^n = 0 \\
& (n+2)(n+1)a_{n+2} - n(n+1)a_{n+1} + (n-2)a_n = 0 \\
& a_{n+2} = \frac{n(n+1)a_{n+1} - (n-2)a_n}{(n+1)(n+2)} \quad \Big|
\end{aligned}$$

Given:  $y(0) = 0 = a_0$ ,  $y'(0) = 1 = a_1$

$$2a_2 - 2a_0 = 0 \rightarrow \underline{a_2 = a_0 = 0}$$

$$n=1 \rightarrow a_3 = \frac{2a_2 + a_1}{6} = \frac{1}{6}$$

$$n=2 \rightarrow a_4 = \frac{6a_3}{12} = \frac{1}{12}$$

$$n=3 \rightarrow a_5 = \frac{1}{20} \left( 12a_4 - a_3 \right) = \frac{1}{20} \left( 1 - \frac{1}{6} \right) = \frac{1}{24}$$

$$n=4 \rightarrow a_6 = \frac{1}{30} \left( 20a_5 - 2a_4 \right) = \frac{1}{30} \left( \frac{5}{6} - \frac{1}{6} \right) = \frac{1}{45}$$

$$n=5 \rightarrow a_7 = \frac{1}{42} \left( 30a_6 - 3a_5 \right) = \frac{1}{30} \left( \frac{2}{3} - \frac{1}{8} \right) = \frac{13}{1008}$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\underline{y(x) = x + \frac{1}{6}x^3 + \frac{1}{12}x^4 + \frac{1}{24}x^5 + \frac{1}{45}x^6 + \frac{13}{1008}x^7 + \dots}$$

### Exercise

Find the series solution to the initial value problem

$$(x^2 + 1)y'' + 2xy' = 0; \quad y(0) = 0, \quad y'(0) = 1$$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} na_n x^{n-1} = \sum_{n=0}^{\infty} (n+1)a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1)a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n$$

$$(x^2 + 1)y'' + 2xy' = 0$$

$$(x^2 + 1) \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n + 2x \sum_{n=0}^{\infty} (n+1)a_{n+1} x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^{n+2} + \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n + \sum_{n=0}^{\infty} 2(n+1)a_{n+1} x^{n+1} = 0$$

$$\sum_{n=2}^{\infty} n(n-1)a_n x^n + \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2} x^n + \sum_{n=1}^{\infty} 2na_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1)a_n x^n + 2a_2 + 6a_3 x + \sum_{n=2}^{\infty} (n+2)(n+1)a_{n+2} x^n + 2a_1 x + \sum_{n=2}^{\infty} 2na_n x^n = 0$$

$$2a_2 + (6a_3 + 2a_1)x + \sum_{n=2}^{\infty} [(n(n-1) + 2n)a_n + (n+2)(n+1)a_{n+2}] x^n = 0$$

$$\text{Given: } y(0) = 0 = a_0, \quad y'(0) = 1 = a_1$$

$$2a_2 + (6a_3 + 2a_1)x = 0 \rightarrow \begin{cases} a_2 = 0 \\ a_3 = -\frac{1}{3}a_1 = -\frac{1}{3} \end{cases}$$

$$n(n+1)a_n + (n+2)(n+1)a_{n+2} = 0$$

$$\rightarrow a_{n+2} = -\frac{n}{n+2}a_n \quad n = 2, 3, \dots$$

$$a_4 = -\frac{1}{2}a_2 = 0$$

$$a_5 = -\frac{3}{5}a_3 = \frac{1}{5}$$

$$a_6 = -\frac{2}{3}a_4 = 0$$

$$a_7 = -\frac{5}{7}a_5 = -\frac{1}{7}$$

$$\underline{y(x) = x - \frac{1}{3}x^3 + \frac{1}{5}x^5 - \frac{1}{7}x^7 + \dots}$$

### Exercise

Find the series solution to the initial value problem

$$(x^2 - 1)y'' + 3xy' + xy = 0; \quad y(0) = 4, \quad y'(0) = 6$$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(x^2 - 1)y'' + 3xy' + xy = 0$$

$$(x^2 - 1) \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + 3x \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n + x \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^{n+2} - \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + \sum_{n=0}^{\infty} 3(n+1) a_{n+1} x^{n+1} + \sum_{n=0}^{\infty} a_n x^{n+1} = 0$$

$$\sum_{n=0}^{\infty} (n+1)(n+2) a_{n+2} x^{n+2} + \sum_{n=0}^{\infty} [(3n+3) a_{n+1} + a_n] x^{n+1} - \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^n + \sum_{n=1}^{\infty} (3n a_n + a_{n-1}) x^n - \sum_{n=0}^{\infty} (n+1)(n+2) a_{n+2} x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^n + (3a_1 + a_0)x + \sum_{n=2}^{\infty} (3n a_n + a_{n-1}) x^n - 2a_2 - 6a_3 x - \sum_{n=2}^{\infty} (n+1)(n+2) a_{n+2} x^n = 0$$

$$-2a_2 + (3a_1 + a_0 - 6a_3)x + \sum_{n=2}^{\infty} [(n^2 + 2n) a_n + a_{n-1} - (n+2)(n+1) a_{n+2}] x^n = 0$$

$$\text{Given: } y(0) = 4 = a_0, \quad y'(0) = 6 = a_1$$

$$\begin{cases} -2a_2 = 0 & \rightarrow \underline{a_2 = 0} \\ 3a_1 + a_0 - 6a_3 = 0 & \rightarrow \underline{a_3 = \frac{22}{6} = \frac{11}{3}} \end{cases}$$

$$(n^2 + 2n) a_n + a_{n-1} - (n+2)(n+1) a_{n+2} = 0$$

$$\underline{a_{n+2} = \frac{(n^2 + 2n) a_n + a_{n-1}}{(n+1)(n+2)}}$$

$$n = 2 \rightarrow a_4 = \frac{8a_2 + a_1}{12} = \frac{6}{12} = \frac{1}{2}$$

$$n = 3 \rightarrow a_5 = \frac{1}{20}(15a_3 + a_2) = \frac{1}{20}(55) = \frac{11}{4}$$

$$n = 4 \rightarrow a_6 = \frac{1}{30}(24a_4 + a_3) = \frac{1}{30}\left(12 + \frac{11}{3}\right) = \frac{47}{90}$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\underline{y(x) = 4 + 6x + \frac{11}{3}x^3 + \frac{1}{2}x^4 + \frac{11}{4}x^5 + \frac{47}{90}x^6 + \dots}$$

### Exercise

Find the series solution to the initial value problem

$$(2 + x^2)y'' - xy' + 4y = 0 \quad y(0) = -1 \quad y'(0) = 3$$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(2 + x^2)y'' - xy' + 4y = 0$$

$$(2 + x^2) \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} - x \sum_{n=1}^{\infty} n a_n x^{n-1} + 4 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$2 \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} + x^2 \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} - \sum_{n=0}^{\infty} n a_n x^{n-1} + 4 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} 2(n+2)(n+1) a_{n+2} x^n + \sum_{n=0}^{\infty} n(n-1) a_n x^n - \sum_{n=0}^{\infty} n a_n x^{n-1} + \sum_{n=0}^{\infty} 4 a_n x^n = 0$$

$$2(n+2)(n+1) a_{n+2} + n(n-1) a_n - n a_n + 4 a_n = 0$$

$$2(n+2)(n+1) a_{n+2} + (n^2 - 2n + 4) a_n = 0$$

$$\underline{a_{n+2} = -\frac{n^2 - 2n + 4}{2(n+2)(n+1)} a_n}$$

$$a_0 = y(0) = -1$$

$$a_1 = y'(0) = 3$$

$$n=0 \rightarrow a_2 = -\frac{4}{4}a_0 = 1$$

$$n=1 \rightarrow a_3 = -\frac{3}{12}a_1 = -\frac{1}{4}(3) = -\frac{3}{4}$$

$$n=2 \rightarrow a_4 = -\frac{4}{24}a_2 = -\frac{1}{6}$$

$$n=3 \rightarrow a_5 = -\frac{7}{40}a_3 = -\frac{7}{40}\left(-\frac{3}{4}\right) = \frac{21}{160}$$

$$y(x) = \underline{-1 + 3x + x^2 - \frac{3}{4}x^3 - \frac{1}{6}x^4 + \frac{21}{160}x^5 + \dots}$$

### Exercise

Find the series solution to the initial value problem

$$(2 - x^2)y'' - xy' + 4y = 0 \quad y(0) = 1 \quad y'(0) = 0$$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(2 - x^2)y'' - xy' + 4y = 0$$

$$(2 - x^2) \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} - x \sum_{n=1}^{\infty} n a_n x^{n-1} + 4 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$2 \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} - x^2 \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} - \sum_{n=0}^{\infty} n a_n x^{n-1} + 4 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} 2(n+2)(n+1) a_{n+2} x^n - \sum_{n=2}^{\infty} n(n-1) a_n x^n - \sum_{n=0}^{\infty} n a_n x^{n-1} + \sum_{n=0}^{\infty} 4 a_n x^n = 0$$

$$\sum_{n=0}^{\infty} 2(n+2)(n+1) a_{n+2} x^n - \sum_{n=0}^{\infty} n(n-1) a_n x^n - \sum_{n=0}^{\infty} n a_n x^{n-1} + \sum_{n=0}^{\infty} 4 a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ 2(n+1)(n+2) a_{n+2} - (n^2 - n + n - 4) a_n \right] x^n = 0$$

$$2(n+1)(n+2) a_{n+2} - (n-2)(n+2) a_n = 0$$

$$a_{n+2} = \underline{\frac{n-2}{2(n+1)} a_n}$$

$$a_0 = y(0) = 1$$

$$a_1 = y'(0) = 0$$

$$n=0 \rightarrow a_2 = \frac{-2}{2}a_0 = -1$$

$$n=1 \rightarrow a_3 = -\frac{1}{4}a_1 = 0$$

$$n=2 \rightarrow a_4 = 0$$

$$n=3 \rightarrow a_5 = *a_3 = 0$$

$$\vdots \quad \vdots \quad \vdots$$

$$\vdots \quad \vdots \quad \vdots$$

$$\boxed{y(x) = 1 - x^2}$$

### Exercise

Find the series solution to the initial value problem

$$(4 - x^2)y'' + 2y = 0 \quad y(0) = 0 \quad y'(0) = 1$$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(4 - x^2)y'' + 2y = 0$$

$$(4 - x^2) \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} + 2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=2}^{\infty} 4n(n-1) a_n x^{n-2} - \sum_{n=2}^{\infty} n(n-1) a_n x^n + 2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} 4(n+1)(n+2) a_{n+2} x^n - \sum_{n=0}^{\infty} n(n-1) a_n x^n + 2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ 4(n+1)(n+2) a_{n+2} - (n^2 - n - 2) a_n \right] x^n = 0$$

$$4(n+1)(n+2) a_{n+2} - (n+1)(n-2) a_n = 0$$

$$\boxed{a_{n+2} = \frac{n-2}{4(n+2)} a_n}$$



$$a_0 = y(0) = 0$$

$$a_1 = y'(0) = 1$$

$$n=0 \rightarrow a_2 = \frac{-2}{8}a_0 = 0$$

$$n=1 \rightarrow a_3 = -\frac{1}{12}a_1 = -\frac{1}{12}$$

$$n=2 \rightarrow a_4 = 0$$

$$n=3 \rightarrow a_5 = \frac{1}{20}a_3 = -\frac{1}{240}$$

$$\vdots \quad \vdots \quad \vdots$$

$$n=5 \rightarrow a_7 = \frac{3}{28}a_5 = -\frac{1}{2,240}$$

$$\vdots \quad \vdots \quad \vdots$$

$$y(x) = x - \frac{1}{12}x^3 - \frac{1}{240}x^5 - \frac{1}{2240}x^7 - \frac{1}{16,128}x^9 - \dots$$

### Exercise

Find a power series solution.  $(x^2 - 4)y'' + 3xy' + y = 0$ ;  $y(0) = 4$ ,  $y'(0) = 1$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(x^2 - 4)y'' + 3xy' + y = 0$$

$$x^2 \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} - 4 \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} + 3x \sum_{n=1}^{\infty} n a_n x^{n-1} + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^n - \sum_{n=2}^{\infty} 4n(n-1) a_n x^{n-2} + \sum_{n=1}^{\infty} 3n a_n x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} n(n-1) a_n x^n - \sum_{n=0}^{\infty} 4(n+2)(n+1) a_{n+2} x^n + \sum_{n=0}^{\infty} 3n a_n x^n + \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ (n^2 - n + 3n + 1) a_n - 4(n+2)(n+1) a_{n+2} \right] x^n = 0$$

$$(n^2 + 2n + 1) a_n - 4(n+2)(n+1) a_{n+2} = 0$$

$$4(n+2)(n+1)a_{n+2} = (n+1)^2 a_n$$

$$\boxed{a_{n+2} = \frac{n+1}{4(n+2)} a_n}$$

$$a_0 = y(0) = 4$$

$$a_1 = y'(0) = 1$$

$$n=0 \rightarrow a_2 = \frac{1}{8}a_0 = \frac{1}{2}$$

$$n=1 \rightarrow a_3 = \frac{2}{4 \cdot 3}a_1 = \frac{1}{6}$$

$$n=2 \rightarrow a_4 = \frac{3}{16}a_2 = \frac{3}{32}$$

$$n=3 \rightarrow a_5 = \frac{1}{5}a_3 = \frac{1}{30}$$

$$n=4 \rightarrow a_6 = \frac{5}{24}a_4 = \frac{5}{256}$$

$$n=5 \rightarrow a_7 = \frac{6}{4 \cdot 7}a_5 = \frac{1}{140}$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\boxed{y(x) = 4 + x + \frac{1}{2}x^2 + \frac{1}{6}x^3 + \frac{3}{32}x^4 + \frac{1}{30}x^5 + \frac{5}{256}x^6 + \frac{1}{140}x^7 + \cdots}$$

### Exercise

Find a power series solution.  $(x^2 + 1)y'' + 2xy' - 2y = 0$ ;  $y(0) = 0$ ,  $y'(0) = 1$

### Solution

$$y(x) = \sum_{n=0}^{\infty} a_n x^n$$

$$y'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} x^n$$

$$y''(x) = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n$$

$$(x^2 + 1)y'' + 2xy' - 2y = 0$$

$$x^2 \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} + \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} + 2x \sum_{n=1}^{\infty} n a_n x^{n-1} - 2 \sum_{n=0}^{\infty} a_n x^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n x^n + \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} + \sum_{n=1}^{\infty} 2n a_n x^n - \sum_{n=0}^{\infty} 2a_n x^n = 0$$

$$\sum_{n=0}^{\infty} n(n-1) a_n x^n + \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} x^n + \sum_{n=0}^{\infty} 2n a_n x^n - \sum_{n=0}^{\infty} 2a_n x^n = 0$$

$$\sum_{n=0}^{\infty} \left[ (n+1)(n+2)a_{n+2} + (n^2 + n - 2)a_n \right] x^n = 0$$

$$(n+1)(n+2)a_{n+2} + (n-1)(n+2)a_n = 0$$

$$\underline{a_{n+2} = -\frac{n-1}{n+1}a_n}$$

$$a_0 = y(0) = 0$$

$$a_1 = y'(0) = 1$$

$$n=0 \rightarrow a_2 = a_0$$

$$n=1 \rightarrow a_3 = 0$$

$$n=2 \rightarrow a_4 = -\frac{1}{3}a_2 = -\frac{1}{3}a_0$$

$$n=3 \rightarrow a_5 = 0$$

$$n=4 \rightarrow a_6 = -\frac{3}{5}a_4 = \frac{1}{5}a_0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$n=6 \rightarrow a_8 = -\frac{5}{7}a_6 = -\frac{1}{7}a_0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$a_{2n} = \frac{(-1)^n}{2n-1}a_0$$

$$\underline{y(x) = a_1x + a_0 \left( 1 + x^2 - \frac{1}{3}x^4 + \frac{1}{5}x^6 - \frac{1}{7}x^8 + \dots \right)}$$

$$\begin{aligned} y(x) &= a_1x + a_0 \left( 1 + x \left( x - \frac{1}{3}x^3 + \frac{1}{5}x^5 - \frac{1}{7}x^7 + \dots \right) \right) \\ &= \underline{a_1x + a_0 \left( 1 + x \tan^{-1}x \right)} \end{aligned}$$

### Exercise

Find a power series solution.  $(2x - x^2)y'' - 6(x-1)y' - 4y = 0$ ;  $y(1)=0, \quad y'(1)=1$

### Solution

$$\text{Let } z = x - 1 \Rightarrow \begin{cases} x = z + 1 \\ dz = dx \end{cases}$$

$$(2x - x^2)y'' - 6(x-1)y' - 4y = 0$$

$$(2z + 2 - z^2 - 2z - 1)y'' - 6zy' - 4y = 0$$

$$(1 - z^2)y'' - 6zy' - 4y = 0$$

$$y(z) = \sum_{n=0}^{\infty} a_n z^n$$

$$y'(z) = \sum_{n=1}^{\infty} n a_n z^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} z^n$$

$$y''(z) = \sum_{n=2}^{\infty} n(n-1) a_n z^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} z^n$$

$$(1 - z^2)y'' - 6zy' - 4y = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n z^{n-2} - z^2 \sum_{n=2}^{\infty} n(n-1) a_n z^{n-2} - 6z \sum_{n=1}^{\infty} n a_n z^{n-1} - 4 \sum_{n=0}^{\infty} a_n z^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n z^{n-2} - \sum_{n=2}^{\infty} n(n-1) a_n z^n - \sum_{n=1}^{\infty} 6n a_n z^n - 4 \sum_{n=0}^{\infty} a_n z^n = 0$$

$$\sum_{n=0}^{\infty} (n+1)(n+2) a_{n+2} z^n - \sum_{n=0}^{\infty} n(n-1) a_n z^n - \sum_{n=0}^{\infty} 6n a_n z^n - 4 \sum_{n=0}^{\infty} a_n z^n = 0$$

$$\sum_{n=0}^{\infty} \left[ (n+1)(n+2) a_{n+2} - (n^2 + 5n + 4) a_n \right] z^n = 0$$

$$(n+1)(n+2) a_{n+2} - (n+1)(n+4) a_n = 0$$

$$\boxed{a_{n+2} = \frac{n+4}{n+2} a_n}$$

$$\text{Given: } y(1) = 0 = a_0, \quad y'(1) = 1 = a_1$$

$$a_0 = 0$$

$$n=0 \rightarrow a_2 = 2a_0 = 0$$

$$n=2 \rightarrow a_4 = \frac{3}{2} a_2 = 0$$

$$n=4 \rightarrow a_6 = 0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots$$

$$a_1 = 1$$

$$n=1 \rightarrow a_3 = \frac{5}{3} a_1 = \frac{5}{3}$$

$$n=3 \rightarrow a_5 = \frac{7}{5} a_3 = \frac{7}{3}$$

$$n=5 \rightarrow a_7 = \frac{9}{7} a_5 = \frac{9}{3}$$

$$n=7 \rightarrow a_9 = \frac{11}{9} a_7 = \frac{11}{3}$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots$$

$$a_{2n+1} = \frac{2n+3}{3}$$

$$y(z) = z + \frac{5}{3} z^3 + \frac{7}{3} z^5 + 3z^7 + \frac{11}{3} z^9 + \dots + \frac{2n+3}{3} z^{2n+1} + \dots$$

$$\boxed{y(x) = (x-1) + \frac{5}{3}(x-1)^3 + \frac{7}{3}(x-1)^5 + 3(x-1)^7 + \frac{11}{3}(x-1)^9 + \dots}$$

### Exercise

Find a power series solution.  $(x^2 - 6x + 10)y'' - 4(x - 3)y' + 6y = 0$  ;  $y(3) = 2$ ,  $y'(3) = 0$

### Solution

$$\text{Let } z = x - 3 \Rightarrow \begin{cases} x = z + 3 \\ dz = dx \end{cases}$$

$$(x^2 - 6x + 10)y'' - 4(x - 3)y' + 6y = 0$$

$$(z^2 + 6z + 9 - 6z - 18 + 10)y'' - 4zy' + 6y = 0$$

$$(z^2 + 1)y'' - 4zy' + 6y = 0$$

$$y(z) = \sum_{n=0}^{\infty} a_n z^n$$

$$y'(z) = \sum_{n=1}^{\infty} n a_n z^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} z^n$$

$$y''(z) = \sum_{n=2}^{\infty} n(n-1) a_n z^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} z^n$$

$$(z^2 + 1)y'' - 4zy' + 6y = 0$$

$$z^2 \sum_{n=2}^{\infty} n(n-1) a_n z^{n-2} + \sum_{n=2}^{\infty} n(n-1) a_n z^{n-2} - 4z \sum_{n=1}^{\infty} n a_n z^{n-1} + 6 \sum_{n=0}^{\infty} a_n z^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1) a_n z^n + \sum_{n=2}^{\infty} n(n-1) a_n z^{n-2} - \sum_{n=1}^{\infty} 4n a_n z^n + \sum_{n=0}^{\infty} 6a_n z^n = 0$$

$$\sum_{n=0}^{\infty} n(n-1) a_n z^n + \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} z^n - \sum_{n=0}^{\infty} 4n a_n z^n + \sum_{n=0}^{\infty} 6a_n z^n = 0$$

$$\sum_{n=0}^{\infty} \left[ (n+1)(n+2) a_{n+2} + (n^2 - 5n + 6) a_n \right] z^n = 0$$

$$(n+1)(n+2) a_{n+2} + (n-2)(n-3) a_n = 0$$

$$a_{n+2} = -\frac{(n-2)(n-3)}{(n+1)(n+2)} a_n$$

$$\text{Given: } y(3) = 2 = a_0, \quad y'(3) = 0 = a_1$$

$$\begin{aligned}
 a_0 &= 2 \\
 n=0 &\rightarrow a_2 = -3a_0 = -6 \\
 n=2 &\rightarrow a_4 = 0 \\
 n=4 &\rightarrow a_6 = 0 \\
 &\vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots
 \end{aligned}$$

$$\begin{aligned}
 a_1 &= 0 \\
 n=1 &\rightarrow a_3 = -\frac{2}{4}a_1 = 0 \\
 n=3 &\rightarrow a_5 = 0 \\
 n=5 &\rightarrow a_7 = 0 \\
 &\vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots
 \end{aligned}$$

$$y(z) = 2 - 6z^2$$

$$\underline{y(x) = 2 - 6(x-3)^2}$$

### Exercise

Find a power series solution.  $(4x^2 + 16x + 17)y'' - 8y = 0$ ;  $y(-2) = 1$ ,  $y'(-2) = 0$

### Solution

$$\text{Let } z = x + 2 \Rightarrow \begin{cases} x = z - 2 \\ dz = dx \end{cases}$$

$$(4z^2 - 16z + 16 + 16z - 32 + 17)y'' - 8y = 0$$

$$\underline{(4z^2 + 1)y'' - 8y = 0}$$

$$y(z) = \sum_{n=0}^{\infty} a_n z^n$$

$$y'(z) = \sum_{n=1}^{\infty} n a_n z^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} z^n$$

$$y''(z) = \sum_{n=2}^{\infty} n(n-1) a_n z^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} z^n$$

$$(4z^2 + 1)y'' - 8y = 0$$

$$4z^2 \sum_{n=2}^{\infty} n(n-1) a_n z^{n-2} + \sum_{n=2}^{\infty} n(n-1) a_n z^{n-2} - 8 \sum_{n=0}^{\infty} a_n z^n = 0$$

$$\sum_{n=2}^{\infty} 4n(n-1) a_n z^n + \sum_{n=2}^{\infty} n(n-1) a_n z^{n-2} - 8 \sum_{n=0}^{\infty} a_n z^n = 0$$

$$\sum_{n=0}^{\infty} 4n(n-1) a_n z^n + \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} z^n - 8 \sum_{n=0}^{\infty} a_n z^n = 0$$

$$\sum_{n=0}^{\infty} \left[ (n+1)(n+2)a_{n+2} + (4n^2 - 4n - 8)a_n \right] z^n = 0$$

$$(n+1)(n+2)a_{n+2} + 4(n+1)(n-2)a_n = 0$$

$$\underline{a_{n+2} = -\frac{4(n-2)}{n+2}a_n}$$

$$\text{Given: } y(-2) = 1 = a_0, \quad y'(-2) = 0 = a_1$$

$$a_0 = 1$$

$$a_1 = 0$$

$$n=0 \rightarrow a_2 = \frac{8}{2}a_0 = 4$$

$$n=1 \rightarrow a_3 = \frac{4}{3}a_1 = 0$$

$$n=2 \rightarrow a_4 = -0a_2 = 0$$

$$n=3 \rightarrow a_5 = 0$$

$$n=4 \rightarrow a_6 = 0$$

$$n=5 \rightarrow a_7 = 0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots$$

$$y(z) = 1 + 4z^2$$

$$\underline{y(x) = 1 + 4(x+2)^2}$$

### Exercise

Find a power series solution.  $(x^2 + 6x)y'' + (3x + 9)y' - 3y = 0$ ;  $y(-3) = 0, \quad y'(-3) = 2$

### Solution

$$\text{Let } z = x + 3 \Rightarrow \begin{cases} x = z - 3 \\ dz = dx \end{cases}$$

$$(x^2 + 6x)y'' + (3x + 9)y' - 3y = 0$$

$$(z^2 - 6z + 9 + 6z - 18)y'' + 3zy' - 3y = 0$$

$$\underline{(z^2 - 9)y'' + 3zy' - 3y = 0}$$

$$y(z) = \sum_{n=0}^{\infty} a_n z^n$$

$$y'(z) = \sum_{n=1}^{\infty} n a_n z^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} z^n$$

$$y''(z) = \sum_{n=2}^{\infty} n(n-1) a_n z^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} z^n$$

$$(z^2 - 9)y'' + 3zy' - 3y = 0$$

$$z^2 \sum_{n=2}^{\infty} n(n-1)a_n z^{n-2} - 9 \sum_{n=2}^{\infty} n(n-1)a_n z^{n-2} + 3z \sum_{n=1}^{\infty} na_n z^{n-1} - 3 \sum_{n=0}^{\infty} a_n z^n = 0$$

$$\sum_{n=2}^{\infty} n(n-1)a_n z^n - \sum_{n=2}^{\infty} 9n(n-1)a_n z^{n-2} + \sum_{n=1}^{\infty} 3na_n z^n - \sum_{n=0}^{\infty} 3a_n z^n = 0$$

$$\sum_{n=0}^{\infty} n(n-1)a_n z^n - \sum_{n=0}^{\infty} 9(n+2)(n+1)a_{n+2} z^n + \sum_{n=0}^{\infty} 3na_n z^n - \sum_{n=0}^{\infty} 3a_n z^n = 0$$

$$\sum_{n=0}^{\infty} \left[ -9(n+2)(n+1)a_{n+2} + (n^2 - n + 3n - 3)a_n \right] z^n = 0$$

$$-9(n+2)(n+1)a_{n+2} + (n+3)(n-1)a_n = 0$$

$$a_{n+2} = \frac{(n+3)(n-1)}{9(n+1)(n+2)} a_n$$

$$\text{Given: } y(-3) = 0 = a_0, \quad y'(-3) = 2 = a_1$$

$$a_0 = 0$$

$$n=0 \rightarrow a_2 = \frac{-3}{18} a_0 = 0$$

$$n=2 \rightarrow a_4 = \frac{5}{108} a_2 = 0$$

$$n=4 \rightarrow a_6 = 0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots$$

$$a_1 = 2$$

$$n=1 \rightarrow a_3 = 0a_1 = 0$$

$$n=3 \rightarrow a_5 = 0$$

$$n=5 \rightarrow a_7 = 0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots$$

$$y(z) = 2z$$

$$y(x) = 2x + 6$$

## Exercise

Find the series solution near the given value  $y'' - (x-2)y' + 2y = 0$  ; near  $x = 2$

### Solution

$$y = \sum_{n=0}^{\infty} a_n (x-2)^n$$

$$y' = \sum_{n=1}^{\infty} na_n (x-2)^{n-1} = \sum_{n=0}^{\infty} (n+1)a_{n+1} (x-2)^n$$



$$y'' = \sum_{n=2}^{\infty} n(n-1)a_n (x-2)^{n-2} = \sum_{n=0}^{\infty} (n+1)(n+2)a_{n+2} (x-2)^n$$

$$y'' - (x-2)y' + 2y = 0$$

$$\sum_{n=0}^{\infty} (n+1)(n+2)a_{n+2} (x-2)^n - (x-2) \sum_{n=0}^{\infty} (n+1)a_{n+1} (x-2)^n + 2 \sum_{n=0}^{\infty} a_n (x-2)^n = 0$$

$$\sum_{n=0}^{\infty} (n+1)(n+2)a_{n+2} (x-2)^n - \sum_{n=0}^{\infty} (n+1)a_{n+1} (x-2)^{n+1} + \sum_{n=0}^{\infty} 2a_n (x-2)^n = 0$$

$$\sum_{n=0}^{\infty} [(n+1)(n+2)a_{n+2} + 2a_n] (x-2)^n - \sum_{n=0}^{\infty} (n+1)a_{n+1} (x-2)^{n+1} = 0$$

$$\sum_{n=0}^{\infty} [(n+1)(n+2)a_{n+2} + 2a_n] (x-2)^n - \sum_{n=1}^{\infty} na_n (x-2)^n = 0$$

$$2a_2 + 2a_0 + \sum_{n=1}^{\infty} [(n+1)(n+2)a_{n+2} + 2a_n] (x-2)^n - \sum_{n=1}^{\infty} na_n (x-2)^n = 0$$

$$2a_2 + 2a_0 + \sum_{n=1}^{\infty} [(n+1)(n+2)a_{n+2} - (n-2)a_n] (x-2)^n = 0$$

$$\text{For } n=0 \rightarrow 2a_2 + 2a_0 = 0 \Rightarrow \underline{a_2 = -a_0}$$

$$(n+1)(n+2)a_{n+2} - (n-2)a_n = 0$$

$$\underline{a_{n+2} = \frac{n-2}{(n+1)(n+2)} a_n}$$

$$a_0$$

$$n=0 \rightarrow a_2 = -a_0$$

$$n=2 \rightarrow a_4 = 0$$

$$n=4 \rightarrow a_6 = 0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$a_1$$

$$n=1 \rightarrow a_3 = -\frac{1}{6}a_1$$

$$n=3 \rightarrow a_5 = \frac{1}{20}a_3 = -\frac{1}{120}a_1$$

$$n=5 \rightarrow a_7 = \frac{3}{42}a_5 = -\frac{1}{1680}a_1$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\underline{y(x) = a_0 \left(1 - (x-2)^2\right) + a_1 \left((x-2) - \frac{1}{6}(x-2)^3 - \frac{1}{120}(x-2)^5 - \frac{1}{1680}(x-2)^7 - \dots\right)}$$

### Exercise

Find the series solution near the given value  $y'' + (x-1)^2 y' - 4(x-1)y = 0$  ; *near  $x=1$*

### Solution

$$y = \sum_{n=0}^{\infty} a_n (x-1)^n$$

$$y' = \sum_{n=1}^{\infty} n a_n (x-1)^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} (x-1)^n$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n (x-1)^{n-2} = \sum_{n=0}^{\infty} (n+1)(n+2) a_{n+2} (x-1)^n$$

$$y'' + (x-1)^2 y' - 4(x-1)y = 0$$

$$\sum_{n=0}^{\infty} (n+1)(n+2) a_{n+2} (x-1)^n + (x-1)^2 \sum_{n=0}^{\infty} (n+1) a_{n+1} (x-1)^n - 4(x-1) \sum_{n=0}^{\infty} a_n (x-1)^n = 0$$

$$\sum_{n=0}^{\infty} (n+1)(n+2) a_{n+2} (x-1)^n + \sum_{n=0}^{\infty} (n+1) a_{n+1} (x-1)^{n+2} - \sum_{n=0}^{\infty} 4a_n (x-1)^{n+1} = 0$$

$$\sum_{n=0}^{\infty} (n+1)(n+2) a_{n+2} (x-1)^n + \sum_{n=2}^{\infty} (n-1) a_{n-1} (x-1)^n - \sum_{n=1}^{\infty} 4a_{n-1} (x-1)^n = 0$$

$$2a_2 + 6a_3(x-1) + \sum_{n=2}^{\infty} (n+1)(n+2) a_{n+2} (x-1)^n + \sum_{n=2}^{\infty} (n-1) a_{n-1} (x-1)^n$$

$$- 4a_0(x-1) - \sum_{n=2}^{\infty} 4a_{n-1} (x-1)^n = 0$$

$$2a_2 + (6a_3 - 4a_0)(x-1) + \sum_{n=2}^{\infty} [(n+1)(n+2) a_{n+2} + (n-5) a_{n-1}] (x-1)^n = 0$$

$$\begin{cases} 2a_2 = 0 & \rightarrow \underline{a_2 = 0} \\ 6a_3 - 4a_0 = 0 & \rightarrow \underline{a_3 = \frac{2}{3}a_0} \end{cases}$$

$$(n+1)(n+2) a_{n+2} + (n-5) a_{n-1} = 0$$

$$\underline{a_{n+2} = -\frac{(n-5)}{(n+1)(n+2)} a_{n-1}}$$

$a_0$	$a_1$	$a_2 = 0$
$n = 1 \rightarrow a_3 = \frac{2}{3}a_0$	$n = 2 \rightarrow a_4 = \frac{1}{4}a_1$	$n = 3 \rightarrow a_5 = \frac{2}{20}a_2 = 0$
$n = 4 \rightarrow a_6 = \frac{1}{30}a_3 = \frac{1}{45}a_0$	$n = 5 \rightarrow a_7 = 0$	$n = 6 \rightarrow a_8 = 0$
$n = 7 \rightarrow a_9 = -\frac{2}{8 \cdot 9}a_6 = -\frac{1}{1,620}a_0$	$n = 8 \rightarrow a_{10} = 0$	$n = 9 \rightarrow a_{11} = 0$
$\vdots \quad \vdots \quad \vdots \quad \vdots$	$\vdots \quad \vdots \quad \vdots \quad \vdots$	$\vdots \quad \vdots \quad \vdots \quad \vdots$

---


$$y(x) = a_0 \left( 1 + \frac{2}{3}(x-1)^3 + \frac{1}{45}(x-1)^6 - \frac{1}{1,620}(x-1)^9 + \dots \right) + a_1 \left( (x-1) + \frac{1}{4}(x-1)^4 \right)$$


---


$$y(x) = a_0 \sum_{n=0}^{\infty} \frac{(-1)^n 4(x-1)^{3n}}{3^n (3n-1)(3n-4)n!} + a_1 \left( (x-1) + \frac{1}{4}(x-1)^4 \right)$$


---

### Exercise

Find the series solution near the given value  $y'' + (x-1)y = e^x$  ; near  $x = 1$

### Solution

$$y = \sum_{n=0}^{\infty} a_n (x-1)^n$$

$$y' = \sum_{n=1}^{\infty} n a_n (x-1)^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} (x-1)^n$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n (x-1)^{n-2} = \sum_{n=0}^{\infty} (n+1)(n+2) a_{n+2} (x-1)^n$$

$$\textcolor{red}{y'' + (x-1)y = e^{x-1+1}}$$

$$\sum_{n=0}^{\infty} (n+1)(n+2) a_{n+2} (x-1)^n + (x-1) \sum_{n=0}^{\infty} a_n (x-1)^n = e \cdot e^{x-1}$$

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$$

$$\sum_{n=0}^{\infty} (n+1)(n+2) a_{n+2} (x-1)^n + \sum_{n=0}^{\infty} a_n (x-1)^{n+1} = e \cdot \sum_{n=0}^{\infty} \frac{(x-1)^n}{n!}$$

$$\sum_{n=0}^{\infty} (n+1)(n+2) a_{n+2} (x-1)^n + \sum_{\textcolor{red}{n}=1}^{\infty} a_{n-1} (x-1)^n = e \cdot \sum_{n=0}^{\infty} \frac{(x-1)^n}{n!}$$

$$2a_2 + \sum_{n=1}^{\infty} (n+1)(n+2)a_{n+2} (x-1)^n + \sum_{n=1}^{\infty} a_{n-1} (x-1)^n = e + e \cdot \sum_{n=1}^{\infty} \frac{(x-1)^n}{n!}$$

$$2a_2 + \sum_{n=1}^{\infty} [(n+1)(n+2)a_{n+2} + a_{n-1}] (x-1)^n = e + e \cdot \sum_{n=1}^{\infty} \frac{(x-1)^n}{n!}$$

$$2a_2 = e \rightarrow a_2 = \frac{e}{2}$$

$$(n+1)(n+2)a_{n+2} + a_{n-1} = \frac{e}{n!}$$

$$a_{n+2} = \frac{e}{(n+1)(n+2)n!} - \frac{1}{(n+1)(n+2)} a_{n-1}$$

$a_0$

$a_1$

$$n=1 \rightarrow a_3 = \frac{e}{6} - \frac{1}{6}a_0$$

$$n=2 \rightarrow a_4 = \frac{e}{24} - \frac{1}{12}a_1$$

$$n=4 \rightarrow a_6 = \frac{e}{720} - \frac{1}{30}a_3 = -\frac{11e}{720} + \frac{1}{180}a_0 \quad n=5 \rightarrow a_7 = \frac{e}{5040} - \frac{1}{42}a_4 = \frac{e}{1260} + \frac{1}{504}a_0$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots$$

$$a_2 = \frac{e}{2}$$

$$n=3 \rightarrow a_5 = \frac{e}{120} - \frac{1}{20}a_2 = \frac{e}{120} - \frac{e}{40} = -\frac{e}{60}$$

$$n=6 \rightarrow a_8 = \frac{e}{40,320} - \frac{1}{56}a_5 = \frac{e}{40,320} + \frac{e}{3360} = \frac{13e}{40,320}$$

$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$\begin{aligned} y(x) &= a_0 + (x-1)a_1 + \frac{e}{2}(x-1)^2 + \left(\frac{e}{6} - \frac{1}{6}a_0\right)(x-1)^3 + \left(\frac{e}{24} - \frac{1}{12}a_1\right)(x-1)^4 - \frac{e}{60}(x-1)^5 + \dots \\ &= a_0 + (x-1)a_1 + \frac{e}{2}(x-1)^2 + \frac{e}{6}(x-1)^3 + -\frac{1}{6}a_0(x-1)^3 + \frac{e}{24}(x-1)^4 \\ &\quad - \frac{1}{12}a_1(x-1)^4 - \frac{e}{60}(x-1)^5 + \dots \end{aligned}$$

$$\begin{aligned} y(x) &= e \left( \frac{1}{2}(x-1)^2 + \frac{1}{6}(x-1)^3 + \frac{1}{24}(x-1)^4 - \frac{1}{60}(x-1)^5 + \dots \right) \\ &\quad + a_0 \left( 1 - \frac{1}{6}(x-1)^3 + \dots \right) + a_1 \left( (x-1) - \frac{1}{12}(x-1)^4 + \dots \right) \end{aligned}$$

## Exercise

Find the series solution near the given value

$$y'' + xy' + (2x-1)y = 0 ; \quad \text{near } x = -1 \quad y(-1) = 2, \quad y'(-1) = -2$$

## Solution

$$t = x+1 \rightarrow x = t-1$$

$$y = \sum_{n=0}^{\infty} a_n t^n$$

$$y' = \sum_{n=1}^{\infty} n a_n t^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} t^n$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n t^{n-2} = \sum_{n=0}^{\infty} (n+1)(n+2) a_{n+2} t^n$$

$$y'' + xy' + (2x-1)y = 0$$

$$y'' + (t-1)y' + (2t-3)y = 0$$

$$\sum_{n=0}^{\infty} (n+1)(n+2) a_{n+2} t^n + (t-1) \sum_{n=0}^{\infty} (n+1) a_{n+1} t^n + (2t-3) \sum_{n=0}^{\infty} a_n t^n = 0$$

$$\sum_{n=0}^{\infty} (n+1)(n+2) a_{n+2} t^n + \sum_{n=0}^{\infty} (n+1) a_{n+1} t^{n+1} - \sum_{n=0}^{\infty} (n+1) a_{n+1} t^n + \sum_{n=0}^{\infty} 2a_n t^{n+1} - \sum_{n=0}^{\infty} 3a_n t^n = 0$$

$$\sum_{n=0}^{\infty} [(n+1)(n+2) a_{n+2} - (n+1) a_{n+1} - 3a_n] t^n + \sum_{n=0}^{\infty} [(n+1) a_{n+1} + 2a_n] t^{n+1} = 0$$

$$\sum_{n=0}^{\infty} [(n+1)(n+2) a_{n+2} - (n+1) a_{n+1} - 3a_n] t^n + \sum_{n=1}^{\infty} [n a_n + 2a_{n-1}] t^n = 0$$

$$2a_2 - a_1 - 3a_0 + \sum_{n=1}^{\infty} [(n+1)(n+2) a_{n+2} - (n+1) a_{n+1} - 3a_n] t^n + \sum_{n=1}^{\infty} [n a_n + 2a_{n-1}] t^n = 0$$

$$2a_2 - a_1 - 3a_0 + \sum_{n=1}^{\infty} [(n+1)(n+2) a_{n+2} - (n+1) a_{n+1} + (n-3) a_n + 2a_{n-1}] t^n = 0$$

$$2a_2 - a_1 - 3a_0 = 0 \rightarrow \underline{a_2 = \frac{1}{2}(a_1 + 3a_0)}$$

$$(n+1)(n+2) a_{n+2} - (n+1) a_{n+1} + (n-3) a_n + 2a_{n-1} = 0$$

$$\underline{a_{n+2} = \frac{1}{n+2} a_{n+1} - \frac{n-3}{(n+1)(n+2)} a_n - \frac{2}{(n+1)(n+2)} a_{n-1}}$$

**Given:**  $t = x+1$

$$y(x=-1) = y(t=0) = 2 = a_0, \quad y'(x=-1) = y'(t=0) = -2 = a_1$$

$$\underline{a_2} = \frac{1}{2}(a_1 + 3a_0) = \frac{1}{2}(-2 + 6) = \underline{2}$$

$$n=1 \rightarrow a_3 = \frac{1}{3}a_2 + \frac{1}{3}a_1 - \frac{1}{3}a_0 = \frac{2}{3} - \frac{2}{3} - \frac{2}{3} = -\frac{2}{3}$$

$$n=2 \rightarrow a_4 = \frac{1}{4}a_3 + \frac{1}{12}a_2 - \frac{1}{6}a_1 = -\frac{1}{6} + \frac{1}{6} + \frac{1}{3} = \frac{1}{3}$$


$$\vdots \quad \vdots \quad \vdots \quad \vdots$$

$$y(t) = 2 - 2t + 3t^2 - \frac{1}{3}t^3 + \frac{1}{3}t^4 + \dots$$

$$y(x) = 2 - 2(x+1) + 3(x+1)^2 - \frac{1}{3}(x+1)^3 + \frac{1}{3}(x+1)^4 + \dots$$

### Exercise

As a spring is heated, its spring “constant” decreases. Suppose the spring is heated so that the spring “constant” at time  $t$  is  $k(t) = 6 - t$  N/m.



If the unforced mass-spring system has mass  $m = 2$  kg and a damping constant  $b = 1$  N-sec/m with initial conditions  $x(0) = 3$  m and  $x'(0) = 0$  m/sec, then the displacement  $x(t)$  is governed by the initial value problem

$$2x''(t) + x'(t) + (6 - t)x(t) = 0 ; \quad x(0) = 3, \quad x'(0) = 0$$

Find at least the first four nonzero terms in a power series expansion about  $t = 0$  for the displacement.

### Solution

$$x(t) = \sum_{n=0}^{\infty} a_n t^n$$

$$x'(t) = \sum_{n=1}^{\infty} n a_n t^{n-1} = \sum_{n=0}^{\infty} (n+1) a_{n+1} t^n$$

$$x''(t) = \sum_{n=2}^{\infty} n(n-1) a_n t^{n-2} = \sum_{n=0}^{\infty} (n+2)(n+1) a_{n+2} t^n$$

$$2x'' + x' + (6 - t)x = 0$$

$$2 \sum_{n=0}^{\infty} (n+2)(n+1)a_{n+2}t^n + \sum_{n=0}^{\infty} (n+1)a_{n+1}t^n + (6-t) \sum_{n=0}^{\infty} a_n t^n = 0$$

$$\sum_{n=0}^{\infty} 2(n+2)(n+1)a_{n+2}t^n + \sum_{n=0}^{\infty} (n+1)a_{n+1}t^n + \sum_{n=0}^{\infty} 6a_n t^n - \sum_{n=0}^{\infty} a_n t^{n+1} = 0$$

$$\sum_{n=0}^{\infty} \left[ 2(n+2)(n+1)a_{n+2} + (n+1)a_{n+1} + 6a_n \right] t^n - \sum_{n=1}^{\infty} a_{n-1} t^n = 0$$

$$4a_2 + a_1 + 6a_0 + \sum_{n=1}^{\infty} \left[ 2(n+2)(n+1)a_{n+2} + (n+1)a_{n+1} + 6a_n \right] t^n - \sum_{n=1}^{\infty} a_{n-1} t^n = 0$$

$$4a_2 + a_1 + 6a_0 + \sum_{n=1}^{\infty} \left[ 2(n+1)(n+2)a_{n+2} + (n+1)a_{n+1} + 6a_n - a_{n-1} \right] t^n = 0$$

**Given:**  $x(0) = 3 = a_0$ ,  $x'(0) = 0 = a_1$

$$4a_2 + a_1 + 6a_0 = 0 \rightarrow \underline{a_2 = -\frac{9}{2}}$$

$$2(n+1)(n+2)a_{n+2} + (n+1)a_{n+1} + 6a_n - a_{n-1} = 0$$

$$\underline{a_{n+2} = \frac{a_{n-1} - 6a_n - (n+1)a_{n+1}}{2(n+1)(n+2)}}$$

$$n=1 \rightarrow a_3 = \frac{1}{12}(a_0 - 6a_1 - 2a_2) = \frac{1}{12}(3+9) = 1$$

$$n=2 \rightarrow a_4 = \frac{1}{24}(a_1 - 6a_2 - 3a_3) = \frac{1}{24}(27-3) = 1$$

$$\underline{x(t) = 3 - \frac{9}{2}t^2 + t^3 + t^4 + \dots}$$