

## ***Solution***      **Section 2.1 – Introducing the Derivative**

### ***Exercise***

Use the definition of the derivative to determine the slope of the curve  $y = f(x)$ . Find an equation of the line tangent to the curve  $y = f(x)$  at  $P$ ; then graph the curve and the tangent line.

$$y = 4 - x^2; \quad P(-1, 3)$$

### **Solution**


$$\begin{aligned} m &= \lim_{h \rightarrow 0} \frac{4 - (x+h)^2 - (4 - x^2)}{h} \\ &= \lim_{h \rightarrow 0} \frac{4 - (-1+h)^2 - (4 - (-1)^2)}{h} \\ &= \lim_{h \rightarrow 0} \frac{4 - (1 - 2h + h^2) - (4 - 1)}{h} \\ &= \lim_{h \rightarrow 0} \frac{4 - 1 + 2h - h^2 - 3}{h} \\ &= \lim_{h \rightarrow 0} \frac{2h - h^2}{h} \\ &= \lim_{h \rightarrow 0} (2 - h) \\ &= 2 \end{aligned}$$

$$y - y_1 = m(x - x_1)$$

$$\text{At } (-1, 3) \Rightarrow y - 3 = 2(x - (-1))$$

$$y - 3 = 2x + 2$$

$$\underline{y = 2x + 5}$$



### Exercise

Use the definition of the derivative to determine the slope of the curve  $y = f(x)$ . Find an equation of the line tangent to the curve  $y = f(x)$  at  $P$ ; then graph the curve and the tangent line.

$$y = \frac{1}{x^2}; \quad P(-1, 1)$$


### Solution

$$\begin{aligned} m &= \lim_{h \rightarrow 0} \frac{\frac{1}{(x+h)^2} - \frac{1}{x^2}}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{1}{(-1+h)^2} - \frac{1}{1} \right) \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{1 - (1 - 2h + h^2)}{(-1+h)^2} \right) \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{1 - 1 + 2h - h^2}{(-1+h)^2} \right) \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{2h - h^2}{(-1+h)^2} \right) \\ &= \lim_{h \rightarrow 0} \frac{h}{h} \left( \frac{2 - h}{(-1+h)^2} \right) \\ &= \lim_{h \rightarrow 0} \left( \frac{2 - h}{(-1+h)^2} \right) \\ &= \frac{2 + 0}{(-1 + 0)^2} \\ &= 2 \end{aligned}$$

$$\text{At } (-1, 3) \Rightarrow y - 1 = 2(x - (-1))$$

$$y - 1 = 2x + 2$$

$$\underline{y = 2x + 3}$$



### Exercise

Use the definition of the derivative to determine the slope of the curve  $y = f(x)$ . Find an equation of the line tangent to the curve  $y = f(x)$  at  $P$ ; then graph the curve and the tangent line.

$$f(x) = 2\sqrt{x}; \quad P(1, 2)$$

### Solution


$$\begin{aligned} m &= \lim_{h \rightarrow 0} \frac{2\sqrt{x+h} - 2\sqrt{x}}{h} \\ &= \lim_{h \rightarrow 0} \frac{2\sqrt{1+h} - 2\sqrt{x}}{h} \cdot \frac{2\sqrt{1+h} + 2}{2\sqrt{1+h} + 2} \\ &= \lim_{h \rightarrow 0} \frac{4(1+h) - 4}{h(2\sqrt{1+h} + 2)} \\ &= \lim_{h \rightarrow 0} \frac{4 + 4h - 4}{h(2\sqrt{1+h} + 2)} \\ &= \lim_{h \rightarrow 0} \frac{4h}{h(2\sqrt{1+h} + 2)} \\ &= \lim_{h \rightarrow 0} \frac{4}{2\sqrt{1+h} + 2} \\ &= \frac{4}{2+2} \\ &= 1 \end{aligned}$$

$$\text{At } (1, 2) \Rightarrow y - 2 = (x - 1)$$

$$y - y_1 = m(x - x_1)$$

$$y - 2 = x - 1$$

$$\underline{y = x + 1}$$



### Exercise

Use the definition of the derivative to determine the slope of the curve  $y = f(x)$ . Find an equation of the line tangent to the curve  $y = f(x)$  at  $P$ ; then graph the curve and the tangent line.

$$f(x) = x^3 + 3x; \quad P(1, 4)$$

### Solution

$$\begin{aligned} m &= \lim_{h \rightarrow 0} \frac{(x+h)^3 + 3(x+h) - (x^3 + 3x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{1 + 3h + 3h^2 + h^3 + 3 + 3h - (4)}{h} \\ &= \lim_{h \rightarrow 0} \frac{3h + 3h^2 + h^3 + 3h}{h} \\ &= \lim_{h \rightarrow 0} \frac{3h^2 + h^3 + 6h}{h} \\ &= 6 \end{aligned}$$


At  $(1, 4)$

$$y - 4 = 6(x - 1)$$

$$y - y_1 = m(x - x_1)$$

$$y - 4 = 6x - 6$$

$$y = 6x - 2$$



### Exercise

Use the definition of the derivative to determine the slope of the curve  $y = f(x)$ . Find an equation of the line tangent to the curve  $y = f(x)$  at  $P$ ; then graph the curve and the tangent line.

$$f(x) = 4x^2 - 7x + 5; \quad P(2, 7)$$

### Solution


$$\begin{aligned} m &= \lim_{h \rightarrow 0} \frac{4(x+h)^2 - 7(x+h) + 5 - 4x^2 + 7x - 5}{h} \\ &= \lim_{h \rightarrow 0} \frac{4(x^2 + 2xh + h^2) - 7x - 7h - 4x^2 + 7x}{h} \\ &= \lim_{h \rightarrow 0} \frac{4x^2 + 8xh + 4h^2 - 7h - 4x^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{8xh + 4h^2 - 7h}{h} \\ &= \lim_{h \rightarrow 0} (8xh + 4h - 7) \end{aligned}$$

$$= 8x - 7]$$

$$\text{At } (2, 7) \rightarrow m = 9]$$

$$y = 9(x - 2) + 7 \quad y = m(x - x_1) + y_1$$

$$= 9x - 11]$$



### Exercise

Use the definition of the derivative to determine the slope of the curve  $y = f(x)$ . Find an equation of the line tangent to the curve  $y = f(x)$  at  $P$ ; then graph the curve and the tangent line.


$$f(x) = 5x^3 + x; \quad P(1, 6)$$

### Solution

$$\begin{aligned} m &= \lim_{h \rightarrow 0} \frac{5(x+h)^3 + (x+h) - 5x^3 - x}{h} \\ &= \lim_{h \rightarrow 0} \frac{5(x^3 + 3x^2h + 3xh^2 + h^3) + h - 5x^3}{h} \\ &= \lim_{h \rightarrow 0} \frac{15x^2h + 15xh^2 + 5h^3 + h}{h} \\ &= \lim_{h \rightarrow 0} (15x^2 + 15xh + 5h^2 + 1) \\ &= 15x^2 + 1 \Big|_{(1, 6)} \\ &= 16] \end{aligned}$$

$$y = 16(x - 1) + 6 \quad y = m(x - x_1) + y_1$$

$$= 16x - 10]$$



## Exercise

Use the definition of the derivative to determine the slope of the curve  $y = f(x)$ . Find an equation of the line tangent to the curve  $y = f(x)$  at  $P$ ; then graph the curve and the tangent line.


$$f(x) = \frac{x+3}{2x+1}; \quad P(0, 3)$$

## Solution

$$\begin{aligned} m &= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{x+h+3}{2x+2h+1} - \frac{x+3}{2x+1} \right] \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{2x^2 + 2hx + 6x + x + h + 3 - 2x^2 - 2hx - x - 6x - 6h - 3}{(2x+2h+1)(2x+1)} \right) \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{-5h}{(2x+2h+1)(2x+1)} \right) \\ &= \lim_{h \rightarrow 0} \left( \frac{-5}{(2x+2h+1)(2x+1)} \right) \\ &= \frac{-5}{(2x+1)^2} \Big|_{(0, 3)} \\ &= -5 \end{aligned}$$

$$y = -5x + 3$$

$$y = m(x - x_1) + y_1$$



## Exercise

Use the definition of the derivative to determine the slope of the curve  $y = f(x)$ . Find an equation of the line tangent to the curve  $y = f(x)$  at  $P$ ; then graph the curve and the tangent line.

$$f(x) = \frac{1}{2\sqrt{3x+1}}; \quad P\left(0, \frac{1}{2}\right)$$


## Solution

$$\begin{aligned} m &= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{1}{2\sqrt{3x+3h+1}} - \frac{1}{2\sqrt{3x+1}} \right] \\ &= \frac{1}{2} \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{\sqrt{3x+1} - \sqrt{3x+3h+1}}{\sqrt{3x+3h+1} \sqrt{3x+1}} \right) = \frac{0}{0} \\ &= \frac{1}{2} \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{\sqrt{3x+1} - \sqrt{3x+3h+1}}{\sqrt{3x+3h+1} \sqrt{3x+1}} \right) \frac{\sqrt{3x+1} + \sqrt{3x+3h+1}}{\sqrt{3x+1} + \sqrt{3x+3h+1}} \\ &= \frac{1}{2} \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{3x+1-3x-3h-1}{\sqrt{3x+3h+1} \sqrt{3x+1} (\sqrt{3x+1} + \sqrt{3x+3h+1})} \right) \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2} \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{-3h}{\sqrt{3x+3h+1} \sqrt{3x+1} (\sqrt{3x+1} + \sqrt{3x+3h+1})} \right) \\
&= -\frac{3}{2} \lim_{h \rightarrow 0} \left( \frac{1}{\sqrt{3x+3h+1} \sqrt{3x+1} (\sqrt{3x+1} + \sqrt{3x+3h+1})} \right) \\
&= -\frac{3}{2} \frac{1}{(3x+1)(2\sqrt{3x+1})} \\
&= -\frac{3}{4} \frac{1}{(3x+1)^{3/2}} \left| \left(0, \frac{1}{2}\right) \right. \\
&= -\frac{3}{4} \left| \right.
\end{aligned}$$

$$y = -\frac{3}{4}x + \frac{1}{2} \left| \right.$$

$$y = m(x - x_1) + y_1$$



### Exercise

Find the slope of the curve  $y = 1 - x^2$  at the point  $x = 2$

### Solution

$$\begin{aligned}
m &= \lim_{h \rightarrow 0} \frac{1 - (x+h)^2 - (1 - x^2)}{h} \\
&= \lim_{h \rightarrow 0} \frac{1 - (2+h)^2 - (1 - 2^2)}{h} \\
&= \lim_{h \rightarrow 0} \frac{1 - (4 + 4h + h^2) - (-3)}{h}
\end{aligned}$$

$$\begin{aligned}
&= \lim_{h \rightarrow 0} \frac{1 - 4 - 4h - h^2 + 3}{h} \\
&= \lim_{h \rightarrow 0} \frac{-4h - h^2}{h} \\
&= \lim_{h \rightarrow 0} (-4 - h) \\
&= \underline{-4}
\end{aligned}$$

### ***Exercise***

Find the slope of the curve  $y = \frac{1}{x-1}$  at the point  $x = 3$

### **Solution**

$$\begin{aligned}
m &= \lim_{h \rightarrow 0} \frac{\frac{1}{x+h-1} - \frac{1}{x-1}}{h} \\
&= \lim_{h \rightarrow 0} \frac{\frac{1}{3+h-1} - \frac{1}{3-1}}{h} \\
&= \lim_{h \rightarrow 0} \frac{\frac{1}{2+h} - \frac{1}{2}}{h} \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{2-2-h}{2+h} \right) \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{-h}{2+h} \right) \\
&= \lim_{h \rightarrow 0} \left( \frac{-1}{2+h} \right) \\
&= \underline{-\frac{1}{2}}
\end{aligned}$$

### ***Exercise***

Find the slope of the curve  $y = \frac{x-1}{x+1}$  at the point  $x = 0$

### **Solution**

$$\begin{aligned}
m &= \lim_{h \rightarrow 0} \frac{\frac{x+h-1}{x+h+1} - \frac{x-1}{x+1}}{h} \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{0+h-1}{0+h+1} - \frac{0-1}{0+1} \right) \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{h-1}{h+1} + 1 \right)
\end{aligned}$$



$$\begin{aligned}
&= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{h-1+h+1}{h+1} \right) \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{2h}{h+1} \right) \\
&= \lim_{h \rightarrow 0} \left( \frac{2}{h+1} \right) \\
&= 2
\end{aligned}$$

### Exercise

Find equations of all lines having slope  $-1$  that are tangent to the curve  $y = \frac{1}{x-1}$

### Solution

$$\begin{aligned}
m &= \lim_{h \rightarrow 0} \frac{\frac{1}{x+h-1} - \frac{1}{x-1}}{h} \\
-1 &= \lim_{h \rightarrow 0} \frac{\frac{1}{x+h-1} - \frac{1}{x-1}}{h} \\
-1 &= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{x-1-(x+h-1)}{x+h-1} \right) \\
-1 &= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{x-1-x-h+1}{x+h-1} \right) \\
-1 &= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{-h}{x+h-1} \right) \\
-1 &= \lim_{h \rightarrow 0} \left( \frac{-1}{x+h-1} \right) \\
-1 &= \frac{-1}{x-1} \\
-x+1 &= -1 \\
x &= 2
\end{aligned}$$

*Cross multiplication*

$$y = \frac{1}{x-1} = \frac{1}{2-1} = 1$$

$$\text{At } (2, 1) \Rightarrow y-1 = -1(x-2)$$

$$y-1 = -x+2$$

$$y = -x+3$$

### Exercise

What is the rate of change of the area of a circle  $(A = \pi r^2)$  with respect to the radius when the radius is  $r = 3$ ?

### Solution

$$\begin{aligned} m &= \lim_{h \rightarrow 0} \frac{\pi(3+h)^2 - \pi(3)^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{\pi(9 + 6h + h^2) - 9\pi}{h} \\ &= \lim_{h \rightarrow 0} \frac{9\pi + 6\pi h + \pi h^2 - 9\pi}{h} \\ &= \lim_{h \rightarrow 0} \frac{6\pi h + \pi h^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{\pi h(6 + h)}{h} \\ &= \lim_{h \rightarrow 0} \pi(6 + h) \\ &= \underline{6\pi} \end{aligned}$$

### Exercise

Find the slope of the tangent to the curve  $y = \frac{1}{\sqrt{x}}$  at the point where  $x = 4$

### Solution

$$\begin{aligned} m &= \lim_{h \rightarrow 0} \frac{\frac{1}{\sqrt{x+h}} - \frac{1}{\sqrt{x}}}{h} \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{\sqrt{4} - \sqrt{4+h}}{2\sqrt{4+h}} \right) \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{2 - \sqrt{4+h}}{2\sqrt{4+h}} \right) \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{2 - \sqrt{4+h}}{2\sqrt{4+h}} \cdot \frac{2 + \sqrt{4+h}}{2 + \sqrt{4+h}} \right) \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{4 - (4+h)}{2\sqrt{4+h}(2 + \sqrt{4+h})} \right) \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \left( \frac{-h}{2\sqrt{4+h}(2 + \sqrt{4+h})} \right) \end{aligned}$$

$$\begin{aligned}
&= \lim_{h \rightarrow 0} \left( \frac{-1}{2\sqrt{4+h}(2+\sqrt{4+h})} \right) \\
&= \frac{-1}{2\sqrt{4}(2+\sqrt{4})} \\
&= \frac{-1}{2(2)(2+2)} \\
&= \underline{\underline{\frac{-1}{16}}}
\end{aligned}$$

### Exercise

Find the values of the derivatives of the function  $f(x) = 4 - x^2$ . Then find the values of  $f'(-3)$ ,  $f'(0)$ ,  $f'(1)$

### Solution

$$\begin{aligned}
\frac{f(x+h) - f(x)}{h} &= \frac{4 - (x+h)^2 - (4 - x^2)}{h} \\
&= \frac{4 - (x^2 + 2xh + h^2) - (4 - x^2)}{h} \\
&= \frac{4 - x^2 - 2xh - h^2 - 4 + x^2}{h} \\
&= \frac{-2xh - h^2}{h} \\
&= \underline{\underline{-2x - h}}
\end{aligned}$$

$$f'(x) = \lim_{h \rightarrow 0} (-2x - h) = -2x$$

$$f'(-3) = \underline{\underline{-6}} \quad f'(0) = \underline{\underline{0}} \quad f'(1) = \underline{\underline{-2}}$$

### Exercise

Find the values of the derivatives of the function  $r(s) = \sqrt{2s+1}$ . Then find the values of  $r'(0)$ ,  $r'\left(\frac{1}{2}\right)$ ,  $r'(1)$

### Solution

$$\begin{aligned}
r'(s) &= \lim_{h \rightarrow 0} \frac{r(s+h) - r(s)}{h} \\
&= \lim_{h \rightarrow 0} \frac{\sqrt{2(s+h)+1} - \sqrt{2s+1}}{h}
\end{aligned}$$

$$\begin{aligned}
&= \lim_{h \rightarrow 0} \frac{\sqrt{2s+2h+1} - \sqrt{2s+1}}{h} \cdot \frac{\sqrt{2s+2h+1} + \sqrt{2s+1}}{\sqrt{2s+2h+1} + \sqrt{2s+1}} \\
&= \lim_{h \rightarrow 0} \frac{2s+2h+1 - (2s+1)}{h(\sqrt{2s+2h+1} + \sqrt{2s+1})} \\
&= \lim_{h \rightarrow 0} \frac{2s+2h+1-2s-1}{h(\sqrt{2s+2h+1} + \sqrt{2s+1})} \\
&= \lim_{h \rightarrow 0} \frac{2h}{h(\sqrt{2s+2h+1} + \sqrt{2s+1})} \\
&= \lim_{h \rightarrow 0} \frac{2}{\sqrt{2s+2h+1} + \sqrt{2s+1}} \\
&= \frac{2}{\sqrt{2s+1} + \sqrt{2s+1}} \\
&= \frac{2}{2\sqrt{2s+1}} \\
&= \frac{1}{\sqrt{2s+1}}
\end{aligned}$$

$$r'(0) = \frac{1}{\sqrt{2(0)+1}} = \underline{1}$$

$$r'\left(\frac{1}{2}\right) = \frac{1}{\sqrt{2\frac{1}{2}+1}} = \underline{\frac{1}{\sqrt{2}}}$$

$$r'(1) = \frac{1}{\sqrt{2(1)+1}} = \underline{\frac{1}{\sqrt{3}}}$$

## Exercise

Find the derivative of  $f(x) = 3x^2 - 2x$

### Solution

$$\begin{aligned}
f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\
&= \lim_{\Delta x \rightarrow 0} \frac{3(x + \Delta x)^2 - 2(x + \Delta x) - (3x^2 - 2x)}{\Delta x} \\
&= \lim_{\Delta x \rightarrow 0} \frac{3(x^2 + \Delta x^2 + 2x\Delta x) - 2x - 2\Delta x - 3x^2 + 2x}{\Delta x} \\
&= \lim_{\Delta x \rightarrow 0} \frac{3x^2 + 3\Delta x^2 + 6x\Delta x - 2x - 2\Delta x - 3x^2 + 2x}{\Delta x} \\
&= \lim_{\Delta x \rightarrow 0} \frac{3\Delta x^2 + 6x\Delta x - 2\Delta x}{\Delta x}
\end{aligned}$$

$$\begin{aligned}
 &= \lim_{\Delta x \rightarrow 0} 3\Delta x + 6x - 2 \\
 &= \underline{6x - 2}
 \end{aligned}$$

### Exercise

Find the derivative of  $y$  with the respect to  $t$  for the function  $y = \frac{4}{t}$

### Solution

$$\begin{aligned}
 f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\
 &= \lim_{\Delta t \rightarrow 0} \frac{\frac{4}{t + \Delta t} - \frac{4}{t}}{\Delta t} \\
 &= \lim_{\Delta t \rightarrow 0} \frac{\frac{4t - 4(t + \Delta t)}{t(t + \Delta t)}}{\Delta t} \\
 &= \lim_{\Delta t \rightarrow 0} \frac{1}{\Delta t} \frac{4t - 4(t + \Delta t)}{t(t + \Delta t)} \\
 &= \lim_{\Delta t \rightarrow 0} \frac{-4\Delta t}{t(t + \Delta t)\Delta t} \\
 &= \lim_{\Delta t \rightarrow 0} \frac{-4}{t(t + \Delta t)} \\
 &= \underline{-\frac{4}{t^2}}
 \end{aligned}$$

### Exercise

Find the derivative of  $\frac{dy}{dx}$  if  $y = 2x^3$

### Solution

$$\begin{aligned}
 f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{2(x + \Delta x)^3 - 2x^3}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{2\left(x^3 + 3x^2\Delta x + 3x(\Delta x)^2 + (\Delta x)^3\right) - 2x^3}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{2x^3 + 6x^2\Delta x + 6x(\Delta x)^2 + 3(\Delta x)^3 - 2x^3}{\Delta x} \\
 &= \lim_{\Delta x \rightarrow 0} \frac{\Delta x\left(6x^2 + 6x(\Delta x) + 3(\Delta x)^2\right)}{\Delta x}
 \end{aligned}$$

$$= \lim_{\Delta x \rightarrow 0} \left( 6x^2 + 6x(\Delta x) + 3(\Delta x)^2 \right)$$

$$\underline{= 6x^2}$$

### Exercise

Differentiate the function  $y = \frac{x+3}{1-x}$  and find the slope of the tangent line at the given value of the independent variable.

### Solution

$$f'(x) = \lim_{\Delta x \rightarrow 0} \frac{\frac{x+\Delta x+3}{1-x-\Delta x} - \frac{x+3}{1-x}}{\Delta x}$$

$$= \lim_{\Delta x \rightarrow 0} \left( \frac{1}{\Delta x} \right) \left( \frac{(x+\Delta x+3)(1-x) - (x+3)(1-x-\Delta x)}{(1-x-\Delta x)(1-x)} \right)$$

$$= \lim_{\Delta x \rightarrow 0} \left( \frac{1}{\Delta x} \right) \left( \frac{x+\Delta x+3-x^2-x\Delta x-3x - (x-x^2-x\Delta x+3-3x-3\Delta x)}{(1-x-\Delta x)(1-x)} \right)$$

$$= \lim_{\Delta x \rightarrow 0} \left( \frac{1}{\Delta x} \right) \left( \frac{x+\Delta x+3-x^2-x\Delta x-3x-x+x^2+x\Delta x-3+3x+3\Delta x}{(1-x-\Delta x)(1-x)} \right)$$

$$= \lim_{\Delta x \rightarrow 0} \left( \frac{1}{\Delta x} \right) \left( \frac{4\Delta x}{(1-x-\Delta x)(1-x)} \right)$$

$$= \lim_{\Delta x \rightarrow 0} \frac{4}{(1-x-\Delta x)(1-x)}$$

$$= \frac{4}{(1-x)(1-x)}$$

$$\underline{= \frac{4}{(1-x)^2}}$$

### Exercise

Find the equation of the tangent line to  $f(x) = x^2 + 1$  that is parallel to  $2x + y = 0$

### Solution

$$2x + y = 0 \Rightarrow y = -2x \Rightarrow \text{slope} = -2$$

$$f'(x) = \lim_{\Delta x \rightarrow 0} \frac{f(x+\Delta x) - f(x)}{\Delta x}$$

$$\begin{aligned}
&= \lim_{\Delta x \rightarrow 0} \frac{(x + \Delta x)^2 + 1 - (x^2 + 1)}{\Delta x} \\
&= \lim_{\Delta x \rightarrow 0} \frac{x^2 + \Delta x^2 + 2x\Delta x + 1 - x^2 - 1}{\Delta x} \\
&= \lim_{\Delta x \rightarrow 0} \frac{\Delta x^2 + 2x\Delta x}{\Delta x} \\
&= \lim_{\Delta x \rightarrow 0} \Delta x + 2x = 2x
\end{aligned}$$

$$f' = 2x = -2 \Rightarrow x = -1 \Rightarrow f(-1) = (-1)^2 + 1 = 2 \rightarrow (-1, 2)$$

The line equation is given by  $y = m(x - x_1) + y_1$

$$y = -2(x + 1) + 2$$

$$\underline{y = -2x}$$

### ***Exercise***

Use the definition of limits to find the derivative:  $f(x) = \frac{3}{\sqrt{x}} \quad y - 2 = -2x - 2$

### ***Solution***

$$\begin{aligned}
f'(x) &= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} \\
&= \lim_{\Delta x \rightarrow 0} \frac{\left(\frac{3}{\sqrt{x + \Delta x}}\right) - \left(\frac{3}{\sqrt{x}}\right)}{\Delta x} \cdot \frac{\sqrt{x} \cdot \sqrt{x + \Delta x}}{\sqrt{x} \cdot \sqrt{x + \Delta x}} \\
&= \lim_{\Delta x \rightarrow 0} \frac{3\sqrt{x} - 3\sqrt{x + \Delta x}}{\Delta x (\sqrt{x} \cdot \sqrt{x + \Delta x})} \\
&= \lim_{\Delta x \rightarrow 0} \frac{3(\sqrt{x} - \sqrt{x + \Delta x})}{\Delta x (\sqrt{x} \cdot \sqrt{x + \Delta x})} \cdot \frac{\sqrt{x} + \sqrt{x + \Delta x}}{\sqrt{x} + \sqrt{x + \Delta x}} \\
&= \lim_{\Delta x \rightarrow 0} \frac{3(x - (x + \Delta x))}{\Delta x (\sqrt{x} \cdot \sqrt{x + \Delta x}) (\sqrt{x} + \sqrt{x + \Delta x})} \\
&= \lim_{\Delta x \rightarrow 0} \frac{-3\Delta x}{\Delta x (\sqrt{x} \cdot \sqrt{x + \Delta x}) (\sqrt{x} + \sqrt{x + \Delta x})}
\end{aligned}$$

$$= \frac{-3}{x(2\sqrt{x})}$$

$$= \frac{-3}{2x^{3/2}}$$

### Exercise

Use the definition of limits to find the derivative:  $f(x) = \sqrt{x+2}$

### Solution

$$f'(x) = \lim_{\Delta x \rightarrow 0} \frac{f(x+\Delta x) - f(x)}{\Delta x}$$

$$= \lim_{\Delta x \rightarrow 0} \frac{\sqrt{x+\Delta x+2} - \sqrt{x+2}}{\Delta x} \cdot \frac{\sqrt{x+\Delta x+2} + \sqrt{x+2}}{\sqrt{x+\Delta x+2} + \sqrt{x+2}}$$

$$= \lim_{\Delta x \rightarrow 0} \frac{x+\Delta x+2 - (x+2)}{\Delta x(\sqrt{x+\Delta x+2} + \sqrt{x+2})}$$

$$= \lim_{\Delta x \rightarrow 0} \frac{\Delta x}{\Delta x(\sqrt{x+\Delta x+2} + \sqrt{x+2})}$$

$$= \lim_{\Delta x \rightarrow 0} \frac{1}{\sqrt{x+\Delta x+2} + \sqrt{x+2}}$$

$$= \frac{1}{2\sqrt{x+2}}$$

### Exercise

Suppose the height  $s$  of an object (in  $m$ ) above the ground after  $t$  seconds is approximated by the function

$$s = -4.9t^2 + 25t + 1$$

- Make a table showing the average velocities of the object from time  $t = 1$  to  $t = 1 + h$ , for  $h = 0.01, 0.001, 0.0001$ , and  $0.00001$ .
- Use the table in part (a) to estimate the instantaneous velocity of the object at  $t = 1$ .
- Use limits to verify your estimate in part (b).

### Solution

$$a) \frac{f(1+h) - f(1)}{h} = \frac{1}{h} \left[ -4.9(1+h)^2 + 25(1+h) + 1 + 4.9 - 25 - 1 \right]$$

$$= \frac{1}{h} \left[ -4.9 - 9.8h - 4.9h^2 + 25h + 4.9 \right]$$

$$= \frac{1}{h} \left( -4.9h^2 + 15.2h \right)$$



$$= 15.2 - 4.9h \quad |$$

$h$	$\frac{f(1+h) - f(1)}{h}$
0.01	15.151
0.001	15.1951
0.0001	15.1995
0.00001	15.2
0.000001	15.2

$$b) \quad f'(1) = \lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h}$$


$$\approx 15.2 \text{ m/sec} \quad |$$

$$c) \quad f'(1) = \lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h}$$

$$= 15.2 \text{ m/sec} \quad |$$

### Exercise

Suppose the following graph represents the number of bacteria in a culture  $t$  hours after the start of an experiment.



- At approximately what time is the instantaneous growth rate the greatest, for  $0 \leq t \leq 36$ ? Estimate the growth rate at this time.
- At approximately what time is the instantaneous growth rate the least, for  $0 \leq t \leq 36$ ? Estimate the growth rate at this time.
- What is the average growth rate over the interval  $0 \leq t \leq 36$ ?

### Solution

$$a) \quad t = \frac{36}{2} = 18$$

$$\text{Point the rate} = \frac{N(20) - N(16)}{20 - 16}$$

$$= \frac{2500 - 1900}{4}$$

$$= 400 \text{ bacteria/hr}$$

**b)** It is smallest at  $t = 0$  or  $t = 36$

$$\frac{N(36) - N(32)}{4} = \frac{4900 - 4800}{4}$$

$$= 25 \text{ bacteria/hr}$$

**c)** Growth rate  $= \frac{N(36) - N(0)}{36}$

$$\approx \frac{4900 - 400}{36}$$

$$= 125 \text{ bacteria/hr}$$