

## ***Solution***      **Section 2.5 – Polynomial Functions**

### ***Exercise***

Determine the end behavior of the graph of the polynomial function  $f(x) = 5x^3 + 7x^2 - x + 9$

### **Solution**

Leading term:  $5x^3$  with  $3^{rd}$  degree ( $n$  is **odd**)

$$x \rightarrow -\infty \Rightarrow f(x) \rightarrow -\infty \quad f(x) \text{ falls left}$$

$$x \rightarrow \infty \Rightarrow f(x) \rightarrow \infty \quad f(x) \text{ rises right}$$

### ***Exercise***

Determine the end behavior of the graph of the polynomial function  $f(x) = 11x^3 - 6x^2 + x + 3$

### **Solution**

Leading term:  $11x^3$  with  $3^{rd}$  degree ( $n$  is **odd**)

$$x \rightarrow -\infty \Rightarrow f(x) \rightarrow -\infty \quad f(x) \text{ falls left}$$

$$x \rightarrow \infty \Rightarrow f(x) \rightarrow \infty \quad \text{rises right}$$

### ***Exercise***

Determine the end behavior of the graph of the polynomial function  $f(x) = -11x^3 - 6x^2 + x + 3$

### **Solution**

Leading term:  $-11x^3$  with  $3^{rd}$  degree ( $n$  is **odd**)

$$x \rightarrow -\infty \Rightarrow f(x) \rightarrow \infty \quad f(x) \text{ rises left}$$

$$x \rightarrow \infty \Rightarrow f(x) \rightarrow -\infty \quad f(x) \text{ falls right}$$

### ***Exercise***

Determine the end behavior of the graph of the polynomial function  $f(x) = 2x^3 + 3x^2 - 23x - 42$

### **Solution**

Leading term:  $2x^3$  with  $3^{rd}$  degree ( $n$  is **odd**)

$$x \rightarrow -\infty \Rightarrow f(x) \rightarrow -\infty \quad f(x) \text{ falls left}$$

$$x \rightarrow \infty \Rightarrow f(x) \rightarrow \infty \quad f(x) \text{ rises right}$$

### Exercise

Determine the end behavior of the graph of the polynomial function  $f(x) = 5x^4 + 7x^2 - x + 9$

#### Solution

Leading term:  $5x^4$  with  $4^{rd}$  degree ( $n$  is *even*)

$$x \rightarrow -\infty \Rightarrow f(x) \rightarrow \infty \quad f(x) \text{ rises left}$$

$$x \rightarrow \infty \Rightarrow f(x) \rightarrow \infty \quad f(x) \text{ rises right}$$

### Exercise

Determine the end behavior of the graph of the polynomial function  $f(x) = 11x^4 - 6x^2 + x + 3$

#### Solution

Leading term:  $11x^4$  with  $4^{rd}$  degree ( $n$  is *even*)

$$x \rightarrow -\infty \Rightarrow f(x) \rightarrow \infty \quad f(x) \text{ rises left}$$

$$x \rightarrow \infty \Rightarrow f(x) \rightarrow \infty \quad f(x) \text{ rises right}$$

### Exercise

Determine the end behavior of the graph of the polynomial function  $f(x) = -5x^4 + 7x^2 - x + 9$

#### Solution

Leading term:  $-5x^4$  with  $4^{rd}$  degree ( $n$  is *even*)

$$x \rightarrow -\infty \Rightarrow f(x) \rightarrow -\infty \quad f(x) \text{ falls left}$$

$$x \rightarrow \infty \Rightarrow f(x) \rightarrow -\infty \quad f(x) \text{ falls right}$$

### Exercise

Determine the end behavior of the graph of the polynomial function  $f(x) = -11x^4 - 6x^2 + x + 3$

#### Solution

Leading term:  $-11x^4$  with  $4^{rd}$  degree ( $n$  is *even*)

$$x \rightarrow -\infty \Rightarrow f(x) \rightarrow -\infty \quad f(x) \text{ falls left}$$

$$x \rightarrow \infty \Rightarrow f(x) \rightarrow -\infty \quad f(x) \text{ falls right}$$

### ***Exercise***

Determine the end behavior of the graph of the polynomial function  $f(x) = 5x^5 - 16x^2 - 20x + 64$

#### **Solution**

Leading term:  $5x^5$  with  $5^{th}$  degree ( $n$  is **odd**)

$$x \rightarrow -\infty \Rightarrow f(x) \rightarrow -\infty \quad f(x) \text{ falls left}$$

$$x \rightarrow \infty \Rightarrow f(x) \rightarrow \infty \quad f(x) \text{ rises right}$$

### ***Exercise***

Determine the end behavior of the graph of the polynomial function  $f(x) = -5x^5 - 16x^2 - 20x + 64$

#### **Solution**

Leading term:  $-5x^5$  with  $5^{th}$  degree ( $n$  is **odd**)

$$x \rightarrow -\infty \Rightarrow f(x) \rightarrow \infty \quad f(x) \text{ rises left}$$

$$x \rightarrow \infty \Rightarrow f(x) \rightarrow -\infty \quad f(x) \text{ falls right}$$

### ***Exercise***

Determine the end behavior of the graph of the polynomial function  $f(x) = -3x^6 - 16x^3 + 64$

#### **Solution**

Leading term:  $-3x^6$  with  $6^{th}$  degree ( $n$  is **even**)

$$x \rightarrow -\infty \Rightarrow f(x) \rightarrow -\infty \quad f(x) \text{ falls left}$$

$$x \rightarrow \infty \Rightarrow f(x) \rightarrow -\infty \quad f(x) \text{ falls right}$$

### ***Exercise***

Determine the end behavior of the graph of the polynomial function  $f(x) = 3x^6 - 16x^3 + 4$

#### **Solution**

Leading term:  $3x^6$  with  $6^{th}$  degree ( $n$  is **even**)

$$x \rightarrow -\infty \Rightarrow f(x) \rightarrow \infty \quad f(x) \text{ rises left}$$

$$x \rightarrow \infty \Rightarrow f(x) \rightarrow \infty \quad f(x) \text{ rises right}$$

### Exercise

Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.  $f(x) = x^3 - x - 1$ ; between 1 and 2

### Solution

$$\begin{aligned} f(1) &= (1)^3 - (1) - 1 \\ &= -1 \end{aligned}$$

$$\begin{aligned} f(2) &= (2)^3 - (2) - 1 \\ &= 5 \end{aligned}$$

Since  $f(1)$  and  $f(2)$  have opposite signs.

Therefore, the polynomial *has a real zero* between 1 and 2.

### Exercise

Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.  $f(x) = x^3 - 4x^2 + 2$ ; between 0 and 1

### Solution

$$\begin{aligned} f(0) &= (0)^3 - 4(0)^2 + 2 \\ &= 2 \end{aligned}$$

$$\begin{aligned} f(1) &= (1)^3 - 4(1)^2 + 2 \\ &= -1 \end{aligned}$$

Since  $f(0)$  and  $f(1)$  have opposite signs.

Therefore, the polynomial *has a real zero* between 0 and 1.

### Exercise

Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.  $f(x) = 2x^4 - 4x^2 + 1$ ; between -1 and 0

### Solution

$$\begin{aligned} f(-1) &= 2(-1)^4 - 4(-1)^2 + 1 \\ &= -1 \end{aligned}$$

$$\begin{aligned} f(0) &= 2(0)^4 - 4(0)^2 + 1 \\ &= 1 \end{aligned}$$

Since  $f(0)$  and  $f(-1)$  have opposite signs.

Therefore, the polynomial *has a real zero* between  $-1$  and  $0$ .

### ***Exercise***

Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.  $f(x) = x^4 + 6x^3 - 18x^2$ ; between  $2$  and  $3$

### **Solution**

$$f(2) = (2)^4 + 6(2)^3 - 18(2)^2$$
$$= -8$$

$$f(3) = (3)^4 + 6(3)^3 - 18(3)^2$$
$$= 81$$

Since  $f(2)$  and  $f(3)$  have opposite signs.

Therefore, the polynomial *has a real zero* between  $2$  and  $3$ .

### ***Exercise***

Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.  $f(x) = x^3 + x^2 - 2x + 1$ ; between  $-3$  and  $-2$

### **Solution**

$$f(-3) = (-3)^3 + (-3)^2 - 2(-3) + 1$$
$$= -11$$

$$f(-2) = (-2)^3 + (-2)^2 - 2(-2) + 1$$
$$= 1$$

Since  $f(-3)$  and  $f(-2)$  have opposite signs.

Therefore, the polynomial *has a real zero* between  $-2$  and  $-3$ .

### ***Exercise***

Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.  $f(x) = x^5 - x^3 - 1$ ; between  $1$  and  $2$

### **Solution**

$$f(1) = (1)^5 - (1)^3 - 1$$

$$\underline{= -1}$$

$$f(2) = (2)^5 - (2)^3 - 1$$

$$\underline{= 23}$$

Since  $f(1)$  and  $f(2)$  have opposite signs.

Therefore, the polynomial *has a real zero* between 1 and 2.

### Exercise

Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.  $f(x) = 3x^3 - 10x + 9$ ; between  $-3$  and  $-2$

### Solution

$$f(-3) = 3(-3)^3 - 10(-3) + 9$$

$$\underline{= -42}$$

$$f(-2) = 3(-2)^3 - 10(-2) + 9$$

$$\underline{= 5}$$

Since  $f(-3)$  and  $f(-2)$  have opposite signs.

Therefore, the polynomial *has a real zero* between  $-3$  and  $-2$ .

### Exercise

Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.  $f(x) = 3x^3 - 8x^2 + x + 2$ ; between 2 and 3

### Solution

$$f(2) = 3(2)^3 - 8(2)^2 + (2) + 2$$

$$\underline{= -4}$$

$$f(3) = 3(3)^3 - 8(3)^2 + (3) + 2$$

$$\underline{= 14}$$

Since  $f(2)$  and  $f(3)$  have opposite signs.

Therefore, the polynomial *has a real zero* between 2 and 3.

### Exercise

Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.  $f(x) = 3x^3 - 8x^2 + x + 2$ ; between 1 and 2

### Solution

$$\begin{aligned} f(1) &= 3(1)^3 - 8(1)^2 + (1) + 2 \\ &= -2 \end{aligned}$$

$$\begin{aligned} f(2) &= 3(2)^3 - 8(2)^2 + (2) + 2 \\ &= -4 \end{aligned}$$

Since  $f(1)$  and  $f(2)$  have same signs.

Therefore, *cannot be determined*.

### Exercise

Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.  $f(x) = x^5 + 2x^4 - 6x^3 + 2x - 3$ ; between 0 and 1

### Solution

$$\begin{aligned} f(0) &= (0)^5 + 2(0)^4 - 6(0)^3 + 2(0) - 3 \\ &= -3 \end{aligned}$$

$$\begin{aligned} f(1) &= (1)^5 + 2(1)^4 - 6(1)^3 + 2(1) - 3 \\ &= -4 \end{aligned}$$

Since  $f(0)$  and  $f(1)$  have same signs.

Therefore, *cannot be determined*.

### Exercise

Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.  $P(x) = 2x^3 + 3x^2 - 23x - 42$ ,  $a = 3$ ,  $b = 4$

### Solution

$$\begin{aligned} P(3) &= 54 + 27 - 69 - 42 \\ &= -30 \end{aligned}$$

$$\begin{aligned} P(4) &= 128 + 48 - 92 - 42 \\ &= 90 \end{aligned}$$

Since  $P(3)$  and  $P(4)$  have opposite signs.

Therefore, the polynomial *has a real zero* between 3 and 4.

### **Exercise**

Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.  $P(x) = 4x^3 - x^2 - 6x + 1$ ,  $a = 0$ ,  $b = 1$

### **Solution**

$$P(0) = \underline{1}$$

$$P(1) = 4 - 1 - 6 + 1 \\ = \underline{-2}$$

Since  $P(0)$  and  $P(1)$  have opposite signs.

Therefore, the polynomial *has a real zero* between 0 and 1.

### **Exercise**

Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.  $P(x) = 3x^3 + 7x^2 + 3x + 7$ ,  $a = -3$ ,  $b = -2$

### **Solution**

$$P(-3) = -81 + 63 - 9 + 7 \\ = \underline{-20}$$

$$P(-2) = -24 + 28 - 6 + 7 \\ = \underline{5}$$

Since  $P(-3)$  and  $P(-2)$  have opposite signs.

Therefore, the polynomial *has a real zero* between  $-3$  and  $-2$ .

### **Exercise**

Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.  $P(x) = 2x^3 - 21x^2 - 2x + 25$ ,  $a = 1$ ,  $b = 2$

### **Solution**

$$P(1) = 2 - 21 - 2 + 25 \\ = \underline{4}$$



$$P(2) = 16 - 84 - 4 + 25$$

$$= -47$$

Since  $P(1)$  and  $P(2)$  have opposite signs.

Therefore, the polynomial *has a real zero* between 1 and 2.

### Exercise

Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.  $P(x) = 4x^4 + 7x^3 - 11x^2 + 7x - 15$ ,  $a = 1$ ,  $b = \frac{3}{2}$

### Solution

$$P(1) = 4 + 7 - 11 + 7 - 15$$

$$= -8$$

$$P\left(\frac{3}{2}\right) = 81 + \frac{189}{8} - \frac{99}{4} + \frac{21}{2} - 15$$

$$= 66 + \frac{189 - 198 + 84}{8}$$

$$= 66 + \frac{75}{8}$$

$$= \frac{603}{8}$$

Since  $P(1)$  and  $P\left(\frac{3}{2}\right)$  have opposite signs.

Therefore, the polynomial *has a real zero* between 1 and  $\frac{3}{2}$ .

### Exercise

Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.  $P(x) = 5x^3 - 16x^2 - 20x + 64$ ,  $a = 3$ ,  $b = \frac{7}{2}$

### Solution

$$P(3) = 135 - 144 - 60 + 64$$

$$= -5$$

$$P\left(\frac{7}{2}\right) = \frac{1715}{8} - 196 - 70 + 64$$

$$= \frac{1715}{8} - 202$$

$$= \frac{99}{8}$$

Since  $P(3)$  and  $P\left(\frac{7}{2}\right)$  have opposite signs.

Therefore, the polynomial *has a real zero* between 3 and  $\frac{7}{2}$ .

### Exercise

Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.  $P(x) = x^4 - x^2 - x - 4$ ,  $a = 1$ ,  $b = 2$

### Solution

$$\begin{aligned} P(1) &= 1 - 1 - 1 - 4 \\ &= -5 \end{aligned}$$

$$\begin{aligned} P(2) &= 16 - 4 - 2 - 4 \\ &= 6 \end{aligned}$$

Since  $P(1)$  and  $P(2)$  have opposite signs.

Therefore, the polynomial *has a real zero* between 1 and 2.

### Exercise

Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.  $P(x) = x^3 - x - 8$ ,  $a = 2$ ,  $b = 3$

### Solution

$$\begin{aligned} P(2) &= 8 - 2 - 8 \\ &= -2 \end{aligned}$$

$$\begin{aligned} P(3) &= 27 - 3 - 8 \\ &= 16 \end{aligned}$$

Since  $P(2)$  and  $P(3)$  have opposite signs.

Therefore, the polynomial *has a real zero* between 2 and 3.

### Exercise

Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.  $P(x) = x^3 - x - 8$ ,  $a = 0$ ,  $b = 1$

### Solution

$$\underline{P(0) = -8}$$

$$P(1) = 1 - 1 - 8$$

$$= -8$$

Since  $P(0)$  and  $P(1)$  have same sign.

Therefore, *cannot be determined*.

### Exercise

Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.  $P(x) = x^3 - x - 8$ ,  $a = 2.1$ ,  $b = 2.2$

### Solution

$$P(2.1) = P\left(\frac{21}{10}\right)$$

$$= \frac{9261}{1000} - \frac{21}{10} - 8$$

$$= \frac{9261 - 2100 - 8000}{1000}$$

$$= -\frac{839}{1,000}$$

$$P(2.2) = P\left(\frac{22}{10}\right)$$

$$= \frac{10,648}{1000} - \frac{22}{10} - 8$$

$$= \frac{10,648 - 2,200 - 8,000}{1000}$$

$$= \frac{448}{1,000}$$

$$= 0.448$$

Since  $P(2.1)$  and  $P(2.2)$  have opposite signs.

Therefore, the polynomial *has a real zero* between 2.1 and 2.2.