

Solution ***Section 1.5 – Length of Curves***

Exercise


Find the length of the curve $y = \frac{1}{3}(x^2 + 2)^{3/2}$ from $x = 0$ to $x = 3$.

Solution

$$\frac{dy}{dx} = \frac{1}{3} \cdot \frac{3}{2} (x^2 + 2)^{1/2} (2x) = x(x^2 + 2)^{1/2}$$

$$\begin{aligned}\sqrt{1 + \left(\frac{dy}{dx}\right)^2} &= \sqrt{1 + x^2(x^2 + 2)} \\ &= \sqrt{1 + x^4 + 2x^2} \\ &= \sqrt{(x^2 + 1)^2} \\ &= x^2 + 1\end{aligned}$$

$$\begin{aligned}L &= \int_0^3 (x^2 + 1) \, dx \\ &= \left[\frac{1}{3}x^3 + x \right]_0^3 \\ &= \frac{1}{3}(3)^3 + (3) - 0 \\ &= \underline{12 \text{ unit}}\end{aligned}$$



Exercise

Find the length of the curve $y = (x)^{3/2}$ from $x = 0$ to $x = 4$.


Solution

$$\frac{dy}{dx} = \frac{3}{2}x^{1/2}$$

$$\begin{aligned}\sqrt{1 + \left(\frac{dy}{dx}\right)^2} &= \sqrt{1 + \frac{9}{4}x} \\ &= \sqrt{\frac{4 + 9x}{4}} \\ &= \frac{1}{2}\sqrt{4 + 9x}\end{aligned}$$

$$L = \int_0^4 \frac{1}{2}(4 + 9x)^{1/2} \, dx \qquad u = 4 + 9x \Rightarrow du = 9dx \rightarrow \frac{1}{9}du = dx \quad \begin{cases} x = 4 & \rightarrow u = 40 \\ x = 0 & \rightarrow u = 4 \end{cases}$$

$$\begin{aligned}
&= \frac{1}{2} \int_4^{40} \frac{1}{9} u^{1/2} du \\
&= \frac{1}{18} \left[\frac{2}{3} u^{3/2} \right]_4^{40} \\
&= \frac{1}{27} \left(40^{3/2} - 4^{3/2} \right) \\
&= \frac{1}{27} \left(\sqrt{40^3} - \sqrt{4^3} \right) \\
&= \frac{1}{27} (80\sqrt{10} - 8) \\
&= \frac{8}{27} (10\sqrt{10} - 1) \text{ unit}
\end{aligned}$$



Exercise

Find the length of the curve $x = \frac{y^{3/2}}{3} - y^{1/2}$ from $y = 1$ to $y = 9$.

Solution

$$\frac{dx}{dy} = \frac{1}{2} y^{1/2} - \frac{1}{2} y^{-1/2} = \frac{1}{2} \left(y^{1/2} - \frac{1}{y^{1/2}} \right)$$

$$\sqrt{1 + \left(\frac{dx}{dy} \right)^2} = \sqrt{1 + \frac{1}{4} \left(y^{1/2} - \frac{1}{y^{1/2}} \right)^2}$$

$$= \sqrt{1 + \frac{1}{4} \left(y - 2 + \frac{1}{y} \right)}$$

$$= \sqrt{1 + \frac{1}{4} y - \frac{1}{2} + \frac{1}{4y}}$$

$$= \sqrt{\frac{1}{4} y + \frac{1}{2} + \frac{1}{4y}}$$

$$= \sqrt{\frac{1}{4} \left(y + 2 + \frac{1}{y} \right)}$$

$$= \frac{1}{2} \sqrt{\left(\sqrt{y} + \frac{1}{\sqrt{y}} \right)^2}$$

$$= \frac{1}{2} \left(\sqrt{y} + \frac{1}{\sqrt{y}} \right)$$

$$L = \frac{1}{2} \int_1^9 \left(y^{1/2} + y^{-1/2} \right) dy$$

$$a = \frac{1}{3}, \quad m = \frac{3}{2}, \quad b = -1, \quad n = \frac{1}{2}$$

$$1. \quad m + n = \frac{3}{2} + \frac{1}{2} = 2 \quad \checkmark$$

$$2. \quad abmn = \frac{1}{3}(-1)\left(\frac{3}{2}\right)\left(\frac{1}{2}\right) = -\frac{1}{4} \quad \checkmark$$

$$L = \left(\frac{1}{3} y^{3/2} + y^{1/2} \right)_1^9$$

$$= 9 + 3 - \frac{4}{3}$$

$$= \frac{32}{3} \text{ unit}$$

$$\begin{aligned}
&= \frac{1}{2} \left[\frac{2}{3} y^{3/2} + 2y^{1/2} \right]_1^9 \\
&= \left[\frac{1}{3} y^{3/2} + y^{1/2} \right]_1^9 \\
&= \left[\frac{1}{3} 9^{3/2} + 9^{1/2} - \left(\frac{1}{3} 1^{3/2} + 1^{1/2} \right) \right] \\
&= \frac{1}{3} 3^3 + 3 - \left(\frac{1}{3} + 1 \right) \\
&= 9 + 3 - \frac{4}{3} \\
&= \frac{32}{3} \text{ unit}
\end{aligned}$$

Exercise

Find the length of the curve $x = \frac{y^3}{6} + \frac{1}{2y}$ from $y = 2$ to $y = 3$.

Solution

$$\begin{aligned}
\frac{dx}{dy} &= \frac{1}{2} y^2 - \frac{1}{2y^2} = \frac{1}{2} (y^2 - y^{-2}) \\
\sqrt{1 + \left(\frac{dx}{dy} \right)^2} &= \sqrt{1 + \frac{1}{4} (y^2 - y^{-2})^2} \\
&= \frac{1}{2} \sqrt{4 + (y^4 - 2 + y^{-4})} \\
&= \frac{1}{2} \sqrt{y^4 + 2 + y^{-4}} \\
&= \frac{1}{2} \sqrt{(y^2 + y^{-2})^2} \\
&= \frac{1}{2} (y^2 + y^{-2})
\end{aligned}$$


$$\begin{aligned}
L &= \frac{1}{2} \int_2^3 (y^2 + y^{-2}) dy \\
&= \frac{1}{2} \left[\frac{1}{3} y^3 - y^{-1} \right]_2^3 \\
&= \frac{1}{2} \left[\left(\frac{1}{3} 3^3 - 3^{-1} \right) - \left(\frac{1}{3} 2^3 - 2^{-1} \right) \right] \\
&= \frac{1}{2} \left[9 - \frac{1}{3} - \left(\frac{8}{3} - \frac{1}{2} \right) \right] \\
&= \frac{1}{2} \left(\frac{26}{3} - \frac{13}{6} \right) \\
&= \frac{13}{4} \text{ unit}
\end{aligned}$$

$$a = \frac{1}{6}, \quad m = 3, \quad b = \frac{1}{2}, \quad n = -1$$

$$1. \quad m + n = 3 - 1 = 2 \quad \checkmark$$

$$2. \quad abmn = \frac{1}{6} \left(\frac{1}{2} \right) (3) (-1) = -\frac{1}{4} \quad \checkmark$$

$$\begin{aligned}
L &= \left(\frac{y^3}{6} - \frac{1}{2y} \right) \Big|_2^3 \\
&= \frac{1}{2} \left[9 - \frac{1}{3} - \left(\frac{8}{3} - \frac{1}{2} \right) \right] \\
&= \frac{13}{4} \text{ unit}
\end{aligned}$$



Exercise

Find the length of the curve $f(x) = x^3 + \frac{1}{12x}$ for $\frac{1}{2} \leq x \leq 2$

Solution

$$a = 1, \quad m = 3, \quad b = \frac{1}{12}, \quad n = -1$$

$$1. \quad m + n = 2 \quad \checkmark$$

$$2. \quad abmn = -\frac{1}{4} \quad \checkmark$$

$$\begin{aligned} L &= \left(x^3 - \frac{1}{12x} \right) \Big|_{1/2}^2 \\ &= 8 - \frac{1}{24} - \frac{1}{8} + \frac{1}{6} \\ &= 8 \text{ unit} \end{aligned}$$

Exercise

Find the length of the curve of $f(x) = \frac{1}{5}x^5 + \frac{1}{12x^3}$ $1 \leq x \leq 2$

Solution

$$a = \frac{1}{5}, \quad m = 5, \quad b = \frac{1}{12}, \quad n = -3$$

$$1. \quad m + n = 5 - 3 = 2 \quad \checkmark$$

$$2. \quad abmn = \frac{1}{5} \left(\frac{1}{12} \right) (5)(-3) = -\frac{1}{4} \quad \checkmark$$

$$\begin{aligned} L &= \frac{1}{5}x^5 - \frac{1}{12x^3} \Big|_1^2 \\ &= \frac{32}{5} - \frac{1}{96} - \frac{1}{5} + \frac{1}{12} \\ &= \frac{31}{5} + \frac{7}{96} \\ &= \frac{3011}{480} \end{aligned}$$

Exercise

Find the length of the curve of $y = \frac{1}{3}x^{1/2} - x^{3/2}$, $0 \leq x \leq \frac{1}{3}$

Solution

$$a = \frac{1}{3}, \quad m = \frac{1}{2}, \quad b = -1, \quad n = \frac{3}{2}$$

$$3. \quad m + n = \frac{1}{2} + \frac{3}{2} = 2 \quad \checkmark$$

$$4. \quad abmn = \frac{1}{3}(-1)\left(\frac{1}{2}\right)\left(\frac{3}{2}\right) = -\frac{1}{4} \quad \checkmark$$

$$L = \frac{1}{3}x^{1/2} + x^{3/2} \Big|_0^{1/3}$$

$$= \frac{1}{3\sqrt{3}} + \frac{1}{3\sqrt{3}}$$

$$= \frac{2}{3\sqrt{3}} \quad = \frac{2\sqrt{3}}{9}$$

Exercise

Find the length of the curve of $y = \frac{1}{3}x^3 + \frac{1}{4x}, \quad 1 \leq x \leq 2$

Solution

$$a = \frac{1}{3}, \quad m = 3, \quad b = \frac{1}{4}, \quad n = -1$$

$$5. \quad m + n = 3 - 1 = 2 \quad \checkmark$$

$$6. \quad abmn = \frac{1}{3} \left(\frac{1}{4} \right) (3) (-1) = -\frac{1}{4} \quad \checkmark$$

$$L = \frac{1}{3}x^3 - \frac{1}{4x} \Big|_1^2$$

$$= \frac{8}{3} - \frac{1}{8} - \frac{1}{3} + \frac{1}{4} = \frac{7}{3} + \frac{1}{8}$$

$$= \frac{59}{24}$$

Exercise

Find the length of the curve of $y = 2e^x + \frac{1}{8}e^{-x} \quad 0 \leq x \leq \ln 2$

Solution

$$a = 2, \quad m = 1, \quad b = \frac{1}{8}, \quad n = -1$$

$$7. \quad m - n = 1 - (-1) = 2 \quad \checkmark$$

$$8. \quad abmn = 2 \left(\frac{1}{8} \right) (1) (-1) = -\frac{1}{4} \quad \checkmark$$

$$L = 2e^x - \frac{1}{8}e^{-x} \Big|_0^{\ln 2}$$

$$= 2e^{\ln 2} - \frac{1}{8}e^{-\ln 2} - 2 + \frac{1}{8}$$

$$= 4 - \frac{1}{16} - \frac{15}{8}$$

$$= \frac{33}{16}$$

Exercise

Find the length of the curve of $y = e^{2x} + \frac{1}{16}e^{-2x}, \quad 0 \leq x \leq \ln 3$

Solution

$$a = 1, \quad m = 2, \quad b = \frac{1}{16}, \quad n = -2$$

$$9. \quad m = -n = 2 \quad \checkmark$$

$$10. \quad abmn = 1\left(\frac{1}{16}\right)(2)(-2) = -\frac{1}{4} \quad \checkmark$$

$$\begin{aligned} L &= e^{2x} - \frac{1}{16}e^{-2x} \Big|_0^{\ln 3} \\ &= e^{2\ln 3} - \frac{1}{16}e^{-2\ln 3} - 1 + \frac{1}{16} \\ &= 9 - \frac{1}{16}\left(\frac{1}{9}\right) - \frac{15}{16} \\ &= \frac{1,160}{144} \\ &= \frac{145}{18} \end{aligned}$$

Exercise

Find the length of the curve $y = \ln(\cos x)$ $0 \leq x \leq \frac{\pi}{4}$

Solution

$$\begin{aligned} \frac{dy}{dx} &= \frac{-\sin x}{\cos x} = -\tan x \\ L &= \int_0^{\pi/4} \sqrt{1 + \tan^2 x} \, dx \\ &= \int_0^{\pi/4} \sqrt{\sec^2 x} \, dx \\ &= \int_0^{\pi/4} \sec x \, dx \\ &= \left[\ln|\sec x + \tan x| \right]_0^{\pi/4} \\ &= \ln\left|\sec \frac{\pi}{4} + \tan \frac{\pi}{4}\right| - \ln|\sec 0 + \tan 0| \\ &= \ln|\sqrt{2} + 1| - \ln|1 + 0| \\ &= \ln|\sqrt{2} + 1| - 0 \\ &= \ln(\sqrt{2} + 1) \text{ unit} \end{aligned}$$

Exercise

Find the length of the curve $f(y) = 2e^{\sqrt{2}y} + \frac{1}{16}e^{-\sqrt{2}y}$ for $0 \leq y \leq \frac{\ln 2}{\sqrt{2}}$

Solution

$$a = 2, \quad m = \sqrt{2}, \quad b = \frac{1}{16}, \quad n = -\sqrt{2}$$

$$1. \quad m = -n \quad \checkmark$$

$$2. \quad abmn = 2(\sqrt{2})\left(\frac{1}{16}\right)(-\sqrt{2}) = -\frac{1}{4} \quad \checkmark$$


$$\begin{aligned} L &= \left(2e^{\sqrt{2}y} + \frac{1}{16}e^{-\sqrt{2}y} \right) \Big|_0^{\ln 2/\sqrt{2}} \\ &= 2e^{\ln 2} + \frac{1}{16}e^{-\ln 2} - 2 - \frac{1}{16} \\ &= 4 + \frac{1}{32} - \frac{33}{16} \\ &= \frac{63}{32} \text{ unit} \end{aligned}$$

Exercise

Find the length of the curve $y = \frac{x^3}{3} + x^2 + x + 1 + \frac{1}{4x+4}$ $0 \leq x \leq 2$

Solution

$$\begin{aligned} \frac{dy}{dx} &= x^2 + 2x + 1 - \frac{1}{4} \frac{1}{(x+1)^2} = (x+1)^2 - \frac{1}{4} \frac{1}{(x+1)^2} \\ \sqrt{1 + \left(\frac{dy}{dx}\right)^2} &= \sqrt{1 + \left((x+1)^2 - \frac{1}{4} \frac{1}{(x+1)^2} \right)^2} \\ &= \sqrt{1 + (x+1)^4 - \frac{1}{2} - \frac{1}{16} \frac{1}{(x+1)^4}} \\ &= \sqrt{(x+1)^4 + \frac{1}{2} - \frac{1}{16} \frac{1}{(x+1)^4}} \\ &= \sqrt{\left((x+1)^2 + \frac{1}{4} \frac{1}{(x+1)^2} \right)^2} \\ &= (x+1)^2 + \frac{1}{4} (x+1)^{-2} \end{aligned}$$



$$\begin{aligned} L &= \int_0^2 \left((x+1)^2 + \frac{1}{4} (x+1)^{-2} \right) dx \\ &= \int_1^3 \left(u^2 + \frac{1}{4} u^{-2} \right) du \\ &= \left[\frac{1}{3} u^3 - \frac{1}{4} u^{-1} \right]_1^3 \\ &= 9 - \frac{1}{12} - \left(\frac{1}{3} - \frac{1}{4} \right) \\ &= \frac{53}{6} \text{ unit} \end{aligned}$$

$$u = x + 1 \Rightarrow du = dx \quad \begin{cases} x = 2 & \rightarrow u = 3 \\ x = 0 & \rightarrow u = 1 \end{cases}$$

Exercise

Find the length of the curve $y = \ln(e^x - 1) - \ln(e^x + 1)$ $\ln 2 \leq x \leq \ln 3$

Solution

$$\begin{aligned} y = \ln(e^x - 1) - \ln(e^x + 1) &\Rightarrow \frac{dy}{dx} = \frac{e^x}{e^x - 1} - \frac{e^x}{e^x + 1} \\ &= \frac{e^{2x} + e^x - e^{2x} - e^x}{e^{2x} - 1} \\ &= \frac{2e^x}{e^{2x} - 1} \end{aligned}$$

$$\begin{aligned} L &= \int_{\ln 2}^{\ln 3} \sqrt{1 + \left(\frac{2e^x}{e^{2x} - 1}\right)^2} dx \\ &= \int_{\ln 2}^{\ln 3} \sqrt{1 + \frac{4e^{2x}}{e^{4x} - 2e^{2x} + 1}} dx \\ &= \int_{\ln 2}^{\ln 3} \sqrt{\frac{e^{4x} - 2e^{2x} + 1 + 4e^{2x}}{(e^{2x} - 1)^2}} dx \\ &= \int_{\ln 2}^{\ln 3} \sqrt{\frac{e^{4x} + 2e^{2x} + 1}{(e^{2x} - 1)^2}} dx \\ &= \int_{\ln 2}^{\ln 3} \sqrt{\frac{(e^{2x} + 1)^2}{(e^{2x} - 1)^2}} dx \\ &= \int_{\ln 2}^{\ln 3} \frac{e^{2x} + 1}{e^{2x} - 1} dx \\ &= \int_{\ln 2}^{\ln 3} \frac{e^{2x} + 1}{\frac{e^{2x}}{e^x} - 1} dx \\ &= \int_{\ln 2}^{\ln 3} \frac{e^{2x}}{\frac{e^{2x}}{e^x} - 1} + \frac{1}{\frac{e^{2x}}{e^x} - 1} dx \\ &= \int_{\ln 2}^{\ln 3} \frac{e^x + e^{-x}}{e^x - e^{-x}} dx \end{aligned}$$

$$\text{or Let } u = e^x - e^{-x} \Rightarrow du = (e^x + e^{-x}) dx$$

$$\begin{aligned}
&= \int_{\ln 2}^{\ln 3} \frac{1}{e^x - e^{-x}} d(e^x - e^{-x}) & d(e^x - e^{-x}) = (e^x + e^{-x}) dx \\
&= \left[\ln |e^x - e^{-x}| \right]_{\ln 2}^{\ln 3} \\
&= \ln \left(3 - \frac{1}{3} \right) - \ln \left(2 - \frac{1}{2} \right) \\
&= \ln \left(\frac{8}{3} \right) - \ln \left(\frac{3}{2} \right) \\
&= \ln \left(\frac{16}{9} \right) \text{ unit}
\end{aligned}$$

Exercise

Find the length of the curve $f(x) = \frac{2}{3}x^{3/2} - \frac{1}{2}x^{1/2}$ $1 \leq x \leq 4$

Solution

$$a = \frac{2}{3}, \quad m = \frac{3}{2}, \quad b = -\frac{1}{2}, \quad n = \frac{1}{2}$$

$$1. \quad m + n = \frac{3}{2} + \frac{1}{2} = 2 \quad \checkmark$$

$$2. \quad abmn = \frac{2}{3} \left(\frac{3}{2} \right) \left(-\frac{1}{2} \right) \left(\frac{1}{2} \right) = -\frac{1}{4} \quad \checkmark$$

$$\begin{aligned}
L &= \left(\frac{2}{3}x^{3/2} + \frac{1}{2}x^{1/2} \right) \Big|_1^4 \\
&= \frac{2}{3}4^{3/2} + 1 - \frac{2}{3} - \frac{1}{2} \\
&= \frac{16}{3} - \frac{2}{3} + \frac{1}{2} \\
&= \frac{31}{6} \text{ unit}
\end{aligned}$$

Exercise

Find the length of the curve $f(x) = x^3 + \frac{1}{12x}$ $1 \leq x \leq 4$

Solution

$$a = 1, \quad m = 3, \quad b = \frac{1}{12}, \quad n = -1$$

$$1. \quad m + n = 3 - 1 = 2 \quad \checkmark$$

$$2. \quad abmn = (1) \left(\frac{1}{12} \right) (3) (-1) = -\frac{1}{4} \quad \checkmark$$

$$\begin{aligned}
L &= \left(x^3 - \frac{1}{12x} \right) \Big|_1^4 \\
&= 4^3 - \frac{1}{48} - 1 + \frac{1}{12} \\
&= 63 + \frac{3}{48}
\end{aligned}$$

$$= \frac{3,027}{48} \text{ unit}$$

Exercise

Find the length of the curve $f(x) = \frac{1}{8}x^4 + \frac{1}{4x^2}$ $1 \leq x \leq 10$

Solution

$$a = \frac{1}{8}, \quad m = 4, \quad b = \frac{1}{4}, \quad n = -2$$

$$1. \quad m + n = 4 - 2 = 2 \quad \checkmark$$

$$2. \quad abmn = \left(\frac{1}{8}\right)\left(\frac{1}{4}\right)(4)(-2) = -\frac{1}{4} \quad \checkmark$$

$$\begin{aligned} L &= \left(\frac{1}{8}x^4 - \frac{1}{4x^2} \right) \Big|_1^{10} \\ &= \frac{10^4}{8} - \frac{1}{400} - \frac{1}{8} + \frac{1}{4} \\ &= \frac{9,999}{8} + \frac{99}{400} \\ &= \frac{9}{8} \left(1111 + \frac{11}{50} \right) \\ &= \frac{9}{8} \left(\frac{55,561}{50} \right) \\ &= \frac{500,049}{400} \text{ unit} \end{aligned}$$

Exercise

Find the length of the curve $f(x) = \frac{1}{4}x^4 + \frac{1}{8x^2}$ $3 \leq x \leq 8$

Solution

$$a = \frac{1}{4}, \quad m = 4, \quad b = \frac{1}{8}, \quad n = -2$$

$$1. \quad m + n = 4 - 2 = 2 \quad \checkmark$$

$$2. \quad abmn = \left(\frac{1}{4}\right)\left(\frac{1}{8}\right)(4)(-2) = -\frac{1}{4} \quad \checkmark$$

$$\begin{aligned} L &= \left(\frac{1}{4}x^4 - \frac{1}{8x^2} \right) \Big|_3^8 \\ &= \frac{8^4}{4} - \frac{1}{8^3} - \frac{81}{4} + \frac{1}{72} \\ &= \frac{4,015}{4} - \frac{1}{512} + \frac{1}{72} \\ &= \frac{1}{4} \left(4,015 - \frac{1}{128} + \frac{1}{18} \right) \end{aligned}$$

$$= \frac{1}{4} \left(4,015 + \frac{55}{1152} \right)$$

$$= \frac{4,625,335}{4,608} \text{ unit}$$

Exercise

Find the length of the curve $f(x) = \frac{1}{10}x^5 + \frac{1}{6x^3}$ $1 \leq x \leq 7$

Solution

$$a = \frac{1}{10}, \quad m = 5, \quad b = \frac{1}{6}, \quad n = -3$$

$$1. \quad m + n = 5 - 3 = 2 \quad \checkmark$$

$$2. \quad abmn = \left(\frac{1}{10}\right)\left(\frac{1}{6}\right)(5)(-3) = -\frac{1}{4} \quad \checkmark$$

$$L = \left(\frac{1}{10}x^5 - \frac{1}{6x^3} \right) \Big|_1^7$$

$$= \frac{7^5}{10} - \frac{1}{2058} - \frac{1}{10} + \frac{1}{6}$$

$$= \frac{8403}{5} + \frac{57}{343}$$

$$= \frac{2,882,514}{1,715} \text{ unit}$$

Exercise

Find the length of the curve $f(x) = \frac{3}{10}x^{1/3} - \frac{3}{2}x^{5/3}$ $0 \leq x \leq 12$

Solution

$$a = \frac{1}{10}, \quad m = 5, \quad b = \frac{1}{6}, \quad n = -3$$

$$1. \quad m + n = 5 - 3 = 2 \quad \checkmark$$

$$2. \quad abmn = \left(\frac{1}{10}\right)\left(\frac{1}{6}\right)(5)(-3) = -\frac{1}{4} \quad \checkmark$$

$$L = \left(\frac{3}{10}x^{1/3} + \frac{3}{2}x^{5/3} \right) \Big|_0^{12}$$

$$= \frac{3}{10}\sqrt[3]{12} + \frac{3}{2}12\sqrt[3]{144}$$

$$= \frac{3}{10}\sqrt[3]{12} + 18\sqrt[3]{144} \text{ unit}$$

$$= \frac{3}{10}\sqrt[3]{12} \left(1 + 600\sqrt[3]{12} \right)$$

Exercise

Find the length of the curve $f(x) = x^{1/2} - \frac{1}{3}x^{3/2}$ $2 \leq x \leq 9$

Solution

$$a = 1, \quad m = \frac{1}{2}, \quad b = -\frac{1}{3}, \quad n = \frac{3}{2}$$

$$1. \quad m + n = \frac{1}{2} + \frac{3}{2} = 2 \quad \checkmark \qquad 2. \quad abmn = (1)\left(-\frac{1}{3}\right)\left(\frac{1}{2}\right)\left(\frac{3}{2}\right) = -\frac{1}{4} \quad \checkmark$$

$$\begin{aligned} L &= \left(x^{1/2} + \frac{1}{3}x^{3/2} \right) \Big|_2^9 \\ &= 3 + 9 - \sqrt{2} - \frac{2\sqrt{2}}{3} \\ &= \frac{1}{3}(36 - 5\sqrt{2}) \text{ unit} \qquad = \frac{3}{10}\sqrt[3]{12}(1 + 600\sqrt[3]{12}) \end{aligned}$$

Exercise


Find the length of the curve $x = \int_0^y \sqrt{\sec^4 t - 1} \, dt$ $-\frac{\pi}{4} \leq y \leq \frac{\pi}{4}$

Solution

$$\frac{dx}{dy} = \sqrt{\sec^4 y - 1}$$

$$\begin{aligned} \sqrt{1 + \left(\frac{dx}{dy}\right)^2} &= \sqrt{1 + \sec^4 y - 1} \\ &= \sqrt{\sec^4 y} \\ &= \sec^2 y \end{aligned}$$

$$\begin{aligned} L &= \int_{-\pi/4}^{\pi/4} \sec^2 y \, dy \\ &= \tan y \Big|_{-\pi/4}^{\pi/4} \\ &= 1 - (-1) \\ &= 2 \text{ unit} \end{aligned}$$



Exercise

Find the length of the curve $y = 3 - 2x$ $0 \leq x \leq 2$. Check your answer by finding the length of the segment as the hypotenuse of a right triangle.

Solution

$$\frac{dy}{dx} = -2$$

$$\sqrt{1 + \left(\frac{dy}{dx}\right)^2} = \sqrt{1 + 4} = \sqrt{5}$$


$$L = \int_0^2 \sqrt{5} \, dx$$

$$= \sqrt{5}x \Big|_0^2$$

$$= \underline{2\sqrt{5} \text{ unit}}$$

$$\begin{cases} x = 0 & \rightarrow y = 3 \\ x = 2 & \rightarrow y = -1 \end{cases}$$

$$d = \sqrt{(2-0)^2 + (3+1)^2} = \sqrt{4+16} = \sqrt{20} = \underline{2\sqrt{5} \text{ unit}}$$



Exercise

Find a curve through the origin in the xy -plane whose length from $x = 0$ to $x = 1$ is $L = \int_0^1 \sqrt{1 + \frac{1}{4}e^x} \, dx$

Solution

$$L = \int_0^1 \sqrt{1 + \frac{1}{4}e^x} \, dx \Rightarrow \frac{dy}{dx} = \frac{e^{x/2}}{2} \quad \rightarrow \quad dy = \frac{e^{x/2}}{2} \, dx$$

$$y = \int \frac{e^{x/2}}{2} \, dx = e^{x/2} + C$$

$$0 = e^{0/2} + C$$

$$0 = 1 + C \quad \Rightarrow \quad \underline{C = -1}$$

$$\underline{y = e^{x/2} - 1}$$

Exercise


Confirm that the circumference of a circle of radius a is $2\pi a$.

Solution

$$f(x) = \sqrt{a^2 - x^2} \quad \text{for } -a \leq x \leq a$$

$$f'(x) = -\frac{x}{\sqrt{a^2 - x^2}} \quad \text{but } x \neq \pm a$$

$$\begin{aligned} \sqrt{1 + f'(x)^2} &= \sqrt{1 + \frac{x^2}{a^2 - x^2}} \\ &= \frac{a}{\sqrt{a^2 - x^2}} \end{aligned}$$




Let's compute the length of $\frac{1}{8}$ of the circle on $\left[0, \frac{a}{\sqrt{2}}\right]$

$$\begin{aligned} L &= 8a \int_0^{a/\sqrt{2}} \frac{dx}{\sqrt{a^2 - x^2}} \\ &= 8a \sin^{-1}\left(\frac{x}{a}\right) \Big|_0^{a/\sqrt{2}} \\ &= 8a \sin^{-1}\left(\frac{1}{\sqrt{2}}\right) \\ &= 8a \left(\frac{\pi}{4}\right) \\ &= 2\pi a \text{ unit} \end{aligned}$$

Exercise

The profile of the cables on a suspension bridge may be modeled by a parabola. The central span of the Golden Gate Bridge is 1280 m long and 152 m high. The parabola $y = 0.00037x^2$ gives a good fit to the shape of the cables, where $|x| \leq 640$, and x and y are measured in *meters*. Approximate the length of the cables that stretch between the tops of the two towers.



Solution

$$y' = 0.00074x$$

$$\begin{aligned}
 L &= \int_{-640}^{640} \sqrt{1 + (.00074x)^2} \, dx & \int \sqrt{a^2 + x^2} \, dx &= \frac{x}{2} \sqrt{a^2 + x^2} + \frac{a^2}{2} \ln \left| x + \sqrt{a^2 + x^2} \right| \\
 &= \left(\frac{x}{2} \sqrt{1 + x^2} + \frac{1}{2} \ln \left| x + \sqrt{1 + x^2} \right| \right) \bigg|_{-640}^{640} \\
 &= 320 \sqrt{1 + 640^2} + \frac{1}{2} \ln \left| 640 + \sqrt{1 + 640^2} \right| + 320 \sqrt{1 + x^2} - \frac{1}{2} \ln \left| -640 + \sqrt{1 + 640^2} \right| \\
 &\approx \underline{1326.4 \, m}
 \end{aligned}$$

Exercise

Electrical wires suspended between two towers form a catenary modeled by the equation

$$y = 20 \cosh \frac{x}{20}, \quad -20 \leq x \leq 20$$


Where x and y are measured in *meters*. The towers are 40 *meters* apart. Find the length of the suspended cable.

Solution

$$y = 20 \cosh \frac{x}{20} \rightarrow y' = \sinh \frac{x}{20}$$

$$\begin{aligned}
 \sqrt{1 + (y')^2} &= \sqrt{1 + \sinh^2 \frac{x}{20}} \\
 &= \sqrt{\cosh^2 \frac{x}{20}} \\
 &= \cosh \frac{x}{20}
 \end{aligned}$$

$$\begin{aligned}
 L &= \int_{-20}^{20} \cosh \frac{x}{20} \, dx \\
 &= 2(20) \sinh \frac{x}{20} \bigg|_0^{20} \\
 &= 40(\sinh 1 - \sinh 0) \\
 &= \underline{40 \sinh 1}
 \end{aligned}$$



Exercise

A barn is 100 *feet* long and 40 *feet* wide. A cross section of the roof is the inverted catenary

$y = 31 - 10(e^{x/20} + e^{-x/20})$. Find the number of **square feet** of roofing on the barn.


Solution

$$a = 10, \quad m = \frac{1}{20}, \quad b = 10, \quad n = -\frac{1}{20}$$

$$1. \quad m = -n \quad \checkmark$$

$$2. \quad abmn = 10(10)\left(\frac{1}{20}\right)\left(-\frac{1}{20}\right) = -\frac{1}{4} \quad \checkmark$$

$$\begin{aligned}
 L &= 10 \left(e^{x/20} - e^{-x/20} \right) \Big|_{-20}^{20} \\
 &= 10 \left(e - \frac{1}{e} - \frac{1}{e} + e \right) \\
 &= 20 \left(e - \frac{1}{e} \right) \approx 47 \text{ ft}
 \end{aligned}$$



\therefore There are $100(47) = 4,700 \text{ ft}^2$ of roofing on the barn

Exercise

A cable for a suspension bridge has the shape of a parabola with equation $y = kx^2$. Let h represent the height of the cable from its lowest point to its highest point and let $2w$ represent the total span of the bridge.

Show that the length C of the cable is given by $C = 2 \int_0^w \sqrt{1 + \frac{4h^2}{w^4} x^2} dx$

Solution


$$y' = 2kx$$

$$\sqrt{1 + (y')^2} = \sqrt{1 + 4k^2 x^2}$$

$$\text{At } (w, h) \rightarrow h = kw^2 \Rightarrow k = \frac{h}{w^2}$$

$$\sqrt{1 + (y')^2} = \sqrt{1 + \frac{4h^2}{w^4} x^2}$$

$$\therefore \text{By symmetry: } C = 2 \int_0^w \sqrt{1 + \frac{4h^2}{w^4} x^2} dx$$



Exercise


Find the total length of the graph of the astroid $x^{2/3} + y^{2/3} = 4$

Solution

$$x^{2/3} + y^{2/3} = 4 \Rightarrow y = (4 - x^{2/3})^{3/2}$$

$$\begin{aligned}
 y' &= \frac{3}{2} \left(-\frac{2}{3} x^{-1/3} \right) (4 - x^{2/3})^{1/2} \\
 &= -\frac{1}{x^{1/3}} (4 - x^{2/3})^{1/2}
 \end{aligned}$$

$$1 + (y')^2 = 1 + \frac{1}{x^{2/3}} (4 - x^{2/3})$$



$$= \frac{4}{x^{2/3}}$$

$$y = 0 \rightarrow x^{2/3} = 4 \Rightarrow \underline{x = 4^{3/2} = 8}$$

$$L = 4 \int_0^8 \sqrt{\frac{4}{x^{2/3}}} dx$$

$$= 8 \int_0^8 x^{-1/3} dx$$

$$= 12x^{2/3} \Big|_0^8$$

$$= 12(4 - 0)$$

$$\underline{= 48}$$

Exercise

Find the arc length from $(0, 3)$ clockwise to $(2, \sqrt{5})$ along the circle $x^2 + y^2 = 9$

Solution

$$y = \sqrt{9 - x^2} \Rightarrow y' = -\frac{x}{\sqrt{9 - x^2}}$$

$$\sqrt{1 + (y')^2} = \sqrt{1 + \frac{x^2}{9 - x^2}}$$


$$= \sqrt{\frac{9}{9 - x^2}}$$

$$= \frac{3}{\sqrt{9 - x^2}}$$

$$L = \int_0^2 \frac{3}{\sqrt{9 - x^2}} dx$$

$$= 3 \arcsin \frac{x}{3} \Big|_0^2$$

$$\underline{= 3 \arcsin \frac{2}{3}} \approx \underline{2.1892}$$



Exercise

Find the arc length from $(-3, 4)$ clockwise to $(4, 3)$ along the circle $x^2 + y^2 = 25$. Show that the result is one-fourth the circumference of the circle.

Solution

$$y = \sqrt{25 - x^2} \Rightarrow y' = -\frac{x}{\sqrt{25 - x^2}}$$

$$\begin{aligned}\sqrt{1 + (y')^2} &= \sqrt{1 + \frac{x^2}{25 - x^2}} \\ &= \sqrt{\frac{25}{25 - x^2}} \\ &= \frac{5}{\sqrt{25 - x^2}}\end{aligned}$$

$$\begin{aligned}L &= \int_{-3}^4 \frac{5}{\sqrt{25 - x^2}} dx \\ &= 5 \arcsin \frac{x}{5} \Big|_{-3}^4 \\ &= 5 \left(\arcsin \frac{4}{5} + \arcsin \frac{3}{5} \right) \Big| \approx 7.854\end{aligned}$$

