

Solution **Section 3.9 – Eigenvalues and Eigenvectors**

Exercise

Find the eigenvalues and eigenvectors of A , A^2 , A^{-1} , and $A + 4I$:

$$A = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix} \quad \text{and} \quad A^2 = \begin{bmatrix} 5 & -4 \\ -4 & 5 \end{bmatrix}$$

Check the trace $\lambda_1 + \lambda_2$ and the determinant $\lambda_1 \lambda_2$ for A and also A^2 .

Solution

For A:

$$\begin{aligned} \det(A - \lambda I) &= \begin{vmatrix} 2-\lambda & -1 \\ -1 & 2-\lambda \end{vmatrix} \\ &= (2-\lambda)^2 - 1 \\ &= \lambda^2 - 4\lambda + 3 = 0 \end{aligned}$$

The eigenvalues of A are $\lambda_1 = 1$ and $\lambda_2 = 3$.

The trace of a square matrix A is the sum of the elements on the main diagonal: $2 + 2$ agrees with $1 + 3$. The $\det(A) = 3$ agrees with the product $\lambda_1 \lambda_2$.

The eigenvectors for A are:

$$\lambda_1 = 1: (A - \lambda_1 I)V_1 = 0$$

$$\begin{pmatrix} 2-1 & -1 \\ -1 & 2-1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} x - y = 0 \\ -x + y = 0 \end{cases} \Rightarrow x = y$$

$$\text{Therefore the eigenvector } V_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$\lambda_2 = 3: (A - \lambda_2 I)V_2 = 0$$

$$\begin{pmatrix} -1 & -1 \\ -1 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} -x - y = 0 \\ -x - y = 0 \end{cases} \Rightarrow x = -y$$

$$\text{Therefore the eigenvector } V_2 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

For A^2 :

$$\det(A^2 - \lambda I) = \begin{vmatrix} 5-\lambda & -4 \\ -4 & 5-\lambda \end{vmatrix} = (5-\lambda)^2 - 16 = \lambda^2 - 10\lambda + 9 = 0$$

The eigenvalues of A^2 are $\lambda_1 = 1$ and $\lambda_2 = 9$. **Or** $\lambda_1 = 1^2 = 1$ and $\lambda_2 = 3^2 = 9$

$$\begin{cases} \text{tr}(A) = 5 + 5 = 10 \\ \lambda_1 + \lambda_2 = 1 + 9 = 10 \end{cases} \Rightarrow \text{tr}(A) = \lambda_1 + \lambda_2$$

$$\begin{cases} |A^2| = \begin{vmatrix} 5 & -4 \\ -4 & 5 \end{vmatrix} = 9 \\ \lambda_1 \lambda_2 = 1(9) = 9 \end{cases} \Rightarrow |A^2| = \lambda_1 \lambda_2$$

$$\lambda_1 = 1: (A^2 - \lambda_1 I)V_1 = 0$$

$$\begin{pmatrix} 4 & -4 \\ -4 & 4 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} 4x - 4y = 0 \\ -4x + 4y = 0 \end{cases} \Rightarrow x = y$$

Therefore the eigenvector $V_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$

$$\lambda_2 = 9: (A^2 - \lambda_2 I)V_2 = 0$$

$$\begin{pmatrix} -4 & -4 \\ -4 & -4 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} -4x - 4y = 0 \\ -4x - 4y = 0 \end{cases} \Rightarrow x = y$$

Therefore the eigenvector $V_2 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$

For A^{-1} :

$$A^{-1} = \frac{1}{3} \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} = \begin{pmatrix} \frac{2}{3} & \frac{1}{3} \\ \frac{1}{3} & \frac{2}{3} \end{pmatrix}$$

$$\det(A^{-1} - \lambda I) = \begin{vmatrix} \frac{2}{3} - \lambda & \frac{1}{3} \\ \frac{1}{3} & \frac{2}{3} - \lambda \end{vmatrix} = \left(\frac{2}{3} - \lambda\right)^2 - \frac{1}{9} = \lambda^2 - \frac{4}{3}\lambda + \frac{1}{3} = 0$$

The eigenvalues of A^{-1} are $\lambda_1 = 1$ and $\lambda_2 = \frac{1}{3}$.

$$\lambda_1 = 1: (A^{-1} - \lambda_1 I)V_1 = 0$$

$$\begin{pmatrix} \frac{2}{3}-1 & \frac{1}{3} \\ \frac{1}{3} & \frac{2}{3}-1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} -\frac{1}{3} & \frac{1}{3} \\ \frac{1}{3} & -\frac{1}{3} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Rightarrow \begin{cases} -\frac{1}{3}x + \frac{1}{3}y = 0 \\ \frac{1}{3}x - \frac{1}{3}y = 0 \end{cases} \rightarrow \boxed{x = y}$$

Therefore the eigenvector $V_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$

$$\lambda_2 = \frac{1}{3} : (A^{-1} - \lambda_2 I)V_2 = 0$$

$$\begin{pmatrix} \frac{2}{3}-\frac{1}{3} & \frac{1}{3} \\ \frac{1}{3} & \frac{2}{3}-\frac{1}{3} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} \frac{1}{3} & \frac{1}{3} \\ \frac{1}{3} & \frac{1}{3} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Rightarrow \begin{cases} \frac{1}{3}x + \frac{1}{3}y = 0 \\ \frac{1}{3}x + \frac{1}{3}y = 0 \end{cases} \rightarrow \boxed{x = -y}$$

Therefore the eigenvector $V_2 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$

For $A + 4I$:

$$A + 4I = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} + \begin{pmatrix} 4 & 0 \\ 0 & 4 \end{pmatrix} = \begin{pmatrix} 6 & 1 \\ 1 & 6 \end{pmatrix}$$

$$\det(A^{-1} - \lambda I) = \begin{vmatrix} 6-\lambda & 1 \\ 1 & 6-\lambda \end{vmatrix} = (6-\lambda)^2 - 1 = \lambda^2 - 12\lambda + 35 = 0$$

The eigenvalues of A^{-1} are $\lambda_1 = 5$ and $\lambda_2 = 7$.

$$\lambda_1 = 5 : (A + 4I - \lambda_1 I)V_1 = 0$$

$$\begin{pmatrix} 6-5 & 1 \\ 1 & 6-5 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Rightarrow \begin{cases} x + y = 0 \\ x + y = 0 \end{cases} \rightarrow \boxed{x = -y}$$

Therefore the eigenvector $V_1 = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$

$$\lambda_2 = 7 : (A + 4I - \lambda_2 I)V_2 = 0$$

$$\begin{pmatrix} -1 & 1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Rightarrow \begin{cases} -x + \frac{1}{3}y = 0 \\ x - y = 0 \end{cases} \rightarrow \boxed{x = y}$$

Therefore the eigenvector $V_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$

The eigenvalues $(A) = \lambda$

The eigenvalues $(A^2) = \lambda^2$

The eigenvalues $(A^{-1}) = \frac{1}{\lambda}$

Exercise

Show directly that the given vectors are eigenvectors of the given matrix. What are the corresponding eigenvalues

$$v_1 = \begin{bmatrix} 1 \\ -3 \end{bmatrix}, \quad v_2 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \quad A = \begin{bmatrix} 1 & -2 \\ -3 & 6 \end{bmatrix}$$

Solution

$$Av_1 = \begin{bmatrix} 1 & -2 \\ -3 & 6 \end{bmatrix} \begin{bmatrix} 1 \\ -3 \end{bmatrix} = \begin{bmatrix} 7 \\ -21 \end{bmatrix} = 7 \begin{bmatrix} 1 \\ -3 \end{bmatrix} = 7v_1$$

$v_1 = \begin{bmatrix} 1 \\ -3 \end{bmatrix}$ is an eigenvectors corresponding to the eigenvalue 7.

$$Av_2 = \begin{bmatrix} 1 & -2 \\ -3 & 6 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} = 0 \begin{bmatrix} 2 \\ 1 \end{bmatrix} = 0v_2$$

$v_2 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ is an eigenvectors corresponding to the eigenvalue 0.

$$\begin{aligned} |A - \lambda I| &= \begin{vmatrix} 1-\lambda & -2 \\ -3 & 6-\lambda \end{vmatrix} \\ &= (1-\lambda)(6-\lambda) - 6 \\ &= 6 - 7\lambda + \lambda^2 - 6 \\ &= \lambda^2 - 7\lambda = 0 \end{aligned}$$

The eigenvalues are: $\lambda_1 = 0$ and $\lambda_2 = 7$

Exercise

For which real numbers c does this matrix A have

$$A = \begin{pmatrix} 2 & -c \\ -1 & 2 \end{pmatrix}$$

- a) Two real eigenvalues and eigenvectors.
- b) A repeated eigenvalue with only one eigenvector
- c) Two complex eigenvalues and eigenvectors.

Solution

$$\begin{aligned} |A - \lambda I| &= \begin{vmatrix} 2-\lambda & -c \\ -1 & 2-\lambda \end{vmatrix} \\ &= (2-\lambda)^2 - c \\ &= \lambda^2 - 4\lambda + 4 - c = 0 \end{aligned}$$

$$\lambda = \frac{-(-4) \pm \sqrt{(-4)^2 - 4(1)(-c)}}{2(1)}$$

$$\lambda_{1,2} = \frac{4 \pm \sqrt{16 + 4c}}{2}$$

- a) Two real eigenvalues and eigenvectors, when $16 + 4c > 0 \rightarrow 4c > -16 \Rightarrow \boxed{c > -4}$
- b) A repeated eigenvalue with only one eigenvector, when $16 + 4c = 0 \Rightarrow \boxed{c = -4}$
- c) Two complex eigenvalues and eigenvectors, when $16 + 4c < 0 \Rightarrow \boxed{c < -4}$

Exercise

Find the eigenvalues of A , B , AB , and BA :

$$A = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \quad B = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \quad AB = \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} \quad BA = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}$$

- a) The eigenvalues of AB (are equal to) (are not equal to) eigenvalues of A times eigenvalues of B .
- b) The eigenvalues of AB (are equal to) (are not equal to) eigenvalues of BA .

Solution

Since A is a lower triangular, then $\lambda_1 = \lambda_2 = 1$

Since B is a upper triangular, then $\lambda_1 = \lambda_2 = 1$

$$\det(AB - I) = \begin{vmatrix} 1-\lambda & 1 \\ 1 & 2-\lambda \end{vmatrix} = (1-\lambda)(2-\lambda) - 1 = \lambda^2 - 3\lambda + 1 = 0 \quad \lambda_1 = \frac{3-\sqrt{5}}{2} \quad \lambda_2 = \frac{3+\sqrt{5}}{2}$$

$$\det(BA - I) = \begin{vmatrix} 2-\lambda & 1 \\ 1 & 1-\lambda \end{vmatrix} = (2-\lambda)(1-\lambda) - 1 = \lambda^2 - 3\lambda + 1 = 0 \quad \lambda_1 = \frac{3-\sqrt{5}}{2} \quad \lambda_2 = \frac{3+\sqrt{5}}{2}$$

- a) The eigenvalues of \mathbf{AB} are not equal to eigenvalues of \mathbf{A} times eigenvalues of \mathbf{B} .
b) The eigenvalues of \mathbf{AB} are equal to the eigenvalues of \mathbf{BA} .

Exercise

When $a + b = c + d$ show that $(1, 1)$ is an eigenvector and find both eigenvalues of

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

Solution

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} a+b \\ c+d \end{pmatrix} = \begin{pmatrix} a+b \\ a+b \end{pmatrix} = (a+b) \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$\text{If } a+b = c+d = \lambda_1$$

$$\text{tr}(A) = a+d = \lambda_1 + \lambda_2$$

$$\lambda_2 = (a+d) - \lambda_1$$

$$= a+d - (a+b)$$

$$= a+d - a - b$$

$$= d-b \quad \text{or} \quad = a-c$$

The eigenvalues for λ_2 :

$$\begin{pmatrix} a-\lambda_2 & b \\ c & d-\lambda_2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} a-(a-c) & b \\ c & d-(d-b) \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} c & b \\ c & b \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \rightarrow \{cx + by = 0 \Rightarrow \boxed{cx = -by}\}$$

$$\text{The eigenvector: } \mathbf{V}_2 = \begin{pmatrix} b \\ -c \end{pmatrix}$$

Exercise

The eigenvalues of A equal to the eigenvalues of A^T . This is because $\det(A - \lambda I)$ equals $\det(A^T - \lambda I)$.

That is true because _____. Show by an example that the eigenvectors of A and A^T are not the same.

Solution

$$\det(A - \lambda I) = \det(A - \lambda I)^T = \det(A^T - (\lambda I)^T) = \det(A^T - \lambda I)$$

Therefore, A and A^T have the same eigenvalues.

Let consider the matrix: $A = \begin{pmatrix} 0 & 1 \\ 4 & 0 \end{pmatrix} \Rightarrow A^T = \begin{pmatrix} 0 & 4 \\ 1 & 0 \end{pmatrix}$

$$|A - \lambda I| = \begin{vmatrix} -\lambda & 1 \\ 4 & -\lambda \end{vmatrix} = \lambda^2 - 4 = 0$$

The eigenvalues are: $\lambda = \pm 2$

For $\lambda = 2$

$$(A - \lambda_1 I)V_1 = \begin{pmatrix} -2 & 1 \\ 4 & -2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} -2x + y = 0 \\ 4x - 2y = 0 \end{cases} \Rightarrow y = 2x$$

$$V_1 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$$

$$(A^T - \lambda_1 I)V_1 = \begin{pmatrix} -2 & 4 \\ 1 & -2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} -2x + 4y = 0 \\ x - 2y = 0 \end{cases} \Rightarrow x = 2y$$

$$V_1 = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$$

Exercise

Let $A = \begin{bmatrix} 2 & -1 \\ 1 & 2 \end{bmatrix}$. Compute the eigenvalues and eigenvectors of A .

Solution

$$\det(A - \lambda I) = \begin{vmatrix} 2 - \lambda & -1 \\ 1 & 2 - \lambda \end{vmatrix} = (2 - \lambda)^2 + 1 = 0$$

$$(2 - \lambda)^2 = -1$$

$$2 - \lambda = \pm \sqrt{-1} = \pm i$$

The eigenvalues of A are: $\lambda = 2 \pm i$

For $\lambda_1 = 2 - i \Rightarrow (A - \lambda_1 I)V_1 = 0$

$$\begin{pmatrix} 2-(2-i) & -1 \\ 1 & 2-(2-i) \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} i & -1 \\ 1 & i \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} ix - y = 0 \\ x + iy = 0 \end{cases} \Rightarrow x = -iy$$

The eigenvector is: $V_1 = \begin{pmatrix} 1 \\ -i \end{pmatrix}$

For $\lambda_2 = 2+i \Rightarrow (A - \lambda_2 I)V_2 = 0$

$$\begin{pmatrix} -i & -1 \\ 1 & -i \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} -ix - y = 0 \\ x - iy = 0 \end{cases} \Rightarrow x = iy$$

The eigenvector is: $V_2 = \begin{pmatrix} i \\ 1 \end{pmatrix}$

Exercise

Let $A = \begin{bmatrix} 2 & -2 & 3 \\ 1 & 1 & 1 \\ 1 & 3 & -1 \end{bmatrix}$

- What is the characteristic polynomial for A (i.e. compute $\det(A - \lambda I)$)?
- Verify that 1 is an eigenvalue of A . What is a corresponding eigenvector?
- What are the other eigenvalues of A ?

Solution

$$\begin{aligned} a) \quad \det(A - \lambda I) &= \begin{vmatrix} 2-\lambda & -2 & 3 \\ 1 & 1-\lambda & 1 \\ 1 & 3 & -1-\lambda \end{vmatrix} \\ &= (2-\lambda)(1-\lambda)(-1-\lambda) - 2 + 9 - 3(1-\lambda) - 3(2-\lambda) + 2(-1-\lambda) \\ &= (2-3\lambda+\lambda^2)(-1-\lambda) + 7 - 3 + 3\lambda - 6 + 3\lambda - 2 - 2\lambda \\ &= -2 + 3\lambda - \lambda^2 - 2\lambda + 3\lambda^2 - \lambda^3 + 4\lambda - 4 \\ &= \underline{-\lambda^3 + 2\lambda^2 + 5\lambda - 6} \end{aligned}$$

$$\begin{aligned} b) \quad \text{If } \lambda = 1 &\rightarrow -\lambda^3 + 2\lambda^2 + 5\lambda - 6 = 0 \\ &\quad -1^3 + 2(1)^2 + 5(1) - 6 = 0 \\ &\quad -1 + 2 + 5 - 6 = 0 \\ &\quad \boxed{0 = 0} \end{aligned}$$

1 is an eigenvalue of A .

$$\begin{pmatrix} 2-\lambda & -2 & 3 \\ 1 & 1-\lambda & 1 \\ 1 & 3 & -1-\lambda \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = 0$$

$$\begin{pmatrix} 1 & -2 & 3 \\ 1 & 0 & 1 \\ 1 & 3 & -2 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = 0 \rightarrow \begin{cases} x - 2y + 3z = 0 \\ x + z = 0 \\ x + 3y - 2z = 0 \end{cases}$$

$$\Rightarrow \begin{cases} \boxed{x = -z} \\ 3y = 2z - x = 2z + z = 3z \Rightarrow \boxed{y = z} \end{cases}$$

The eigenvector for $\lambda = 1$ is $V = \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix}$

c) $-\lambda^3 + 2\lambda^2 + 5\lambda - 6 = 0 \Rightarrow \underline{\lambda_1 = 1 \quad \lambda_2 = -2 \quad \lambda_3 = 3}$

Exercise

For the matrix:

$$a) \begin{pmatrix} 3 & 0 \\ 8 & -1 \end{pmatrix}$$

$$b) \begin{pmatrix} 10 & -9 \\ 4 & -2 \end{pmatrix}$$

$$c) \begin{pmatrix} 0 & 3 \\ 4 & 0 \end{pmatrix}$$

$$d) \begin{bmatrix} -2 & -7 \\ 1 & 2 \end{bmatrix}$$

$$e) \begin{pmatrix} 4 & 0 & 1 \\ -2 & 1 & 0 \\ -2 & 0 & 1 \end{pmatrix}$$

$$f) \begin{pmatrix} 3 & 0 & -5 \\ \frac{1}{5} & -1 & 0 \\ 1 & 1 & -2 \end{pmatrix}$$

$$g) \begin{pmatrix} -2 & 0 & 1 \\ -6 & -2 & 0 \\ 19 & 5 & -4 \end{pmatrix}$$

$$h) \begin{pmatrix} 0 & 0 & 2 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$i) \begin{pmatrix} 10 & -9 & 0 & 0 \\ 4 & -2 & 0 & 0 \\ 0 & 0 & -2 & -7 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

- i. Find the characteristic equation
- ii. Find the eigenvalues
- iii. Find the eigenvectors

Solution

a)

$$\begin{aligned} \text{i. } \det(A - \lambda I) &= \begin{vmatrix} 3-\lambda & 0 \\ 8 & -1-\lambda \end{vmatrix} \\ &= (3-\lambda)(-1-\lambda) - 0 \\ &= \lambda^2 - 2\lambda - 3 \end{aligned}$$

The characteristic equation: $\lambda^2 - 2\lambda - 3$

$$\text{ii. } \lambda^2 - 2\lambda - 3 = 0$$

The eigenvalues are $\lambda_1 = -1$ and $\lambda_2 = 3$

$$\text{iii. } \lambda_1 = -1 \rightarrow (A - \lambda_1 I)V_1 = 0$$

$$\begin{pmatrix} 4 & 0 \\ 8 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} 4x = 0 \\ 8x = 0 \end{cases} \Rightarrow x = 0$$

Therefore the eigenvector $V_1 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$

$$\lambda_2 = 3 \rightarrow (A - \lambda_2 I)V_2 = 0$$

$$\begin{pmatrix} 0 & 0 \\ 8 & -4 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} 0 = 0 \\ 8x - 4y = 0 \end{cases} \Rightarrow 8x = 4y \rightarrow \boxed{2x = y}$$

Therefore the eigenvector $V_2 = \begin{pmatrix} x \\ 2x \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$

b) For the matrix: $\begin{pmatrix} 10 & -9 \\ 4 & -2 \end{pmatrix}$

$$\begin{aligned} \text{i. } \det(A - \lambda I) &= \begin{vmatrix} 10 - \lambda & -9 \\ 4 & -2 - \lambda \end{vmatrix} \\ &= (10 - \lambda)(-2 - \lambda) + 36 \\ &= \lambda^2 - 8\lambda + 16 \end{aligned}$$

\Rightarrow The characteristic equation: $\lambda^2 - 8\lambda + 16$

$$\text{ii. } \lambda^2 - 8\lambda + 16 = 0$$

\Rightarrow The eigenvalues are $\lambda_{1,2} = 4$

$$\text{iii. } \lambda_1 = 4 \rightarrow (A - \lambda_1 I)V_1 = 0$$

$$\begin{pmatrix} 6 & -9 \\ 4 & -6 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} 6x - 9y = 0 \\ 4x - 6y = 0 \end{cases} \Rightarrow \boxed{2x = 3y}$$

Therefore the eigenvector $V_1 = \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix}$

c) For the matrix: $\begin{pmatrix} 0 & 3 \\ 4 & 0 \end{pmatrix}$

$$\begin{aligned} \text{i. } \det(A - \lambda I) &= \begin{vmatrix} -\lambda & 3 \\ 4 & -\lambda \end{vmatrix} \\ &= \lambda^2 - 12 \end{aligned}$$

\Rightarrow The characteristic equation: $\lambda^2 - 12$

$$\text{ii. } \lambda^2 - 12 = 0 \Rightarrow \lambda = \pm\sqrt{12}$$

The eigenvalues are $\lambda_{1,2} = \pm\sqrt{12}$

$$\text{iii. } \text{For } \lambda_1 = \sqrt{12} \rightarrow \begin{pmatrix} -\sqrt{12} & 3 \\ 4 & -\sqrt{12} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} -\sqrt{12} & 3 \\ 4 & -\sqrt{12} \end{pmatrix} \xrightarrow{rref} \begin{pmatrix} 1 & -\frac{3}{\sqrt{12}} \\ 0 & 0 \end{pmatrix} \Rightarrow x - \frac{3}{\sqrt{12}}y = 0$$

Therefore the eigenvector $V_1 = \begin{pmatrix} \frac{3}{\sqrt{12}} \\ 1 \end{pmatrix}$

For $\lambda_2 = -\sqrt{12} \rightarrow \begin{pmatrix} \sqrt{12} & 3 \\ 4 & \sqrt{12} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$

$$\begin{pmatrix} \sqrt{12} & 3 \\ 4 & \sqrt{12} \end{pmatrix} \xrightarrow{rref} \begin{pmatrix} 1 & \frac{3}{\sqrt{12}} \\ 0 & 0 \end{pmatrix} \Rightarrow x + \frac{3}{\sqrt{12}} y = 0$$

Therefore the eigenvector $V_2 = \begin{pmatrix} -\frac{3}{\sqrt{12}} \\ 1 \end{pmatrix}$

d) For the matrix $\begin{bmatrix} -2 & -7 \\ 1 & 2 \end{bmatrix}$

$$\begin{aligned} i. \quad \begin{vmatrix} -2-\lambda & -7 \\ 1 & 2-\lambda \end{vmatrix} &= (-2-\lambda)(2-\lambda) + 7 \\ &= -4 + \lambda^2 + 7 \\ &= \lambda^2 + 3 \end{aligned}$$

The characteristic equation: $\lambda^2 + 3 = 0$

ii. $\lambda^2 = -3 \rightarrow$ The eigenvalues $\lambda_{1,2} = \pm i\sqrt{3}$

iii. For $\lambda_1 = -i\sqrt{3}$, we have: $(A - \lambda_1 I)V_1 = 0$

$$\begin{pmatrix} -2+i\sqrt{3} & -7 \\ 1 & 2+i\sqrt{3} \end{pmatrix} \begin{pmatrix} x_1 \\ y_1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} (-2+i\sqrt{3})x_1 - 7y_1 = 0 \\ x_1 + (2+i\sqrt{3})y_1 = 0 \end{cases}$$

$$x_1 = -(2+i\sqrt{3})y_1$$

Therefore the eigenvector $V_1 = \begin{pmatrix} 2+i\sqrt{3} \\ -1 \end{pmatrix}$

For $\lambda_2 = i\sqrt{3}$, we have: $(A - \lambda_2 I)V_2 = 0$

$$\begin{pmatrix} -2-i\sqrt{3} & -7 \\ 1 & 2-i\sqrt{3} \end{pmatrix} \begin{pmatrix} x_2 \\ y_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} (-2-i\sqrt{3})x_2 - 7y_2 = 0 \\ x_2 + (2-i\sqrt{3})y_2 = 0 \end{cases}$$

$$x_2 = -(2-i\sqrt{3})y_2$$

Therefore the eigenvector $V_2 = \begin{pmatrix} 2-i\sqrt{3} \\ -1 \end{pmatrix}$

e) For the matrix:

$$\begin{pmatrix} 4 & 0 & 1 \\ -2 & 1 & 0 \\ -2 & 0 & 1 \end{pmatrix}$$

i. $\det(A - \lambda I) = \begin{vmatrix} 4-\lambda & 0 & 1 \\ -2 & 1-\lambda & 0 \\ -2 & 0 & 1-\lambda \end{vmatrix} = (4-\lambda)(1-\lambda)(1-\lambda) + 2(1-\lambda)$

$$= (1-\lambda)[(4-\lambda)(1-\lambda) + 2]$$

$$= (1-\lambda)(\lambda^2 - 5\lambda + 6) \Rightarrow \text{The characteristic equation: } \underline{-\lambda^3 + 6\lambda^2 - 11\lambda + 6}$$

ii. $-\lambda^3 + 6\lambda^2 - 11\lambda + 6 = 0 \Rightarrow$ The eigenvalues are $\boxed{\lambda = 1, 2, 3}$

iii. $\lambda_1 = 1 \rightarrow \begin{pmatrix} 3 & 0 & 1 \\ -2 & 0 & 0 \\ -2 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} 3x_1 + x_3 = 0 \\ x_1 = 0 \end{cases} \Rightarrow x_1 = x_3 = 0$

Therefore the eigenvector $V_1 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$

$$\lambda_2 = 2 \rightarrow \begin{pmatrix} 2 & 0 & 1 \\ -2 & -1 & 0 \\ -2 & 0 & -1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} 2x_1 + x_3 = 0 \\ -2x_1 - x_2 = 0 \\ -2x_1 - x_3 = 0 \end{cases} \Rightarrow \begin{cases} x_3 = -2x_1 \\ x_2 = -2x_1 \end{cases}$$

Therefore the eigenvector $V_2 = \begin{pmatrix} 1 \\ -2 \\ -2 \end{pmatrix}$

$$\lambda_3 = 3 \rightarrow \begin{pmatrix} 1 & 0 & 1 \\ -2 & -2 & 0 \\ -2 & 0 & -2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} x_1 + x_3 = 0 \\ -2x_1 - 2x_2 = 0 \\ -2x_1 - 2x_3 = 0 \end{cases} \Rightarrow \begin{cases} x_3 = -x_1 \\ x_2 = -x_1 \end{cases}$$

Therefore the eigenvector $V_3 = \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix}$

f) For the matrix:

$$\begin{pmatrix} 3 & 0 & -5 \\ \frac{1}{5} & -1 & 0 \\ 1 & 1 & -2 \end{pmatrix}$$

$$\begin{aligned}
 \text{i. } \det(A - \lambda I) &= \begin{vmatrix} 3-\lambda & 0 & -5 \\ \frac{1}{5} & -1-\lambda & 0 \\ 1 & 1 & -2-\lambda \end{vmatrix} \\
 &= (3-\lambda)(-1-\lambda)(-2-\lambda) - 1 + 5(-1-\lambda) \\
 &= (3-\lambda)(\lambda^2 + 3\lambda + 2) - 1 - 5 - 5\lambda \\
 &= 3\lambda^2 + 9\lambda + 6 - \lambda^3 - 3\lambda^2 - 2\lambda - 6 - 5\lambda \\
 &= -\lambda^3 + 2\lambda
 \end{aligned}$$

\Rightarrow The characteristic equation: $-\lambda^3 + 2\lambda$

$$\text{ii. } -\lambda^3 + 2\lambda = 0 \Rightarrow \text{The eigenvalues are } \boxed{\lambda = 0, \pm\sqrt{2}}$$

$$\text{iii. } \lambda_1 = -\sqrt{2} \rightarrow \begin{pmatrix} 3+\sqrt{2} & 0 & -5 \\ \frac{1}{5} & -1+\sqrt{2} & 0 \\ 1 & 1 & -2+\sqrt{2} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} (3+\sqrt{2})x_1 - 5x_3 = 0 \\ \frac{1}{5}x_1 + (-1+\sqrt{2})x_2 = 0 \\ x_1 + x_2 + (-2+\sqrt{2})x_3 = 0 \end{cases}$$

$$\rightarrow \begin{cases} x_3 = \frac{3+\sqrt{2}}{5}x_1 \\ (-1+\sqrt{2})x_2 = -\frac{1}{5}x_1 \Rightarrow x_2 = -\frac{1}{5(-1+\sqrt{2})}x_1 \end{cases}$$

$$\text{Therefore the eigenvector } V_1 = \begin{pmatrix} 1 \\ \frac{1}{5(1-\sqrt{2})} \\ \frac{3+\sqrt{2}}{5} \end{pmatrix}$$

$$\lambda_2 = 0 \rightarrow \begin{pmatrix} 3 & 0 & -5 \\ \frac{1}{5} & -1 & 0 \\ 1 & 1 & -2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} 3x_1 - 5x_3 = 0 \\ \frac{1}{5}x_1 - x_2 = 0 \\ x_1 + x_2 - 2x_3 = 0 \end{cases} \Rightarrow \begin{cases} x_3 = \frac{3}{5}x_1 \\ x_2 = \frac{1}{5}x_1 \end{cases}$$

$$\text{Therefore the eigenvector } V_2 = \begin{pmatrix} 5 \\ 1 \\ 3 \end{pmatrix}$$

$$\lambda_3 = \sqrt{2} \rightarrow \begin{pmatrix} 3-\sqrt{2} & 0 & -5 \\ \frac{1}{5} & -1-\sqrt{2} & 0 \\ 1 & 1 & -2-\sqrt{2} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} (3-\sqrt{2})x_1 - 5x_3 = 0 \\ \frac{1}{5}x_1 + (-1-\sqrt{2})x_2 = 0 \\ x_1 + x_2 + (-2-\sqrt{2})x_3 = 0 \end{cases}$$

$$\rightarrow \begin{cases} x_3 = \frac{3-\sqrt{2}}{5}x_1 \\ (-1-\sqrt{2})x_2 = -\frac{1}{5}x_1 \end{cases} \Rightarrow x_2 = \frac{1}{5(1+\sqrt{2})}x_1$$

$$\text{Therefore the eigenvector } V_3 = \begin{pmatrix} 1 \\ \frac{1}{5(1+\sqrt{2})} \\ \frac{3-\sqrt{2}}{5} \end{pmatrix}$$

g) For the matrix: $\begin{pmatrix} -2 & 0 & 1 \\ -6 & -2 & 0 \\ 19 & 5 & -4 \end{pmatrix}$

$$\begin{aligned} \text{i. } \det(A - \lambda I) &= \begin{vmatrix} -2-\lambda & 0 & 1 \\ -6 & -2-\lambda & 0 \\ 19 & 5 & -4-\lambda \end{vmatrix} \\ &= (-2-\lambda)^2(-4-\lambda) - 30 - 19(-2-\lambda) \\ &= (4 + 4\lambda + \lambda^2)(-4-\lambda) - 30 + 38 + 19\lambda \\ &= -16 - 16\lambda - 4\lambda^2 - 4\lambda - 4\lambda^2 - \lambda^3 + 8 + 19\lambda \\ &= -\lambda^3 - 8\lambda^2 - \lambda - 8 \end{aligned}$$

$$\Rightarrow \text{The characteristic equation: } \underline{-\lambda^3 - 8\lambda^2 - \lambda - 8}$$

$$\text{ii. } \lambda^3 + 8\lambda^2 + \lambda + 8 = 0 \Rightarrow (\lambda + 8)(\lambda^2 + 1) = 0$$

$$\lambda^3 + 8\lambda^2 + \lambda + 8 = 0 \Rightarrow \text{The eigenvalues are } \boxed{\lambda_{1,2,3} = -8, \pm i}$$

$$\text{iii. } \lambda_1 = -8 \rightarrow \begin{pmatrix} 6 & 0 & 1 \\ -6 & 6 & 0 \\ 19 & 5 & 4 \end{pmatrix} \begin{pmatrix} x_1 \\ y_1 \\ z_1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 6 & 0 & 1 \\ -6 & 6 & 0 \\ 19 & 5 & 4 \end{pmatrix} \xrightarrow{rref} \begin{pmatrix} 1 & 0 & \frac{1}{6} \\ 0 & 1 & \frac{1}{6} \\ 0 & 0 & 0 \end{pmatrix} \Rightarrow \begin{cases} x_1 + \frac{1}{6}z_1 = 0 \\ y_1 + \frac{1}{6}z_1 = 0 \end{cases}$$

Therefore the eigenvector $V_1 = \begin{pmatrix} -\frac{1}{6} \\ -\frac{1}{6} \\ 1 \end{pmatrix}$

For $\lambda_2 = -i \rightarrow \begin{pmatrix} -2+i & 0 & 1 \\ -6 & -2+i & 0 \\ 19 & 5 & -4+i \end{pmatrix} \begin{pmatrix} x_2 \\ y_2 \\ z_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$

$$\rightarrow \begin{cases} (-2+i)x_2 + z_2 = 0 \\ -6x_2 + (-2+i)y_2 = 0 \\ 19x_2 + 5y_2 + (-4+i)z_2 = 0 \end{cases}$$

Therefore the eigenvector $V_2 = \begin{pmatrix} 1 \\ -\frac{12}{5} - i\frac{6}{5} \\ 2-i \end{pmatrix}$

For $\lambda_3 = i \rightarrow \begin{pmatrix} -2-i & 0 & 1 \\ -6 & -2-i & 0 \\ 19 & 5 & -4-i \end{pmatrix} \begin{pmatrix} x_2 \\ y_2 \\ z_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$

$$\begin{cases} (-2-i)x_2 + z_2 = 0 \\ -6x_2 + (-2-i)y_2 = 0 \\ 19x_2 + 5y_2 + (-4-i)z_2 = 0 \end{cases}$$

Therefore the eigenvector $V_2 = \begin{pmatrix} 1 \\ -\frac{12}{5} + i\frac{6}{5} \\ 2+i \end{pmatrix}$

h) For the matrix:

$$\begin{pmatrix} 0 & 0 & 2 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$\begin{aligned}
 \text{i. } \det(A - \lambda I) &= \begin{vmatrix} -\lambda & 0 & 2 & 0 \\ 1 & -\lambda & 1 & 0 \\ 0 & 1 & -2-\lambda & 0 \\ 0 & 0 & 0 & 1-\lambda \end{vmatrix} = (1-\lambda) \begin{vmatrix} -\lambda & 0 & 2 \\ 1 & -\lambda & 1 \\ 0 & 1 & -2-\lambda \end{vmatrix} \\
 &= (1-\lambda)(\lambda^2(-2-\lambda) + 2 + \lambda) \\
 &= (1-\lambda)(-\lambda^3 - 2\lambda^2 + \lambda + 2) \\
 &= \lambda^4 + \lambda^3 - 3\lambda^2 - \lambda + 2
 \end{aligned}$$

\Rightarrow The characteristic equation: $\lambda^4 + \lambda^3 - 3\lambda^2 - \lambda + 2$

$$\text{ii. } \lambda^4 + \lambda^3 - 3\lambda^2 - \lambda + 2 = 0 \Rightarrow \text{The eigenvalues are } \boxed{\lambda = -2, -1, 1, 1}$$

$$\begin{aligned}
 \text{iii. } \lambda_1 = -2 \rightarrow \begin{pmatrix} 2 & 0 & 2 & 0 \\ 1 & 2 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} 2x_1 + 2x_3 = 0 \\ x_1 + 2x_2 + x_3 = 0 \\ x_2 = 0 \\ x_4 = 0 \end{cases} \\
 &\rightarrow \begin{cases} x_1 = -x_3 \\ x_1 = -x_3 \\ x_2 = 0 \\ x_4 = 0 \end{cases}
 \end{aligned}$$

Therefore the eigenvector $V_1 = \begin{pmatrix} -1 \\ 0 \\ 1 \\ 0 \end{pmatrix}$

$$\begin{aligned}
 \lambda_2 = -1 \rightarrow \begin{pmatrix} 1 & 0 & 2 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} x_1 + 2x_3 = 0 \\ x_1 + x_2 + x_3 = 0 \\ x_2 - x_3 = 0 \\ x_4 = 0 \end{cases} \\
 &\rightarrow \begin{cases} x_1 = -2x_3 \\ x_1 = -x_2 - x_3 \\ x_2 = x_3 \end{cases}
 \end{aligned}$$

Therefore the eigenvector $V_2 = \begin{pmatrix} -2 \\ 1 \\ 1 \\ 0 \end{pmatrix}$

$$\lambda_3 = 1 \rightarrow \begin{pmatrix} -1 & 0 & 2 & 0 \\ 1 & -1 & 1 & 0 \\ 0 & 1 & -3 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} -x_1 + 2x_3 = 0 \\ x_1 - x_2 + x_3 = 0 \\ x_2 - 3x_3 = 0 \end{cases}$$

$$\rightarrow \begin{cases} x_1 = 2x_3 \\ x_1 = x_2 - x_3 \\ x_2 = 3x_3 \end{cases}$$

Therefore the eigenvector $V_3 = \begin{pmatrix} 2 \\ 3 \\ 1 \\ 0 \end{pmatrix}$

$\lambda_4 = 1 \rightarrow$ Therefore the eigenvector $V_4 = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}$

i) For the matrix:

$$\begin{pmatrix} 10 & -9 & 0 & 0 \\ 4 & -2 & 0 & 0 \\ 0 & 0 & -2 & -7 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

i. $\det(A - \lambda I) = \begin{vmatrix} 10 - \lambda & -9 & 0 & 0 \\ 4 & -2 - \lambda & 0 & 0 \\ 0 & 0 & -2 - \lambda & -7 \\ 0 & 0 & 1 & 2 - \lambda \end{vmatrix}$

$$= (10 - \lambda) \begin{vmatrix} -2 - \lambda & 0 & 0 \\ 0 & -2 - \lambda & -7 \\ 0 & 1 & 2 - \lambda \end{vmatrix} + 9 \begin{vmatrix} 4 & 0 & 0 \\ 0 & -2 - \lambda & -7 \\ 0 & 1 & 2 - \lambda \end{vmatrix}$$

$$= (10 - \lambda) [(-2 - \lambda)^2(2 - \lambda) + 7(-2 - \lambda)] + 9[(4)(-2 - \lambda)(2 - \lambda) + 28]$$

$$= (10 - \lambda)(-2 - \lambda)(3 + \lambda^2) + 9(4\lambda^2 + 12)$$

$$= (3 + \lambda^2)(-8\lambda + \lambda^2 + 16)$$

$$= (3 + \lambda^2)(\lambda - 4)^2$$

\Rightarrow The characteristic equation: $\underline{(3 + \lambda^2)(\lambda - 4)^2}$

ii. $(3 + \lambda^2)(\lambda - 4)^2 = 0 \Rightarrow$ The eigenvalues are $\boxed{\lambda = 4, 4, \pm i\sqrt{3}}$

$$\text{iii. } \lambda_1 = 4 \rightarrow \begin{pmatrix} 6 & -9 & 0 & 0 \\ 4 & -6 & 0 & 0 \\ 0 & 0 & -6 & -7 \\ 0 & 0 & 1 & -2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} 6x_1 - 9x_2 = 0 \\ 4x_1 - 6x_2 = 0 \\ -6x_3 - 7x_4 = 0 \\ x_3 - 2x_4 = 0 \end{cases}$$

$$\begin{cases} 6x_1 = 9x_2 \\ 6x_3 = -7x_4 \\ x_3 = 2x_4 \end{cases} \Rightarrow \begin{cases} x_1 = \frac{3}{2}x_2 \\ x_3 = x_4 = 0 \end{cases}$$

Therefore the eigenvector $V_1 = \begin{pmatrix} \frac{3}{2} \\ 1 \\ 0 \\ 0 \end{pmatrix}$

$$\lambda_2 = 4 \rightarrow \text{Therefore the eigenvector } V_2 = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\lambda_3 = -i\sqrt{3} \rightarrow \begin{pmatrix} 10+i\sqrt{3} & -9 & 0 & 0 \\ 4 & -2+i\sqrt{3} & 0 & 0 \\ 0 & 0 & -2+i\sqrt{3} & -7 \\ 0 & 0 & 1 & 2+i\sqrt{3} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\rightarrow \begin{cases} (10+i\sqrt{3})x_1 - 9x_2 = 0 \\ 4x_1 + (-2+i\sqrt{3})x_2 = 0 \\ (-2+i\sqrt{3})x_3 - 7x_4 = 0 \\ x_3 + (2+i\sqrt{3})x_4 = 0 \end{cases}$$

$$\rightarrow \begin{cases} (10+i\sqrt{3})x_1 = 9x_2 \\ 4x_1 = -(-2+i\sqrt{3})x_2 \\ (-2+i\sqrt{3})x_3 = 7x_4 \\ x_3 = -(2+i\sqrt{3})x_4 \end{cases} \Rightarrow \begin{cases} x_1 = x_2 = 0 \\ x_3 = \frac{7}{-2+i\sqrt{3}}x_4 \left(\frac{-2-i\sqrt{3}}{-2-i\sqrt{3}} \right) = -(2+i\sqrt{3})x_4 \\ x_3 = -(2+i\sqrt{3})x_4 \end{cases}$$

Therefore the eigenvector $V_3 = \begin{pmatrix} 0 \\ 0 \\ -(2+i\sqrt{3}) \\ 1 \end{pmatrix}$

$$\lambda_4 = i\sqrt{3} \rightarrow \begin{pmatrix} 10-i\sqrt{3} & -9 & 0 & 0 \\ 4 & -2-i\sqrt{3} & 0 & 0 \\ 0 & 0 & -2-i\sqrt{3} & -7 \\ 0 & 0 & 1 & 2-i\sqrt{3} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\rightarrow \begin{cases} (10-i\sqrt{3})x_1 - 9x_2 = 0 \\ 4x_1 + (-2-i\sqrt{3})x_2 = 0 \\ (-2-i\sqrt{3})x_3 - 7x_4 = 0 \\ x_3 + (2-i\sqrt{3})x_4 = 0 \end{cases} \Rightarrow \begin{cases} x_1 = x_2 = 0 \\ x_3 = \frac{7}{-2-i\sqrt{3}}x_4 \left(\frac{-2+i\sqrt{3}}{-2+i\sqrt{3}} \right) = (-2+i\sqrt{3})x_4 \\ x_3 = -(2-i\sqrt{3})x_4 \end{cases}$$

Therefore the eigenvector $V_4 = \begin{pmatrix} 0 \\ 0 \\ -2+i\sqrt{3} \\ 1 \end{pmatrix}$

j) For the matrix $\begin{bmatrix} -1 & 0 & 1 \\ -1 & 3 & 0 \\ -4 & 13 & -1 \end{bmatrix}$

i. $\begin{vmatrix} -1-\lambda & 0 & 1 \\ -1 & 3-\lambda & 0 \\ -4 & 13 & -1-\lambda \end{vmatrix} = (-1-\lambda)^2(3-\lambda) - 13 + 4(3-\lambda)$

$$= (\lambda^2 + 2\lambda + 1)(3-\lambda) - 13 + 12 - 4\lambda$$

$$= 3\lambda^2 + 6\lambda + 3 - \lambda^3 - 2\lambda^2 - \lambda - 1 - 4\lambda$$

$$= -\lambda^3 + \lambda^2 + \lambda + 2$$

The characteristic equation: $\underline{-\lambda^3 + \lambda^2 + \lambda + 2 = 0}$

ii. \rightarrow The eigenvalues $\underline{\lambda_{1,2,3} = 2, -\frac{1}{2} \pm i\frac{\sqrt{3}}{2}}$

iii. For $\lambda_1 = 2$, we have: $(A - \lambda_1 I)V_1 = 0$

$$\begin{pmatrix} -3 & 0 & 1 \\ -1 & 1 & 0 \\ -4 & 13 & -3 \end{pmatrix} \begin{pmatrix} x_1 \\ y_1 \\ z_1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} -3x_1 + z_1 = 0 \\ -x_1 + y_1 = 0 \\ -4x_1 + 13y_1 - 3z_1 = 0 \end{cases} \Rightarrow \begin{cases} z_1 = 3x_1 \\ y_1 = x_1 \end{cases}$$

If we let $x_1 = 1$; therefore the eigenvector $V_1 = \begin{pmatrix} 1 \\ 1 \\ 3 \end{pmatrix}$

For $\lambda_2 = -\frac{1}{2} - i\frac{\sqrt{3}}{2}$, we have: $(A - \lambda_2 I)V_2 = 0$

$$\begin{pmatrix} -\frac{1}{2} + i\frac{\sqrt{3}}{2} & 0 & 1 \\ -1 & \frac{7}{2} + i\frac{\sqrt{3}}{2} & 0 \\ -4 & 13 & -\frac{1}{2} + i\frac{\sqrt{3}}{2} \end{pmatrix} \begin{pmatrix} x_2 \\ y_2 \\ z_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} \left(-\frac{1}{2} + i\frac{\sqrt{3}}{2}\right)x_2 + z_2 = 0 \\ -x_2 + \left(\frac{7}{2} + i\frac{\sqrt{3}}{2}\right)y_2 = 0 \\ -4x_2 + 13y_2 + \left(-\frac{1}{2} + i\frac{\sqrt{3}}{2}\right)z_2 = 0 \end{cases}$$

$$\rightarrow \begin{cases} z_2 = -\left(-\frac{1}{2} + i\frac{\sqrt{3}}{2}\right)x_2 \\ y_2 = \left(\frac{2}{7 + i\sqrt{3}}\right)x_2 \end{cases}$$

If we let $x_2 = 1$; therefore the eigenvector $V_2 = \begin{pmatrix} 1 \\ \frac{1-i\sqrt{3}}{2} \\ \frac{2}{7+i\sqrt{3}} \end{pmatrix} = \begin{pmatrix} 1 \\ \frac{1-i\sqrt{3}}{2} \\ \frac{7-i\sqrt{3}}{26} \end{pmatrix}$

For $\lambda_3 = -\frac{1}{2} + i\frac{\sqrt{3}}{2}$, we have: $(A - \lambda_3 I)V_3 = 0$

$$\begin{pmatrix} -\frac{1}{2} - i\frac{\sqrt{3}}{2} & 0 & 1 \\ -1 & \frac{7}{2} - i\frac{\sqrt{3}}{2} & 0 \\ -4 & 13 & -\frac{1}{2} - i\frac{\sqrt{3}}{2} \end{pmatrix} \begin{pmatrix} x_3 \\ y_3 \\ z_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} \left(-\frac{1}{2} - i\frac{\sqrt{3}}{2}\right)x_3 + z_3 = 0 \\ -x_3 + \left(\frac{7}{2} - i\frac{\sqrt{3}}{2}\right)y_3 = 0 \\ -4x_3 + 13y_3 + \left(-\frac{1}{2} - i\frac{\sqrt{3}}{2}\right)z_3 = 0 \end{cases}$$

$$\rightarrow \begin{cases} z_3 = -\left(\frac{1+i\sqrt{3}}{2}\right)x_3 \\ y_3 = \left(\frac{2}{7-i\sqrt{3}}\right)x_3 \end{cases}$$

If we let $x_3 = 1$; therefore the eigenvector $V_3 = \begin{pmatrix} 1 \\ \frac{1+i\sqrt{3}}{2} \\ \frac{2}{7-i\sqrt{3}} \end{pmatrix} = \begin{pmatrix} 1 \\ \frac{1+i\sqrt{3}}{2} \\ \frac{7+i\sqrt{3}}{26} \end{pmatrix}$

Exercise

Find the eigenvalues of A^9 for $A = \begin{pmatrix} 1 & 3 & 7 & 11 \\ 0 & \frac{1}{2} & 3 & 8 \\ 0 & 0 & 0 & 4 \\ 0 & 0 & 0 & 2 \end{pmatrix}$

Solution

The eigenvalues are: $\lambda = 1, \frac{1}{2}, 0, 2$

The eigenvalues of A^9 are: $1^9 = 1$ $\left(\frac{1}{2}\right)^9 = \frac{1}{512}$ $0^9 = 0$ $2^9 = 512$

Exercise

Find the eigenvalues of the matrices

$$A = \begin{pmatrix} 0.7 & 0.4 \\ 0.3 & 0.6 \end{pmatrix}, \quad A^2 = \begin{pmatrix} 0.61 & 0.52 \\ 0.39 & 0.48 \end{pmatrix}, \quad A^\infty = \begin{pmatrix} 0.57143 & 0.57143 \\ 0.42857 & 0.42857 \end{pmatrix}, \quad \text{and} \quad B = \begin{pmatrix} 0.3 & 0.6 \\ 0.7 & 0.4 \end{pmatrix}$$

Solution

The eigenvalues for A :

$$\begin{vmatrix} 0.7 - \lambda & 0.4 \\ 0.3 & 0.6 - \lambda \end{vmatrix} = (0.7 - \lambda)(0.6 - \lambda) - .12 \\ = \lambda^2 - 1.3\lambda + .3 = 0 \quad \lambda_{1,2} = 0.65 \pm 0.35$$

The eigenvalues are: $\lambda_1 = 1$ $\lambda_2 = 0.3$

The eigenvalues for A^2 : $\lambda_1 = 1^2 = 1$ $\lambda_2 = 0.3^2 = 0.09$

The eigenvalues for A^∞ : $\lambda^2 - \lambda = 0$ $\lambda_1 = 1$ $\lambda_2 = 0.3^\infty = 0$

The eigenvalues for B :

$$\begin{vmatrix} 0.3 - \lambda & 0.6 \\ 0.7 & 0.4 - \lambda \end{vmatrix} = \lambda^2 - .7\lambda - .3 = 0 \quad \lambda_{1,2} = 0.35 \pm 0.65$$

The eigenvalues are: $\lambda_1 = 1$ $\lambda_2 = -0.3$

Exercise

Given the matrix $\begin{bmatrix} -1 & -3 \\ -3 & 7 \end{bmatrix}$

- Find the characteristic polynomial.
- Find the eigenvalues
- Find the bases for its eigenspaces
- Graph the eigenspaces
- Verify directly that $A\mathbf{v} = \lambda\mathbf{v}$, for all associated eigenvectors and eigenvalues.

Solution

$$\begin{aligned} a) \quad \begin{vmatrix} -1-\lambda & -3 \\ -3 & 7-\lambda \end{vmatrix} &= (-1-\lambda)(7-\lambda) - 9 \\ &= -7 - 6\lambda + \lambda^2 - 9 \\ &= \lambda^2 - 6\lambda - 16 \end{aligned}$$

The characteristic polynomial is $\lambda^2 - 6\lambda - 16 = 0$

$$b) \quad \lambda^2 - 6\lambda - 16 = 0 \Rightarrow \lambda_1 = -2 \text{ and } \lambda_2 = 8$$

$$c) \quad \text{For } \lambda_1 = -2, \text{ we have: } (A + 2I)V_1 = 0$$

$$\begin{pmatrix} 1 & -3 \\ -3 & 9 \end{pmatrix} \begin{pmatrix} x_1 \\ y_1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} x_1 - 3y_1 = 0 \\ -3x_1 + 9y_1 = 0 \end{cases} \Rightarrow x_1 = 3y_1$$


Therefore the eigenvector $V_1 = \begin{pmatrix} 3 \\ 1 \end{pmatrix}$

$$\text{For } \lambda_2 = 8, \text{ we have: } (A - 8I)V_2 = 0$$

$$\begin{pmatrix} -9 & -3 \\ -3 & -1 \end{pmatrix} \begin{pmatrix} x_2 \\ y_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} -9x_2 - 3y_2 = 0 \\ -3x_2 - y_2 = 0 \end{cases} \Rightarrow y_2 = -3x_2$$

Therefore the eigenvector $V_2 = \begin{pmatrix} 1 \\ -3 \end{pmatrix}$

d)



$$\begin{aligned}
e) \quad AV_1 &= \lambda V_1 \rightarrow \begin{pmatrix} -1 & -3 \\ -3 & 7 \end{pmatrix} \begin{pmatrix} 3 \\ 1 \end{pmatrix} = -2 \begin{pmatrix} 3 \\ 1 \end{pmatrix} \\
&\begin{pmatrix} -6 \\ -2 \end{pmatrix} = \begin{pmatrix} -6 \\ -2 \end{pmatrix} \quad \checkmark \\
AV_2 &= \lambda V_2 \rightarrow \begin{pmatrix} -1 & -3 \\ -3 & 7 \end{pmatrix} \begin{pmatrix} 1 \\ -3 \end{pmatrix} = 8 \begin{pmatrix} 1 \\ -3 \end{pmatrix} \\
&\begin{pmatrix} 8 \\ -24 \end{pmatrix} = \begin{pmatrix} 8 \\ -24 \end{pmatrix} \quad \checkmark
\end{aligned}$$

Exercise

Given the matrix $\begin{bmatrix} 5 & 0 & -4 \\ 0 & -3 & 0 \\ -4 & 0 & -1 \end{bmatrix}$

- Find the characteristic polynomial.
- Find the eigenvalues
- Find the bases for its eigenspaces
- Graph the eigenspaces
- Verify directly that $Av = \lambda v$, for all associated eigenvectors and eigenvalues.

Solution

$$\begin{aligned}
a) \quad \begin{vmatrix} 5-\lambda & 0 & -4 \\ 0 & -3-\lambda & 0 \\ -4 & 0 & -1-\lambda \end{vmatrix} &= (5-\lambda)(-3-\lambda)(-1-\lambda) - 16(-3-\lambda) \\
&= (5-\lambda)(3+4\lambda+\lambda^2) + 48 + 16\lambda \\
&= 15 + 20\lambda + 5\lambda^2 - 3\lambda - 4\lambda^2 - \lambda^3 + 48 + 16\lambda \\
&= -\lambda^3 + \lambda^2 + 33\lambda + 63
\end{aligned}$$

The characteristic polynomial is $\underline{-\lambda^3 + \lambda^2 + 33\lambda + 63 = 0}$

$$b) \quad -\lambda^3 + \lambda^2 + 33\lambda + 63 = 0 \Rightarrow \lambda = -3, -3, 7$$

$$c) \quad \text{For } \lambda_{1,2} = -3, \text{ we have: } (A + 3I)V_1 = 0$$

$$\begin{pmatrix} 8 & 0 & -4 \\ 0 & 0 & 0 \\ -4 & 0 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ y_1 \\ z_1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} 8x_1 - 4z_1 = 0 \\ -4x_1 + 2z_1 = 0 \end{cases} \Rightarrow z_1 = 2x_1$$


Therefore the eigenvector $V_1 = \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix}$ and $V_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$

For $\lambda_3 = 7$, we have: $(A - 7I)V_3 = 0$

$$\begin{pmatrix} -2 & 0 & -4 \\ 0 & -10 & 0 \\ -4 & 0 & -8 \end{pmatrix} \begin{pmatrix} x_1 \\ y_1 \\ z_1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} -2x_1 - 4z_1 = 0 \\ -10y_1 = 0 \\ -4x_1 - 8z_1 = 0 \end{cases} \Rightarrow x_1 = -2z_1 \text{ and } y_1 = 0$$

Therefore the eigenvector $V_3 = \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix}$

d)



$$e) \quad AV_1 = \lambda V_1 \rightarrow \begin{pmatrix} 5 & 0 & -4 \\ 0 & -3 & 0 \\ -4 & 0 & -1 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix} = -3 \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix}$$

$$\begin{pmatrix} -3 \\ 0 \\ -6 \end{pmatrix} = \begin{pmatrix} -3 \\ 0 \\ -6 \end{pmatrix} \checkmark$$

$$AV_2 = \lambda V_2 \rightarrow \begin{pmatrix} 5 & 0 & -4 \\ 0 & -3 & 0 \\ -4 & 0 & -1 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} = -3 \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 0 \\ -3 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ -3 \\ 0 \end{pmatrix} \checkmark$$

$$AV_3 = \lambda V_3 \rightarrow \begin{pmatrix} 5 & 0 & -4 \\ 0 & -3 & 0 \\ -4 & 0 & -1 \end{pmatrix} \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix} = 7 \begin{pmatrix} -2 \\ 0 \\ 1 \end{pmatrix}$$

$$\begin{pmatrix} -14 \\ 0 \\ 7 \end{pmatrix} = \begin{pmatrix} -14 \\ 0 \\ 7 \end{pmatrix} \checkmark$$

Exercise

Given: $A = \begin{bmatrix} -1 & 7 & -1 \\ 0 & 1 & 0 \\ 0 & 15 & -2 \end{bmatrix}$. Compute A^{11}

Solution

$$\det(A - \lambda I) = \begin{vmatrix} -1-\lambda & 7 & -1 \\ 0 & 1-\lambda & 0 \\ 0 & 15 & -2-\lambda \end{vmatrix}$$

$$= (-1-\lambda)(1-\lambda)(-2-\lambda)$$

The eigenvalues are: $-1, 1, -2$

For $\lambda_1 = -1$, we have: $(A - \lambda_1 I)V_1 = 0$

$$\begin{pmatrix} 0 & 7 & -1 \\ 0 & 2 & 0 \\ 0 & 15 & -1 \end{pmatrix} \begin{pmatrix} x_1 \\ y_1 \\ z_1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} 7y_1 - z_1 = 0 \\ 2y_1 = 0 \\ 15y_1 - z_1 = 0 \end{cases} \Rightarrow \begin{cases} z_1 = 7y_1 \\ y_1 = 0 \end{cases}$$

The eigenvector $V_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$

For $\lambda_2 = 1$, we have: $(A - I)V_2 = 0$

$$\begin{pmatrix} -2 & 7 & -1 \\ 0 & 0 & 0 \\ 0 & 15 & -3 \end{pmatrix} \begin{pmatrix} x_2 \\ y_2 \\ z_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} -2x_2 + 7y_2 - z_2 = 0 \\ 15y_2 - 3z_2 = 0 \end{cases} \Rightarrow \begin{cases} 2x_2 = 7y_2 - z_2 \\ 5y_2 = z_2 \end{cases}$$

If we let $y_2 = 1 \rightarrow z_2 = 5$ and $x_2 = \frac{7-5}{2} = 1$;

The eigenvector $V_2 = \begin{pmatrix} 1 \\ 1 \\ 5 \end{pmatrix}$

For $\lambda_3 = -2$, we have: $(A + 2I)V_3 = 0$

$$\begin{pmatrix} 1 & 7 & -1 \\ 0 & 3 & 0 \\ 0 & 15 & 0 \end{pmatrix} \begin{pmatrix} x_3 \\ y_3 \\ z_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \begin{cases} x_3 + 7y_3 - z_3 = 0 \\ 3y_3 = 0 \\ 15y_3 = 0 \end{cases} \Rightarrow \begin{cases} x_3 = -7y_3 + z_3 \\ y_3 = 0 \end{cases}$$

The eigenvector $V_3 = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$

$$D = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix} \Rightarrow D^{11} = \begin{pmatrix} (-1)^{11} & 0 & 0 \\ 0 & 1^{11} & 0 \\ 0 & 0 & (-2)^{11} \end{pmatrix} = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2048 \end{pmatrix}$$

$$P = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 0 & 5 & 1 \end{pmatrix} \Rightarrow P^{-1} = \begin{pmatrix} 1 & 4 & -1 \\ 0 & 1 & 0 \\ 0 & -5 & 1 \end{pmatrix}$$

$$\begin{aligned} A^{11} &= \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 0 & 5 & 1 \end{pmatrix} \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2048 \end{pmatrix} \begin{pmatrix} 1 & 4 & -1 \\ 0 & 1 & 0 \\ 0 & -5 & 1 \end{pmatrix} \\ &= \begin{pmatrix} -1 & 1 & -2048 \\ 0 & 1 & 0 \\ 0 & 5 & -2048 \end{pmatrix} \begin{pmatrix} 1 & 4 & -1 \\ 0 & 1 & 0 \\ 0 & -5 & 1 \end{pmatrix} \\ &= \begin{pmatrix} -1 & 10237 & 2047 \\ 0 & 1 & 0 \\ 0 & 10245 & -2048 \end{pmatrix} \end{aligned}$$