

Lecture Three

Section 3.1 – Inner Products

Definition

An **inner product** on a real vector space V is a function that associates a real number $\langle \mathbf{u}, \mathbf{v} \rangle$ with each pair of vectors in V in such a way that the following axioms are satisfied for all vectors \mathbf{u}, \mathbf{v} , and \mathbf{w} in V and all scalars k .

1. $\langle \mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{v}, \mathbf{u} \rangle$ *Symmetry axiom*
2. $\langle \mathbf{u} + \mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{w} \rangle + \langle \mathbf{v}, \mathbf{w} \rangle$ *Additivity axiom*
3. $\langle k\mathbf{u}, \mathbf{v} \rangle = k \langle \mathbf{u}, \mathbf{v} \rangle$ *Homogeneity axiom*
4. $\langle \mathbf{v}, \mathbf{v} \rangle \geq 0$ and $\langle \mathbf{v}, \mathbf{v} \rangle = 0$ iff $\mathbf{v} = \mathbf{0}$ *Positivity axiom*

A real vector space with an inner product is called a **real inner product space**.

$$\langle \mathbf{u}, \mathbf{v} \rangle = \mathbf{u} \cdot \mathbf{v} = u_1 v_1 + u_2 v_2 + \cdots + u_n v_n$$

This is called the **Euclidean inner product** (or the **standard inner product**)

Definition

If V is a real inner product space, then the norm (or length) of a vector \mathbf{v} in V is denoted by $\|\mathbf{v}\|$ and is defined by

$$\|\mathbf{v}\| = \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle}$$

And the **distance** between two vectors is denoted by $d(\mathbf{u}, \mathbf{v})$ and is defined by

$$d(\mathbf{u}, \mathbf{v}) = \|\mathbf{u} - \mathbf{v}\| = \sqrt{\langle \mathbf{u} - \mathbf{v}, \mathbf{u} - \mathbf{v} \rangle}$$

A vector of norm 1 is called a **unit vector**.

Theorem

If \mathbf{u} and \mathbf{v} are vectors in a real inner product space V , and if k is a scalar, then:

- a) $\|\mathbf{v}\| \geq 0$ with equality iff $\mathbf{v} = \mathbf{0}$
- b) $\|k\mathbf{v}\| = |k| \|\mathbf{v}\|$
- c) $d(\mathbf{u}, \mathbf{v}) = d(\mathbf{v}, \mathbf{u})$
- d) $d(\mathbf{u}, \mathbf{v}) \geq 0$ with equality iff $\mathbf{u} = \mathbf{v}$

Although the Euclidean inner product is the most important inner product on R^n , there are various applications in which is desirable to modify it by weighing each term differently. More precisely, if w_1, w_2, \dots, w_n are positive real numbers, which we will call weighs, and if $\mathbf{u} = (u_1, u_2, \dots, u_n)$ and \mathbf{v} are vectors in R^n , then it can be shown that the formula

$$\langle \mathbf{u}, \mathbf{v} \rangle = w_1 u_1 v_1 + w_2 u_2 v_2 + \dots + w_n u_n v_n$$

Defines an inner product on R^n that we call the **weighted Euclidean inner product** with weights w_1, w_2, \dots, w_n

Example

Let $\mathbf{u} = (u_1, u_2)$ and $\mathbf{v} = (v_1, v_2)$ be vectors in R^2 , verify that the weighted Euclidean inner product $\langle \mathbf{u}, \mathbf{v} \rangle = 3u_1 v_1 + 2u_2 v_2$ satisfies the four inner product axioms.

Solution

$$\text{Axiom 1: } \langle \mathbf{u}, \mathbf{v} \rangle = 3u_1 v_1 + 2u_2 v_2 = 3v_1 u_1 + 2v_2 u_2 = \langle \mathbf{v}, \mathbf{u} \rangle$$

$$\begin{aligned} \text{Axiom 2: } \langle \mathbf{u} + \mathbf{v}, \mathbf{w} \rangle &= 3(u_1 + v_1)w_1 + 2(u_2 + v_2)w_2 \\ &= 3(u_1 w_1 + v_1 w_1) + 2(u_2 w_2 + v_2 w_2) \\ &= 3u_1 w_1 + 3v_1 w_1 + 2u_2 w_2 + 2v_2 w_2 \\ &= (3u_1 w_1 + 2u_2 w_2) + (3v_1 w_1 + 2v_2 w_2) \\ &= \langle \mathbf{u}, \mathbf{w} \rangle + \langle \mathbf{v}, \mathbf{w} \rangle \end{aligned}$$

$$\begin{aligned} \text{Axiom 3: } \langle k\mathbf{u}, \mathbf{v} \rangle &= 3(ku_1)v_1 + 2(ku_2)v_2 \\ &= k(3u_1 v_1 + 2u_2 v_2) \\ &= k \langle \mathbf{u}, \mathbf{v} \rangle \end{aligned}$$

$$\begin{aligned} \text{Axiom 3: } \langle \mathbf{v}, \mathbf{v} \rangle &= 3v_1 v_1 + 2v_2 v_2 \\ &= 3v_1^2 + 2v_2^2 \geq 0 \\ v_1 = v_2 = 0 &\text{ iff } \mathbf{v} = \mathbf{0} \end{aligned}$$

Exercises Section 3.1 – Inner Products

1. Let $\langle \mathbf{u}, \mathbf{v} \rangle$ be the Euclidean inner product on R^2 , and let $\mathbf{u} = (1, 1)$, $\mathbf{v} = (3, 2)$, $\mathbf{w} = (0, -1)$, and $k = 3$. Compute the following.

a) $\langle \mathbf{u}, \mathbf{v} \rangle$	c) $\langle \mathbf{u} + \mathbf{v}, \mathbf{w} \rangle$	e) $d(\mathbf{u}, \mathbf{v})$
b) $\langle k\mathbf{v}, \mathbf{w} \rangle$	d) $\ \mathbf{v}\ $	f) $\ \mathbf{u} - k\mathbf{v}\ $

2. Let $\langle \mathbf{u}, \mathbf{v} \rangle$ be the Euclidean inner product on R^2 , and let $\mathbf{u} = (1, 1)$, $\mathbf{v} = (3, 2)$, $\mathbf{w} = (0, -1)$ and $k = 3$. Compute the following for the weighted Euclidean inner product $\langle \mathbf{u}, \mathbf{v} \rangle = 2u_1v_1 + 3u_2v_2$.

a) $\langle \mathbf{u}, \mathbf{v} \rangle$	c) $\langle \mathbf{u} + \mathbf{v}, \mathbf{w} \rangle$	e) $d(\mathbf{u}, \mathbf{v})$
b) $\langle k\mathbf{v}, \mathbf{w} \rangle$	d) $\ \mathbf{v}\ $	f) $\ \mathbf{u} - k\mathbf{v}\ $

3. Let $\langle \mathbf{u}, \mathbf{v} \rangle$ be the Euclidean inner product on R^2 , and let $\mathbf{u} = (3, -2)$, $\mathbf{v} = (4, 5)$, $\mathbf{w} = (-1, 6)$, and $k = -4$. Verify the following.

a) $\langle \mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{v}, \mathbf{u} \rangle$	d) $\langle k\mathbf{u}, \mathbf{v} \rangle = k\langle \mathbf{u}, \mathbf{v} \rangle$
b) $\langle \mathbf{u} + \mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{w} \rangle + \langle \mathbf{v}, \mathbf{w} \rangle$	e) $\langle \mathbf{0}, \mathbf{v} \rangle = \langle \mathbf{v}, \mathbf{0} \rangle = 0$
c) $\langle \mathbf{u}, \mathbf{v} + \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{v} \rangle + \langle \mathbf{u}, \mathbf{w} \rangle$	

4. Let $\langle \mathbf{u}, \mathbf{v} \rangle$ be the Euclidean inner product on R^2 , and let $\mathbf{u} = (3, -2)$, $\mathbf{v} = (4, 5)$, $\mathbf{w} = (-1, 6)$, and $k = -4$. Verify the following for the weighted Euclidean inner product $\langle \mathbf{u}, \mathbf{v} \rangle = 4u_1v_1 + 5u_2v_2$

a) $\langle \mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{v}, \mathbf{u} \rangle$	d) $\langle k\mathbf{u}, \mathbf{v} \rangle = k\langle \mathbf{u}, \mathbf{v} \rangle$
b) $\langle \mathbf{u} + \mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{w} \rangle + \langle \mathbf{v}, \mathbf{w} \rangle$	e) $\langle \mathbf{0}, \mathbf{v} \rangle = \langle \mathbf{v}, \mathbf{0} \rangle = 0$
c) $\langle \mathbf{u}, \mathbf{v} + \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{v} \rangle + \langle \mathbf{u}, \mathbf{w} \rangle$	

5. Let $\mathbf{u} = (u_1, u_2)$ and $\mathbf{v} = (v_1, v_2)$. Show that the following are inner product on R^3 by verifying that the inner product axioms hold. $\langle \mathbf{u}, \mathbf{v} \rangle = 3u_1v_1 + 5u_2v_2$

6. Show that the following identity holds for the vectors in any inner product space

$$\|\mathbf{u} + \mathbf{v}\|^2 + \|\mathbf{u} - \mathbf{v}\|^2 = 2\|\mathbf{u}\|^2 + 2\|\mathbf{v}\|^2$$