

Section 1.6 – Determinants and Properties

The determinant is a number that contains information about matrix. It is used to find formulas for inverse matrices, pivots, and solutions $A^{-1}b$.

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \text{ has inverse } A^{-1} = \frac{1}{ad-bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$$

Determinant of the matrix $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ is written $\det(A)$ or $|A|$ and is define as

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

The determinant is zero when the matrix has no inverse.

Properties of the Determinants

By using these property rules, we can compute the determinant of any square matrix.

1. **Determinant of the n by n identity matrix is 1.**

$$\begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} = 1 \quad \text{and} \quad \begin{vmatrix} 1 & & \\ & 1 & \\ & & 1 \end{vmatrix} = 1$$

2. **Determinant changes sign when 2 rows are exchanged.**

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc \quad \begin{vmatrix} c & d \\ a & b \end{vmatrix} = bc - ad = -(ad - bc) \Rightarrow \begin{vmatrix} a & b \\ c & d \end{vmatrix} = - \begin{vmatrix} c & d \\ a & b \end{vmatrix}$$

3. **Determinant is a linear function of each row separately.**

$$\text{Multiply row 1 by any number } t: \begin{vmatrix} ta & tb \\ c & d \end{vmatrix} = t \begin{vmatrix} a & b \\ c & d \end{vmatrix}$$

$$\text{Add row 1 of } A \text{ to row 1 of } A': \begin{vmatrix} a+a' & b+b' \\ c & d \end{vmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix} + \begin{vmatrix} a' & b' \\ c & d \end{vmatrix}$$

✚ For 2 by 2 determinants, if you expand to a rectangle, the determinants equal areas.

✚ For n -dimensional, the determinants equal volumes.

4. If 2 rows of A are equal, then $\det A = 0$.

$$\begin{vmatrix} a & b \\ a & b \end{vmatrix} = ab - ab = 0$$

5. Subtracting a multiple of one row from another row leaves $\det A$ unchanged.

$$\begin{vmatrix} a & b \\ c - ta & d - tb \end{vmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix}$$

6. A matrix with a row of zeros has $\det A = 0$.

$$\begin{vmatrix} a & b \\ 0 & 0 \end{vmatrix} = 0 \quad \text{and} \quad \begin{vmatrix} 0 & 0 \\ b & c \end{vmatrix} = 0$$

7. If A is triangular then $\det A = a_{11} a_{22} \dots a_{nn} = \text{product of diagonal entries}$.

$$\begin{vmatrix} a & b \\ 0 & d \end{vmatrix} = ad \quad \text{and} \quad \begin{vmatrix} a_{11} & & 0 \\ & a_{22} & \\ 0 & & a_{nn} \end{vmatrix} = a_{11} a_{22} \dots a_{nn}$$

8. If A is singular then $\det A = 0$.

9. If A is invertible then $\det A \neq 0$.

10. The determinant of AB is times $\det A$: $|AB| = |A||B|$

11. The transpose A^T has the same determinant as A : $\det(A) = \det(A^T)$

➤ $\det(A + B) \neq \det(A) + \det(B)$

Big Formula for Determinants (Diagonal)

Determinant Using Diagonal Method

$$\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

$$\begin{array}{cccccc} a_{11} & a_{12} & a_{13} & a_{11} & a_{12} \\ a_{21} & a_{22} & a_{23} & a_{21} & a_{22} \\ a_{31} & a_{32} & a_{33} & a_{31} & a_{32} \end{array}$$

$$a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} \quad (1)$$

$$\begin{array}{cccccc} a_{11} & a_{12} & a_{13} & a_{11} & a_{12} \\ a_{21} & a_{22} & a_{23} & a_{21} & a_{22} \\ a_{31} & a_{32} & a_{33} & a_{31} & a_{32} \end{array}$$

$$-a_{13}a_{22}a_{31} - a_{11}a_{23}a_{32} - a_{12}a_{21}a_{33} \quad (2)$$

$$\text{Determinant: } D = (1) + (2)$$

$$\det = a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} - a_{13}a_{22}a_{31} - a_{11}a_{23}a_{32} - a_{12}a_{21}a_{33}$$

Example

Evaluate: $\begin{vmatrix} x & 0 & -1 \\ 2 & x & x^2 \\ -3 & x & 1 \end{vmatrix}$

Solution

$$\begin{vmatrix} x & 0 & -1 \\ 2 & x & x^2 \\ -3 & x & 1 \end{vmatrix} \begin{array}{cc} x & 0 \\ 2 & x \\ -3 & x \end{array} = x(x)(1) + 0(x^2)(2) + (-1)(2)(x) - (-1)(x)(-3) - x(x^2)(x) - 0(-3)(1)$$

$$= x^2 - 2x - 3x - x^4$$

$$= x^2 - 5x - x^4$$

Determinant by *Cofactors*

$$A = [a_{ij}] = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

Minor

For a square matrix $A = [a_{ij}]$, the minor M_{ij} of an element a_{ij} is the **determinant** of the matrix formed by deleting the i^{th} row and the j^{th} column of A .

Example

$$\text{Let } A = \begin{bmatrix} 3 & 1 & -4 \\ 2 & 5 & 6 \\ 1 & 4 & 8 \end{bmatrix} \text{ Find } M_{32}$$

Solution

$$\begin{aligned} M_{32} &= \begin{vmatrix} 3 & \cancel{1} & -4 \\ 2 & \cancel{5} & 6 \\ \cancel{1} & \cancel{4} & \cancel{8} \end{vmatrix} \\ &= \begin{vmatrix} 3 & -4 \\ 2 & 6 \end{vmatrix} \\ &= \underline{26} \end{aligned}$$

Theorem

The determinant is the dot product of any row i of A with its cofactors:

$$\text{Cofactor Formula: } C_{ij} = (-1)^{i+j} M_{ij}$$

$$|A| = a_{11}C_{11} + a_{12}C_{12} + a_{13}C_{13}$$

$$= a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}$$

Example

Find the determinant of the matrix:

$$A = \begin{bmatrix} -8 & 0 & 6 \\ 4 & -6 & 7 \\ -1 & -3 & 5 \end{bmatrix}$$

Solution

$$\begin{aligned} |A| &= \begin{vmatrix} -8 & 0 & 6 \\ 4 & -6 & 7 \\ -1 & -3 & 5 \end{vmatrix} \\ &= -8 \begin{vmatrix} -6 & 7 \\ -3 & 5 \end{vmatrix} - 0 \begin{vmatrix} 4 & 7 \\ -1 & 5 \end{vmatrix} + 6 \begin{vmatrix} 4 & -6 \\ -1 & -3 \end{vmatrix} \\ &= -8(-30 - (-21)) - 0 + 6(-12 - 6) \\ &= -8(-9) + 6(-18) \\ &= \underline{-36} \end{aligned}$$

✓ By the property of determinants, If A is triangular then $\det A = a_{11} a_{22} \dots a_{nn}$ = product of diagonal entries.

Example

$$\begin{vmatrix} 2 & 7 & -3 & 8 & 3 \\ 0 & -3 & 7 & 5 & 1 \\ 0 & 0 & 6 & 7 & 6 \\ 0 & 0 & 0 & 9 & 8 \\ 0 & 0 & 0 & 0 & 4 \end{vmatrix} = (2)(-3)(6)(9)(4) \\ = \underline{-1296}$$

Theorem

Let A be any n by n matrix.

- a) If A' is the matrix that results when a single row of A is multiplied by a constant k , then $\det(A') = k \det(A)$.
- b) If A' is the matrix that results when two rows of A are interchanged, then $\det(A') = -\det(A)$
- c) If A' is the matrix that results when a multiple of one row of A is added to another row then $\det(A') = \det(A)$

Example

$$\text{Evaluate } \begin{vmatrix} 0 & 1 & 5 \\ 3 & -6 & 9 \\ 2 & 6 & 1 \end{vmatrix}$$

Solution

$$\begin{vmatrix} 0 & 1 & 5 \\ 3 & -6 & 9 \\ 2 & 6 & 1 \end{vmatrix} = - \begin{vmatrix} 3 & -6 & 9 \\ 0 & 1 & 5 \\ 2 & 6 & 1 \end{vmatrix}$$

Interchanged 1st and 2nd row

$$= -3 \begin{vmatrix} 1 & -2 & 3 \\ 0 & 1 & 5 \\ 2 & 6 & 1 \end{vmatrix}$$

A common factor of 3 from the first row (no need)

$$= -3 \begin{vmatrix} 1 & -2 & 3 \\ 0 & 1 & 5 \\ 2 & 6 & 1 \end{vmatrix} \quad R_3 - 2R_1$$

$$= -3 \begin{vmatrix} 1 & -2 & 3 \\ 0 & 1 & 5 \\ 0 & 10 & -5 \end{vmatrix} \quad R_3 - 10R_2$$

$$= -3 \begin{vmatrix} 1 & -2 & 3 \\ 0 & 1 & 5 \\ 0 & 0 & -55 \end{vmatrix}$$

$$= -3(1)(1)(-55)$$

$$= \underline{165}$$

Exercises Section 1.6 – Determinants and Properties

1. Verify that $\det(AB) = \det(A)\det(B)$ when: $A = \begin{pmatrix} 2 & 1 & 0 \\ 3 & 4 & 0 \\ 0 & 0 & 2 \end{pmatrix}$ and $B = \begin{pmatrix} 1 & -1 & 3 \\ 7 & 1 & 2 \\ 5 & 0 & 1 \end{pmatrix}$
2. For which value(s) of k does A fail to be invertible? $A = \begin{bmatrix} k-3 & -2 \\ -2 & k-2 \end{bmatrix}$
3. Without directly evaluating, show that $\begin{vmatrix} b+c & c+a & b+a \\ a & b & c \\ 1 & 1 & 1 \end{vmatrix} = 0$
4. If the entries in every row of A add to zero, solve $A\mathbf{x} = 0$ to prove $\det(A) = 0$. If those entries add to one, show that $\det(A - I) = 0$. Does this mean $\det(A) = 1$?
5. Does $\det(AB) = \det(BA)$ in general?
 - a) True or false if A and B are square $n \times n$ matrices?
 - b) True or false if A is $m \times n$ and B is $n \times m$ with $m \neq n$?
6. True or false, with a reason if true or a counterexample if false:
 - a) The determinant of $I + A$ is $1 + \det(A)$.
 - b) The determinant of ABC is $|A||B||C|$.
 - c) The determinant of $4A$ is $4|A|$.
 - d) The determinant of $AB - BA$ is zero. (try an example)
 - e) If A is not invertible then AB is not invertible.
 - f) The determinant of $A - B$ equals to $\det(A) - \det(B)$.
7. Use row operations to show the 3 by 3 “Vandermonde determinant” is

$$\det \begin{bmatrix} 1 & a & a^2 \\ 1 & b & b^2 \\ 1 & c & c^2 \end{bmatrix} = (b-a)(c-a)(c-b)$$

8. The inverse of a 2 by 2 matrix seems to have determinant = 1:

$$\det A^{-1} = \det \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} = \frac{ad-bc}{ad-bc} = 1$$

What is wrong with this calculation? What is the correct $\det A^{-1}$

9. A *Hessenberg* matrix is a triangular matrix with one extra diagonal. Use cofactors of row 1 to show that the 4 by 4 determinant satisfies Fibonacci's rule $|H_4| = |H_3| + |H_2|$. The same rule will continue for all sizes $|H_n| = |H_{n-1}| + |H_{n-2}|$. Which Fibonacci number is $|H_n|$?

$$H_2 = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} \quad H_3 = \begin{bmatrix} 2 & 1 & \\ 1 & 2 & 1 \\ 1 & 1 & 2 \end{bmatrix} \quad H_4 = \begin{bmatrix} 2 & 1 & & \\ 1 & 2 & 1 & \\ 1 & 1 & 2 & 1 \\ 1 & 1 & 1 & 2 \end{bmatrix}$$

(10 – 44) Evaluate

10. $\begin{vmatrix} -1 & 3 \\ -2 & 9 \end{vmatrix}$

19. $\begin{vmatrix} -4 & -1 \\ 5 & 6 \end{vmatrix}$

26. $\begin{vmatrix} x & x^2 \\ 4 & x \end{vmatrix}$

11. $\begin{vmatrix} 6 & -4 \\ 0 & -1 \end{vmatrix}$

20. $\begin{vmatrix} \sqrt{3} & -2 \\ -3 & \sqrt{3} \end{vmatrix}$

27. $\begin{vmatrix} x & x^2 \\ x & 9 \end{vmatrix}$

12. $\begin{vmatrix} x & 4x \\ 2x & 8x \end{vmatrix}$

21. $\begin{vmatrix} \sqrt{7} & 6 \\ -3 & \sqrt{7} \end{vmatrix}$

28. $\begin{vmatrix} x^2 & x \\ -3 & 2 \end{vmatrix}$

13. $\begin{vmatrix} x & 2x \\ 4 & 3 \end{vmatrix}$

22. $\begin{vmatrix} \sqrt{5} & 3 \\ -2 & 2 \end{vmatrix}$

29. $\begin{vmatrix} x+2 & 6 \\ x-2 & 4 \end{vmatrix}$

14. $\begin{vmatrix} x^4 & 2 \\ x & -3 \end{vmatrix}$

23. $\begin{vmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{8} & -\frac{3}{4} \end{vmatrix}$

30. $\begin{vmatrix} x+1 & -6 \\ x+3 & -3 \end{vmatrix}$

15. $\begin{vmatrix} -8 & -5 \\ b & a \end{vmatrix}$

24. $\begin{vmatrix} \frac{1}{5} & \frac{1}{6} \\ -6 & -5 \end{vmatrix}$

31. $\begin{vmatrix} 3 & 0 & 0 \\ 2 & 1 & -5 \\ 2 & 5 & -1 \end{vmatrix}$

16. $\begin{vmatrix} 5 & 7 \\ 2 & 3 \end{vmatrix}$

25. $\begin{vmatrix} \frac{2}{3} & \frac{1}{3} \\ -\frac{1}{2} & \frac{3}{4} \end{vmatrix}$

32. $\begin{vmatrix} 4 & 0 & 0 \\ 3 & -1 & 4 \\ 2 & -3 & 6 \end{vmatrix}$

18. $\begin{vmatrix} 5 & 3 \\ -2 & 3 \end{vmatrix}$

$$33. \begin{vmatrix} 3 & 1 & 0 \\ -3 & -4 & 0 \\ -1 & 3 & 5 \end{vmatrix}$$

$$37. \begin{vmatrix} 4 & -7 & 8 \\ 2 & 1 & 3 \\ -6 & 3 & 0 \end{vmatrix}$$

$$41. \begin{vmatrix} 0 & x & x \\ x & x^2 & 5 \\ x & 7 & -5 \end{vmatrix}$$

$$34. \begin{vmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \\ 3 & -4 & 5 \end{vmatrix}$$

$$38. \begin{vmatrix} 2 & 1 & -1 \\ 4 & 7 & -2 \\ 2 & 4 & 0 \end{vmatrix}$$

$$42. \begin{vmatrix} 2 & x & 1 \\ -3 & 1 & 0 \\ 2 & 1 & 4 \end{vmatrix}$$

$$35. \begin{vmatrix} x & 0 & -1 \\ 2 & 1 & x^2 \\ -3 & x & 1 \end{vmatrix}$$

$$39. \begin{vmatrix} 3 & 1 & 2 \\ -2 & 3 & 1 \\ 3 & 4 & -6 \end{vmatrix}$$

$$43. \begin{vmatrix} 1 & x & -2 \\ 3 & 1 & 1 \\ 0 & -2 & 2 \end{vmatrix}$$

$$36. \begin{vmatrix} x & 1 & -1 \\ x^2 & x & x \\ 0 & x & 1 \end{vmatrix}$$

$$40. \begin{vmatrix} 2x & 1 & -1 \\ 0 & 4 & x \\ 3 & 0 & 2 \end{vmatrix}$$

$$44. \begin{vmatrix} 1 & 1 & 0 & 0 & 0 \\ -1 & 1 & 1 & 0 & 0 \\ 0 & -1 & 1 & 1 & 0 \\ 0 & 0 & -1 & 1 & 1 \\ 0 & 0 & 0 & -1 & 1 \end{vmatrix}$$

45. Find all the values of λ for which $\det(A) = 0$

$$a) A = \begin{bmatrix} \lambda - 1 & -2 \\ 1 & \lambda - 4 \end{bmatrix} \quad b) A = \begin{bmatrix} \lambda - 6 & 0 & 0 \\ 0 & \lambda & -1 \\ 0 & 4 & \lambda - 4 \end{bmatrix}$$

46. Prove that if a square matrix A has a column of zeros, then $\det(A) = 0$

47. With 2 by 2 blocks in 4 by 4 matrices, you cannot always use block determinants:

$$\begin{vmatrix} A & B \\ 0 & D \end{vmatrix} = |A||D| \quad \text{but} \quad \begin{vmatrix} A & B \\ C & D \end{vmatrix} \neq |A||D| - |C||B|$$

a) Why is the first statement true? Somehow B doesn't enter.

b) Show by example that equality fails (as shown) when C enters.

c) Show by example that the answer $\det(AD - CB)$ is also wrong.

48. Show that the value of the following determinant is independent of θ .

$$\begin{vmatrix} \sin \theta & \cos \theta & 0 \\ -\cos \theta & \sin \theta & 0 \\ \sin \theta - \cos \theta & \sin \theta + \cos \theta & 1 \end{vmatrix}$$

49. Show that the matrices $A = \begin{bmatrix} a & b \\ 0 & c \end{bmatrix}$ and $B = \begin{bmatrix} d & e \\ 0 & f \end{bmatrix}$

commute if and only if $\begin{vmatrix} b & a-c \\ e & d-f \end{vmatrix} = 0$

50. Show that $\det(A) = \frac{1}{2} \begin{vmatrix} \text{tr}(A) & 1 \\ \text{tr}(A^2) & \text{tr}(A) \end{vmatrix}$ for every 2×2 matrix A .

51. What is the maximum number of zeros that a 4×4 matrix can have without a zero determinant? Explain your reasoning.

52. Evaluate $\det(A)$, $\det(E)$, and $\det(AE)$. Then verify that $\det(A) \cdot \det(E) = \det(AE)$

$$A = \begin{bmatrix} 4 & 1 & 3 \\ 0 & -2 & 0 \\ 3 & 1 & 5 \end{bmatrix}, \quad E = \begin{bmatrix} 1 & & \\ & 3 & \\ & & 1 \end{bmatrix}$$

53. Show that $\begin{bmatrix} \sin^2 \alpha & \sin^2 \beta & \sin^2 \gamma \\ \cos^2 \alpha & \cos^2 \beta & \cos^2 \gamma \\ 1 & 1 & 1 \end{bmatrix}$ is not invertible for any values of α, β, γ

54. The determinant of a 2×2 matrix $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ is $\det(A) = ad - bc$.

Assuming no rows swaps are required, perform elimination on A and show explicitly that $ad - bc$ is the product of the pivots.

55. If A is a 7×7 matrix and let $\det(A) = 17$. What is $\det(3A^2)$?

56. Explain without computations why the following determinant is equal to zero

$$\begin{vmatrix} a_1 & a_2 & a_3 & a_4 & a_5 \\ b_1 & b_2 & b_3 & b_4 & b_5 \\ c_1 & c_2 & 0 & 0 & 0 \\ d_1 & d_2 & 0 & 0 & 0 \\ e_1 & e_2 & 0 & 0 & 0 \end{vmatrix}$$

57. Let A be $n \times n$ real matrix.

- Show that if $A^t = -A$ and n is odd, then $|A| = 0$.
- Show that if $A^2 + I = 0$, then n must be even.
- Does part (b) remain true for complex matrices?

58. Let A and C be $m \times m$ and $n \times n$ matrices, respectively.

a) Show that $\begin{vmatrix} A & B \\ 0 & C \end{vmatrix} = \begin{vmatrix} A & 0 \\ B & C \end{vmatrix} = |A||C|$

b) Evaluate

i. $\begin{vmatrix} I_m & 0 \\ 0 & I_n \end{vmatrix}$

ii. $\begin{vmatrix} 0 & I_m \\ I_n & 0 \end{vmatrix}$

iii. $\begin{vmatrix} I_m & B \\ 0 & I_n \end{vmatrix}$

iv. $\begin{vmatrix} 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & \cdots & 1 & 0 \\ \vdots & \vdots & & \vdots & \vdots \\ 0 & 1 & \cdots & 0 & 0 \\ 1 & 0 & \cdots & 0 & 0 \end{vmatrix}_{n \times n}$

c) Find a formula for $\begin{vmatrix} 0 & A \\ C & B \end{vmatrix}_{n \times n}$

59. Let $f(x) = (p_1 - x)(p_2 - x) \cdots (p_n - x)$ and let

$$\Delta_n = \begin{vmatrix} p_1 & a & a & \cdots & a & a \\ b & p_2 & a & \cdots & a & a \\ b & b & p_3 & \cdots & a & a \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ b & b & b & \cdots & p_{n-1} & a \\ b & b & b & \cdots & b & p_n \end{vmatrix}$$

a) Show that, if $a \neq b$,

$$\Delta_n = \frac{bf(a) - af(b)}{b - a}$$

b) Show that, if $a = b$,

$$\Delta_n = a \sum_{i=1}^n f_i(a) + p_n f_n(a)$$

Where $f_i(a)$ means $f(a)$ with factor $(p_i - a)$ missing.

c) Use part (b) to evaluate

$$\begin{vmatrix} a & b & b & \dots & b & b \\ b & a & b & \dots & b & b \\ b & b & a & \dots & a & a \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ b & b & b & \dots & a & a \\ b & b & b & \dots & b & a \end{vmatrix}_{n \times n}$$

60. Let $A, B, C, D \in M_n(\mathbb{C})$

a) Show that when A is invertible: $\begin{vmatrix} A & B \\ C & D \end{vmatrix} = |A| |D - CA^{-1}B|$

b) Show that when $AC = CA$: $\begin{vmatrix} A & B \\ C & D \end{vmatrix} = |AD - CB|$

c) Can B and C on the right-hand side of the identity be switched?

d) Does part (b) remain true if the condition $AC = CA$ is dropped?