


Section 2.6 – Improper Integrals

Definition

Integrals with infinite limits of integration are *improper integrals*.


1. If $f(x)$ is continuous on $[a, \infty)$, then

$$\int_a^{\infty} f(x) dx = \lim_{b \rightarrow \infty} \int_a^b f(x) dx$$




2. If $f(x)$ is continuous on $(-\infty, b]$, then

$$\int_{-\infty}^b f(x) dx = \lim_{a \rightarrow -\infty} \int_a^b f(x) dx$$



3. If $f(x)$ is continuous on $(-\infty, \infty)$, then

$$\int_{-\infty}^{\infty} f(x) dx = \int_{-\infty}^c f(x) dx + \int_c^{\infty} f(x) dx$$



In each case, if the limit is finite we say that the improper integral *converges* and that the limit is the *value* of the improper integral. If the limit fails to exist, the improper integral *diverges*.

Example

Is the area under the curve $y = \frac{\ln x}{x^2}$ from $x = 1$ to $x = \infty$ finite? If so, what is its value?

Solution

$$\begin{aligned} \int_1^b \frac{\ln x}{x^2} dx &= -\frac{1}{x} \ln x \Big|_1^b - \int_1^b \left(-\frac{1}{x}\right) \left(\frac{1}{x}\right) dx \\ &= -\left(\frac{1}{b} \ln b - \ln 1\right) + \int_1^b \frac{1}{x^2} dx \\ &= -\frac{1}{b} \ln b + \left[-\frac{1}{x}\right]_1^b \\ &= -\frac{1}{b} \ln b - \left(\frac{1}{b} - 1\right) \\ &= -\frac{1}{b} \ln b - \frac{1}{b} + 1 \end{aligned}$$

$$\begin{aligned} u &= \ln x & dv &= \frac{dx}{x^2} \\ du &= \frac{1}{x} dx & v &= -\frac{1}{x} \end{aligned}$$

$$\begin{aligned}
\int_1^{\infty} \frac{\ln x}{x^2} dx &= \lim_{b \rightarrow \infty} \int_1^b \frac{\ln x}{x^2} dx \\
&= \lim_{b \rightarrow \infty} \left(-\frac{1}{b} \ln b - \frac{1}{b} + 1 \right) \\
&= -\lim_{b \rightarrow \infty} \left(\frac{\frac{1}{b}}{1} \right) - 0 + 1 \\
&= -0 + 1 \\
&= \underline{1}
\end{aligned}$$

L'Hôpital Rule


Example

Evaluate $\int_{-\infty}^{\infty} \frac{dx}{1+x^2}$

Solution

$$\begin{aligned}
\int_{-\infty}^{\infty} \frac{dx}{1+x^2} &= \int_{-\infty}^0 \frac{dx}{1+x^2} + \int_0^{\infty} \frac{dx}{1+x^2} \\
\int_{-\infty}^0 \frac{dx}{1+x^2} &= \lim_{a \rightarrow -\infty} \int_a^0 \frac{dx}{1+x^2} \\
&= \lim_{a \rightarrow -\infty} \tan^{-1} x \Big|_a^0 \\
&= \lim_{a \rightarrow -\infty} \left(\tan^{-1} 0 - \tan^{-1} a \right) \\
&= 0 - \left(-\frac{\pi}{2} \right) \\
&= \underline{\frac{\pi}{2}}
\end{aligned}$$

$$\begin{aligned}
\int_0^{\infty} \frac{dx}{1+x^2} &= \lim_{b \rightarrow \infty} \int_0^b \frac{dx}{1+x^2} \\
&= \lim_{b \rightarrow \infty} \tan^{-1} x \Big|_0^b \\
&= \lim_{b \rightarrow \infty} \left(\tan^{-1} b - \tan^{-1} 0 \right) \\
&= \frac{\pi}{2} - 0 \\
&= \underline{\frac{\pi}{2}}
\end{aligned}$$



$$\begin{aligned}
 \int_{-\infty}^{\infty} \frac{dx}{1+x^2} &= \int_{-\infty}^0 \frac{dx}{1+x^2} + \int_0^{\infty} \frac{dx}{1+x^2} \\
 &= \frac{\pi}{2} + \frac{\pi}{2} \\
 &= \pi
 \end{aligned}$$

Example


For what value of p does the integral $\int_1^{\infty} \frac{dx}{x^p}$ converge? When the integral does converge, what is its value?

Solution

$$\text{If } p \neq 1 \quad \int_1^b \frac{dx}{x^p} = \left. \frac{x^{-p+1}}{-p+1} \right|_1^b = \frac{1}{1-p} (b^{1-p} - 1)$$

$$\begin{aligned}
 \int_1^{\infty} \frac{dx}{x^p} &= \lim_{b \rightarrow \infty} \int_1^b \frac{dx}{x^p} \\
 &= \lim_{b \rightarrow \infty} \left[\frac{1}{1-p} (b^{1-p} - 1) \right] \\
 &= \begin{cases} \frac{1}{p-1}, & p > 1 \\ \infty, & p < 1 \end{cases}
 \end{aligned}$$

$$\begin{aligned}
 \text{If } p = 1 \quad \int_1^{\infty} \frac{dx}{x^p} &= \int_1^{\infty} \frac{dx}{x} \\
 &= \lim_{b \rightarrow \infty} \int_1^b \frac{dx}{x} \\
 &= \lim_{b \rightarrow \infty} [\ln x]_1^b \\
 &= \lim_{b \rightarrow \infty} (\ln b - \ln 1) \\
 &= \infty
 \end{aligned}$$



Integrands with Vertical Asymptotes


Definition

Integrals of functions that become infinite at a point within the interval of integration are **improper integrals**.

If the limit is finite we say that the improper integral **converges** and that the limit is the **value** of the improper integral. If the limit does not exist, the integral **diverges**.


1. If $f(x)$ is continuous on $(a, b]$, then

$$\int_a^b f(x) dx = \lim_{c \rightarrow a^+} \int_c^b f(x) dx$$




2. If $f(x)$ is continuous on $[a, b)$, then

$$\int_a^b f(x) dx = \lim_{c \rightarrow b^-} \int_a^c f(x) dx$$



3. If $f(x)$ is continuous on $[a, b]$, then

$$\int_a^b f(x) dx = \lim_{c \rightarrow p^-} \int_a^c f(x) dx + \lim_{d \rightarrow p^+} \int_d^b f(x) dx$$



Example

Investigate the convergence of $\int_0^1 \frac{1}{1-x} dx$

Solution

$$\begin{aligned}\int_0^1 \frac{1}{1-x} dx &= \lim_{b \rightarrow 1^-} \int_0^b \frac{1}{1-x} dx \\&= \lim_{b \rightarrow 1^-} \left[-\ln|1-x| \right]_0^b \\&= \lim_{b \rightarrow 1^-} \left[-\ln|1-b| + 0 \right] \\&= \underline{\infty}\end{aligned}$$

The limit is infinite, so the integral diverges.


Example

Evaluate $\int_0^3 \frac{dx}{(x-1)^{2/3}}$

Solution

The integrand has a vertical asymptote at $x = 1$ and is continuous on $[0, 1)$ and $(1, 3]$.

$$\begin{aligned}\int \frac{dx}{(x-1)^{2/3}} &= \int (x-1)^{-2/3} d(x-1) = 3(x-1)^{1/3} \\ \int_0^3 \frac{dx}{(x-1)^{2/3}} &= \int_0^1 \frac{dx}{(x-1)^{2/3}} + \int_1^3 \frac{dx}{(x-1)^{2/3}} \\&= \left[3(x-1)^{1/3} \right]_0^{1^-} + \left[3(x-1)^{1/3} \right]_{1^+}^3 \\&= 3(0+1) + 3(\sqrt[3]{2} - 0) \\&= \underline{3 + 3\sqrt[3]{2}}\end{aligned}$$



Example


Does the integral $\int_1^{\infty} e^{-x^2} dx$ converge?

Solution

$$\int_1^{\infty} e^{-x^2} dx = \lim_{b \rightarrow \infty} \int_1^b e^{-x^2} dx$$

$$\int_1^b e^{-x^2} dx \leq \int_1^b e^{-x} dx = -e^{-b} + e^{-1} < e^{-1} \approx 0.36788$$

The integral converges



Theorem – Direct Comparison Test

Let f and g be continuous on $[a, \infty)$ with $0 \leq f(x) \leq g(x)$ for all $x \geq a$. Then

1. $\int_a^{\infty} f(x) dx$ converges if $\int_a^{\infty} g(x) dx$ converges
2. $\int_a^{\infty} g(x) dx$ diverges if $\int_a^{\infty} f(x) dx$ diverges

Theorem – Limit Comparison Test

If the positive functions f and g are continuous on $[a, \infty)$, and if

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = L, \quad 0 < L < \infty$$

Then

$$\int_a^{\infty} f(x) dx \quad \text{and} \quad \int_a^{\infty} g(x) dx$$

Both converge or both diverge

Example

Show that $\int_1^{\infty} \frac{dx}{1+x^2}$ converges by comparison with $\int_1^{\infty} \frac{dx}{x^2}$. Find and compare the two integral values.


Solution

The functions $f(x) = \frac{1}{x^2}$ and $g(x) = \frac{1}{1+x^2}$ are positive and continuous on $[1, \infty)$. Also,

$$\begin{aligned}\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} &= \lim_{x \rightarrow \infty} \frac{\frac{1}{x^2}}{\frac{1}{1+x^2}} \\ &= \lim_{x \rightarrow \infty} \frac{1+x^2}{x^2} \\ &= 1\end{aligned}$$

Therefore, $\int_1^{\infty} \frac{dx}{1+x^2}$ converges because $\int_1^{\infty} \frac{dx}{x^2}$ converges.

$$\begin{aligned}\int_1^{\infty} \frac{dx}{1+x^2} &= \lim_{b \rightarrow \infty} \int_1^b \frac{dx}{1+x^2} \\ &= \lim_{b \rightarrow \infty} \left(\tan^{-1} b - \tan^{-1} 1 \right) \\ &= \frac{\pi}{2} - \frac{\pi}{4} \\ &= \frac{\pi}{4}\end{aligned}$$



Example

Let R be the region bounded by the graph of $y = x^{-1}$ and the x -axis, for $x \geq 1$.

- What is the volume of the solid generated when R is revolved about the x -axis?
- What is the surface area of the solid generated when R is revolved about the x -axis?
- What is the volume of the solid generated when R is revolved about the y -axis?

Solution


$$a) \quad V = \pi \int_1^{\infty} \frac{1}{x^2} dx$$

$$= -\pi \frac{1}{x} \Big|_1^{\infty}$$

$$= -\pi(0 - 1)$$

$$= \pi \text{ unit}^3$$

$$V = \pi \int_a^b (f(x))^2 dx$$



$$b) \quad S = 2\pi \int_1^{\infty} \frac{1}{x} \sqrt{1 + \left(-\frac{1}{x^2}\right)^2} dx$$

$$= 2\pi \int_1^{\infty} \frac{1}{x^3} \sqrt{x^4 + 1} dx$$

$$> 2\pi \int_1^{\infty} \frac{x^2}{x^3} dx \quad \sqrt{x^4 + 1} > \sqrt{x^4} = x^2$$

$$= 2\pi \int_1^{\infty} \frac{1}{x} dx$$

$$= 2\pi (\ln x) \Big|_1^{\infty}$$

$$= \infty \text{ unit}^2$$

$$S = 2\pi \int_a^b f(x) \sqrt{1 + f'(x)^2} dx$$

$$c) \quad V = 2\pi \int_1^{\infty} x \frac{1}{x} dx$$

$$= 2\pi x \Big|_1^{\infty}$$

$$= \infty \text{ unit}^3$$

$$V = 2\pi \int_a^b x \cdot f(x) dx \quad (\text{Shell method})$$

Exercises Section 2.6 – Improper Integrals

Evaluate the integrals

1. $\int_0^{\infty} \frac{dx}{x^2 + 1}$

2. $\int_0^4 \frac{dx}{\sqrt{4-x}}$

3. $\int_{-\infty}^2 \frac{2dx}{x^2 + 4}$

4. $\int_{-\infty}^{\infty} \frac{xdx}{(x^2 + 4)^{3/2}}$

5. $\int_1^{\infty} \frac{dx}{x\sqrt{x^2 - 1}}$

6. $\int_{-\infty}^{\infty} 2xe^{-x^2} dx$

7. $\int_0^1 (-\ln x) dx$

8. $\int_{-1}^4 \frac{dx}{\sqrt{|x|}}$

9. $\int_0^{\infty} e^{-3x} dx$

10. $\int_{-\infty}^{\infty} \frac{dx}{1+x^2}$

11. $\int_1^{10} \frac{dx}{(x-2)^{1/3}}$

12. $\int_1^{\infty} \frac{dx}{x^2}$

13. $\int_0^{\infty} \frac{dx}{(x+1)^3}$

14. $\int_{-\infty}^0 e^x dx$

15. $\int_1^{\infty} 2^{-x} dx$

16. $\int_{-\infty}^0 \frac{dx}{\sqrt[3]{2-x}}$

17. $\int_{4/\pi}^{\infty} \frac{1}{x^2} \sec^2\left(\frac{1}{x}\right) dx$

18. $\int_{e^2}^{\infty} \frac{dx}{x \ln^p x} \quad p > 1$

19. $\int_0^{\infty} \frac{p}{\sqrt[5]{p^2 + 1}} dp$

20. $\int_{-1}^1 \ln y^2 dy$

21. $\int_{-2}^6 \frac{dx}{\sqrt{|x-2|}}$

22. $\int_0^{\infty} xe^{-x} dx$

23. $\int_0^1 x \ln x dx$

24. $\int_1^{\infty} \frac{\ln x}{x^2} dx$

25. $\int_1^{\infty} (1-x)e^x dx$

26. $\int_{-\infty}^{\infty} \frac{e^x}{1+e^{2x}} dx$

27. $\int_0^1 \frac{dx}{\sqrt[3]{x}}$

28. $\int_1^{\infty} \frac{3}{\sqrt[3]{x}} dx$

29. $\int_1^{\infty} \frac{4}{\sqrt[4]{x}} dx$

30. $\int_0^2 \frac{dx}{x^3}$

31. $\int_1^{\infty} \frac{dx}{x^3}$

32. $\int_1^{\infty} \frac{6}{x^4} dx$

33. $\int_0^{\infty} \frac{dx}{\sqrt{x}(x+1)}$

34. $\int_{-\infty}^0 xe^{-4x} dx$

35. $\int_0^{\infty} xe^{-x/3} dx$

36. $\int_0^{\infty} x^2 e^{-x} dx$

37. $\int_0^{\infty} e^{-x} \cos x dx$

38. $\int_4^{\infty} \frac{1}{x(\ln x)^3} dx$

39. $\int_1^{\infty} \frac{\ln x}{x} dx$

40. $\int_{-\infty}^{\infty} \frac{4}{16+x^2} dx$

41. $\int_0^{\infty} \frac{x^3}{(x^2+1)^2} dx$

43. $\int_0^{\infty} \frac{e^x}{1+e^x} dx$


45. $\int_0^{\infty} \sin \frac{x}{2} dx$

42. $\int_0^{\infty} \frac{1}{e^x + e^{-x}} dx$


44. $\int_0^{\infty} \cos \pi x dx$

Find the area of the unbounded shaded region


46. $y = e^x, \quad -\infty < x \leq 1$




47. $y = -\ln x$



48. $y = \frac{1}{x^2 + 1}$



49. $y = \frac{8}{x^2 + 4}$



50. Find the area of the region R between the graph of $f(x) = \frac{1}{\sqrt{9-x^2}}$ and the x -axis on the interval $(-3, 3)$ (if it exists)

51. Find the volume of the region bounded by $f(x) = (x^2 + 1)^{-1/2}$ and the x -axis on the interval $[2, \infty)$ is revolved about the x -axis.

52. Find the volume of the region bounded by $f(x) = \sqrt{\frac{x+1}{x^3}}$ and the x -axis on the interval $[1, \infty)$ is revolved about the x -axis.

53. Find the volume of the region bounded by $f(x) = (x+1)^{-3}$ and the x -axis on the interval $[0, \infty)$ is revolved about the y -axis.

54. Find the volume of the region bounded by $f(x) = \frac{1}{\sqrt{x} \ln x}$ and the x -axis on the interval $[2, \infty)$ is revolved about the x -axis .
55. Find the volume of the region bounded by $f(x) = \frac{\sqrt{x}}{\sqrt[3]{x^2 + 1}}$ and the x -axis on the interval $[0, \infty)$ is revolved about the x -axis .
56. Find the volume of the region bounded by $f(x) = (x^2 - 1)^{-1/4}$ and the x -axis on the interval $(1, 2]$ is revolved about the y -axis .
57. Find the volume of the region bounded by $f(x) = \tan x$ and the x -axis on the interval $\left[0, \frac{\pi}{2}\right)$ is revolved about the x -axis .
58. Find the volume of the region bounded by $f(x) = -\ln x$ and the x -axis on the interval $(0, 1]$ is revolved about the x -axis .
59. Find the volume of the solid generated by revolving the region bounded by the graphs of $y = xe^{-x}$, $y = 0$, and $x = 0$ about the x -axis .

Consider the region satisfying the inequalities

- Find the area of the region
 - Find the volume of the solid generated by revolving the region about the x -axis .
 - Find the volume of the solid generated by revolving the region about the y -axis .
60. $y \leq e^{-x}$, $y \geq 0$, $x \geq 0$
61. $y \leq \frac{1}{x^2}$, $y \geq 0$, $x \geq 1$
62. Find the perimeter of the hypocycloid of four cusps $x^{2/3} + y^{2/3} = 4$
63. Find the arc length of the graph $y = \sqrt{16 - x^2}$ over the interval $[0, 4]$
64. The region bounded by $(x - 2)^2 + y^2 = 1$ is revolved about the y -axis to form a torus. Find the surface area of the torus.
65. Find the surface area formed by revolving the graph $y = 2e^{-x}$ on the interval $[0, \infty)$ about the x -axis
66. The magnetic potential P at a point on the axis of a circular coil is given by

$$P = \frac{2\pi N I r}{k} \int_c^\infty \frac{1}{(r^2 + x^2)^{3/2}} dx$$


Where N , I , r , k , and c are constants. Find P .

67. A “semi-infinite” uniform rod occupies the nonnegative x -axis. The rod has a linear density δ , which means that a segment of length dx has a mass of δdx . A particle of mass M is located at the point $(-a, 0)$. The gravitational force F that the rod exerts on the mass is given by

$$F = \int_0^{\infty} \frac{GM\delta}{(a+x)^2} dx$$

Where G is the gravitational constant. Find F .

68. Let R be the region bounded by the graph of $f(x) = x^{-p}$ and the x -axis
- Let S be the solid generated when R is revolved about the x -axis. For what values of p is the volume of S finite for $0 < x \leq 1$?
 - Let S be the solid generated when R is revolved about the y -axis. For what values of p is the volume of S finite for $0 < x \leq 1$?
 - Let S be the solid generated when R is revolved about the x -axis. For what values of p is the volume of S finite for $x \geq 1$?
 - Let S be the solid generated when R is revolved about the y -axis. For what values of p is the volume of S finite for $x \geq 1$?
69. The solid formed by revolving (about the x -axis) the unbounded region lying between the graph of $f(x) = \frac{1}{x}$ and the x -axis ($x \geq 1$) is called **Gabriel's Horn**.



Show that this solid has a finite volume and an infinite surface area

70. Water is drained from a 3000-gal tank at a rate that starts at 100 gal/hr. and decreases continuously by 5% /hr. If the drain left open indefinitely, how much water drains from the tank? Can a full tank be emptied at this rate?