

Solution

Section 4.2 – Calculus with Parametric Curves

Exercise

Find all the points at which the curve has the given slope. $x = 4\cos t$, $y = 4\sin t$; $\text{slope} = \frac{1}{2}$


Solution

$$\begin{aligned} \frac{dy}{dx} &= \frac{4\cos t}{-4\sin t} & \frac{dy}{dx} &= \frac{dy/dt}{dx/dt} \\ &= -\cot t = \frac{1}{2} \end{aligned}$$

$$\cot t = -\frac{1}{2} \Rightarrow t = \cot^{-1}\left(-\frac{1}{2}\right) \quad t \in QII \text{ \& } QIV$$

$$\begin{cases} x = -\cos\left(\cot^{-1}\frac{1}{2}\right) = -\frac{1}{\sqrt{5}} \\ y = 4\sin\left(\cot^{-1}\frac{1}{2}\right) = \frac{8}{\sqrt{5}} \end{cases} \rightarrow \left(-\frac{\sqrt{5}}{5}, \frac{8\sqrt{5}}{5}\right)$$

$$\begin{cases} x = \cos\left(\cot^{-1}\frac{1}{2}\right) = \frac{1}{\sqrt{5}} \\ y = -4\sin\left(\cot^{-1}\frac{1}{2}\right) = -\frac{8}{\sqrt{5}} \end{cases} \rightarrow \left(\frac{\sqrt{5}}{5}, -\frac{8\sqrt{5}}{5}\right)$$



Exercise

Find all the points at which the curve has the given slope. $x = 2\cos t$, $y = 8\sin t$; $\text{slope} = -1$


Solution

$$\begin{aligned} \frac{dy}{dx} &= \frac{8\cos t}{-2\sin t} & \frac{dy}{dx} &= \frac{dy/dt}{dx/dt} \\ &= -4\cot t = -1 \end{aligned}$$

$$\cot t = \frac{1}{4} \Rightarrow t = \cot^{-1}\left(\frac{1}{4}\right) \quad t \in QI \text{ \& } QIII$$

$$\begin{cases} x = 2\cos\left(\cot^{-1}\frac{1}{4}\right) = \frac{2}{\sqrt{17}} \\ y = 8\sin\left(\cot^{-1}\frac{1}{4}\right) = \frac{32}{\sqrt{17}} \end{cases} \rightarrow \left(\frac{2\sqrt{17}}{17}, \frac{32\sqrt{17}}{17}\right)$$

$$\begin{cases} x = -2\cos\left(\cot^{-1}\frac{1}{4}\right) = -\frac{2}{\sqrt{17}} \\ y = -8\sin\left(\cot^{-1}\frac{1}{4}\right) = -\frac{32}{\sqrt{17}} \end{cases} \rightarrow \left(-\frac{2\sqrt{17}}{17}, -\frac{32\sqrt{17}}{17}\right)$$



Exercise

Find all the points at which the curve has the given slope. $x = t + \frac{1}{t}$, $y = t - \frac{1}{t}$; slope = 1

Solution

$$\begin{aligned}\frac{dy}{dx} &= \frac{1 + \frac{1}{t^2}}{1 - \frac{1}{t^2}} & \frac{dy}{dx} &= \frac{dy/dt}{dx/dt} \\ &= \frac{t^2 + 1}{t^2 - 1} = 1\end{aligned}$$

$t^2 + 1 \neq 1 \therefore$ There are **no** points on this curve with slope 1.


Exercise

Find all the points at which the curve has the given slope. $x = 2 + \sqrt{t}$, $y = 2 - 4t$; slope = -8

Solution

$$\begin{aligned}\frac{dy}{dx} &= \frac{-4}{\frac{1}{2\sqrt{t}}} & \frac{dy}{dx} &= \frac{dy/dt}{dx/dt} \\ &= -8\sqrt{t} \\ &= -8\end{aligned}$$

$$t = 1 \rightarrow (3, -2)$$



Exercise

Find an equation of the line tangent to the curve at the point corresponding to the given value of t .

$$x = \sin t, \quad y = \cos t, \quad t = \frac{\pi}{4}$$

Solution

$$\begin{aligned}\frac{dy}{dx} &= \frac{-\sin t}{\cos t} & \frac{dy}{dx} &= \frac{dy/dt}{dx/dt} \\ &= -\tan t \Big|_{t=\frac{\pi}{4}} \\ &= -1\end{aligned}$$

$$\text{At } t = \frac{\pi}{4}$$

$$\begin{cases} x = \sin \frac{\pi}{4} = \frac{\sqrt{2}}{2} \\ y = \cos \frac{\pi}{4} = \frac{\sqrt{2}}{2} \end{cases} \rightarrow \left(\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2} \right)$$

The equation of the tangent line is

$$y = -\left(x - \frac{\sqrt{2}}{2}\right) + \frac{\sqrt{2}}{2}$$

$$\underline{= -x + \sqrt{2} \quad |}$$

Exercise

Find an equation of the line tangent to the curve at the point corresponding to the given value of t .

$$x = t^2 - 1, \quad y = t^3 + t, \quad t = 2$$

Solution

$$\frac{dy}{dx} = \frac{3t^2 + 1}{2t} \Big|_{t=2}$$

$$\frac{dy}{dx} = \frac{dy / dt}{dx / dt}$$

$$\underline{= \frac{13}{4} \quad |}$$

At $t = 2$

$$\begin{cases} x = 3 \\ y = 10 \end{cases} \rightarrow (3, 10)$$

The equation of the tangent line is

$$y = \frac{13}{4}(x - 3) + 10$$

$$\underline{= \frac{13}{4}x + \frac{1}{4} \quad |}$$

Exercise

Find an equation of the line tangent to the curve at the point corresponding to the given value of t .

$$x = e^t, \quad y = \ln(t + 1), \quad t = 0$$

Solution

$$\frac{dy}{dx} = \frac{1}{t+1} \Big|_{t=0}$$

$$\frac{dy}{dx} = \frac{dy / dt}{dx / dt}$$

$$\underline{= 1 \quad |}$$

At $t = 0$

$$\begin{cases} x = 1 \\ y = 0 \end{cases} \rightarrow (1, 0)$$

The equation of the tangent line is

$$\underline{y = x - 1}$$

Exercise

Find an equation of the line tangent to the curve at the point corresponding to the given value of t .

$$x = \cos t + t \sin t, \quad y = \sin t - t \cos t, \quad t = \frac{\pi}{4}$$

Solution

$$\begin{aligned} \frac{dy}{dx} &= \frac{\cos t - \cos t + t \sin t}{-\sin t + \sin t + t \cos t} & \frac{dy}{dx} &= \frac{dy / dt}{dx / dt} \\ &= \tan t \Big|_{t=\frac{\pi}{4}} \\ &= 1 \end{aligned}$$

$$\text{At } t = \frac{\pi}{4}$$

$$\begin{cases} x = \frac{\sqrt{2}}{2} + \frac{\pi}{4} \frac{\sqrt{2}}{2} \\ y = \frac{\sqrt{2}}{2} - \frac{\pi}{4} \frac{\sqrt{2}}{2} \end{cases} \rightarrow \left(\frac{4\sqrt{2} + \pi\sqrt{2}}{8}, \frac{4\sqrt{2} - \pi\sqrt{2}}{8} \right)$$

The equation of the tangent line is

$$\begin{aligned} y &= x - \frac{4\sqrt{2} + \pi\sqrt{2}}{8} + \frac{4\sqrt{2} - \pi\sqrt{2}}{8} & y &= m(x - x_0) + y_0 \\ &= x - \frac{\pi\sqrt{2}}{4} \end{aligned}$$

Exercise

Find an equation of the line tangent to the curve at the point corresponding to the given value of t .

$$x = 6t, \quad y = t^2 + 4, \quad t = 1$$

Solution

$$\begin{aligned} \frac{dy}{dx} &= \frac{2t}{6} \Big|_{t=1} & \frac{dy}{dx} &= \frac{dy / dt}{dx / dt} \\ &= \frac{1}{3} \end{aligned}$$

$$\text{At } t = 1:$$

$$\begin{cases} x = 6 \\ y = 5 \end{cases} \rightarrow (6, 5)$$

The equation of the tangent line is

$$y = \frac{1}{3}(x - 6) + 5$$

$$y = m(x - x_0) + y_0$$

$$\underline{= \frac{1}{3}x + 3}$$

Exercise

Find an equation of the line tangent to the curve at the point corresponding to the given value of t .

$$x = t - 2, \quad y = \frac{1}{t} + 3, \quad t = 1$$

Solution

$$\frac{dy}{dx} = -\frac{1}{t^2} \Big|_{t=1}$$

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt}$$

$$\underline{= -1}$$

At $t = 1$:

$$\begin{cases} x = -1 \\ y = 4 \end{cases} \rightarrow (-1, 4)$$

The equation of the tangent line is

$$y = -(x + 1) + 4$$

$$y = m(x - x_0) + y_0$$

$$\underline{= -x + 3}$$

Exercise

Find an equation of the line tangent to the curve at the point corresponding to the given value of t .

$$x = t^2 - t + 2, \quad y = t^3 - 3t, \quad t = -1$$

Solution

$$\frac{dy}{dx} = \frac{3t^2 - 3}{2t - 1} \Big|_{t=-1}$$

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt}$$

$$\underline{= 0}$$

At $t = -1$

$$\begin{cases} x = 4 \\ y = 2 \end{cases} \rightarrow (4, 2)$$

The equation of the tangent line is

$$\underline{y = 2}$$

$$y = m(x - x_0) + y_0$$

Exercise

Find an equation of the line tangent to the curve at the point corresponding to the given value of t .

$$x = -t^2 + 3t, \quad y = 2t^{3/2}, \quad t = \frac{1}{4}$$

Solution

$$\begin{aligned} \frac{dy}{dx} &= \frac{3t^{1/2}}{-2t+3} \Big|_{t=1/4} & \frac{dy}{dx} &= \frac{dy/dt}{dx/dt} \\ &= \frac{3}{2} \frac{1}{-\frac{1}{2}+3} \\ &= \frac{3}{5} \end{aligned}$$

At $t = \frac{1}{4}$:

$$\begin{cases} x = -\frac{1}{16} + \frac{3}{4} = \frac{11}{16} \\ y = \frac{1}{4} \end{cases} \rightarrow \left(\frac{11}{16}, \frac{1}{4} \right)$$

The equation of the tangent line is

$$\begin{aligned} y &= \frac{3}{5} \left(x - \frac{11}{16} \right) + \frac{1}{4} & y &= m(x - x_0) + y_0 \\ &= \frac{3}{5}x - \frac{13}{80} \end{aligned}$$

Exercise

Find the tangent to the curve at the point defined by the given value of t . Also find the value of $\frac{d^2y}{dx^2}$ at

this point $x = \sin 2\pi t, \quad y = \cos 2\pi t, \quad t = -\frac{1}{6}$

Solution

$$x = \sin 2\pi \left(-\frac{1}{6} \right) = -\sin \left(\frac{\pi}{3} \right) = -\frac{\sqrt{3}}{2}$$

$$y = \cos 2\pi \left(-\frac{1}{6} \right) = \cos \left(\frac{\pi}{3} \right) = \frac{1}{2}$$

The point $\left(-\frac{\sqrt{3}}{2}, \frac{1}{2} \right)$

$$\frac{dx}{dt} = 2\pi \cos 2\pi t$$

$$\frac{dy}{dt} = -2\pi \sin 2\pi t$$

$$\begin{aligned}\frac{dy}{dx} &= \frac{-2\pi \sin 2\pi t}{2\pi \cos 2\pi t} \\ &= -\tan 2\pi t\end{aligned}$$

$$\frac{dy}{dx} = \frac{dy / dt}{dx / dt}$$

$$\begin{aligned}\left. \frac{dy}{dx} \right|_{t=-\frac{1}{6}} &= -\tan 2\pi \left(-\frac{1}{6}\right) \\ &= -\tan \left(-\frac{\pi}{3}\right) \\ &= \sqrt{3}\end{aligned}$$

The tangent to the curve at the point $\left(-\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$ is:

$$\begin{aligned}y &= \sqrt{3} \left(x + \frac{\sqrt{3}}{2}\right) + \frac{1}{2} \\ &= \sqrt{3}x + 2\end{aligned} \qquad y = m(x - x_0) + y_0$$

$$\frac{dy'}{dt} = \frac{d}{dt}(-\tan 2\pi t) = -2\pi \sec^2 2\pi t$$

$$\begin{aligned}\frac{d^2y}{dx^2} &= \frac{-2\pi \sec^2 2\pi t}{2\pi \cos 2\pi t} \\ &= -\frac{1}{\cos^3 2\pi t}\end{aligned} \qquad \frac{d^2y}{dx^2} = \frac{dy' / dt}{dx / dt}$$

$$\begin{aligned}\left. \frac{d^2y}{dx^2} \right|_{t=-\frac{1}{6}} &= -\frac{1}{\cos^3 \left(-\frac{\pi}{3}\right)} \\ &= -8\end{aligned}$$

Exercise

Find the tangent to the curve at the point defined by the given value of t . Also find the value of $\frac{d^2y}{dx^2}$ at

this point $x = \cos t, \quad y = \sqrt{3} \cos t, \quad t = \frac{2\pi}{3}$

Solution

$$x = \cos \left(\frac{2\pi}{3}\right) = -\frac{1}{2}$$

$$y = \sqrt{3} \cos \left(\frac{2\pi}{3}\right) = -\frac{\sqrt{3}}{2}$$

The point $\left(-\frac{1}{2}, -\frac{\sqrt{3}}{2}\right)$

$$\frac{dx}{dt} = -\sin t, \quad \frac{dy}{dt} = -\sqrt{3} \sin t$$

$$\frac{dy}{dx} = \frac{-\sqrt{3} \sin t}{-\sin t} = \sqrt{3} \qquad \frac{dy}{dx} = \frac{dy / dt}{dx / dt}$$

$$\left. \frac{dy}{dx} \right|_{t=\frac{2\pi}{3}} = \sqrt{3}$$

The tangent to the curve at the point $\left(-\frac{1}{2}, -\frac{\sqrt{3}}{2}\right)$ is:

$$y = \sqrt{3}\left(x + \frac{1}{2}\right) - \frac{\sqrt{3}}{2} \qquad y = m(x - x_0) + y_0$$

$$= \sqrt{3}x$$

$$\frac{dy'}{dt} = \frac{d}{dt}(\sqrt{3}) = 0$$

$$\frac{d^2y}{dx^2} = \frac{0}{-\sin t} \qquad \frac{d^2y}{dx^2} = \frac{dy' / dt}{dx / dt}$$

$$= 0$$

$$\left. \frac{d^2y}{dx^2} \right|_{t=\frac{2\pi}{3}} = 0$$

Exercise

Find the tangent to the curve at the point defined by the given value of t . Also find the value of $\frac{d^2y}{dx^2}$ at

this point $x = t, \quad y = \sqrt{t}, \quad t = \frac{1}{4}$

Solution

$$x = \frac{1}{4}$$

$$y = \sqrt{\frac{1}{4}} = \frac{1}{2}$$

The point $\left(\frac{1}{4}, \frac{1}{2}\right)$

$$\frac{dx}{dt} = 1, \quad \frac{dy}{dt} = \frac{1}{2\sqrt{t}}$$

$$\frac{dy}{dx} = \frac{1}{2\sqrt{t}} \cdot 1 \qquad \frac{dy}{dx} = \frac{dy / dt}{dx / dt}$$

$$= \frac{1}{2\sqrt{t}}$$

$$\left. \frac{dy}{dx} \right|_{t=\frac{1}{4}} = \frac{1}{2\sqrt{\frac{1}{4}}} \\ = 1$$

The tangent is:

$$y = \left(x - \frac{1}{4} \right) + \frac{1}{2} \\ = x + \frac{1}{4}$$

$$y = m(x - x_0) + y_0$$

$$\frac{dy'}{dt} = \frac{d}{dt} \left(\frac{1}{2\sqrt{t}} \right) \\ = -\frac{1}{4} t^{-3/2}$$

$$\frac{d^2y}{dx^2} = \frac{-\frac{1}{4} t^{-3/2}}{1} \\ = -\frac{1}{4} t^{-3/2}$$

$$\frac{d^2y}{dx^2} = \frac{dy' / dt}{dx / dt}$$

$$\left. \frac{d^2y}{dx^2} \right|_{t=\frac{1}{4}} = -\frac{1}{4} \left(\frac{1}{4} \right)^{-3/2} \\ = -2$$

Exercise

Find the tangent to the curve at the point defined by the given value of t . Also find the value of $\frac{d^2y}{dx^2}$ at

this point $x = \sec^2 t - 1$, $y = \tan t$, $t = -\frac{\pi}{4}$

Solution

$$= -\frac{1}{2}x + \frac{1}{2}$$

$$y = \tan\left(-\frac{\pi}{4}\right) = -1$$

The point $(1, -1)$

$$\frac{dx}{dt} = 2\sec^2 t \tan t, \quad \frac{dy}{dt} = \sec^2 t$$

$$\begin{aligned}\frac{dy}{dx} &= \frac{\sec^2 t}{2 \sec^2 t \tan t} \\ &= \frac{1}{2 \tan t}\end{aligned}$$

$$\frac{dy}{dx} = \frac{dy / dt}{dx / dt}$$

$$\begin{aligned}\left. \frac{dy}{dx} \right|_{t=-\frac{\pi}{4}} &= \frac{1}{2 \tan\left(-\frac{\pi}{4}\right)} \\ &= \underline{-\frac{1}{2}}\end{aligned}$$

The tangent is:

$$y = -\frac{1}{2}(x-1) - 1$$

$$y = m(x - x_0) + y_0$$

$$\underline{= -\frac{1}{2}x + \frac{1}{2}}$$

$$\begin{aligned}\frac{dy'}{dt} &= \frac{d}{dt}\left(\frac{1}{2 \tan t}\right) \\ &= \frac{1 - \sec^2 t}{2 \tan^2 t} \\ &= -\frac{1}{2} \frac{\frac{1}{\cos^2 t}}{\frac{\sin^2 t}{\cos^2 t}} \\ &= -\frac{1}{2} \csc^2 t\end{aligned}$$

$$\begin{aligned}\frac{d^2 y}{dx^2} &= \frac{-\frac{1}{2} \csc^2 t}{2 \sec^2 t \tan t} \\ &= -\frac{1}{4} \frac{\frac{\sin^2 t}{\cos^2 t}}{\frac{1}{\cos^2 t} \frac{\sin t}{\cos t}} \\ &= -\frac{1}{4} \frac{\cos^3 t}{\sin^3 t} \\ &= \underline{-\frac{1}{4} \cot^3 t}\end{aligned}$$

$$\frac{d^2 y}{dx^2} = \frac{dy' / dt}{dx / dt}$$

$$\begin{aligned}\left. \frac{d^2 y}{dx^2} \right|_{t=-\frac{\pi}{4}} &= -\frac{1}{4} \cot^3\left(-\frac{\pi}{4}\right) \\ &= \underline{\frac{1}{4}}\end{aligned}$$

Exercise

Find the tangent to the curve at the point defined by the given value of t . Also find the value of d^2y/dx^2 at this point $x = \frac{1}{t+1}$, $y = \frac{t}{t-1}$, $t = 2$

Solution

$$x = \frac{1}{2+1} = \frac{1}{3}$$

$$y = \frac{2}{2-1} = 2$$

The point $\left(\frac{1}{3}, 2\right)$

$$\frac{dx}{dt} = \frac{-1}{(t+1)^2}$$

$$\begin{aligned}\frac{dy}{dt} &= \frac{t-1-t}{(t-1)^2} \\ &= \frac{-1}{(t-1)^2}\end{aligned}$$

$$\begin{aligned}\frac{dy}{dx} &= \frac{\frac{-1}{(t-1)^2}}{\frac{-1}{(t+1)^2}} \\ &= \frac{(t+1)^2}{(t-1)^2}\end{aligned}$$

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt}$$

$$\begin{aligned}\left.\frac{dy}{dx}\right|_{t=2} &= \frac{(2+1)^2}{(2-1)^2} \\ &= 9\end{aligned}$$

The tangent is:

$$\begin{aligned}y &= 9\left(x - \frac{1}{3}\right) + 2 \\ &= 9x - 1\end{aligned}$$

$$y = m(x - x_0) + y_0$$

$$\begin{aligned}\frac{dy'}{dt} &= \frac{d}{dt}\left(\frac{t+1}{t-1}\right)^2 \\ &= 2\left(\frac{t+1}{t-1}\right)\left(\frac{t-1-t-1}{(t-1)^2}\right) \\ &= -4\frac{t+1}{(t-1)^3}\end{aligned}$$

$$\frac{d^2y}{dx^2} = -4 \frac{t+1}{(t-1)^3} \frac{(t+1)^2}{-1} \quad \frac{d^2y}{dx^2} = \frac{dy' / dt}{dx / dt}$$

$$= 4 \frac{(t+1)^3}{(t-1)^3} \Bigg|$$

$$\frac{d^2y}{dx^2} \Bigg|_{t=2} = 4 \frac{(2+1)^3}{(2-1)^3}$$

$$= 108 \Bigg|$$

Exercise

Find the tangent to the curve at the point defined by the given value of t . Also find the value of d^2y / dx^2 at this point $x = t + e^t$, $y = 1 - e^t$, $t = 0$

Solution

$$x = 0 + e^0 = 1$$

$$y = 1 - e^0 = 0$$

The point (1, 0)

$$\frac{dx}{dt} = 1 + e^t$$

$$\frac{dy}{dt} = -e^t$$

$$\frac{dy}{dx} = \frac{-e^t}{1 + e^t} \quad \frac{dy}{dx} = \frac{dy / dt}{dx / dt}$$

$$\frac{dy}{dx} \Bigg|_{t=0} = -\frac{e^0}{1 + e^0}$$

$$= -\frac{1}{2} \Bigg|$$

The tangent is:

$$y = -\frac{1}{2}(x - 1)$$

$$= -\frac{1}{2}x + \frac{1}{2} \Bigg|$$

$$\frac{dy'}{dt} = \frac{d}{dt} \left(\frac{-e^t}{1 + e^t} \right)$$

$$\begin{aligned}
&= \frac{-e^t(1+e^t) - e^t(-e^t)}{(1+e^t)^2} \\
&= \frac{-e^t - e^{2t} + e^{2t}}{(1+e^t)^2} \\
&= \frac{-e^t}{(1+e^t)^2}
\end{aligned}$$

$$\begin{aligned}
\frac{d^2y}{dx^2} &= \frac{-e^t}{(1+e^t)^2} \cdot \frac{1}{1+e^t} & \frac{d^2y}{dx^2} &= \frac{dy' / dt}{dx / dt} \\
&= \frac{-e^t}{(1+e^t)^3}
\end{aligned}$$

$$\begin{aligned}
\left. \frac{d^2y}{dx^2} \right|_{t=0} &= \frac{-e^0}{(1+e^0)^3} \\
&= -\frac{1}{8}
\end{aligned}$$

Exercise

Find the tangent to the curve at the point defined by the given value of t . Also find the value of $\frac{d^2y}{dx^2}$ at this point $x = 4t, \quad y = 3t - 2, \quad t = 3$

Solution

$$\frac{dx}{dt} = 4$$

$$\frac{dy}{dt} = 3$$

$$\frac{dy}{dx} = \frac{3}{4} \Big|_{t=3} = \frac{3}{4}$$

$$\frac{dy}{dx} = \frac{dy / dt}{dx / dt}$$

$$t = 3 \Rightarrow \begin{cases} x = 12 \\ y = 7 \end{cases}$$

The tangent to the curve at the point (12, 7)

$$y = \frac{3}{4}(x - 12) + 7$$

$$y = m(x - x_0) + y_0$$

$$= \frac{3}{4}x - 2 \Big|$$

$$\frac{dy'}{dt} = \frac{d}{dt} \left(\frac{3}{4} \right) = 0$$

$$\frac{d^2y}{dx^2} = \underline{0}$$

$$\frac{d^2y}{dx^2} = \frac{dy' / dt}{dx / dt}$$

Exercise

Find the tangent to the curve at the point defined by the given value of t . Also find the value of $\frac{d^2y}{dx^2}$ at this point $x = \sqrt{t}$, $y = 3t - 1$, $t = 1$

Solution

$$\frac{dx}{dt} = \frac{1}{2\sqrt{t}}$$

$$\frac{dy}{dt} = 3$$

$$\frac{dy}{dx} = 6\sqrt{t} \Big|_{t=1}$$

$$\frac{dy}{dx} = \frac{dy / dt}{dx / dt}$$

$$= 6 \Big|$$

$$t = 1 \Rightarrow \begin{cases} x = 1 \\ y = 2 \end{cases}$$

The tangent to the curve at the point (1, 2)

$$y = 6(x - 1) + 2$$

$$y = m(x - x_0) + y_0$$

$$= 6x - 4 \Big|$$

$$\begin{aligned} \frac{dy'}{dt} &= \frac{d}{dt} (6\sqrt{t}) \\ &= \frac{3}{\sqrt{t}} \end{aligned}$$

$$\begin{aligned} \frac{d^2y}{dx^2} &= \frac{3}{\sqrt{t}} \cdot 2\sqrt{t} \\ &= 6 \Big| \end{aligned}$$

$$\frac{d^2y}{dx^2} = \frac{dy' / dt}{dx / dt}$$

Exercise

Find the tangent to the curve at the point defined by the given value of t . Also find the value of $\frac{d^2y}{dx^2}$ at this point $x = t + 1, \quad y = t^2 + 3t, \quad t = -1$

Solution

$$\frac{dx}{dt} = 1$$

$$\frac{dy}{dt} = 2t + 3$$

$$\frac{dy}{dx} = 2t + 3 \Big|_{t=-1}$$

$$= 1 \Big|$$

$$\frac{dy}{dx} = \frac{dy / dt}{dx / dt}$$

$$t = -1 \Rightarrow \begin{cases} x = 2 \\ y = 4 \end{cases}$$

The tangent to the curve at the point $(2, 4)$

$$y = (x - 2) + 4$$

$$y = m(x - x_0) + y_0$$

$$= x + 2 \Big|$$

$$\frac{dy'}{dt} = \frac{d}{dt}(2t + 3) = 2$$

$$\frac{d^2y}{dx^2} = 2 \Big|$$

$$\frac{d^2y}{dx^2} = \frac{dy' / dt}{dx / dt}$$

Exercise

Find the tangent to the curve at the point defined by the given value of t . Also find the value of $\frac{d^2y}{dx^2}$ at this point $x = t^2 + 5t + 4, \quad y = 4t, \quad t = 0$

Solution

$$\frac{dx}{dt} = 2t + 5$$

$$\frac{dy}{dt} = 4$$

$$\frac{dy}{dx} = \frac{4}{2t + 5} \Big|_{t=0}$$

$$= \frac{4}{5} \Big|$$

$$\frac{dy}{dx} = \frac{dy / dt}{dx / dt}$$

$$t = 0 \Rightarrow \begin{cases} x = 4 \\ y = 0 \end{cases}$$

The tangent to the curve at the point (4, 0)

$$y = \frac{4}{5}(x - 4) \qquad y = m(x - x_0) + y_0$$

$$\underline{= \frac{4}{5}x - \frac{16}{5} \mid}$$

$$\frac{dy'}{dt} = \frac{d}{dt} \left(\frac{4}{2t+5} \right)$$

$$= \frac{-8}{(2t+5)^2}$$

$$\frac{d^2y}{dx^2} = \frac{-8}{(2t+5)^2} \cdot \frac{1}{2t+5} \qquad \frac{d^2y}{dx^2} = \frac{dy' / dt}{dx / dt}$$

$$= \frac{-8}{(2t+5)^3} \Big|_{t=0}$$

$$\underline{= -\frac{8}{125} \mid}$$

Exercise

Find the tangent to the curve at the point defined by the given value of t . Also find the value of $\frac{d^2y}{dx^2}$ at

this point $x = 4 \cos \theta, \quad y = 4 \sin \theta, \quad \theta = \frac{\pi}{4}$

Solution

$$\frac{dx}{d\theta} = -4 \sin \theta$$

$$\frac{dy}{d\theta} = 4 \cos \theta$$

$$\frac{dy}{dx} = \frac{4 \cos \theta}{-4 \sin \theta} \qquad \frac{dy}{dx} = \frac{dy / d\theta}{dx / d\theta}$$

$$= -\cot \theta \Big|_{\theta = \frac{\pi}{4}}$$

$$\underline{= -1 \mid}$$

$$\theta = \frac{\pi}{4} \Rightarrow \begin{cases} x = 2\sqrt{2} \\ y = 2\sqrt{2} \end{cases}$$

The tangent to the curve at the point $(2\sqrt{2}, 2\sqrt{2})$:

$$y = -(x - 2\sqrt{2}) + 2\sqrt{2}$$

$$= -x + 4\sqrt{2}$$

$$y = m(x - x_0) + y_0$$

$$\frac{dy'}{d\theta} = \frac{d}{d\theta}(-\cot \theta)$$

$$= \csc^2 \theta$$

$$\frac{d^2y}{dx^2} = \frac{\csc^2 \theta}{-4 \sin \theta}$$

$$= -\frac{1}{4} \csc^3 \theta \Big|_{\theta=\frac{\pi}{4}}$$

$$= -\frac{\sqrt{2}}{2}$$

$$\frac{d^2y}{dx^2} = \frac{dy' / d\theta}{dx / d\theta}$$

Exercise

Find the tangent to the curve at the point defined by the given value of t . Also find the value of $\frac{d^2y}{dx^2}$ at

this point $x = \cos \theta$, $y = 3 \sin \theta$, $\theta = 0$

Solution

$$\frac{dx}{d\theta} = -\sin \theta$$

$$\frac{dy}{d\theta} = 3 \cos \theta$$

$$\frac{dy}{dx} = \frac{3 \cos \theta}{-\sin \theta}$$

$$= -3 \cot \theta \Big|_{\theta=0}$$

$$= \infty$$

$$\frac{dy}{dx} = \frac{dy / d\theta}{dx / d\theta}$$

$$\theta = 0 \Rightarrow \begin{cases} x = 1 \\ y = 0 \end{cases}$$

The tangent to the curve at the point $(1, 0)$: $x = 1$

$$\frac{dy'}{d\theta} = \frac{d}{d\theta}(-3 \cot \theta)$$

$$= 3 \csc^2 \theta$$

$$\frac{d^2y}{dx^2} = \frac{3 \csc^2 \theta}{-\sin \theta}$$

$$\frac{d^2y}{dx^2} = \frac{dy' / d\theta}{dx / d\theta}$$

$$= -3 \csc^3 \theta \Big|_{\theta=0}$$

$$= \underline{\infty} \quad \text{undefined}$$

Exercise

Find the tangent to the curve at the point defined by the given value of t . Also find the value of $\frac{d^2y}{dx^2}$ at

this point $x = 2 + \sec \theta$, $y = 1 + 2 \tan \theta$, $\theta = \frac{\pi}{6}$

Solution

$$\frac{dx}{d\theta} = \sec \theta \tan \theta$$

$$\frac{dy}{d\theta} = 2 \sec^2 \theta$$

$$\frac{dy}{dx} = \frac{2 \sec^2 \theta}{\sec \theta \tan \theta}$$

$$\frac{dy}{dx} = \frac{dy / d\theta}{dx / d\theta}$$

$$= 2 \csc \theta \Big|_{\theta=\frac{\pi}{6}}$$

$$= \underline{4}$$

$$\theta = \frac{\pi}{6} \Rightarrow \begin{cases} x = 2 + \frac{2}{\sqrt{3}} \\ y = 1 + \frac{2\sqrt{3}}{3} \end{cases}$$

The tangent to the curve at the point $\left(2 + \frac{2\sqrt{3}}{3}, 1 + \frac{2\sqrt{3}}{3}\right)$:

$$y = 2 \left(x - 2 - \frac{2\sqrt{3}}{3} \right) + 1 + \frac{2\sqrt{3}}{3}$$

$$y = m(x - x_0) + y_0$$

$$= \underline{2x - 3 - \frac{2\sqrt{3}}{3}}$$

$$\frac{dy'}{d\theta} = \frac{d}{d\theta} (2 \csc \theta)$$

$$= -2 \csc \theta \cot \theta$$

$$\frac{d^2y}{dx^2} = \frac{-2 \csc \theta \cot \theta}{\sec \theta \tan \theta}$$

$$\frac{d^2y}{dx^2} = \frac{dy' / d\theta}{dx / d\theta}$$

$$= -2 \cot^3 \theta \Big|_{\theta=\frac{\pi}{6}}$$

$$= \underline{-6\sqrt{3}}$$

Exercise

Find the tangent to the curve at the point defined by the given value of t . Also find the value of $\frac{d^2y}{dx^2}$ at this point $x = \sqrt{t}, \quad y = \sqrt{t-1}, \quad t = 2$

Solution

$$\frac{dx}{dt} = \frac{1}{2\sqrt{t}}$$

$$\frac{dy}{dt} = \frac{1}{2\sqrt{t-1}}$$

$$\frac{dy}{dx} = \frac{2\sqrt{t}}{2\sqrt{t-1}} \Big|_{t=2}$$

$$= \sqrt{2} \Big|$$

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt}$$

$$t = 2 \Rightarrow \begin{cases} x = \sqrt{2} \\ y = 1 \end{cases}$$

The tangent to the curve at the point $(\sqrt{2}, 1)$

$$y = \sqrt{2}(x - \sqrt{2}) + 1 \\ = \sqrt{2}x - 1 \Big|$$

$$y = m(x - x_0) + y_0$$

$$\frac{dy'}{dt} = \frac{d}{dt} \left(\frac{\sqrt{t}}{\sqrt{t-1}} \right)$$

$$= \frac{\frac{1}{2}t - \frac{1}{2} - \frac{1}{2}t}{(t-1)^{3/2} \sqrt{t}}$$

$$= -\frac{1}{2} \frac{1}{(t-1)^{3/2} \sqrt{t}}$$

$$(U^n V^m)' = U^{n-1} V^{m-1} (nU'V + mUV')$$

$$\frac{d^2y}{dx^2} = -\frac{1}{2} \frac{1}{(t-1)^{3/2} \sqrt{t}} \cdot 2\sqrt{t}$$

$$= -\frac{1}{(t-1)^{3/2}} \Big|_{t=2}$$

$$= -1 \Big|$$

$$\frac{d^2y}{dx^2} = \frac{dy'/dt}{dx/dt}$$

Exercise

Find the tangent to the curve at the point defined by the given value of t . Also find the value of $\frac{d^2y}{dx^2}$ at

this point $x = \cos^3 \theta$, $y = \sin^3 \theta$, $\theta = \frac{\pi}{4}$

Solution

$$\frac{dx}{d\theta} = -3 \sin \theta \cos^2 \theta$$

$$\frac{dy}{d\theta} = 3 \cos \theta \sin^2 \theta$$

$$\frac{dy}{dx} = \frac{3 \cos \theta \sin^2 \theta}{-3 \sin \theta \cos^2 \theta} \qquad \frac{dy}{dx} = \frac{dy / d\theta}{dx / d\theta}$$

$$= -\tan \theta \Big|_{\theta=\frac{\pi}{4}}$$

$$= -1$$

$$\theta = \frac{\pi}{4} \Rightarrow \begin{cases} x = \frac{\sqrt{2}}{4} \\ y = \frac{\sqrt{2}}{4} \end{cases}$$

The tangent to the curve at the point $\left(\frac{\sqrt{2}}{4}, \frac{\sqrt{2}}{4}\right)$:

$$y = -\left(x - \frac{\sqrt{2}}{4}\right) + \frac{\sqrt{2}}{4} \qquad y = m(x - x_0) + y_0$$

$$= -x + \frac{\sqrt{2}}{2}$$

$$\frac{dy'}{d\theta} = \frac{d}{d\theta}(-\tan \theta)$$

$$= -\sec^2 \theta$$

$$\frac{d^2y}{dx^2} = \frac{-\sec^2 \theta}{-3 \sin \theta \cos^2 \theta} \qquad \frac{d^2y}{dx^2} = \frac{dy' / d\theta}{dx / d\theta}$$

$$= \frac{1}{3 \sin \theta \cos^4 \theta} \Big|_{\theta=\frac{\pi}{4}}$$

$$= \frac{4\sqrt{2}}{3}$$

Exercise

Find the tangent to the curve at the point defined by the given value of t . Also find the value of $\frac{d^2y}{dx^2}$ at this point $x = \theta - \sin \theta, \quad y = 1 - \cos \theta, \quad \theta = \pi$

Solution

$$\frac{dx}{d\theta} = 1 - \cos \theta$$

$$\frac{dy}{d\theta} = \sin \theta$$

$$\frac{dy}{dx} = \frac{\sin \theta}{1 - \cos \theta} \Big|_{\theta=\pi} \\ = 0$$

$$\frac{dy}{dx} = \frac{dy / d\theta}{dx / d\theta}$$

$$\theta = \pi \Rightarrow \begin{cases} x = \pi \\ y = 2 \end{cases}$$

The tangent to the curve at the point $(\pi, 2)$:

$$y = 2$$

$$y = m(x - x_0) + y_0$$

$$\begin{aligned} \frac{dy'}{d\theta} &= \frac{d}{d\theta} \left(\frac{\sin \theta}{1 - \cos \theta} \right) \\ &= \frac{\cos \theta - \cos^2 \theta - \sin^2 \theta}{(1 - \cos \theta)^2} \\ &= \frac{\cos \theta - 1}{(1 - \cos \theta)^2} \\ &= \frac{-1}{1 - \cos \theta} \end{aligned}$$

$$\begin{aligned} \frac{d^2y}{dx^2} &= \left(\frac{-1}{1 - \cos \theta} \right) \frac{1}{1 - \cos \theta} \Big|_{\theta=\pi} \\ &= \frac{-1}{(1 - \cos \theta)^2} \Big|_{\theta=\pi} \end{aligned}$$

$$\frac{d^2y}{dx^2} = \frac{dy' / d\theta}{dx / d\theta}$$

$$= -\frac{1}{4}$$

Exercise

Find the equations of the tangent lines at the point where the curve crosses itself

$$x = 2 \sin 2t, \quad y = 3 \sin t$$

Solution

$$x = y \rightarrow 2 \sin 2t = 3 \sin t$$

$$\Rightarrow t = 0, \pi$$

$$\frac{dx}{dt} = 4 \cos 2t, \quad \frac{dy}{dt} = 3 \cos t$$

$$\frac{dy}{dx} = \frac{3 \cos t}{4 \cos 2t}$$

$$\text{At } t = 0 \quad \frac{dy}{dx} = \frac{3}{4}$$


The point at $t = 0$ is $(0, 0)$

$$\text{The tangent line: } y = \frac{3}{4}x$$

$$\text{At } t = \pi \quad \frac{dy}{dx} = -\frac{3}{4}$$

The point at $t = \pi$ is $(0, 0)$

$$\text{The tangent line: } y = -\frac{3}{4}x$$



Exercise

Find the equations of the tangent lines at the point where the curve crosses itself

$$x = 2 - \pi \cos t, \quad y = 2t - \pi \sin t$$

Solution

The graph crosses itself at the point $(2, 0)$

$$x = 2 - \pi \cos t = 2$$

$$\cos t = 0$$


$$\Rightarrow t = \pm \frac{\pi}{2}$$

$$\frac{dx}{dt} = \pi \sin t$$

$$\frac{dy}{dt} = 2 - \pi \cos t$$

$$\frac{dy}{dx} = \frac{2 - \pi \cos t}{\pi \sin t}$$

$$\text{At } t = \frac{\pi}{2} \quad \frac{dy}{dx} = \frac{2}{\pi}$$



The tangent line: $y = \frac{2}{\pi}(x - 2)$

$$= \frac{2}{\pi}x - \frac{4}{\pi}$$

At $t = -\frac{\pi}{2}$ $\frac{dy}{dx} = -\frac{2}{\pi}$

The tangent line: $y = -\frac{2}{\pi}(x - 2)$

$$= -\frac{2}{\pi}x + \frac{4}{\pi}$$

Exercise

Find the equations of the tangent lines at the point where the curve crosses itself

$$x = t^2 - t, \quad y = t^3 - 3t - 1$$

Solution

The graph crosses itself at the point $(2, 1)$

$$x = t^2 - t = 2$$

$$t^2 - t - 2 = 0$$

$$\Rightarrow t = -1, 2$$

$$\frac{dx}{dt} = 2t - 1$$

$$\frac{dy}{dt} = 3t^2 - 3$$

$$\frac{dy}{dx} = \frac{3t^2 - 3}{2t - 1}$$


At $t = -1$ $\frac{dy}{dx} = 0$

The tangent line: $y = 1$

At $t = 2$ $\frac{dy}{dx} = 3$

The tangent line: $y = 3(x - 2) + 1$

$$= 3x - 5$$



Exercise

Find the equations of the tangent lines at the point where the curve crosses itself

$$x = t^3 - 6t, \quad y = t^2$$

Solution

The graph crosses itself at the point (0, 6)

$$y = t^2 = 6$$

$$\Rightarrow t = \pm\sqrt{6}$$

$$\frac{dx}{dt} = 3t^2 - 6, \quad \frac{dy}{dt} = 2t$$

$$\frac{dy}{dx} = \frac{2t}{3t^2 - 6}$$

$$\text{At } t = -\sqrt{6}$$


$$\begin{aligned} \frac{dy}{dx} &= \frac{-2\sqrt{6}}{12} \\ &= -\frac{\sqrt{6}}{6} \end{aligned}$$

$$\text{The tangent line: } y = -\frac{\sqrt{6}}{6}x + 6$$

$$\text{At } t = \sqrt{6}$$

$$\begin{aligned} \frac{dy}{dx} &= \frac{2\sqrt{6}}{12} \\ &= \frac{\sqrt{6}}{6} \end{aligned}$$

$$\text{The tangent line: } y = \frac{\sqrt{6}}{6}x + 6$$



Exercise

Find the slope of the curve $x = f(t)$, $y = g(t)$ at the given value of t . Define x and y as differentiable functions. $x^3 + 2t^2 = 9$, $2y^3 - 3t^2 = 4$, $t = 2$

Solution

$$x^3 + 2(2)^2 = 9$$

$$x^3 = 9 - 8 = 1$$

$$\rightarrow x = 1$$

$$2y^3 - 3(2)^2 = 4$$

$$2y^3 = 4 + 12 = 16$$

$$y^3 = 8$$

$$\Rightarrow \underline{y = 2}$$

$$x^3 + 2t^2 = 9 \Rightarrow 3x^2 \frac{dx}{dt} + 4t = 0$$

$$3x^2 \frac{dx}{dt} = -4t$$

$$\frac{dx}{dt} = -\frac{4t}{3x^2}$$

$$2y^3 - 3t^2 = 4 \Rightarrow 6y^2 \frac{dy}{dt} - 6t = 0$$

$$y^2 \frac{dy}{dt} = t$$

$$\frac{dy}{dt} = \frac{t}{y^2}$$

$$\frac{dy}{dx} = \frac{\frac{t}{y^2}}{-\frac{4t}{3x^2}}$$

$$\frac{dy}{dx} = \frac{dy / dt}{dx / dt}$$

$$= -\frac{3x^2}{4y^2}$$

$$\left. \frac{dy}{dx} \right|_{t=2} = -\frac{3(1)^2}{4(2)^2}$$

$$\underline{= -\frac{3}{16}}$$

Exercise

Find the slope of the curve $x = f(t)$, $y = g(t)$ at the given value of t . Define x and y as differentiable functions. $x + 2x^{3/2} = t^2 + t$, $y\sqrt{t+1} + 2t\sqrt{y} = 4$, $t = 0$

Solution

$$x + 2x^{3/2} = 0^2 + 0$$

$$x(1 + 2x^{1/2}) = 0$$

$$\rightarrow \boxed{x=0} \quad \cancel{x^{1/2} = -\frac{1}{2}} \quad (False)$$

$$y\sqrt{0+1} + 2(0)\sqrt{y} = 4$$

$$\Rightarrow \underline{y = 4}$$

$$x + 2x^{3/2} = t^2 + t$$

$$\frac{dx}{dt} + 3x^{1/2} \frac{dx}{dt} = 2t + 1$$

$$\frac{dx}{dt} (1 + 3x^{1/2}) = 2t + 1$$

$$\frac{dx}{dt} = \frac{2t + 1}{1 + 3x^{1/2}}$$

$$y\sqrt{t+1} + 2t\sqrt{y} = 4$$

$$\frac{dy}{dt} \sqrt{t+1} + \frac{1}{2} y (t+1)^{-1/2} + 2\sqrt{y} + 2t \left(\frac{1}{2} y^{-1/2} \right) \frac{dy}{dt} = 0$$

$$\frac{dy}{dt} \left(\sqrt{t+1} + \frac{t}{\sqrt{y}} \right) = -\frac{y}{2\sqrt{t+1}} - 2\sqrt{y}$$

$$\frac{dy}{dt} \left(\frac{\sqrt{t+1}\sqrt{y} + t}{\sqrt{y}} \right) = \frac{-y - 4\sqrt{t+1}\sqrt{y}}{2\sqrt{t+1}}$$

$$\frac{dy}{dt} = \frac{-y - 4\sqrt{t+1}\sqrt{y}}{2\sqrt{t+1}} \cdot \frac{\sqrt{y}}{\sqrt{t+1}\sqrt{y} + t}$$

$$\frac{dy}{dt} = \frac{-y\sqrt{y} - 4y\sqrt{t+1}}{2(t+1)\sqrt{y} + 2t\sqrt{t+1}}$$

$$\frac{dy}{dx} = \frac{-y\sqrt{y} - 4y\sqrt{t+1}}{2(t+1)\sqrt{y} + 2t\sqrt{t+1}} \cdot \frac{1 + 3\sqrt{x}}{2t + 1}$$

$$\frac{dy}{dx} = \frac{dy / dt}{dx / dt}$$

$$\left. \frac{dy}{dx} \right|_{t=0} = \frac{-4\sqrt{4} - 4(4)\sqrt{0+1}}{2(0+1)\sqrt{4} + 2(0)\sqrt{0+1}} \cdot \frac{1 + 3\sqrt{0}}{2(0) + 1} = -6$$

Exercise

Find the slope of the curve $x = f(t)$, $y = g(t)$ at the given value of t . Define x and y as differentiable functions. $t = \ln(x-t)$, $y = te^t$, $t = 0$

Solution

$$0 = \ln(x - 0)$$

$$\ln x = 0 \rightarrow \underline{x = 1}$$

$$y = (0)e^0 \Rightarrow \boxed{y = 0}$$

$$t = \ln(x - t)$$

$$1 = \frac{\frac{dx}{dt} - 1}{x - t}$$

$$\frac{dx}{dt} - 1 = x - t$$

$$\frac{dx}{dt} = x - t + 1$$

$$y = te^t$$

$$\begin{aligned} \frac{dy}{dt} &= e^t + te^t \\ &= e^t(1 + t) \end{aligned}$$

$$\frac{dy}{dx} = \frac{e^t(1+t)}{x-t+1}$$

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt}$$

$$\begin{aligned} \left. \frac{dy}{dx} \right|_{t=0} &= \frac{e^0(1+0)}{1-0+1} \\ &= \frac{1}{2} \end{aligned}$$

Exercise

Find $\frac{d^2y}{dx^2}$ for $x(t) = t - t^2$ $y(t) = t - t^3$

Solution

$$\frac{dx}{dt} = 1 - 2t$$

$$\frac{dy}{dt} = 1 - 3t^2$$

$$\left. \frac{dy}{dx} = \frac{1-3t^2}{1-2t} \right|$$

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt}$$

$$\begin{aligned} \frac{dy'}{dt} &= \frac{d}{dt} \left(\frac{-3t^2 + 1}{-2t + 1} \right) \\ &= \frac{-6t + 2}{(1 - 2t)^2} \end{aligned}$$

$$\frac{d}{dx} \left(\frac{ax^2 + bx + c}{dx^2 + ex + f} \right) = \frac{(ae - bd)x^2 + 2(af - cd)x + bf - ce}{(dx^2 + ex + f)^2}$$

$$\begin{aligned}\frac{d^2y}{dx^2} &= \frac{-6t+2}{(1-2t)^2} \cdot \frac{1}{1-2t} \\ &= \frac{-6t+2}{(1-2t)^3} \quad | \end{aligned}$$

$$\frac{d^2y}{dx^2} = \frac{dy' / dt}{dx / dt}$$

Exercise

Find $\frac{d^2y}{dx^2}$ for $x(t) = 2 \sec t$ $y(t) = 4 \tan t + 2$

Solution

$$\frac{dx}{dt} = 2 \sec t \tan t$$

$$\frac{dy}{dt} = 4 \sec^2 t$$

$$\frac{dy}{dx} = \frac{4 \sec^2 t}{2 \sec t \tan t}$$

$$= \frac{2 \sec t}{\tan t}$$

$$= \frac{2}{\sin t}$$

$$= 2 \csc t \quad |$$

$$\frac{dy}{dx} = \frac{dy / dt}{dx / dt}$$

$$\begin{aligned}\frac{dy'}{dt} &= \frac{d}{dt}(2 \csc t) \\ &= -2 \csc t \cot t \quad | \end{aligned}$$

$$\frac{d^2y}{dx^2} = \frac{-2 \csc t \cot t}{2 \sec t \tan t}$$

$$= -\frac{\csc t \cot t}{\frac{1}{\csc t \cot t}}$$

$$= -\csc^2 t \cot^2 t \quad |$$

$$\frac{d^2y}{dx^2} = \frac{dy' / dt}{dx / dt}$$

Exercise

Find $\frac{d^2y}{dx^2}$ for $x(t) = t^2 + 1$ $y(t) = 2t - 1$

Solution

$$\frac{dx}{dt} = 2t$$

$$\frac{dy}{dt} = 2$$

$$\frac{dy}{dx} = \frac{2t}{2}$$

$$= t \quad |$$

$$\frac{dy'}{dt} = \frac{d}{dt}(t)$$

$$= 1 \quad |$$

$$\frac{d^2y}{dx^2} = \frac{1}{2t} \quad |$$

$$\frac{dy}{dx} = \frac{dy / dt}{dx / dt}$$

$$\frac{d^2y}{dx^2} = \frac{dy' / dt}{dx / dt}$$

Exercise

Find $\frac{d^2y}{dx^2}$ for $x(t) = 2t^2 - 1$ $y(t) = 2t^3 + t$

Solution

$$\frac{dx}{dt} = 4t$$

$$\frac{dy}{dt} = 6t^2$$

$$\frac{dy}{dx} = \frac{6t^2}{4t}$$

$$= \frac{3}{2}t \quad |$$

$$\frac{dy'}{dt} = \frac{d}{dt}\left(\frac{3}{2}t\right)$$

$$= \frac{3}{2} \quad |$$

$$\frac{d^2y}{dx^2} = \frac{3}{2} \cdot \frac{1}{4t}$$

$$= \frac{3}{8t} \quad |$$

$$\frac{dy}{dx} = \frac{dy / dt}{dx / dt}$$

$$\frac{d^2y}{dx^2} = \frac{dy' / dt}{dx / dt}$$

Exercise

Find an equation of the line tangent to cycloid $x(t) = t - \sin t$, $y(t) = 2 - \cos t$ at the points corresponding to $t = \frac{\pi}{6}$ and $t = \frac{2\pi}{3}$.

Solution

$$\frac{dx}{dt} = 1 - \cos t$$

$$\frac{dy}{dt} = \sin t$$

$$\frac{dy}{dx} = \frac{\sin t}{1 - \cos t} \qquad \frac{dy}{dx} = \frac{dy / dt}{dx / dt}$$

$$@ \quad t = \frac{\pi}{6} \rightarrow \begin{cases} x = \frac{\pi}{6} - \sin \frac{\pi}{6} = \frac{\pi}{6} - \frac{1}{2} \\ y = 2 - \cos \frac{\pi}{6} = 2 - \frac{\sqrt{3}}{2} \end{cases}$$

$$\left. \frac{dy}{dx} = \frac{\sin t}{1 - \cos t} \right|_{t=\frac{\pi}{6}}$$

$$m = \frac{\sin \frac{\pi}{6}}{1 - \cos \frac{\pi}{6}}$$

$$= \frac{\frac{1}{2}}{1 - \frac{\sqrt{3}}{2}}$$

$$= \frac{1}{2 - \sqrt{3}} \cdot \frac{2 + \sqrt{3}}{2 + \sqrt{3}}$$

$$= 2 + \sqrt{3}$$

$$\begin{aligned} y &= (2 + \sqrt{3}) \left(x - \frac{\pi}{6} + \frac{1}{2} \right) + 2 - \frac{\sqrt{3}}{2} \\ &= (2 + \sqrt{3})x - (2 + \sqrt{3})\frac{\pi}{6} + 1 + \frac{\sqrt{3}}{2} + 2 - \frac{\sqrt{3}}{2} \\ &= (2 + \sqrt{3})x - (2 + \sqrt{3})\frac{\pi}{6} + 3 \end{aligned}$$

$$@ \quad t = \frac{2\pi}{3} \rightarrow \begin{cases} x = \frac{2\pi}{3} - \sin \frac{2\pi}{3} = \frac{2\pi}{3} - \frac{\sqrt{3}}{2} \\ y = 2 - \cos \frac{2\pi}{3} = 2 + \frac{1}{2} = \frac{5}{2} \end{cases}$$

$$\begin{aligned}
 m &= \frac{\sin \frac{2\pi}{3}}{1 - \cos \frac{2\pi}{3}} \\
 &= \frac{\frac{\sqrt{3}}{2}}{1 + \frac{1}{2}} \\
 &= \frac{\sqrt{3}}{3} \Big| \\
 y &= \frac{\sqrt{3}}{3} \left(x - \frac{2\pi}{3} + \frac{\sqrt{3}}{2} \right) + \frac{5}{2} \\
 &= \frac{\sqrt{3}}{3} x - \frac{2\pi\sqrt{3}}{9} + \frac{1}{2} + \frac{5}{2} \\
 &= \frac{\sqrt{3}}{3} x - \frac{2\pi\sqrt{3}}{9} + 3 \Big|
 \end{aligned}$$

Exercise

Consider Lissajous curve, estimate the coordinates of the points on the curve at which there is

$$x = \sin 2t, \quad y = 2 \sin t; \quad 0 \leq t \leq 2\pi$$

- a) A horizontal tangent line
- b) A vertical tangent line.

Solution

$$a) \quad \frac{dy}{dx} = \frac{2 \cos t}{2 \cos 2t} = 0$$

$$\frac{dy}{dx} = \frac{dy / dt}{dx / dt}$$


$$\cos t = 0 \rightarrow t = \frac{\pi}{2}, \frac{3\pi}{2}$$

$$t = \frac{\pi}{2} \rightarrow \begin{cases} x = \sin \pi = 0 \\ y = 2 \sin \frac{\pi}{2} = 2 \end{cases}$$

$$(0, 2) \Big|$$

$$t = \frac{3\pi}{2} \rightarrow \begin{cases} x = \sin 3\pi = 0 \\ y = 2 \sin \frac{3\pi}{2} = -2 \end{cases}$$

$$(0, -2) \Big|$$



- b) Vertical tangent line: $\cos 2t = 0 \quad \cos t \neq 0$

$$\cos 2t = 0$$

$$2t = \frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}, \frac{7\pi}{2}$$

$$t = \frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4} \Big|$$

$$t = \frac{\pi}{4} \quad \begin{cases} x = 1 \\ y = \sqrt{2} \end{cases} \rightarrow \underline{(1, \sqrt{2})}$$

$$t = \frac{3\pi}{4} \quad \begin{cases} x = -1 \\ y = \sqrt{2} \end{cases} \rightarrow \underline{(-1, \sqrt{2})}$$

$$t = \frac{5\pi}{4} \quad \begin{cases} x = -1 \\ y = -\sqrt{2} \end{cases} \rightarrow \underline{(-1, -\sqrt{2})}$$

$$t = \frac{7\pi}{4} \quad \begin{cases} x = 1 \\ y = -\sqrt{2} \end{cases} \rightarrow \underline{(1, -\sqrt{2})}$$

Exercise

Consider Lissajous curve, estimate the coordinates of the points on the curve at which there is

$$x = \sin 4t, \quad y = \sin 3t; \quad 0 \leq t \leq 2\pi$$

a) A horizontal tangent line

b) A vertical tangent line.

Solution

$$a) \quad \frac{dy}{dx} = \frac{3 \cos 3t}{4 \cos 4t} = 0 \quad \frac{dy}{dx} = \frac{dy/dt}{dx/dt}$$

$$\cos 3t = 0 \rightarrow 3t = \frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}, \frac{7\pi}{2}, \frac{9\pi}{2}, \frac{11\pi}{2}$$


$$t = \frac{\pi}{6}, \frac{\pi}{2}, \frac{5\pi}{6}, \frac{7\pi}{6}, \frac{3\pi}{2}, \frac{11\pi}{6} \Big|$$

$$t = \frac{\pi}{6} \Rightarrow \begin{cases} x = -\frac{\sqrt{3}}{2} \\ y = 1 \end{cases} \rightarrow \underline{\left(-\frac{\sqrt{3}}{2}, 1\right)}$$

$$t = \frac{\pi}{2} \Rightarrow \begin{cases} x = 0 \\ y = -1 \end{cases} \rightarrow \underline{(0, -1)}$$

$$t = \frac{5\pi}{6} \Rightarrow \begin{cases} x = \frac{\sqrt{3}}{2} \\ y = -1 \end{cases} \rightarrow \underline{\left(\frac{\sqrt{3}}{2}, -1\right)}$$

$$t = \frac{7\pi}{6} \Rightarrow \begin{cases} x = -\frac{\sqrt{3}}{2} \\ y = 1 \end{cases} \rightarrow \underline{\left(-\frac{\sqrt{3}}{2}, 1\right)}$$



$$t = \frac{3\pi}{2} \Rightarrow \begin{cases} x=0 \\ y=1 \end{cases} \rightarrow \underline{(0, 1)}$$

$$t = \frac{11\pi}{6} \Rightarrow \begin{cases} x = \frac{\sqrt{3}}{2} \\ y=1 \end{cases} \rightarrow \underline{\left(\frac{\sqrt{3}}{2}, 1\right)}$$

b) Vertical tangent line: $\cos 4t = 0$ $\cos 3t \neq 0$

$$\cos 4t = 0 \rightarrow 4t = \frac{(n+1)\pi}{2}$$

$$\underline{t = \frac{(n+1)\pi}{8}}$$

$$t = \frac{(n+1)\pi}{8} \Rightarrow \begin{cases} x = \pm 1 \\ y = \pm \sin \frac{3\pi}{8} \end{cases}$$

$$\underline{\left(\pm 1, \pm \sin \frac{3\pi}{8}\right)}$$

Exercise

Find the area of the region $x = 2\sin^2 \theta$, $y = 2\sin^2 \theta \tan \theta$, $0 \leq \theta < \frac{\pi}{2}$

Solution

$$dx = 4\sin \theta \cos \theta d\theta$$

$$A = \int_0^{\pi/2} 2\sin^2 \theta \tan \theta (4\sin \theta \cos \theta) d\theta$$

$$A = \int_a^b y dx$$


$$= 8 \int_0^{\pi/2} \sin^4 \theta d\theta$$

$$= 2 \int_0^{\pi/2} (1 - \cos 2\theta)^2 d\theta$$

$$= 2 \int_0^{\pi/2} \left(1 - 2\cos 2\theta + \cos^2 2\theta\right) d\theta$$

$$= 2 \int_0^{\pi/2} \left(1 - 2\cos 2\theta + \frac{1}{2} + \frac{1}{2}\cos 4\theta\right) d\theta$$

$$= 2 \left(\frac{3}{2}\theta - \sin 2\theta + \frac{1}{8}\sin 4\theta \right) \Big|_0^{\pi/2}$$



$$= 2 \left(\frac{3\pi}{4} \right)$$

$$= \frac{3\pi}{2} \text{ unit}^2$$

Exercise

Find the area of the region $x = 2 \cot \theta$, $y = 2 \sin^2 \theta$, $0 \leq \theta < \pi$

Solution

$$dx = -2 \csc^2 \theta \, d\theta$$


$$A = -4 \int_0^\pi \sin^2 \theta \csc^2 \theta \, d\theta$$

$$= -4 \int_0^\pi d\theta$$

$$= \left| -4\theta \right|_0^\pi$$

$$= 4\pi \text{ unit}^2$$

$$A = \int_a^b y \, dx$$



Exercise

Find the area under one arch of the cycloid $x = a(t - \sin t)$, $y = a(1 - \cos t)$

Solution

$$A = \int_0^{2\pi} y \, dx$$

$$= \int_0^{2\pi} a(1 - \cos t) \, d[a(t - \sin t)]$$

$$d[a(t - \sin t)] = a(1 - \cos t) \, dt$$

$$= \int_0^{2\pi} a^2 (1 - \cos t)^2 \, dt$$

$$= a^2 \int_0^{2\pi} (1 - 2 \cos t + \cos^2 t) \, dt$$

$$= a^2 \int_0^{2\pi} \left(1 - 2 \cos t + \frac{1 + \cos 2t}{2} \right) dt$$

$$\begin{aligned}
&= a^2 \int_0^{2\pi} \left(\frac{3}{2} - 2 \cos t + \frac{1}{2} \cos 2t \right) dt \\
&= a^2 \left(\frac{3}{2}t - 2 \sin t + \frac{1}{4} \sin 2t \right) \Big|_0^{2\pi} \\
&= a^2 \left(\frac{3}{2}(2\pi) - 2 \sin(2\pi) + \frac{1}{4} \sin 2(2\pi) - 0 \right) \\
&= \underline{3\pi a^2 \text{ unit}^2}
\end{aligned}$$

Exercise

Find the area enclosed by the y-axis and the curve $x = t - t^2$, $y = 1 + e^{-t}$

Solution

$$x = t - t^2 = 0 \Rightarrow \underline{t = 0, 1}$$

$$\begin{aligned}
A &= \int_0^1 x dy \\
&= \int_0^1 (t - t^2) d(1 + e^{-t}) \\
&= \int_0^1 (t - t^2)(-e^{-t}) dt
\end{aligned}$$

$\int e^{-t}$		
+	$t - t^2$	$-e^{-t}$
-	$1 - 2t$	e^{-t}
+	-2	$-e^{-t}$

$$\begin{aligned}
&= - \int_0^1 (t - t^2) e^{-t} dt \\
&= - \left((t - t^2)(-e^{-t}) - (1 - 2t)(e^{-t}) - 2(-e^{-t}) \right) \Big|_0^1 \\
&= - \left(e^{-t}(t^2 - t) - e^{-t}(1 - 2t) + 2e^{-t} \right) \Big|_0^1
\end{aligned}$$

$$\begin{aligned}
&= -\left[e^{-1}(1^2 - 1) - e^{-1}(1 - 2(1)) + 2e^{-1} - \left(e^{-0}(0^2 - 0) - e^{-0}(1 - 2(0)) + 2e^{-0} \right) \right] \\
&= -\left[e^{-1} + 2e^{-1} - (-1 + 2) \right] \\
&= -(3e^{-1} - 1) \\
&= 1 - 3e^{-1} \\
&= \underline{1 - \frac{3}{e} \text{ unit}^2}
\end{aligned}$$

Exercise

Find the area enclosed by the ellipse $x = a \cos t$, $y = b \sin t$, $0 \leq t \leq 2\pi$

Solution


$$\begin{aligned}
A &= \int_0^{2\pi} y dx \\
&= 2 \left| \int_0^{\pi} y dx \right| \\
&= 2 \int_0^{\pi} b \sin t \, d(a \cos t) \\
&= 2 \int_0^{\pi} b \sin t (-a \sin t) \, dt \\
&= -2ab \int_0^{\pi} \sin^2 t \, dt \\
&= -2ab \int_0^{\pi} \left(\frac{1 - \cos 2t}{2} \right) dt \\
&= -ab \left(t - \frac{1}{2} \sin 2t \right) \Big|_0^{\pi} \\
&= -ab \left(\pi - \frac{1}{2} \sin 2\pi - 0 \right) \\
&= |-\pi ab| \\
&= \underline{\pi ab \text{ unit}^2}
\end{aligned}$$

Exercise

Find the area of the closed curve *Ellipse* $\begin{cases} x = b \cos t \\ y = a \sin t \end{cases} \quad 0 \leq t \leq 2\pi$

Solution

$$\begin{aligned} A &= \int_0^{2\pi} y dx \\ &= 2 \left| \int_0^{\pi} y dx \right| \\ &= 2 \int_0^{\pi} a \sin t \, d(b \cos t) \\ &= -2ab \int_0^{\pi} \sin^2 t \, dt \\ &= -2ab \int_0^{\pi} \left(\frac{1 - \cos 2t}{2} \right) dt \\ &= -ab \left(t - \frac{1}{2} \sin 2t \right) \Big|_0^{\pi} \\ &= -ab \left(\pi - \frac{1}{2} \sin 2\pi - 0 \right) \\ &= |-\pi ab| \\ &= \underline{\pi ab \text{ unit}^2} \end{aligned}$$




Exercise

Find the area of the closed curve *Astroid* $\begin{cases} x = a \cos^3 t \\ y = a \sin^3 t \end{cases} \quad 0 \leq t \leq 2\pi$

Solution

$$\begin{aligned} A &= \int_0^{2\pi} y dx \\ &= 4 \int_0^{\pi/2} a \sin^3 t \, \left| d(a \cos^3 t) \right| \\ &= 12a^2 \int_0^{\pi/2} \sin^4 t \cos^2 t \, dt \end{aligned}$$




$$\begin{aligned}
&= 12a^2 \int_0^{\pi/2} \left(\frac{1 - \cos 2t}{2} \right)^2 \left(\frac{1 + \cos 2t}{2} \right) dt \\
&= \frac{3}{2} a^2 \int_0^{\pi/2} (1 - 2 \cos 2t + \cos^2 2t)(1 + \cos 2t) dt \\
&= \frac{3}{2} a^2 \int_0^{\pi/2} (1 - \cos 2t - \cos^2 2t + \cos^3 2t) dt \\
&= \frac{3}{2} a^2 \int_0^{\pi/2} \left(\frac{1}{2} - \cos 2t - \cos 4t \right) dt + \frac{3}{2} a^2 \int_0^{\pi/2} \cos^2 2t \cos 2t dt \\
&= \frac{3}{2} a^2 \left(\frac{1}{2} t - \frac{1}{2} \sin 2t - \frac{1}{4} \sin 4t \right) \Big|_0^{\pi/2} + \frac{3}{4} a^2 \int_0^{\pi/2} (1 - \sin^2 2t) d(\sin 2t) \\
&= \frac{3}{2} a^2 \left(\frac{\pi}{4} \right) + \frac{3}{4} a^2 \left(\sin 2t - \frac{1}{3} \sin^3 2t \right) \Big|_0^{\pi/2} \\
&= \frac{3}{8} \pi a^2 \text{ unit}^2
\end{aligned}$$

Exercise

Find the area of the closed curve *Cardioid* $\begin{cases} x = 2a \cos t - a \cos 2t \\ y = 2a \sin t - a \sin 2t \end{cases} \quad 0 \leq t \leq 2\pi$

Solution

$$\begin{aligned}
A &= \int_0^{2\pi} y dx \\
&= 2 \left| \int_0^{\pi} (2a \sin t - a \sin 2t) d(2a \cos t - a \cos 2t) \right| \\
&= 2 \left| \int_0^{\pi} (2a \sin t - a \sin 2t)(-2a \sin t + 2a \sin 2t) dt \right| \\
&= 4a^2 \left| \int_0^{\pi} (2 \sin t - \sin 2t)(-\sin t + \sin 2t) dt \right| \\
&= 4a^2 \left| \int_0^{\pi} (-2 \sin^2 t + 3 \sin t \sin 2t - \sin^2 2t) dt \right|
\end{aligned}$$



$$\sin \alpha \sin \beta = \frac{1}{2} [\cos(\alpha - \beta) - \cos(\alpha + \beta)]$$


$$\begin{aligned}
&= 4a^2 \left| \int_0^\pi \left(-1 + \cos 2t + \frac{3}{2} \cos t - \frac{3}{2} \cos 3t - \frac{1}{2} + \frac{1}{2} \cos 4t \right) dt \right| \\
&= 4a^2 \left| \int_0^\pi \left(-\frac{3}{2} + \cos 2t + \frac{3}{2} \cos t - \frac{3}{2} \cos 3t + \frac{1}{2} \cos 4t \right) dt \right| \\
&= 4a^2 \left| \left(-\frac{3}{2}t + \frac{1}{2} \sin 2t + \frac{3}{2} \sin t - \frac{1}{2} \sin 3t + \frac{1}{8} \sin 4t \right) \right|_0^\pi \\
&= 4a^2 \left| \left(-\frac{3\pi}{2} \right) \right| \\
&= \underline{6\pi a^2 \text{ unit}^2}
\end{aligned}$$

Exercise

Find the area of the closed curve *Deltoid* $\begin{cases} x = 2a \cos t + a \cos 2t \\ y = 2a \sin t - a \sin 2t \end{cases} \quad 0 \leq t \leq 2\pi$

Solution

$$\begin{aligned}
A &= \int_0^{2\pi} y dx \\
&= 2 \left| \int_0^\pi (2a \sin t - a \sin 2t) d(2a \cos t + a \cos 2t) \right| \\
&= 2 \left| \int_0^\pi (2a \sin t - a \sin 2t)(-2a \sin t - 2a \sin 2t) dt \right| \\
&= 4a^2 \left| \int_0^\pi (2 \sin t - \sin 2t)(\sin t + \sin 2t) dt \right| \\
&= 4a^2 \left| \int_0^\pi (2 \sin^2 t + \sin t \sin 2t - \sin^2 2t) dt \right| \\
&= 4a^2 \left| \int_0^\pi \left(1 - \cos 2t + \frac{1}{2} \cos t - \frac{1}{2} \cos 3t - \frac{1}{2} + \frac{1}{2} \cos 4t \right) dt \right| \\
&= 4a^2 \left| \int_0^\pi \left(\frac{1}{2} - \cos 2t + \frac{1}{2} \cos t - \frac{1}{2} \cos 3t + \frac{1}{2} \cos 4t \right) dt \right|
\end{aligned}$$



$$\sin \alpha \sin \beta = \frac{1}{2} [\cos(\alpha - \beta) - \cos(\alpha + \beta)]$$

$$= 4a^2 \left| \left(\frac{1}{2}t - \frac{1}{2}\sin 2t + \frac{1}{2}\sin t - \frac{1}{6}\sin 3t + \frac{1}{8}\sin 4t \right) \right|_0^\pi$$

$$= \underline{2\pi a^2 \text{ unit}^2}$$

Exercise

Find the area of the closed curve *Hourglass* $\begin{cases} x = a \sin 2t \\ y = b \sin t \end{cases} \quad 0 \leq t \leq 2\pi$

Solution

$$A = \int_0^{2\pi} y dx$$

$$= 2 \left| \int_0^\pi (b \sin t) d(a \sin 2t) \right|$$

$$= 4ab \left| \int_0^\pi (\sin t \cos 2t) dt \right|$$

$$\sin \alpha \cos \beta = \frac{1}{2} [\sin(\alpha + \beta) + \sin(\alpha - \beta)]$$


$$= 2ab \left| \int_0^\pi (\sin 3t + \sin(-t)) dt \right|$$

$$= 2ab \left| \int_0^\pi (\sin 3t - \sin t) dt \right|$$

$$= 2ab \left| \left(-\frac{1}{3} \cos 3t + \cos t \right) \right|_0^\pi$$

$$= 2ab \left| \frac{1}{3} - 1 + \frac{1}{3} - 1 \right|$$

$$= \underline{\frac{8}{3} ab \text{ unit}^2}$$




Exercise

Find the area of the closed curve *Teardrop* $\begin{cases} x = 2a \cos t - a \sin 2t \\ y = b \sin t \end{cases} \quad 0 \leq t \leq 2\pi$

Solution

$$\begin{aligned}
 A &= \int_0^{2\pi} y dx \\
 &= 2 \left| \int_{-\pi/2}^{\pi/2} (b \sin t) d(2a \cos t - a \sin 2t) \right| \\
 &= 2 \left| \int_{-\pi/2}^{\pi/2} (b \sin t)(-2a \sin t - 2a \cos 2t) dt \right| \\
 &= 4ab \left| \int_{-\pi/2}^{\pi/2} (\sin^2 t + \sin t \cos 2t) dt \right| \\
 &= 2ab \left| \int_{-\pi/2}^{\pi/2} (1 - \cos 2t + \sin 3t - \sin t) dt \right| \\
 &= 2ab \left| \left(t - \frac{1}{2} \sin 2t - \frac{1}{3} \cos 3t + \cos t \right) \right|_{-\pi/2}^{\pi/2} \\
 &= 2ab \left| \left(\frac{\pi}{2} + \frac{\pi}{2} \right) \right| \\
 &= \underline{2\pi ab \text{ unit}^2}
 \end{aligned}$$



$$\sin \alpha \cos \beta = \frac{1}{2} [\sin(\alpha + \beta) + \sin(\alpha - \beta)]$$

Exercise

Find the lengths of the curves $x = \cos t, \quad y = t + \sin t, \quad 0 \leq t \leq \pi$

Solution

$$x = \cos t \Rightarrow \frac{dx}{dt} = -\sin t$$

$$y = t + \sin t \Rightarrow \frac{dy}{dt} = 1 + \cos t$$

$$\begin{aligned}
 \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} &= \sqrt{\sin^2 t + (1 + \cos t)^2} \\
 &= \sqrt{\sin^2 t + 1 + 2\cos t + \cos^2 t} \\
 &= \sqrt{2 + 2\cos t}
 \end{aligned}$$

$$\begin{aligned}
L &= \int_0^{\pi} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt \\
&= \int_0^{\pi} \sqrt{2 + 2\cos t} dt \\
&= \sqrt{2} \int_0^{\pi} \sqrt{(1 + \cos t) \frac{1 - \cos t}{1 - \cos t}} dt \\
&= \sqrt{2} \int_0^{\pi} \sqrt{\frac{1 - \cos^2 t}{1 - \cos t}} dt \\
&= \sqrt{2} \int_0^{\pi} \sqrt{\frac{\sin^2 t}{1 - \cos t}} dt \\
&= \sqrt{2} \int_0^{\pi} \frac{\sin t}{\sqrt{1 - \cos t}} dt \quad d(1 - \cos t) = \sin t dt \\
&= \sqrt{2} \int_0^{\pi} \frac{d(1 - \cos t)}{\sqrt{1 - \cos t}} \\
&= \sqrt{2} \left(2\sqrt{1 - \cos t} \right) \Big|_0^{\pi} \\
&= 2\sqrt{2} (\sqrt{1 - \cos \pi} - \sqrt{1 - \cos 0}) \\
&= 2\sqrt{2} (\sqrt{2} - 0) \\
&= 4 \text{ unit}
\end{aligned}$$

$$\begin{aligned}
L &= \int_0^{\pi} \sqrt{4 \sin^2 \frac{t}{2}} dt \quad 2 \sin^2 \frac{t}{2} = 1 - \cos t \\
&= 2 \int_0^{\pi} \sin \frac{t}{2} dt \\
&= -4 \cos \frac{t}{2} \Big|_0^{\pi} \\
&= -4(0 - 1) \\
&= 4 \text{ unit}
\end{aligned}$$

Exercise

Find the lengths of the curves $x = t^3, \quad y = \frac{3}{2}t^2, \quad 0 \leq t \leq \sqrt{3}$

Solution

$$x = t^3 \Rightarrow \frac{dx}{dt} = 3t^2$$

$$y = \frac{3}{2}t^2 \Rightarrow \frac{dy}{dt} = 3t$$

$$\begin{aligned}\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} &= \sqrt{9t^4 + 9t^2} \\ &= 3t\sqrt{t^2 + 1}\end{aligned}$$

$$L = \int_0^{\sqrt{3}} 3t \sqrt{t^2 + 1} \, dt$$

$$L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} \, dt$$

$$u = t^2 + 1 \Rightarrow du = 2t \, dt \rightarrow \begin{cases} t = \sqrt{3} \Rightarrow u = 4 \\ t = 0 \Rightarrow u = 1 \end{cases}$$

$$= \frac{3}{2} \int_1^4 \sqrt{u} \, du$$

$$= \frac{3}{2} \frac{2}{3} \left(u^{3/2} \right) \Big|_1^4$$

$$= 4^{3/2} - 1$$

$$= \underline{7 \text{ unit}}$$

Exercise

Find the lengths of the curves $x = 8 \cos t + 8t \sin t, \quad y = 8 \sin t - 8t \cos t, \quad 0 \leq t \leq \frac{\pi}{2}$

Solution

$$x = 8 \cos t + 8t \sin t$$

$$\frac{dx}{dt} = -8 \sin t + 8 \sin t + 8t \cos t = \underline{8t \cos t}$$

$$y = 8 \sin t - 8t \cos t$$

$$\frac{dy}{dt} = 8 \cos t - 8 \cos t + 8t \sin t = \underline{8t \sin t}$$

$$\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} = \sqrt{(8t \cos t)^2 + (8t \sin t)^2}$$

$$\begin{aligned}
&= \sqrt{(8t)^2 \cos^2 t + (8t)^2 \sin^2 t} \\
&= 8t \sqrt{\cos^2 t + \sin^2 t} & \cos^2 t + \sin^2 t = 1 \\
&= \underline{8t}
\end{aligned}$$

$$\begin{aligned}
L &= \int_0^{\pi/2} 8t \, dt \\
&= 4t^2 \Big|_0^{\pi/2} \\
&= 4 \left(\frac{\pi^2}{4} - 0 \right) \\
&= \underline{\pi^2 \text{ unit}}
\end{aligned}$$

$$L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

Exercise

Find the lengths of the curves $x = \ln(\sec t + \tan t) - \sin t$, $y = \cos t$, $0 \leq t \leq \frac{\pi}{3}$

Solution

$$x = \ln(\sec t + \tan t) - \sin t$$

$$\begin{aligned}
\frac{dx}{dt} &= \frac{\sec t \tan t + \sec^2 t}{\sec t + \tan t} - \cos t \\
&= \frac{\sec t (\tan t + \sec t)}{\sec t + \tan t} - \cos t \\
&= \underline{\sec t - \cos t}
\end{aligned}$$

$$y = \cos t \Rightarrow \underline{\frac{dy}{dt} = -\sin t}$$

$$\begin{aligned}
\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} &= \sqrt{(\sec t - \cos t)^2 + (-\sin t)^2} \\
&= \sqrt{\sec^2 t - 2\sec t \cos t + \cos^2 t + \sin^2 t} \\
&= \sqrt{\sec^2 t - 2\frac{1}{\cos t} \cos t + 1} \\
&= \sqrt{\sec^2 t - 2 + 1} \\
&= \sqrt{\sec^2 t - 1} \\
&= \sqrt{\tan^2 t} \\
&= \tan t
\end{aligned}$$

$$\begin{aligned}
L &= \int_0^{\pi/3} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt \\
&= \int_0^{\pi/3} \tan t \, dt \\
&= \int_0^{\pi/3} \frac{\sin t}{\cos t} dt \\
&= \int_0^{\pi/3} -\frac{d(\cos t)}{\cos t} \\
&= -\ln|\cos t| \Big|_0^{\pi/3} \\
&= -\ln \cos \frac{\pi}{3} + \ln \cos 0 \\
&= -\ln \frac{1}{2} + \ln 1 \\
&= \ln 2 \text{ unit}
\end{aligned}$$

Exercise

Find the arc length of the Hypocycloid perimeter curve: $x = a \cos \theta$, $y = a \sin \theta$

Solution

$$x = a \cos \theta \rightarrow \frac{dx}{d\theta} = -a \sin \theta$$

$$y = a \sin \theta \rightarrow \frac{dy}{d\theta} = a \cos \theta$$

$$L = 4 \int_0^{\pi/2} \sqrt{a^2 \sin^2 \theta + a^2 \cos^2 \theta} \, d\theta$$

$$= 4a \int_0^{\pi/2} d\theta$$

$$= 4a \theta \Big|_0^{\pi/2}$$

$$= 2\pi a \text{ unit}$$

$$L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

Exercise

Find the arc length of the circle circumference: $x = a \cos^3 \theta$, $y = a \sin^3 \theta$

Solution

$$\frac{dx}{d\theta} = -3a \sin \theta \cos^2 \theta$$

$$\frac{dy}{d\theta} = 3a \cos \theta \sin^2 \theta$$

$$\begin{aligned}\sqrt{\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2} &= \sqrt{9a^2 \sin^2 \theta \cos^4 \theta + 9a^2 \cos^2 \theta \sin^4 \theta} \\ &= 3a \sin \theta \cos \theta \sqrt{\cos^2 \theta + \sin^2 \theta} \\ &= 3a \sin \theta \cos \theta\end{aligned}$$

$$L = 4 \int_0^{\pi/2} 3a \sin \theta \cos \theta \, d\theta$$

$$= 6a \int_0^{\pi/2} \sin 2\theta \, d\theta$$

$$= -3a \cos 2\theta \Big|_0^{\pi/2}$$

$$= -3a(-1-1)$$

$$= \underline{6a \text{ unit}}$$

$$L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} \, dt$$

Exercise

Find the arc length of the Cycloid arch: $x = a(\theta - \sin \theta)$, $y = a(1 - \cos \theta)$

Solution

$$x = a(\theta - \sin \theta) \rightarrow \frac{dx}{d\theta} = a(1 - \cos \theta)$$

$$y = a(1 - \cos \theta) \rightarrow \frac{dy}{d\theta} = a \sin \theta$$

$$\begin{aligned}\sqrt{\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2} &= \sqrt{a^2(1 - 2\cos \theta + \cos^2 \theta) + a^2 \sin^2 \theta} \\ &= a\sqrt{1 - 2\cos \theta + \cos^2 \theta + \sin^2 \theta} \\ &= a\sqrt{2 - 2\cos \theta}\end{aligned}$$

$$L = 2a\sqrt{2} \int_0^{\pi} \sqrt{1 - \cos \theta} \, d\theta$$

$$L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} \, dt$$

$$\begin{aligned}
&= 2a\sqrt{2} \int_0^{\pi} \sqrt{1-\cos\theta} \frac{\sqrt{1+\cos\theta}}{\sqrt{1+\cos\theta}} d\theta \\
&= 2a\sqrt{2} \int_0^{\pi} \frac{\sin\theta}{\sqrt{1+\cos\theta}} d\theta \\
&= -2a\sqrt{2} \int_0^{\pi} (1+\cos\theta)^{-1/2} d(1+\cos\theta) \\
&= -4a\sqrt{2} \sqrt{1+\cos\theta} \Big|_0^{\pi} \\
&= -4a\sqrt{2} (0 - \sqrt{2}) \\
&= \underline{8a \text{ unit}}
\end{aligned}$$

Exercise

Find the arc length of the involute of a circle: $x = \cos\theta + \theta\sin\theta$, $y = \sin\theta - \theta\cos\theta$

Solution

$$x = \cos\theta + \theta\sin\theta \rightarrow \frac{dx}{d\theta} = -\sin\theta + \sin\theta + \theta\cos\theta = \theta\cos\theta$$

$$y = \sin\theta - \theta\cos\theta \rightarrow \frac{dy}{d\theta} = \theta\sin\theta$$

$$\begin{aligned}
\sqrt{\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2} &= \sqrt{\theta^2 \cos^2\theta + \theta^2 \sin^2\theta} \\
&= \theta\sqrt{\cos^2\theta + \sin^2\theta} \\
&= \theta
\end{aligned}$$

$$\begin{aligned}
L &= \int_0^{2\pi} \theta d\theta \\
&= \frac{1}{2} \theta^2 \Big|_0^{2\pi} \\
&= \underline{2\pi^2 \text{ unit}}
\end{aligned}$$

$$L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

Exercise

Find the arc length of $x = t^2$, $y = t^3$, $0 \leq t \leq 2$

Solution

$$\frac{dx}{dt} = 2t$$

$$\frac{dy}{dt} = 3t^2$$

$$\begin{aligned}\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} &= \sqrt{4t^2 + 9t^4} \\ &= t \sqrt{4 + 9t^2}\end{aligned}$$

$$\begin{aligned}L &= \int_0^2 t \sqrt{4 + 9t^2} \, dt & L &= \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} \, dt \\ &= \frac{1}{18} \int_0^2 (4 + 9t^2)^{1/2} d(4 + 9t^2) \\ &= \frac{1}{27} (4 + 9t^2)^{3/2} \Big|_0^2 \\ &= \frac{1}{27} ((40)^{3/2} - 4^{3/2}) \\ &= \frac{1}{27} (8(10)^{3/2} - 8) \\ &= \frac{8}{27} ((10)^{3/2} - 1) \text{ unit}\end{aligned}$$

Exercise

Find the arc length of $x = 5 \sin t$, $y = 5 \cos t$, $-\frac{\pi}{3} \leq t \leq \frac{\pi}{2}$

Solution

$$\frac{dx}{dt} = 5 \cos t$$

$$\frac{dy}{dt} = -5 \sin t$$

$$\begin{aligned}\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} &= \sqrt{25 \cos^2 t + 25 \sin^2 t} \\ &= \sqrt{25 (\cos^2 t + \sin^2 t)}\end{aligned}$$

$$= 5 \Big|$$

$$\begin{aligned} L &= \int_{-\frac{\pi}{3}}^{\frac{\pi}{2}} 5 \, dt \\ &= 5t \Big|_{-\frac{\pi}{3}}^{\frac{\pi}{2}} \\ &= 5 \left(\frac{\pi}{2} + \frac{\pi}{3} \right) \\ &= \frac{25\pi}{6} \text{ unit} \Big| \end{aligned}$$

$$L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} \, dt$$

Exercise

Find the area of the surface generated by revolving the curve about each given axis.

$$x = \frac{1}{3}t^3, \quad y = t + 1, \quad 1 \leq t \leq 2, \quad y\text{-axis}$$

Solution

$$\frac{dx}{dt} = t^2, \quad \frac{dy}{dt} = 1$$

$$\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} = \sqrt{t^4 + 1}$$

$$\begin{aligned} S &= 2\pi \int_1^2 \frac{1}{3}t^3 \sqrt{t^4 + 1} \, dt \\ &= \frac{1}{6}\pi \int_1^2 \sqrt{t^4 + 1} \, d(t^4 + 1) \\ &= \frac{\pi}{9} (t^4 + 1)^{3/2} \Big|_1^2 \\ &= \frac{\pi}{9} (17^{3/2} - 2^{3/2}) \\ &= \frac{\pi}{9} (17\sqrt{17} - 2\sqrt{2}) \text{ unit}^2 \Big| \end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} \, dt$$

Exercise

Find the areas of the surfaces generated by revolving the curves

$$x = \frac{2}{3}t^{3/2}, \quad y = 2\sqrt{t}, \quad 0 \leq t \leq \sqrt{3}; \quad x\text{-axis}$$

Solution

$$x = \frac{2}{3}t^{3/2} \Rightarrow \frac{dx}{dt} = t^{1/2}$$

$$y = 2\sqrt{t} \Rightarrow \frac{dy}{dt} = t^{-1/2}$$

$$\begin{aligned}\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} &= \sqrt{(t^{1/2})^2 + (t^{-1/2})^2} \\ &= \sqrt{t + t^{-1}} \\ &= \sqrt{\frac{t^2 + 1}{t}}\end{aligned}$$

$$\begin{aligned}A &= 2\pi \int_0^{\sqrt{3}} x ds \\ &= 2\pi \int_0^{\sqrt{3}} \frac{2}{3}t^{3/2} \sqrt{\frac{t^2 + 1}{t}} dt \\ &= \frac{4\pi}{3} \int_0^{\sqrt{3}} t \sqrt{t^2 + 1} dt\end{aligned}$$

$$u = t^2 + 1 \Rightarrow du = 2t dt \rightarrow \begin{cases} t = \sqrt{3} \Rightarrow u = 4 \\ t = 0 \Rightarrow u = 1 \end{cases}$$

$$\begin{aligned}&= \frac{2\pi}{3} \int_1^4 \sqrt{u} du \\ &= \frac{2\pi}{3} \left(\frac{2}{3} u^{3/2} \right) \Big|_1^4 \\ &= \frac{4\pi}{9} (4^{3/2} - 1) \\ &= \underline{\underline{\frac{28\pi}{9} \text{ unit}^2}}\end{aligned}$$

Exercise

Find the areas of the surfaces generated by revolving the curves

$$x = t + \sqrt{2}, \quad y = \frac{t^2}{2} + \sqrt{2}t, \quad -\sqrt{2} \leq t \leq \sqrt{2}; \quad y - \text{axis}$$

Solution

$$x = t + \sqrt{2} \Rightarrow \frac{dx}{dt} = 1$$

$$y = \frac{t^2}{2} + \sqrt{2}t \Rightarrow \frac{dy}{dt} = t + \sqrt{2}$$

$$\begin{aligned} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} &= \sqrt{1^2 + (t + \sqrt{2})^2} \\ &= \sqrt{1 + t^2 + 2\sqrt{2}t + 2} \\ &= \sqrt{t^2 + 2\sqrt{2}t + 3} \end{aligned}$$

$$\begin{aligned} A &= 2\pi \int_{-\sqrt{2}}^{\sqrt{2}} x ds \\ &= 2\pi \int_{-\sqrt{2}}^{\sqrt{2}} (t + \sqrt{2}) \sqrt{t^2 + 2\sqrt{2}t + 3} dt \end{aligned}$$

$$\begin{aligned} u &= t^2 + 2\sqrt{2}t + 3 \\ du &= (2t + 2\sqrt{2}) dt = 2(t + \sqrt{2}) dt \\ &\rightarrow \begin{cases} t = \sqrt{2} \Rightarrow u = 9 \\ t = -\sqrt{2} \Rightarrow u = 1 \end{cases} \end{aligned}$$

$$= \pi \int_1^9 \sqrt{u} du$$

$$= \pi \left(\frac{2}{3} u^{3/2} \right) \Big|_1^9$$

$$= \frac{2\pi}{3} (9^{3/2} - 1)$$

$$= \frac{52\pi}{9} \text{ unit}^2$$

Exercise

Find the areas of the surfaces generated by revolving the curves $x = 2t$, $y = 3t$; $0 \leq t \leq 3$ x -axis

Solution

$$x = 2t \rightarrow \frac{dx}{dt} = 2$$

$$y = 3t \rightarrow \frac{dy}{dt} = 3$$

$$\begin{aligned}\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} &= \sqrt{4 + 9} \\ &= \sqrt{13}\end{aligned}$$

$$\begin{aligned}S &= 2\pi \int_0^3 (3t) \sqrt{13} \, dt \\ &= 3\pi \sqrt{13} \left(t^2 \right) \Big|_0^3 \\ &= \underline{27\pi \sqrt{13} \text{ unit}^2}\end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} \, dt$$

Exercise

Find the areas of the surfaces generated by revolving the curves $x = 2t$, $y = 3t$; $0 \leq t \leq 3$ y -axis

Solution

$$x = 2t \rightarrow \frac{dx}{dt} = 2$$

$$y = 3t \rightarrow \frac{dy}{dt} = 3$$

$$\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} = \sqrt{13}$$

$$\begin{aligned}S &= 2\pi \int_0^3 (2t) \sqrt{13} \, dt \\ &= 2\pi \sqrt{13} \left(t^2 \right) \Big|_0^3 \\ &= \underline{18\pi \sqrt{13} \text{ unit}^2}\end{aligned}$$

$$S = 2\pi \int_a^b x \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} \, dt$$

Exercise

Find the areas of the surfaces generated by revolving the curves $x = t$, $y = 4 - 2t$; $0 \leq t \leq 2$ x -axis

Solution

$$\frac{dx}{dt} = 1$$

$$\frac{dy}{dt} = -2$$

$$\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} = \sqrt{5}$$

$$\begin{aligned} S &= 2\pi \int_0^2 (4 - 2t) \sqrt{5} \, dt \\ &= 2\pi \sqrt{5} \left(4t - t^2 \right) \Big|_0^2 \\ &= \underline{8\pi\sqrt{5} \text{ unit}^2} \end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} \, dt$$

Exercise

Find the areas of the surfaces generated by revolving the curves $x = t$, $y = 4 - 2t$; $0 \leq t \leq 2$ y -axis

Solution

$$\frac{dx}{dt} = 1$$

$$\frac{dy}{dt} = -2$$

$$\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} = \sqrt{5}$$

$$\begin{aligned} S &= 2\pi \int_0^2 (t) \sqrt{5} \, dt \\ &= \pi \sqrt{5} \left(t^2 \right) \Big|_0^2 \\ &= \underline{4\pi\sqrt{5} \text{ unit}^2} \end{aligned}$$

$$S = 2\pi \int_a^b x \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} \, dt$$

Exercise

Find the area of the surface generated by revolving the curve about each given axis.

$$x = 5 \cos \theta, \quad y = 5 \sin \theta, \quad 0 \leq \theta \leq \frac{\pi}{2}, \quad y\text{-axis}$$

Solution

$$\frac{dx}{d\theta} = -5 \sin \theta$$


$$\frac{dy}{d\theta} = 5 \cos \theta$$

$$\sqrt{\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2} = \sqrt{25 \sin^2 \theta + 25 \cos^2 \theta} \\ = 5$$

$$S = 2\pi \int_0^{\pi/2} 5 \cos \theta (5) d\theta$$

$$= 50\pi \sin \theta \Big|_0^{\pi/2}$$

$$= 50\pi \text{ unit}^2$$



$$S = 2\pi \int_a^b x \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

Exercise

Find the area of the surface generated by revolving the curve about each given axis.

$$x = a \cos^3 \theta, \quad y = a \sin^3 \theta, \quad 0 \leq \theta \leq \pi, \quad x\text{-axis}$$

Solution

$$x = a \cos^3 \theta \rightarrow \frac{dx}{d\theta} = -3a \sin \theta \cos^2 \theta$$

$$y = a \sin^3 \theta \rightarrow \frac{dy}{d\theta} = 3a \cos \theta \sin^2 \theta$$

$$\sqrt{\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2} = \sqrt{9a^2 \sin^2 \theta \cos^4 \theta + 9a^2 \cos^2 \theta \sin^4 \theta} \\ = 3a \sin \theta \cos \theta$$

$$S = 2\pi \int_0^{\pi/2} a \sin^3 \theta (3a \sin \theta \cos \theta) d\theta$$

$$= 12a^2 \pi \int_0^{\pi/2} \sin^4 \theta d(\sin \theta)$$

$$= \frac{12a^2 \pi}{5} \sin^5 \theta \Big|_0^{\pi/2}$$

$$S = 2\pi \int_a^b y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$= \frac{12}{5} \pi a^2 \text{ unit}^2$$

Exercise

Find the area of the surface generated by revolving the curve about each given axis.

$$x = a \cos \theta, \quad y = b \sin \theta, \quad 0 \leq \theta \leq 2\pi$$

$$a) \quad x\text{-axis} \qquad b) \quad y\text{-axis}$$

Solution

$$x = a \cos \theta \rightarrow \frac{dx}{d\theta} = -a \sin \theta$$

$$y = b \sin \theta \rightarrow \frac{dy}{d\theta} = b \cos \theta$$

$$\sqrt{\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2} = \sqrt{a^2 \sin^2 \theta + b^2 \cos^2 \theta}$$

$$\begin{aligned} a) \quad S &= 4\pi \int_0^{\pi/2} b \sin \theta \sqrt{a^2 \sin^2 \theta + b^2 \cos^2 \theta} \, d\theta \\ &= 4\pi \int_0^{\pi/2} b \sin \theta \sqrt{a^2 (1 - \cos^2 \theta) + b^2 \cos^2 \theta} \, d\theta \\ &= 4\pi \int_0^{\pi/2} b \sin \theta \sqrt{a^2 + (b^2 - a^2) \cos^2 \theta} \, d\theta \\ &= 4\pi \int_0^{\pi/2} ab \sin \theta \sqrt{1 - \left(\frac{a^2 - b^2}{a^2}\right) \cos^2 \theta} \, d\theta \end{aligned}$$


$$\text{Let: } K^2 = \frac{a^2 - b^2}{a^2}$$

$$= 4\pi \int_0^{\pi/2} ab \sin \theta \sqrt{1 - K^2 \cos^2 \theta} \, d\theta$$

$$\begin{aligned} K \cos \theta &= \sin \alpha & \sqrt{1 - K^2 \cos^2 \theta} &= \cos \alpha \\ -K \sin \theta d\theta &= \cos \alpha d\alpha \end{aligned}$$

$$= -\frac{4ab\pi}{K} \int_0^{\pi/2} \cos^2 \alpha \, d\alpha$$

$$S = \pi \int_a^b y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} \, dt$$



$$\begin{aligned}
&= -\frac{2ab\pi}{K} \int_0^{\pi/2} (1 + \cos 2\alpha) d(\alpha) \\
&= -\frac{2ab\pi}{K} \left(\alpha + \frac{1}{2} \sin 2\alpha \right) \Big|_0^{\pi/2} \\
&= -\frac{2ab\pi}{K} \left(\arcsin(K \cos \theta) + K \cos \theta \sqrt{1 - K^2 \cos^2 \theta} \right) \Big|_0^{\pi/2} \\
&= -\frac{2a^2 b \pi}{\sqrt{a^2 - b^2}} \left(-\arcsin\left(\frac{\sqrt{a^2 - b^2}}{a}\right) - \frac{\sqrt{a^2 - b^2}}{a} \right) \\
&\quad e = \frac{\sqrt{a^2 - b^2}}{a} = \frac{c}{a}: \text{eccentricity} \\
&= \frac{2ab\pi}{e} (e + \arcsin(e)) \text{ unit}^2 \Big| \quad c = \sqrt{a^2 - b^2}
\end{aligned}$$

$$b) \quad S = 4\pi \int_0^{\pi/2} a \cos \theta \sqrt{a^2 \sin^2 \theta + b^2 \cos^2 \theta} d\theta$$

$$S = \pi \int_a^b x \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$= 4a\pi \int_0^{\pi/2} \cos \theta \sqrt{(a^2 - b^2) \sin^2 \theta + b^2} d\theta$$

$$= 4a\pi \int_0^{\pi/2} \cos \theta \sqrt{c^2 \sin^2 \theta + b^2} d\theta$$

$$c \sin \theta = b \tan \alpha \quad \sqrt{c^2 \sin^2 \theta + b^2} = b \sec \alpha$$

$$c \cos \theta d\theta = b \sec^2 \alpha d\alpha$$

$$= 4a\pi \int_0^{\pi/2} \frac{b^2}{c} \sec^3 \alpha d\alpha$$

$$u = \sec x \quad dv = \sec^2 x dx$$

$$du = \sec x \tan x dx \quad v = \tan x$$

$$\int \sec^3 x dx = \sec x \tan x - \int \tan x (\sec x \tan x dx)$$

$$= \sec x \tan x - \int \tan^2 x \sec x dx$$

$$= \sec x \tan x - \int (\sec^2 x - 1) \sec x dx$$

$$\begin{aligned}
&= \sec x \tan x - \int (\sec^3 x - \sec x) dx \\
&= \sec x \tan x - \int \sec^3 x dx + \int \sec x dx \\
2 \int \sec^3 x dx &= \sec x \tan x + \int \sec x dx \\
&= \sec x \tan x + \ln |\sec x + \tan x| \\
\int \sec^3 x dx &= \frac{1}{2} \sec x \tan x + \frac{1}{2} \ln |\sec x + \tan x| \\
&= \frac{2ab^2\pi}{c^2} (\sec \alpha \tan \alpha + \ln |\sec \alpha + \tan \alpha|) \Big|_0^{\pi/2} \\
&= \frac{2ab^2\pi}{c} \left(\frac{c \sin \theta \sqrt{c^2 \sin^2 \theta + b^2}}{b^2} + \ln \left| \frac{c \sin \theta + \sqrt{c^2 \sin^2 \theta + b^2}}{b} \right| \right) \Big|_0^{\pi/2} \\
&= \frac{2ab^2\pi}{c} \left(\frac{c\sqrt{c^2 + b^2}}{b^2} + \ln \left| \frac{c + \sqrt{c^2 + b^2}}{b} \right| \right) \\
&= 2a\pi\sqrt{a^2 - b^2 + b^2} + \frac{2ab^2\pi}{c} \ln \left| \frac{\sqrt{a^2 - b^2} + \sqrt{a^2 - b^2 + b^2}}{b} \right| \\
&= 2a^2\pi + \frac{2ab^2\pi}{\sqrt{a^2 - b^2}} \ln \left| \frac{\sqrt{a^2 - b^2} + a}{b} \right| \\
&= 2a^2\pi + \frac{2b^2\pi}{e} \ln \left| \frac{a(e+1)}{b} \right| \text{ unit}^2
\end{aligned}$$

Exercise

Find the area of the surface generated by revolving the curve about each given axis.

$$x = 2t, \quad y = 3t, \quad 0 \leq t \leq 3$$


a) x -axis

b) y -axis

Solution

$$x = 2t \rightarrow \frac{dx}{dt} = 2$$

$$y = 3t \rightarrow \frac{dy}{dt} = 3$$




$$\begin{aligned}
 a) \quad S &= 2\pi \int_0^3 3t\sqrt{4+9} \, dt \\
 &= 3\pi\sqrt{13} \, t^2 \Big|_0^3 \\
 &= \underline{27\pi\sqrt{13} \, \text{unit}^2}
 \end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} \, dt$$

$$\begin{aligned}
 b) \quad S &= 2\pi \int_0^3 2t\sqrt{13} \, dt \\
 &= 2\pi\sqrt{13} \, t^2 \Big|_0^3 \\
 &= \underline{18\pi\sqrt{13} \, \text{unit}^2}
 \end{aligned}$$

$$S = 2\pi \int_a^b x \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} \, dt$$



Exercise

Find the area of the surface generated by revolving the curve about each given axis.


$$x = t, \quad y = 4 - 2t, \quad 0 \leq t \leq 2$$

a) x -axis

b) y -axis

Solution

$$\begin{aligned}
 x = t &\rightarrow \frac{dx}{dt} = 1 \\
 y = 4 - 2t &\rightarrow \frac{dy}{dt} = -2
 \end{aligned}$$




$$\begin{aligned}
 a) \quad S &= 2\pi \int_0^2 (4 - 2t)\sqrt{1+4} \, dt \\
 &= 2\pi\sqrt{5} \left(4t - t^2\right) \Big|_0^2 \\
 &= \underline{8\pi\sqrt{5} \, \text{unit}^2}
 \end{aligned}$$

$$S = 2\pi \int_a^b y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} \, dt$$

$$\begin{aligned}
 b) \quad S &= 2\pi \int_0^2 t \sqrt{5} \, dt \\
 &= \pi\sqrt{5} \, t^2 \Big|_0^2 \\
 &= \underline{4\pi\sqrt{5} \, \text{unit}^2}
 \end{aligned}$$

$$S = 2\pi \int_a^b x \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} \, dt$$




Exercise

Use the parametric equations $x = t^2\sqrt{3}$ and $y = 3t - \frac{1}{3}t^3$ to

- Graph the curve on the interval $-3 \leq t \leq 3$.
- Find $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$
- Find the equation of the tangent line at the point $\left(\sqrt{3}, \frac{8}{3}\right)$
- Find the length of the curve
- Find the surface area generated by revolving the curve about the x -axis

Solution

a)



$$b) \quad \frac{dy}{dx} = \frac{3-t^2}{2t\sqrt{3}} \quad \Bigg|$$

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt}$$

$$\begin{aligned} \frac{dy'}{dt} &= \frac{1}{2\sqrt{3}} \frac{-2t^2 - 3 + t^2}{t^2} \\ &= -\frac{t^2 + 3}{2\sqrt{3}t^2} \end{aligned}$$

$$\begin{aligned} \frac{d^2y}{dx^2} &= -\frac{t^2 + 3}{2\sqrt{3}t^2} \cdot \frac{1}{2t\sqrt{3}} & \frac{d^2y}{dx^2} &= \frac{dy'/dt}{dx/dt} \\ &= -\frac{t^2 + 3}{12t^3} & & \end{aligned} \quad \Bigg|$$

$$c) \quad \left(\sqrt{3}, \frac{8}{3}\right) \rightarrow x = t^2\sqrt{3} = \sqrt{3}$$

$$\Rightarrow t = 1 \Big|$$

$$m = \frac{dy}{dx} \Big|_{t=1}$$

$$= \frac{3-t^2}{2t\sqrt{3}} \Big|_{t=1}$$

$$= \frac{1}{\sqrt{3}} \Big|$$

$$y = \frac{\sqrt{3}}{3}(x - \sqrt{3}) + \frac{8}{3}$$

$$= \frac{\sqrt{3}}{3}x + \frac{5}{3} \Big|$$

$$d) \quad \frac{dx}{dt} = 2t\sqrt{3} \quad \frac{dy}{dt} = 3 - t^2$$

$$L = \int_{-3}^3 \sqrt{12t^2 + 9 - 6t^2 + t^4} \, dt$$

$$= \int_{-3}^3 \sqrt{(t^2 + 3)^2} \, dt$$

$$= \int_{-3}^3 (t^2 + 3) \, dt$$

$$= \frac{1}{3}t^3 + 3t \Big|_{-3}^3$$

$$= 9 + 9 + 9 + 9$$

$$= 36 \text{ unit} \Big|$$

$$L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} \, dt$$

$$e) \quad S = 2\pi \int_0^3 \left(3t - \frac{1}{3}t^3\right)(t^2 + 3) \, dt$$

$$= 2\pi \int_0^3 \left(2t^3 - \frac{1}{3}t^5 + 9t\right) \, dt$$

$$= 2\pi \left(\frac{1}{2}t^4 - \frac{1}{18}t^6 + \frac{9}{2}t^2 \right) \Big|_0^3$$

$$= 2\pi \left(\frac{81}{2} - \frac{81}{2} + \frac{81}{2} \right)$$

$$= 81\pi \text{ unit}^2 \Big|$$

$$S = 2\pi \int_a^b y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} \, dt$$

Exercise

Use the parametric equations $x = a(\theta - \sin \theta)$ and $y = a(1 - \cos \theta)$ $a > 0$

- Find $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$
- Find the equation of the tangent line at the point where $\theta = \frac{\pi}{6}$
- Find all points (if any) of horizontal tangency.
- Determine where the curve is concave upward or concave downward.
- Find the length of one arc of the curve


Solution

$$a) \quad \frac{dx}{d\theta} = a(1 - \cos \theta) \quad \frac{dy}{d\theta} = a \sin \theta$$

$$\frac{dy}{dx} = \frac{\sin \theta}{1 - \cos \theta} \quad \left| \quad \frac{dy}{dx} = \frac{dy / d\theta}{dx / d\theta} \right.$$

$$\begin{aligned} \frac{dy'}{d\theta} &= \frac{d}{d\theta} \left(\frac{\sin \theta}{1 - \cos \theta} \right) \\ &= \frac{\cos \theta - \cos^2 \theta - \sin^2 \theta}{(1 - \cos \theta)^2} \\ &= \frac{\cos \theta - 1}{(1 - \cos \theta)^2} \\ &= \frac{-1}{1 - \cos \theta} \end{aligned}$$

$$\begin{aligned} \frac{d^2y}{dx^2} &= \left(\frac{-1}{1 - \cos \theta} \right) \frac{1}{a(1 - \cos \theta)} \quad \frac{d^2y}{dx^2} = \frac{dy' / d\theta}{dx / d\theta} \\ &= \frac{-1}{a(1 - \cos \theta)^2} \end{aligned}$$



$$b) \quad \text{At } \theta = \frac{\pi}{6} \quad x = a\left(\frac{\pi}{6} - \frac{1}{2}\right) \quad y = a\left(1 - \frac{\sqrt{3}}{2}\right)$$

$$\begin{aligned} m &= \frac{dy}{dx} = \frac{\sin \theta}{1 - \cos \theta} \bigg|_{\theta = \frac{\pi}{6}} \\ &= \frac{\frac{1}{2}}{1 - \frac{\sqrt{3}}{2}} \\ &= \frac{1}{2 - \sqrt{3}} \end{aligned}$$

Tangent Line:

$$y = \frac{1}{2-\sqrt{3}} \left(x - \frac{\pi a}{6} + \frac{a}{2} \right) + a - \frac{a\sqrt{3}}{2}$$

$$y = m(x - x_0) + y_0$$

$$= \left(2 + \sqrt{3} \right) \left(x - \frac{\pi a}{6} + \frac{a}{2} \right) + a - \frac{a\sqrt{3}}{2} \quad \Big|$$

c) $\frac{dy}{dx} = \frac{\sin \theta}{1 - \cos \theta} = 0$

$$\sin \theta = 0 \rightarrow \theta = (2n+1)\pi \quad \Big|$$

$$1 - \cos \theta \neq 0 \rightarrow \theta = 2\pi n$$

$$x = a(2n+1)\pi, \quad y = 2a$$

$$\text{Points of horizontal tangency: } (x, y) = (a(2n+1)\pi, 2a) \quad \Big|$$

d) Concave downward on all open θ -intervals

$$\dots, (-2\pi, 0), (0, 2\pi), (2\pi, 4\pi), \dots$$

e) $\frac{dx}{d\theta} = a(1 - \cos \theta) \quad \frac{dy}{d\theta} = a \sin \theta$

$$\sqrt{\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2} = \sqrt{a^2(1 - 2\cos \theta + \cos^2 \theta) + a^2 \sin^2 \theta}$$

$$= a\sqrt{1 - 2\cos \theta + \cos^2 \theta + \sin^2 \theta}$$

$$= a\sqrt{2 - 2\cos \theta}$$

$$L = 2a\sqrt{2} \int_0^\pi \sqrt{1 - \cos \theta} \, d\theta$$

$$L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} \, dt$$

$$= 2a\sqrt{2} \int_0^\pi \sqrt{1 - \cos \theta} \frac{\sqrt{1 + \cos \theta}}{\sqrt{1 + \cos \theta}} \, d\theta$$

$$= 2a\sqrt{2} \int_0^\pi \frac{\sin \theta}{\sqrt{1 + \cos \theta}} \, d\theta$$

$$= -2a\sqrt{2} \int_0^\pi (1 + \cos \theta)^{-1/2} \, d(1 + \cos \theta)$$


$$= -4a\sqrt{2} \sqrt{1 + \cos \theta} \Big|_0^\pi$$

$$= -4a\sqrt{2}(0 - \sqrt{2})$$

$$= 8a \text{ unit} \quad \Big|$$

Exercise

The involute of a circle is described by the endpoint P of a string that is held taut as it is unwound from a spool that does not turn.



Show that a parametric representation of the involute is

$$x = r(\cos \theta + \theta \sin \theta) \quad \text{and} \quad y = r(\sin \theta - \theta \cos \theta)$$

Solution

$$\Delta OAC: \quad \cos \theta = \frac{t}{r} \quad \sin \theta = \frac{h}{r}$$

$$\Delta PBC: \quad \cos \theta = \frac{w}{r\theta} \quad \sin \theta = \frac{u}{r\theta}$$

$$x = t + u$$


$$= r \cos \theta + r\theta \sin \theta$$

$$= r(\cos \theta + \theta \sin \theta)$$

$$y = h - w$$


$$= r \sin \theta - r\theta \cos \theta$$

$$= r(\sin \theta - \theta \cos \theta)$$



Exercise

The figure shows a piece of string tied to a circle with a radius of one unit. The string is just long enough to reach the opposite side of the circle.



Find the area that is covered when the string is unwound counterclockwise.

Solution

From previous exercise, we have

$$x = \cos \theta + \theta \sin \theta \quad \text{and} \quad y = \sin \theta - \theta \cos \theta$$

At $(-1, \pi)$, the string is fully extended and has length x .

The area of region **A** is:

$$\frac{1}{4} \pi r^2 = \frac{1}{4} \pi^3 \Big|$$

The area of region **C + D** is:


$$\frac{1}{2} \pi r^2 = \frac{\pi}{2} \Big|$$

$$\frac{dx}{d\theta} = -\sin \theta + \sin \theta + \theta \cos \theta$$

$$= \theta \cos \theta \Big|$$

$$\frac{dx}{d\theta} = \theta \cos \theta = 0$$

$$= \theta \cos \theta \Big|$$



The area of the region **B + C + D** is given by

$$\int_{\pi}^{\pi/2} y dx - \int_0^{\pi/2} y dx = \int_{\pi}^0 y dx$$

$$A_2 = \int_{\pi}^0 (\sin \theta - \theta \cos \theta) \theta \cos \theta d\theta$$

$$= \int_{\pi}^0 \left(\theta \cos \theta \sin \theta - \theta^2 \cos^2 \theta \right) d\theta$$

$$= \int_{\pi}^0 \left(\frac{1}{2} \theta \sin 2\theta - \frac{1}{2} \theta^2 - \frac{1}{2} \theta^2 \cos 2\theta \right) d\theta$$

		$\int \sin 2\theta$
+	θ	$-\frac{1}{2} \cos 2\theta$
-	1	$-\frac{1}{4} \sin 2\theta$

		$\int \cos 2\theta$
+	$\frac{1}{2} \theta^2$	$\frac{1}{2} \sin 2\theta$
-	θ	$-\frac{1}{4} \cos 2\theta$
+	1	$-\frac{1}{8} \sin 2\theta$

$$= -\frac{1}{4} \theta \cos 2\theta + \frac{1}{8} \sin 2\theta - \frac{1}{6} \theta^3 - \frac{1}{4} \theta^2 \sin 2\theta - \frac{1}{4} \theta \cos 2\theta + \frac{1}{8} \sin 2\theta \Big|_{\pi}^0$$


$$= \frac{\pi}{4} + \frac{\pi^3}{6} + \frac{\pi}{4}$$

$$= \frac{\pi^3}{6} + \frac{\pi}{2} \Big|$$

$$\begin{aligned}\text{Total area covered} &= 2 \left(\frac{\pi^3}{4} + \frac{\pi^3}{6} + \frac{\pi}{2} - \frac{\pi}{2} \right) \\ &= \frac{5\pi^3}{6} \text{ unit}^2\end{aligned}$$

Exercise

An Air traffic controller spots two planes at the same altitude flying toward each other.



Their flight paths are 20° and 315° . One plane is 150 miles from point P with a speed of 375 miles per hour. The other is 190 miles from point P with a speed of 450 miles per hour.

- Find parametric equations for the path of each plane where t is the time in hours, with $t = 0$ corresponding to the time at which the air traffic controller spots the planes.
- Use part (a) to write the distance between the planes as a function of t .
- Graph the function in part (b).
- When the distance between the planes be minimum?
- If the planes must keep a separation of at least 3 miles, is the requirement met?


Solution

a) **First Plane:**

Given: $\theta_1 = 90^\circ - 20^\circ = 70^\circ$ $d_1 = 150$ $v_1 = 375$

$$\begin{cases} x_1 = (150 - 375t) \cos 70^\circ \\ y_1 = (150 - 375t) \sin 70^\circ \end{cases}$$

$$\begin{cases} x_1 = 75(2 - 5t) \cos 70^\circ \\ y_1 = 75(2 - 5t) \sin 70^\circ \end{cases}$$



Second Plane:

Given: $\theta_2 = 45^\circ$ $d_2 = 190$ $v_2 = 450$

$$\begin{cases} x_2 = -(190 - 450t) \cos 45^\circ \\ y_2 = (190 - 450t) \sin 45^\circ \end{cases}$$

$$\begin{cases} x_2 = -10(19 - 45t)\left(\frac{\sqrt{2}}{2}\right) \\ y_2 = 10(19 - 45t)\left(\frac{\sqrt{2}}{2}\right) \end{cases}$$

$$\begin{cases} x_2 = -5\sqrt{2}(19 - 45t) \\ y_2 = 5\sqrt{2}(19 - 45t) \end{cases}$$

$$b) \quad d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$


$$= \sqrt{\left(-5\sqrt{2}(19 - 45t) - 75(2 - 5t)\cos 70^\circ\right)^2 + \left(5\sqrt{2}(19 - 45t) - 75(2 - 5t)\sin 70^\circ\right)^2}$$

At $t = 0$


$$d = \sqrt{190^2 + 150^2 - 2(190)(150)\cos 65^\circ}$$

$$\approx 185.77 \quad |$$

c)



d) Using software:



0.4105000000	7.8029045315
0.4110000000	7.7540582513
0.4115000000	7.7114048932
0.4120000000	7.6750477077
0.4125000000	7.6450765223
0.4130000000	7.6215666744
0.4135000000	7.6045780909
0.4140000000	7.5941545371
0.4145000000	7.5903230599
0.4150000000	7.5930936382
0.4155000000	7.6024590542
0.4160000000	7.6183949864
0.4165000000	7.6408603242
0.4170000000	7.6697976925
0.4175000000	7.7051341728
0.4180000000	7.7467821981
0.4185000000	7.7946405981
0.4190000000	7.8485957667
0.4195000000	7.9085229209
0.4200000000	7.9742874225

The minimum distance is 7.59 *miles* when $t = 0.4145$

e) Yes, the planes must keep a separation of at least 3 *miles*.