

## Section 2.5 – Graphing Polynomial Functions

### Polynomial Function

A *Polynomial function*  $P(x)$  in  $x$  is a sum of the form is given by:

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + a_{n-2} x^{n-2} + \dots + a_2 x^2 + a_1 x + a_0$$

Where the coefficients  $a_n, a_{n-1}, \dots, a_2, a_1, a_0$  are real numbers and the exponents are whole numbers.

The diagram shows the term  $a_n x^n$ . An arrow points from the word "Degree" to the exponent  $n$ . Another arrow points from the words "Leading Term" to the entire term  $a_n x^n$ . A third arrow points from the words "Leading Coefficient" to the coefficient  $a_n$ .

Non-polynomial Functions:  $\frac{1}{x} + 2x$ ;  $\sqrt{x^2 - 3} + x$ ;  $\frac{x-5}{x^2+2}$

<i>Degree of <math>f</math></i>	<i>Form of <math>f(x)</math></i>	<i>Graph of <math>f(x)</math></i>
0	$f(x) = a_0$	A horizontal line
1	$f(x) = a_1 x + a_0$	A line with slope $a_1$
2	$f(x) = a_2 x^2 + a_1 x + a_0$	A parabola with a vertical axis

All polynomial functions are *continuous functions*.

## End Behavior ( $a_n x^n$ )


If  $n$  (degree) is **even**:

If  $a_n < 0$  (in front  $x^n$  is negative).

Then the function falls from the left and right side

$$x \rightarrow -\infty \Rightarrow f(x) \rightarrow -\infty$$

$$x \rightarrow \infty \Rightarrow f(x) \rightarrow -\infty$$




If  $a_n > 0$  (in front  $x^n$  is positive).

Then the function rises from the left and right side

$$x \rightarrow -\infty \Rightarrow f(x) \rightarrow \infty$$

$$x \rightarrow \infty \Rightarrow f(x) \rightarrow \infty$$




If  $n$  (degree) is **odd**:

If  $a_n < 0$  (negative).

Then the function rises from the left side and falls from the right side

$$x \rightarrow -\infty \Rightarrow f(x) \rightarrow \infty$$

$$x \rightarrow \infty \Rightarrow f(x) \rightarrow -\infty$$




If  $a_n > 0$  (positive).

Then the function falls from the left side and rises from the right side

$$x \rightarrow -\infty \Rightarrow f(x) \rightarrow -\infty$$

$$x \rightarrow \infty \Rightarrow f(x) \rightarrow \infty$$



## Example

Determine the end behavior of the graph of the polynomial function  $f(x) = -4x^5 + 7x^2 - x + 9$

### Solution

Leading term:  $-4x^5$  with 5th degree ( $n$  is odd)

$$x \rightarrow -\infty \Rightarrow f(x) = -(-)^5 = (-)(-) = + \rightarrow \infty \quad f(x) \text{ rises left}$$

$$x \rightarrow \infty \Rightarrow f(x) \rightarrow -\infty \quad f(x) \text{ falls right}$$



## The Intermediate Value *Theorem*

For any polynomial function  $f(x)$  with real coefficients and  $f(a) \neq f(b)$  for  $a < b$ , then  $f$  takes on every value between  $f(a)$  and  $f(b)$  in the interval  $[a, b]$ .

$\therefore f(a)$  and  $f(b)$  are the **opposite signs**. Then the function has a real zero between  $a$  and  $b$ .

### Example

Using the intermediate value theorem, determine, if possible, whether the function has a real zero between  $a$  and  $b$ .

a)  $f(x) = x^3 + x^2 - 6x$ ;  $a = -4$ ,  $b = -2$

b)  $f(x) = x^3 + x^2 - 6x$ ;  $a = -1$ ,  $b = 3$

### Solution

a)  $f(x) = x^3 + x^2 - 6x$ ;  $a = -4$ ,  $b = -2$

$$f(-4) = (-4)^3 + (-4)^2 - 6(-4) \\ = -24$$

$$f(-2) = (-2)^3 + (-2)^2 - 6(-2) \\ = 8$$

$\therefore f(x)$  has a zero between  $-4$  and  $-2$

b)  $f(x) = x^3 + x^2 - 6x$ ;  $a = -1$ ,  $b = 3$

$$f(-1) = (-1)^3 + (-1)^2 - 6(-1) \\ = 6$$

$$f(3) = (3)^3 + (3)^2 - 6(3) = 18 \\ = 18$$

$\therefore f(x)$  zeros *can't be determined*

### Example

Show that  $f(x) = x^5 + 2x^4 - 6x^3 + 2x - 3$  has a zero between 1 and 2.

### Solution

$$f(1) = 1 + 2 - 6 + 2 - 3 \\ = -4$$

$$f(2) = (2)^5 + 2(2)^4 - 6(2)^3 + 2(2) - 3 \\ = 17$$

Since  $f(1)$  and  $f(2)$  have opposite signs.

Therefore,  $f(c) = 0$  for at least one real number  $c$  between 1 and 2.

## Sketching

### Example


Let  $f(x) = x^3 + x^2 - 4x - 4$ . Find all values of  $x$  such that  $f(x) > 0$  and all  $x$  such that  $f(x) < 0$ , and then sketch the graph of  $f$ .

### Solution

$$\begin{aligned}f(x) &= x^3 + x^2 - 4x - 4 \\&= x^2(x+1) - 4(x+1) \\&= (x+1)(x^2 - 4) \\&= (x+1)(x+2)(x-2)\end{aligned}$$

The zeros of  $f(x)$  ( $x$ -intercepts) are:  $-2$ ,  $-1$ , and  $2$

<i>Interval</i>	$-\infty$	$-2$	$-1$	<b>0</b>	$2$	$\infty$
Sign of $f(x)$		<b>−</b>	<b>+</b>	<b>−</b>	<b>+</b>	
Position		<b>Below <math>x</math>-axis</b>	<b>Above <math>x</math>-axis</b>	<b>Below <math>x</math>-axis</b>	<b>Above <math>x</math>-axis</b>	



We can conclude from the chart and the graph that:

$$f(x) > 0 \quad \text{if } x \text{ is in } (-2, -1) \cup (2, \infty)$$

$$f(x) < 0 \quad \text{if } x \text{ is in } (-\infty, -2) \cup (-1, 2)$$

### Example

Let  $f(x) = x^4 - 4x^3 + 3x^2$ . Find all values of  $x$  such that  $f(x) > 0$  and all  $x$  such that  $f(x) < 0$ , and then sketch the graph of  $f$ .


### Solution

$$\begin{aligned} f(x) &= x^2(x^2 - 4x + 3) \\ &= x^2(x-1)(x-3) \end{aligned}$$

The zeros are: 0, 1, 3.

Since the factor  $x^2$  is always positive, it has no factor

$-\infty$	1	<b>2</b>	3	$\infty$
+		-		+



$$f(x) > 0 \Rightarrow x \text{ is in } (-\infty, 0) \cup (0, 1) \cup (3, \infty)$$

$$f(x) < 0 \Rightarrow x \text{ is in } (1, 3)$$

## Exercises      Section 2.5 – Polynomial Functions

(1 – 12) Determine the end behavior of the graph of the polynomial function

1.  $f(x) = 5x^3 + 7x^2 - x + 9$

7.  $f(x) = -5x^4 + 7x^2 - x + 9$

2.  $f(x) = 11x^3 - 6x^2 + x + 3$

8.  $f(x) = -11x^4 - 6x^2 + x + 3$

3.  $f(x) = -11x^3 - 6x^2 + x + 3$

9.  $f(x) = 5x^5 - 16x^2 - 20x + 64$

4.  $f(x) = 2x^3 + 3x^2 - 23x - 42$

10.  $f(x) = -5x^5 - 16x^2 - 20x + 64$

5.  $f(x) = 5x^4 + 7x^2 - x + 9$

11.  $f(x) = -3x^6 - 16x^3 + 64$

6.  $f(x) = 11x^4 - 6x^2 + x + 3$

12.  $f(x) = 3x^6 - 16x^3 + 4$

(13 – 32) Use the Intermediate Value Theorem to show that each polynomial has a real zero between the given integers.

13.  $f(x) = x^3 - x - 1$ ; between 1 and 2

14.  $f(x) = x^3 - 4x^2 + 2$ ; between 0 and 1

15.  $f(x) = 2x^4 - 4x^2 + 1$ ; between -1 and 0

16.  $f(x) = x^4 + 6x^3 - 18x^2$ ; between 2 and 3

17.  $f(x) = x^3 + x^2 - 2x + 1$ ; between -3 and -2

18.  $f(x) = x^5 - x^3 - 1$ ; between 1 and 2

19.  $f(x) = 3x^3 - 10x + 9$ ; between -3 and -2

20.  $f(x) = 3x^3 - 8x^2 + x + 2$ ; between 2 and 3

21.  $f(x) = 3x^3 - 8x^2 + x + 2$ ; between 1 and 2

22.  $f(x) = x^5 + 2x^4 - 6x^3 + 2x - 3$ ; between 0 and 1

23.  $P(x) = 2x^3 + 3x^2 - 23x - 42$ ,  $a = 3$ ,  $b = 4$

24.  $P(x) = 4x^3 - x^2 - 6x + 1$ ,  $a = 0$ ,  $b = 1$

25.  $P(x) = 3x^3 + 7x^2 + 3x + 7$ ,  $a = -3$ ,  $b = -2$

26.  $P(x) = 2x^3 - 21x^2 - 2x + 25$ ,  $a = 1$ ,  $b = 2$

27.  $P(x) = 4x^4 + 7x^3 - 11x^2 + 7x - 15$ ,  $a = 1$ ,  $b = \frac{3}{2}$

28.  $P(x) = 5x^3 - 16x^2 - 20x + 64$ ,  $a = 3$ ,  $b = \frac{7}{2}$



29.  $P(x) = x^4 - x^2 - x - 4$ ,  $a = 1$ ,  $b = 2$   
 30.  $P(x) = x^3 - x - 8$ ,  $a = 2$ ,  $b = 3$   
 31.  $P(x) = x^3 - x - 8$ ,  $a = 0$ ,  $b = 1$   
 32.  $P(x) = x^3 - x - 8$ ,  $a = 2.1$ ,  $b = 2.2$

(33 – 91) Find all values of  $x$  such that  $f(x) > 0$  and all  $x$  such that  $f(x) < 0$ , and then sketch the graph of  $f$

- |   |   |
|---|---|
| 33. $f(x) = x^4 - 4x^2$                 | 53. $f(x) = 3x^3 + 11x^2 - 6x - 8$          |
| 34. $f(x) = x^4 + 3x^3 - 4x^2$          | 54. $f(x) = 2x^3 + 9x^2 - 2x - 9$           |
| 35. $f(x) = x^3 + 2x^2 - 4x - 8$        | 55. $f(x) = x^3 + 3x^2 - 6x - 8$            |
| 36. $f(x) = x^3 - 3x^2 - 9x + 27$       | 56. $f(x) = 3x^3 - x^2 - 6x + 2$            |
| 37. $f(x) = -x^4 + 12x^2 - 27$          | 57. $f(x) = x^3 - 8x^2 + 8x + 24$           |
| 38. $f(x) = x^2(x+2)(x-1)^2(x-2)$       | 58. $f(x) = x^3 - 7x^2 - 7x + 69$           |
| 39. $f(x) = 2x^3 + 11x^2 - 7x - 6$      | 59. $f(x) = x^3 - 3x - 2$                   |
| 40. $f(x) = x^3 + 2x^2 - 5x - 6$        | 60. $f(x) = x^3 - 2x + 1$                   |
| 41. $f(x) = x^3 + 8x^2 + 11x - 20$      | 61. $f(x) = x^3 - 2x^2 - 11x + 12$          |
| 42. $f(x) = x^4 + x^2 - 2$              | 62. $f(x) = x^3 - 2x^2 - 7x - 4$            |
| 43. $f(x) = x^4 - x^3 - 6x^2 + 4x + 8$  | 63. $f(x) = x^3 - 10x - 12$                 |
| 44. $f(x) = 4x^5 - 8x^4 - x + 2$        | 64. $f(x) = x^3 - 5x^2 + 17x - 13$          |
| 45. $f(x) = 2x^4 - x^3 - 5x^2 + 2x + 2$ | 65. $f(x) = 6x^3 + 25x^2 - 24x + 5$         |
| 46. $f(x) = x^3 - x^2 - 10x - 8$        | 66. $f(x) = 8x^3 + 18x^2 + 45x + 27$        |
| 47. $f(x) = x^3 + x^2 - 14x - 24$       | 67. $f(x) = 3x^3 - x^2 + 11x - 20$          |
| 48. $f(x) = 2x^3 - 3x^2 - 17x + 30$     | 68. $f(x) = x^4 - x^3 - 9x^2 + 3x + 18$     |
| 49. $f(x) = 12x^3 + 8x^2 - 3x - 2$      | 69. $f(x) = 2x^4 - 9x^3 + 9x^2 + x - 3$     |
| 50. $f(x) = x^3 + x^2 - 6x - 8$         | 70. $f(x) = 6x^4 + 5x^3 - 17x^2 - 6x$       |
| 51. $f(x) = x^3 - 19x - 30$             | 71. $f(x) = x^4 - 2x^2 - 16x - 15$          |
|   | 72. $f(x) = x^4 - 2x^3 - 5x^2 + 8x + 4$     |
|   | 73. $f(x) = 2x^4 - 17x^3 + 4x^2 + 35x - 24$ |

52.  $f(x) = 2x^3 + x^2 - 25x + 12$
74.  $f(x) = x^4 + x^3 - 3x^2 - 5x - 2$
75.  $f(x) = 6x^4 - 17x^3 - 11x^2 + 42x$
84.  $f(x) = x^4 + 3x^3 - 30x^2 - 6x + 56$
76.  $f(x) = x^4 - 5x^2 - 2x$
85.  $f(x) = 3x^5 - 10x^4 - 6x^3 + 24x^2 + 11x - 6$
77.  $f(x) = 3x^4 - 4x^3 - 11x^2 + 16x - 4$
86.  $f(x) = 6x^5 + 19x^4 + x^3 - 6x^2$
78.  $f(x) = 6x^4 + 23x^3 + 19x^2 - 8x - 4$
87.  $f(x) = x^5 + 5x^4 + 10x^3 + 10x^2 + 5x + 1$
79.  $f(x) = 4x^4 - 12x^3 + 3x^2 + 12x - 7$
88.  $f(x) = x^5 - x^4 - 7x^3 + 7x^2 + 12x - 12$
80.  $f(x) = 2x^4 - 9x^3 - 2x^2 + 27x - 12$
89.  $f(x) = x^5 - 2x^3 - 8x$
81.  $f(x) = 2x^4 - 19x^3 + 51x^2 - 31x + 5$
90.  $f(x) = x^5 - 7x^4 + 19x^3 - 37x^2 + 60x - 36$
82.  $f(x) = 4x^4 - 35x^3 + 71x^2 - 4x - 6$
91.  $f(x) = 3x^6 - 10x^5 - 29x^4 + 34x^3 + 50x^2 - 24x - 24$
83.  $f(x) = 2x^4 + 3x^3 - 4x^2 - 3x + 2$