

***Solution*****Section 2.9 – Rank and the Fundamental Matrix Spaces*****Exercise***

Verify that  $\text{rank}(A) = \text{rank}(A^T)$

$$A = \begin{bmatrix} 1 & 2 & 4 & 0 \\ -3 & 1 & 5 & 2 \\ -2 & 3 & 9 & 2 \end{bmatrix}$$

**Solution**

$$\begin{bmatrix} 1 & 2 & 4 & 0 \\ -3 & 1 & 5 & 2 \\ -2 & 3 & 9 & 2 \end{bmatrix} \quad \begin{array}{l} \\ R_2 + 3R_1 \\ R_3 + 2R_1 \end{array}$$

$$\begin{bmatrix} 1 & 2 & 4 & 0 \\ 0 & 7 & 17 & 2 \\ 0 & 7 & 17 & 2 \end{bmatrix} \quad R_3 - R_2$$

$$\begin{bmatrix} 1 & 2 & 4 & 0 \\ 0 & 7 & 17 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad 7R_1 - 2R_2$$

$$\begin{bmatrix} 7 & 0 & -6 & -4 \\ 0 & 7 & 17 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \begin{array}{l} \frac{1}{7}R_1 \\ \frac{1}{7}R_2 \\ \end{array}$$

$$\begin{bmatrix} 1 & 0 & -\frac{6}{7} & -\frac{4}{7} \\ 0 & 1 & \frac{17}{7} & \frac{2}{7} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\text{rank}(A) = 2$$

$$A^T = \begin{bmatrix} 1 & -3 & -2 \\ 2 & 1 & 3 \\ 4 & 5 & 9 \\ 0 & 2 & 2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -3 & -2 \\ 2 & 1 & 3 \\ 4 & 5 & 9 \\ 0 & 2 & 2 \end{bmatrix} \quad \begin{array}{l} \\ R_2 - 2R_1 \\ R_3 - 4R_1 \\ \end{array}$$

$$\begin{bmatrix} 1 & -3 & -2 \\ 0 & 7 & 7 \\ 0 & 17 & 17 \\ 0 & 2 & 2 \end{bmatrix} \quad \begin{array}{l} \frac{1}{7}R_2 \\ \\ \\ \frac{1}{2}R_4 \end{array}$$

$$\begin{bmatrix} 1 & -3 & -2 \\ 0 & 1 & 1 \\ 0 & 17 & 17 \\ 0 & 1 & 1 \end{bmatrix} \quad \begin{array}{l} R_1 + 3R_2 \\ \\ R_3 - 17R_2 \\ R_4 - R_2 \end{array}$$

$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\text{rank}(A^T) = 2$$

$$\underline{\text{rank}(A) = \text{rank}(A^T) = 2}$$

### ***Exercise***

Find the rank and nullity of the matrix; then verify that the values obtained satisfy

$$\text{rank}(A) + N(A) = n$$

$$A = \begin{bmatrix} 1 & -1 & 3 \\ 5 & -4 & -4 \\ 7 & -6 & 2 \end{bmatrix}$$

### **Solution**

$$\begin{bmatrix} 1 & -1 & 3 \\ 5 & -4 & -4 \\ 7 & -6 & 2 \end{bmatrix} \quad \begin{array}{l} \\ R_2 - 5R_1 \\ R_3 - 7R_1 \end{array}$$

$$\begin{bmatrix} 1 & -1 & 3 \\ 0 & 1 & -19 \\ 0 & 1 & -19 \end{bmatrix} \quad \begin{array}{l} R_1 + R_2 \\ \\ R_3 - R_2 \end{array}$$

$$\begin{bmatrix} 1 & 0 & -16 \\ 0 & 1 & -19 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\text{rank}(A) = 2$$

$$\text{nullity}(A) = 1$$

$$\text{rank}(A) + \text{nullity}(A) = 2 + 1 = 3 = \textcolor{red}{n} \leftarrow \text{number of columns}$$

### ***Exercise***

Find the rank and nullity of the matrix; then verify that the values obtained satisfy

$$\text{rank}(A) + N(A) = n$$

$$A = \begin{bmatrix} 1 & 4 & 5 & 2 \\ 2 & 1 & 3 & 0 \\ -1 & 3 & 2 & 2 \end{bmatrix}$$

### **Solution**

$$\begin{bmatrix} 1 & 4 & 5 & 2 \\ 2 & 1 & 3 & 0 \\ -1 & 3 & 2 & 2 \end{bmatrix} \begin{array}{l} \\ R_2 - 2R_1 \\ R_3 + R_1 \end{array}$$

$$\begin{bmatrix} 1 & 4 & 5 & 2 \\ 0 & -7 & -7 & -4 \\ 0 & 7 & 7 & 4 \end{bmatrix} \begin{array}{l} 7R_1 + 4R_2 \\ \\ R_3 + R_2 \end{array}$$

$$\begin{bmatrix} 7 & 0 & 7 & -2 \\ 0 & -7 & -7 & -4 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{array}{l} \frac{1}{7}R_1 \\ -\frac{1}{7}R_2 \\ \end{array}$$

$$\begin{bmatrix} 1 & 0 & 1 & -\frac{2}{7} \\ 0 & 1 & 1 & \frac{4}{7} \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\text{rank}(A) = 2$$

$$\text{nullity}(A) = 1$$

$$\text{rank}(A) + \text{nullity}(A) = 2 + 1 = 3 = \textcolor{red}{n}$$

### Exercise

Find the rank and nullity of the matrix; then verify that the values obtained satisfy

$$\text{rank}(A) + N(A) = n$$

$$A = \begin{bmatrix} 1 & 4 & 5 & 6 & 9 \\ 3 & -2 & 1 & 4 & -1 \\ -1 & 0 & -1 & -2 & -1 \\ 2 & 3 & 5 & 7 & 8 \end{bmatrix}$$

### Solution

$$\begin{bmatrix} 1 & 4 & 5 & 6 & 9 \\ 3 & -2 & 1 & 4 & -1 \\ -1 & 0 & -1 & -2 & -1 \\ 2 & 3 & 5 & 7 & 8 \end{bmatrix} \begin{array}{l} \\ R_2 - 3R_1 \\ R_3 + R_1 \\ R_4 - 2R_1 \end{array}$$

$$\begin{bmatrix} 1 & 4 & 5 & 6 & 9 \\ 0 & -14 & -14 & -14 & -28 \\ 0 & 4 & 4 & 4 & 8 \\ 0 & -5 & -5 & -5 & -10 \end{bmatrix} \begin{array}{l} \\ -\frac{1}{14}R_2 \\ \frac{1}{4}R_3 \\ -\frac{1}{5}R_4 \end{array}$$

$$\begin{bmatrix} 1 & 4 & 5 & 6 & 9 \\ 0 & 1 & 1 & 1 & 2 \\ 0 & 1 & 1 & 1 & 2 \\ 0 & 1 & 1 & 1 & 2 \end{bmatrix} \begin{array}{l} R_1 - 4R_2 \\ \\ R_3 - R_2 \\ R_4 - R_2 \end{array}$$

$$\begin{bmatrix} 1 & 0 & 1 & 2 & 1 \\ 0 & 1 & 1 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\text{rank}(A) = 2$$

$$\text{nullity}(A) = 2$$

$$\text{rank}(A) + \text{nullity}(A) = 2 + 2 = 4 = n$$

### Exercise

Find the rank and nullity of the matrix; then verify that the values obtained satisfy

$$\text{rank}(A) + N(A) = n$$

$$A = \begin{bmatrix} 1 & -3 & 2 & 2 & 1 \\ 0 & 3 & 6 & 0 & -3 \\ 2 & -3 & -2 & 4 & 4 \\ 3 & -6 & 0 & 6 & 5 \\ -2 & 9 & 2 & -4 & -5 \end{bmatrix}$$

### Solution

$$\begin{bmatrix} 1 & -3 & 2 & 2 & 1 \\ 0 & 3 & 6 & 0 & -3 \\ 2 & -3 & -2 & 4 & 4 \\ 3 & -6 & 0 & 6 & 5 \\ -2 & 9 & 2 & -4 & -5 \end{bmatrix} \begin{array}{l} \\ R_3 - 2R_1 \\ R_4 - 3R_1 \\ R_5 + 2R_1 \end{array}$$

$$\begin{bmatrix} 1 & -3 & 2 & 2 & 1 \\ 0 & 3 & 6 & 0 & -3 \\ 0 & 3 & -6 & 0 & 2 \\ 0 & 3 & -6 & 0 & 2 \\ 0 & 3 & 6 & 0 & -3 \end{bmatrix} \begin{array}{l} \\ R_4 - R_3 \\ R_5 - R_2 \end{array}$$

$$\begin{bmatrix} 1 & -3 & 2 & 2 & 1 \\ 0 & 3 & 6 & 0 & -3 \\ 0 & 3 & -6 & 0 & 2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{array}{l} R_1 + R_2 \\ \\ R_3 - R_2 \\ \\ \end{array}$$

$$\begin{bmatrix} 1 & 0 & 8 & 2 & -2 \\ 0 & 3 & 6 & 0 & -3 \\ 0 & 0 & -12 & 0 & 5 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{array}{l} \\ \frac{1}{3}R_2 \\ -\frac{1}{12}R_3 \\ \\ \end{array}$$

$$\begin{bmatrix} 1 & 0 & 8 & 2 & -2 \\ 0 & 1 & 2 & 0 & -1 \\ 0 & 0 & 1 & 0 & -\frac{5}{12} \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{array}{l} R_1 - 8R_3 \\ R_2 - 2R_3 \\ \\ \\ \end{array}$$

$$\begin{bmatrix} 1 & 0 & 0 & 2 & \frac{4}{3} \\ 0 & 1 & 0 & 0 & -\frac{1}{6} \\ 0 & 0 & 1 & 0 & -\frac{5}{12} \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\text{rank}(A) = 3$$

$$NS(A) = 2$$

$$\text{Number of columns} = 5$$

$$\text{rank}(A) + NS(A) = 3 + 2 = 5 = n$$

### Exercise

If  $A$  is an  $m \times n$  matrix, what is the largest possible value for its rank and the smallest possible value of the nullity of  $A$ .

### Solution

The largest possible value for the rank of an  $m \times n$  matrix:

- $n$  if  $m \geq n$  (when every column of the  $rref(A)$  contains a leading 1)
- $m$  if  $m < n$  (when every row of the  $rref(A)$  contains a leading 1)

The smallest possible value for the nullity of an  $m \times n$  matrix:

- $0$  if  $m \geq n$  (when every column of the  $rref(A)$  contains a leading 1)
- $n - m$  if  $m < n$  (when every row of the  $rref(A)$  contains a leading 1)

### Exercise

Discuss how the rank of  $A$  varies with  $t$ .

$$a) A = \begin{bmatrix} 1 & 1 & t \\ 1 & t & 1 \\ t & 1 & 1 \end{bmatrix} \quad b) A = \begin{bmatrix} t & 3 & -1 \\ 3 & 6 & -2 \\ -1 & -3 & t \end{bmatrix}$$

### Solution

$$\begin{aligned} a) \begin{vmatrix} 1 & 1 & t \\ 1 & t & 1 \\ t & 1 & 1 \end{vmatrix} &= t + t + t - t^3 - 1 - 1 \\ &= -t^3 + 3t - 2 = 0 \end{aligned}$$

Solve for  $t$ :  $\underline{t=1, -2, -2}$

Therefore,  $\text{rank}(A) = 3$  for  $\forall t - \{1, -2, -2\}$

$$\text{If } t = 1, A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix} \begin{array}{l} \\ R_2 - R_1 \\ R_3 - R_1 \end{array}$$

$$\begin{bmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\text{rank}(A) = 1$$

$$\text{If } t = -2, A = \begin{bmatrix} 1 & 1 & -2 \\ 1 & -2 & 1 \\ -2 & 1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & -2 \\ 0 & -3 & 3 \\ 0 & 3 & -3 \end{bmatrix} \begin{array}{l} \\ \\ R_3 + R_2 \end{array}$$

$$\begin{bmatrix} 1 & 1 & -2 \\ 0 & -3 & 3 \\ 0 & 0 & 0 \end{bmatrix} \begin{array}{l} \\ -\frac{1}{3}R_2 \\ \end{array}$$

$$\begin{bmatrix} 1 & 1 & -2 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{bmatrix} \begin{array}{l} R_1 - R_2 \\ \\ \end{array}$$

$$\begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\text{rank}(A) = 2$$

$$\begin{aligned} b) \quad \begin{vmatrix} t & 3 & -1 \\ 3 & 6 & -2 \\ -1 & -3 & t \end{vmatrix} &= 6t^2 + 6 + 9 - 6 - 6t - 9t \\ &= 6t^2 - 15t + 9 = 0 \end{aligned}$$

Solve for  $t$ :  $t=1, \frac{3}{2}$

Therefore,  $\text{rank}(A)=3$  for  $\forall t - \left\{1, \frac{3}{2}\right\}$

$$\text{If } t=1, A = \begin{bmatrix} 1 & 3 & -1 \\ 3 & 6 & -2 \\ -1 & -3 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 3 & -1 \\ 3 & 6 & -2 \\ -1 & -3 & 1 \end{bmatrix} \begin{array}{l} \\ R_2 - 3R_1 \\ R_3 + R_1 \end{array}$$

$$\begin{bmatrix} 1 & 3 & -1 \\ 0 & -3 & 1 \\ 0 & 0 & 0 \end{bmatrix} R_1 + R_2$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & -3 & 1 \\ 0 & 0 & 0 \end{bmatrix} -\frac{1}{3}R_2$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -\frac{1}{3} \\ 0 & 0 & 0 \end{bmatrix}$$

$$\text{rank}(A)=2$$

$$\text{If } t=\frac{3}{2}, A = \begin{bmatrix} \frac{3}{2} & 3 & -1 \\ 3 & 6 & -2 \\ -1 & -3 & \frac{3}{2} \end{bmatrix}$$

$$\begin{bmatrix} \frac{3}{2} & 3 & -1 \\ 3 & 6 & -2 \\ -1 & -3 & \frac{3}{2} \end{bmatrix} \begin{array}{l} 2R_1 \\ \\ 2R_3 \end{array}$$

$$\begin{bmatrix} 3 & 6 & -2 \\ 3 & 6 & -2 \\ -2 & -6 & 3 \end{bmatrix} \begin{array}{l} \\ R_2 - R_1 \\ 3R_3 + 2R_1 \end{array}$$



$$\begin{bmatrix} 3 & 6 & -2 \\ 0 & 0 & 0 \\ 0 & -6 & 5 \end{bmatrix}$$

$$\begin{bmatrix} 3 & 6 & -2 \\ 0 & -6 & 5 \\ 0 & 0 & 0 \end{bmatrix} \quad R_1 + R_2$$

$$\begin{bmatrix} 3 & 0 & 3 \\ 0 & -6 & 5 \\ 0 & 0 & 0 \end{bmatrix} \quad \begin{matrix} \frac{1}{3}R_1 \\ -\frac{1}{6}R_2 \end{matrix}$$

$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & -\frac{5}{6} \\ 0 & 0 & 0 \end{bmatrix}$$

$$\text{rank}(A) = 2$$

### ***Exercise***

Are there values of  $r$  and  $s$  for which

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & r-2 & 2 \\ 0 & s-1 & r+2 \\ 0 & 0 & 3 \end{bmatrix}$$

Has rank 1? Has rank 2? If so, find those values.

### ***Solution***

Since the third column will always have a nonzero entry, the *rank* will never be 1. (row 1 and row 4 never have a nonzero entry).

If  $r = 2$  and  $s = 1$ , that implies to

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & r-2 & 2 \\ 0 & s-1 & r+2 \\ 0 & 0 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 2 \\ 0 & 0 & 4 \\ 0 & 0 & 3 \end{bmatrix}$$

$$\rightarrow \text{rank} = 2$$

### Exercise

Find the row reduced form  $R$  and the rank  $r$  of  $A$  (those depend on  $c$ ).

Which are the pivot columns of  $A$ ? Which variables are free? What are the special solutions and the nullspace matrix  $N$  (always depending on  $c$ )?

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 3 & 6 & 3 \\ 4 & 8 & c \end{bmatrix} \quad \text{and} \quad A = \begin{bmatrix} c & c \\ c & c \end{bmatrix}$$

### Solution

$$a) \quad c \neq 4 \quad R = \begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix},$$

$\text{rank}(A) = 2$ , the pivot columns are 1 and 3, the second variable  $x_2$  is free.

$$\text{The special solution: } x_2 = 1 \text{ which yields to } N = \begin{pmatrix} -2 \\ 1 \\ 0 \end{pmatrix}$$

$$c = 4 \quad R = \begin{pmatrix} 1 & 2 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

$\text{rank}(A) = 1$ , the pivot column is column 1, the second and third variables  $x_2, x_3$  are free.

$$\text{The special solution goes into } N = \begin{pmatrix} -2 & -1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$b) \quad c \neq 0 \quad R = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix},$$

$\text{rank}(A) = 1$ , the pivot column is the first column, the second variable  $x_2$  is free.

$$\text{The special solution: } x_2 = 1 \text{ which yields to } N = \begin{pmatrix} -1 \\ 1 \end{pmatrix}$$

$$c = 0 \quad R = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$$

$\text{rank}(A) = 0$ , the matrix has no pivot column, and both variables are free.

$$\text{The special solution is: } N = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

### Exercise

Find the row reduced form  $R$  and the rank  $r$  of  $A$  (those depend on  $c$ ).

Which are the pivot columns of  $A$ ? Which variables are free? What are the special solutions and the nullspace matrix  $N$  (always depending on  $c$ )?

$$A = \begin{bmatrix} 1 & 1 & 2 & 2 \\ 2 & 2 & 4 & 4 \\ 1 & c & 2 & 2 \end{bmatrix} \quad \text{and} \quad A = \begin{bmatrix} 1-c & 2 \\ 0 & 2-c \end{bmatrix}$$

### Solution

$$A = \begin{bmatrix} 1 & 1 & 2 & 2 \\ 2 & 2 & 4 & 4 \\ 1 & c & 2 & 2 \end{bmatrix} \quad \begin{array}{l} R_2 - 2R_1 \\ R_3 - R_1 \end{array}$$

$$\begin{bmatrix} 1 & 1 & 2 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & c-1 & 0 & 0 \end{bmatrix} \quad \text{Interchange } R_2 \text{ \& } R_1$$

$$\begin{bmatrix} 1 & 1 & 2 & 2 \\ 0 & c-1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

a) If  $c = 1$ , then

$$\begin{pmatrix} 1 & 1 & 2 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

This has only one pivot (first column) and 3 free variables  $x_2, x_3, x_4$ .

$$\text{The nullspace matrix: } \begin{pmatrix} -1 & -2 & -2 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

b) If  $c \neq 1$ , then

$$\begin{pmatrix} 1 & 1 & 2 & 2 \\ 0 & c-1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad \frac{1}{c-1} R_2$$

$$\begin{pmatrix} 1 & 1 & 2 & 2 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad R_2 - R_1$$

$$\begin{pmatrix} 1 & 0 & 2 & 2 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

There are two pivots  $(C_1, C_2)$  and 2 free variables  $x_3, x_4$

The nullspace matrix:  $\begin{pmatrix} -2 & -2 \\ 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}$

---

$$A = \begin{bmatrix} 1-c & 2 \\ 0 & 2-c \end{bmatrix}$$

**a)** If  $c = 1$

$$A = \begin{bmatrix} 0 & 2 \\ 0 & 1 \end{bmatrix} \quad R_1 - 2R_2$$

$$\begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \rightarrow 0a + b = 0 \Rightarrow b = 0$$

This has a single pivot in the second column and one free variable with the nullspace matrix

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

**b)** If  $c = 2$

$$A = \begin{bmatrix} -1 & 2 \\ 0 & 0 \end{bmatrix} \quad -R_1$$

$$\begin{bmatrix} 1 & -2 \\ 0 & 0 \end{bmatrix}$$

$$\begin{pmatrix} 1 & -2 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \rightarrow a - 2b = 0$$

$$\text{if } b = 1 \quad a = 2$$

This has a single pivot in the first column with the nullspace matrix  $\begin{pmatrix} 2 \\ 1 \end{pmatrix}$

**c)** Otherwise  $c \neq 1, 2$

$$A = \begin{bmatrix} 1-c & 2 \\ 0 & 2-c \end{bmatrix} \quad \frac{1}{1-c} R_1$$

$$\begin{bmatrix} 1 & \frac{2}{1-c} \\ 0 & 2-c \end{bmatrix} \quad \frac{1}{2-c} R_2$$

$$\begin{bmatrix} 1 & \frac{2}{1-c} \\ 0 & 1 \end{bmatrix} \quad R_1 - \frac{2}{1-c} R_2$$

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

The result is the identity matrix with 2 pivots, which has  $(2 - 2) = 0$  null space.

### Exercise

If  $A$  has a rank  $r$ , then it has an  $r$  by  $r$  sub-matrix  $S$  that is invertible. Remove  $m - r$  rows and  $n - r$  columns to find an invertible sub-matrix  $S$  inside each  $A$  (you could keep the pivot rows and pivot columns of  $A$ ).

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 4 \end{pmatrix} \quad A = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \end{pmatrix} \quad A = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

### Solution

If a matrix  $A$  has rank  $r$ , then the

***(dimension of the column space) = (dimension of the row space) =  $r$***

For the invertible sub-matrix  $S$ , we need to find  $r$  linearly independent rows and  $r$  linearly independent columns.

For matrix  $A$ :

$$\begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 4 \end{pmatrix} \quad R_2 - R_1$$

$$\begin{pmatrix} 1 & 2 & 3 \\ 0 & 0 & 1 \end{pmatrix} \quad R_1 - 3R_2$$

$$\begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

The 1<sup>st</sup> and 3<sup>rd</sup> columns are linearly independent, and the 1<sup>st</sup> and 2<sup>nd</sup> rows are also linearly independent.

Rank ( $A$ ) = 2.

The sub matrices are:  $S_A = \begin{pmatrix} 1 & 3 \\ 1 & 4 \end{pmatrix}$   $S_A = \begin{pmatrix} 2 & 3 \\ 2 & 4 \end{pmatrix}$   $S_A = \begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix}$

For matrix **B**:

$$\begin{pmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \end{pmatrix} \quad R_2 - 2R_1$$

$$\begin{pmatrix} 1 & 2 & 3 \\ 0 & 0 & 0 \end{pmatrix}$$

Rank (**B**) = 1.

The submatrix is:  $S_A = (1)$

For matrix **C**:

$$C = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Rank (**C**) = 2.

The submatrix is by disregarding (deleting) 1<sup>st</sup> column and 2<sup>nd</sup> row:  $S_A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$

## ***Exercise***

Suppose that column 3 of 4 x 6 matrix is all zero. Then  $x_3$  must be a \_\_\_\_\_ variable. Give one special solution for this matrix.

## **Solution**

The  $x_3$  must be a ***free variable***.

A special solution for this variable can be taken to be.

$$\begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

### Exercise

Fill in the missing numbers to make  $A$  rank 1, rank 2, rank 3. (your solution should be 3 matrices)

$$A = \begin{pmatrix} & -3 & \\ 1 & 3 & -1 \\ & 9 & -3 \end{pmatrix}$$

### Solution

$$A = \begin{pmatrix} a & -3 & b \\ 1 & 3 & -1 \\ c & 9 & -3 \end{pmatrix}$$

If  $\text{rank}(A) = 1$ , then we need the 1<sup>st</sup> and 3<sup>rd</sup> to be multiple of the 2<sup>nd</sup> row to get zero in these rows.

$$A = \begin{pmatrix} a & -3 & b \\ 1 & 3 & -1 \\ c & 9 & -3 \end{pmatrix} \begin{array}{l} R_1 + R_2 \\ R_3 - 3R_2 \end{array}$$

$$\begin{pmatrix} a+1 & 0 & b-1 \\ 1 & 3 & -1 \\ c-3 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 3 & -1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\begin{cases} a+1=0 \\ b-1=0 \\ c-3=0 \end{cases} \rightarrow \begin{cases} a=-1 \\ b=1 \\ c=3 \end{cases}$$

$$A = \begin{pmatrix} -1 & -3 & 1 \\ 1 & 3 & -1 \\ 3 & 9 & -3 \end{pmatrix}$$

If  $\text{rank}(A) = 2$ , then we need the 1<sup>st</sup> **or** 3<sup>rd</sup> to be multiple of the 2<sup>nd</sup> row to get zero row.

$$A = \begin{pmatrix} a & -3 & b \\ 1 & 3 & -1 \\ c & 9 & -3 \end{pmatrix} \begin{array}{l} R_1 + R_2 \\ R_3 - 3R_2 \end{array}$$

$$\begin{pmatrix} a+1 & 0 & b-1 \\ 1 & 3 & -1 \\ c-3 & 0 & 0 \end{pmatrix} \quad \boxed{c \neq 3}$$

$$A = \begin{pmatrix} -1 & -3 & 1 \\ 1 & 3 & -1 \\ 2 & 9 & -3 \end{pmatrix}$$

If  $\text{rank}(A) = 3$  (full rank), then the appropriate to start using 0's or 1's to fill the blank.

$$A = \begin{pmatrix} 0 & -3 & 0 \\ 1 & 3 & -1 \\ 1 & 9 & -3 \end{pmatrix} \quad \text{Interchange } R_1 \text{ \& } R_2$$

$$\begin{pmatrix} 1 & 3 & -1 \\ 0 & -3 & 0 \\ 1 & 9 & -3 \end{pmatrix} \quad R_3 - R_1$$

$$\begin{pmatrix} 1 & 3 & -1 \\ 0 & -3 & 0 \\ 0 & 6 & -2 \end{pmatrix} \quad -\frac{1}{3}R_2$$

$$\begin{pmatrix} 1 & 3 & -1 \\ 0 & 1 & 0 \\ 0 & 6 & -2 \end{pmatrix} \quad \begin{array}{l} R_1 - 3R_2 \\ R_1 - 6R_2 \end{array}$$

$$\begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix} \quad -\frac{1}{2}R_3$$

$$\begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad R_1 + R_3$$

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Hence, it has *rank* 3.

### ***Exercise***

Fill out these matrices so that they have rank 1:

$$A = \begin{pmatrix} 1 & 2 & 4 \\ 2 & & \\ 4 & & \end{pmatrix} \quad B = \begin{pmatrix} 2 & & \\ 1 & & \\ 2 & 6 & -3 \end{pmatrix} \quad M = \begin{pmatrix} a & b \\ c & \end{pmatrix}$$

### **Solution**

Rank = 1 means that all the rows are multiples of each other.



$$A = \begin{pmatrix} 1 & 2 & 4 \\ 2 & a & b \\ 4 & c & d \end{pmatrix} \xrightarrow[R_3=4R_1]{R_2=2R_1} \begin{matrix} a=2(2) & b=2(4) \\ c=4(2) & d=4(4) \end{matrix}$$

$$A = \begin{pmatrix} 1 & 2 & 4 \\ 2 & 4 & 8 \\ 4 & 8 & 16 \end{pmatrix}$$

$$B = \begin{pmatrix} 2 & a & b \\ 1 & c & d \\ 2 & 6 & -3 \end{pmatrix} \xrightarrow[R_2=\frac{1}{2}R_3]{R_1=R_3} \begin{matrix} a=6 & b=-3 \\ c=3 & d=-\frac{3}{2} \end{matrix}$$

$$B = \begin{pmatrix} 2 & 6 & -3 \\ 1 & 3 & -\frac{3}{2} \\ 2 & 6 & -3 \end{pmatrix}$$

$$M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \xrightarrow{R_2=\frac{c}{a}R_1} d = \frac{c}{a}b$$

$$M = \begin{pmatrix} a & b \\ c & \frac{bc}{a} \end{pmatrix}$$

### ***Exercise***

Suppose  $A$  and  $B$  are  $n$  by  $n$  matrices, and  $AB = I$ . Prove from  $\text{rank}(AB) \leq \text{rank}(A)$  that the  $\text{rank}(A) = n$ . So,  $A$  is invertible and  $B$  must be its two-sided inverse. Therefore  $BA = I$  (which is not so obvious!).

### **Solution**

Since  $A$  is  $n$  by  $n \Rightarrow \text{rank}(A) \leq n$

$$n = \text{rank}(I_n) = \text{rank}(AB) \leq \text{rank}(A)$$

### Exercise

Every  $m$  by  $n$  matrix of rank  $r$  reduces to  $(m$  by  $r)$  times  $(r$  by  $n)$ :

$$A = (\text{pivot columns of } A) (\text{first } r \text{ rows of } R) = (COL)(ROW)^T$$

Write the 3 by 4 matrix  $A = \begin{pmatrix} 1 & 1 & 2 & 4 \\ 1 & 2 & 2 & 5 \\ 1 & 3 & 2 & 6 \end{pmatrix}$  as the product of the 3 by 2 from the pivot columns and the 2 by 4 matrix from  $R$ .

### Solution

$$A = \begin{pmatrix} 1 & 1 & 2 & 4 \\ 1 & 2 & 2 & 5 \\ 1 & 3 & 2 & 6 \end{pmatrix} \begin{array}{l} \\ R_2 - R_1 \\ R_3 - R_1 \end{array}$$

$$\begin{pmatrix} 1 & 1 & 2 & 4 \\ 0 & 1 & 0 & 1 \\ 0 & 2 & 0 & 2 \end{pmatrix} \begin{array}{l} \\ \\ R_3 - 2R_2 \end{array}$$

$$\begin{pmatrix} 1 & 1 & 2 & 4 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{array}{l} R_1 - R_2 \\ \\ \end{array}$$

$$\begin{pmatrix} 1 & 0 & 2 & 3 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

The pivots columns are the 1<sup>st</sup> and 2<sup>nd</sup> column.

$$A = (\text{pivot columns of } A) (\text{first } r \text{ rows of } R) = (COL)(ROW)^T$$

$$\begin{aligned} A &= \begin{pmatrix} 1 & 1 & 2 & 4 \\ 1 & 2 & 2 & 5 \\ 1 & 3 & 2 & 6 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} 1 & 0 & 2 & 3 \\ 0 & 1 & 0 & 1 \end{pmatrix} \end{aligned}$$

### Exercise

Suppose  $R$  is  $m$  by  $n$  matrix of rank  $r$ , with pivot columns first:  $\begin{bmatrix} I & F \\ 0 & 0 \end{bmatrix}$

- What are the shapes of those 4 blocks?
- Find the right-inverse  $B$  with  $RB = I$  if  $r = m$ .
- Find the right-inverse  $C$  with  $CR = I$  if  $r = n$ .
- What is the reduced row echelon form of  $R^T$  (with shapes)?
- What is the reduced row echelon form of  $R^T R$  (with shapes)?

Prove that  $R^T R$  has the same nullspace as  $R$ . Then show that  $A^T A$  always has the same nullspace as  $A$  (a value fact).

- Suppose you allow elementary column operations on  $A$  as well as elementary row operations (which get to  $R$ ). What is the “row-and-column reduced form” for an  $m$  by  $n$  matrix of rank  $r$ ?

### Solution

$$a) \begin{bmatrix} I & F \\ 0 & 0 \end{bmatrix} : \begin{bmatrix} r \times r & r \times (n-r) \\ (m-r) \times r & (m-r) \times (n-r) \end{bmatrix}$$

$$b) R = \begin{bmatrix} I & F \end{bmatrix}$$

$$RB = I \Rightarrow \begin{bmatrix} I & F \end{bmatrix} B = I$$

$$\begin{bmatrix} I & F \end{bmatrix} \begin{pmatrix} M \\ N \end{pmatrix} = I$$

$$IM + FN = I$$

$$\Rightarrow \begin{cases} M = I \\ N = 0 \end{cases} \rightarrow F : r \times (n-r)$$

$$B = \begin{bmatrix} I_{r \times r} \\ 0_{(n-r) \times r} \end{bmatrix}$$

$$c) R = \begin{bmatrix} I & 0 \end{bmatrix}$$

$$CR = I \Rightarrow C \begin{bmatrix} I & 0 \end{bmatrix} = I$$

$$C = \begin{bmatrix} I_{r \times r} & 0_{r \times (m-r)} \end{bmatrix}$$

$$d) R^T = \begin{bmatrix} I_{r \times r} & 0_{(m-r) \times r} \\ F_{r \times (n-r)} & 0_{(m-r) \times (n-r)} \end{bmatrix} \quad R_2 - F_{r \times (n-r)} R_1$$

$$rref(R^T) = \begin{bmatrix} I_{r \times r} & 0_{(m-r) \times r} \\ 0_{(n-r) \times r} & 0_{(m-r) \times (n-r)} \end{bmatrix}$$

$$\begin{aligned} e) \quad R^T R &= \begin{bmatrix} I & 0 \\ F & 0 \end{bmatrix} \begin{bmatrix} I & F \\ 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} I & IF \\ FI & F^2 \end{bmatrix} \end{aligned}$$

$FI: r \times (n-r) \quad r \times r$ , the inner is not equal but to make work, we can use the  $F$  transpose.

$$(n-r) \times r \quad r \times r \Rightarrow F^T I = F^T$$

$$\begin{aligned} R^T R &= \begin{bmatrix} I & 0 \\ F & 0 \end{bmatrix} \begin{bmatrix} I & F \\ 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} I & F \\ F & F^2 \end{bmatrix} \end{aligned}$$

$$\begin{bmatrix} I & F \\ F & F^2 \end{bmatrix} \quad R_2 - FR_1$$

$$\begin{bmatrix} I & F \\ 0 & 0 \end{bmatrix}$$

$$rref(R^T R) = \begin{bmatrix} I_{r \times r} & F_{r \times (n-r)} \\ 0_{(m-r) \times r} & 0_{(m-r) \times (n-r)} \end{bmatrix}$$

$$= R \mid$$

So, that  $N(A) = N(rref(A))$  for any matrix  $A$ . So,  $N(A) = N(R^T R)$

**f)** After getting to  $R$  we can use the column operations to get rid of  $F$ .

$$\begin{bmatrix} I_{r \times r} & 0_{r \times (n-r)} \\ 0_{(m-r) \times r} & 0_{(m-r) \times (n-r)} \end{bmatrix}$$

### ***Exercise***

True or False (check addition or give a counterexample)

- a) The symmetric matrices in  $M$  (with  $A^T = A$ ) form a subspace.
- b) The skew-symmetric matrices in  $M$  (with  $A^T = -A$ ) form a subspace.
- c) The un-symmetric matrices in  $M$  (with  $A^T \neq A$ ) form a subspace.
- d) Invertible matrices
- e) Singular matrices

### **Solution**

a) True:  $A^T = A$  and  $B^T = B$  lead to  $(A+B)^T = A^T + B^T = A+B$

b) True:  $A^T = -A$  and  $B^T = -B$  lead to  $(A+B)^T = A^T + B^T = -A-B = -(A+B)$

c) False:  $\begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$

d) False:  $A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  and  $B = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$  are invertible matrices but  $A+B = \begin{pmatrix} 0 & 0 \\ 0 & 2 \end{pmatrix}$  is not invertible.

$\therefore$  The zero matrix is not invertible but any linear subspace should contain the zero matrix. So, invertible matrices do not form a linear subspace.

e) False:  $A = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$  and  $B = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$  are singular matrices

But  $A+B = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = I$  is not singular.

### Exercise

$$\text{Let } A = \begin{pmatrix} 1 & 2 & -2 & 3 & 0 \\ 2 & 4 & -3 & 7 & 0 \\ 3 & 6 & -5 & 10 & -2 \\ 5 & 10 & -9 & 16 & 0 \end{pmatrix}$$

- a) Reduce  $A$  to row-reduced echelon form.
- b) What is the rank of  $A$ ?
- c) What are the pivots?
- d) What are the free variables?
- e) Find the special solutions. What is the nullspace  $N(A)$ ?
- f) Exhibit an  $r \times r$  submatrix of  $A$  which is invertible, where  $r = \text{rank}(A)$ . (An  $r \times r$  submatrix of  $A$  is obtained by keeping  $r$  rows and  $r$  columns of  $A$ )

### Solution

$$a) \quad A = \begin{pmatrix} 1 & 2 & -2 & 3 & 0 \\ 2 & 4 & -3 & 7 & 0 \\ 3 & 6 & -5 & 10 & -2 \\ 5 & 10 & -9 & 16 & 0 \end{pmatrix} \quad \begin{array}{l} R_2 - 2R_1 \\ R_3 - 3R_1 \\ R_4 - 5R_1 \end{array}$$

$$\begin{pmatrix} 1 & 2 & -2 & 3 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 & -2 \\ 0 & 0 & 1 & 1 & 0 \end{pmatrix} \quad \begin{array}{l} R_3 - R_2 \\ R_4 - R_2 \end{array}$$

$$\begin{pmatrix} 1 & 2 & -2 & 3 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & -2 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

- b)  $\text{Rank}(A) = 3$
- c) The pivots are  $x_1, x_3, x_5$
- d) The free variables are  $x_2, x_4$

$$e) \quad \begin{pmatrix} 1 & 2 & -2 & 3 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & -2 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad -\frac{1}{2}R_3$$

$$\begin{pmatrix} 1 & 2 & -2 & 3 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad R_1 + 2R_2$$

$$\begin{pmatrix} 1 & 2 & 0 & 5 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Let  $x = x_2 s_2 + x_4 s_4$

$$Rx = \begin{pmatrix} 1 & 2 & 0 & 5 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\rightarrow \begin{cases} x_1 + 2x_2 + 5x_4 = 0 \\ x_3 + x_4 = 0 \\ x_5 = 0 \end{cases}$$

$$1. \text{ Set } x_2 = 1, \quad x_4 = 0 \rightarrow \begin{cases} x_1 + 2 = 0 \Rightarrow x_1 = -2 \\ x_3 = 0 \\ x_5 = 0 \end{cases}$$

The special solution:  $\vec{s}_2 = (-2, 1, 0, 0, 0)$

$$2. \text{ Set } x_2 = 0, \quad x_4 = 1 \rightarrow \begin{cases} x_1 + 5 = 0 \Rightarrow x_1 = -5 \\ x_3 + 1 = 0 \Rightarrow x_3 = -1 \\ x_5 = 0 \end{cases}$$

The special solution:  $\vec{s}_3 = (-5, 0, -1, 1, 0)$

The nullspace is the set  $\left\{ x_2 \begin{pmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -5 \\ 0 \\ -1 \\ 1 \\ 0 \end{pmatrix} \right\}$

- f) The pivot rows and columns must be included in a submatrix. To do that, just take the rows and columns of  $A$  containing pivots, which are columns 1, 3, 5 and rows 1, 2, 3. That will give us a 3 by 3 submatrix. Therefore, this submatrix of  $A$  will be invertible.

$$A = \begin{pmatrix} 1 & -2 & 0 \\ 2 & -3 & 0 \\ 3 & -5 & -2 \end{pmatrix}$$

### Exercise

Let  $A = \begin{pmatrix} -1 & 2 & 5 & 0 & 5 \\ 2 & 1 & 0 & 0 & -15 \\ 6 & -1 & -8 & -1 & -47 \\ 0 & 2 & 4 & 3 & 16 \end{pmatrix}$

- Reduce  $A$  to (ordinary) echelon form.
- What the pivots?
- What are the free variables?
- Reduce  $A$  to row-reduced echelon form.
- Find the special solutions. What is the nullspace  $N(A)$ ?
- What is the rank of  $A$ ?

g) Give the complete solution to  $Ax = b$ , where  $b = A \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$

### Solution

a)  $A = \begin{pmatrix} -1 & 2 & 5 & 0 & 5 \\ 2 & 1 & 0 & 0 & -15 \\ 6 & -1 & -8 & -1 & -47 \\ 0 & 2 & 4 & 3 & 16 \end{pmatrix} \quad \begin{matrix} R_2 + 2R_1 \\ R_3 + 6R_1 \end{matrix}$

$$\begin{pmatrix} -1 & 2 & 5 & 0 & 5 \\ 0 & 5 & 10 & 0 & -5 \\ 0 & 11 & 22 & -1 & -17 \\ 0 & 2 & 4 & 3 & 16 \end{pmatrix} \quad \begin{matrix} 5R_3 - 11R_2 \\ 5R_4 - 2R_2 \end{matrix}$$

$$\begin{pmatrix} -1 & 2 & 5 & 0 & 5 \\ 0 & 5 & 10 & 0 & -5 \\ 0 & 0 & 0 & -5 & -30 \\ 0 & 0 & 0 & 15 & 90 \end{pmatrix} \quad R_4 + 3R_3$$



$$\begin{pmatrix} \boxed{-1} & 2 & 5 & 0 & 5 \\ 0 & \boxed{5} & 10 & 0 & -5 \\ 0 & 0 & 0 & \boxed{-5} & -30 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

**b)** The pivots are  $-1$ ,  $5$ , and  $-5$  (Columns 1, 2, 4)

**c)** The free variables are 3<sup>rd</sup> and 5<sup>th</sup> ( $x_3, x_5$ )

$$d) \begin{pmatrix} -1 & 2 & 5 & 0 & 5 \\ 0 & 5 & 10 & 0 & -5 \\ 0 & 0 & 0 & -5 & -30 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \begin{array}{l} -R_1 \\ \frac{1}{5}R_2 \\ -\frac{1}{5}R_3 \end{array}$$

$$\begin{pmatrix} 1 & -2 & -5 & 0 & -5 \\ 0 & 1 & 2 & 0 & -1 \\ 0 & 0 & 0 & 1 & 6 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad R_1 + 2R_2$$

$$R = \begin{pmatrix} 1 & 0 & -1 & 0 & -7 \\ 0 & 1 & 2 & 0 & -1 \\ 0 & 0 & 0 & 1 & 6 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

**e)** Let  $x = x_3 s_1 + x_5 s_2$

$$R\vec{x} = \begin{pmatrix} 1 & 0 & -1 & 0 & -7 \\ 0 & 1 & 2 & 0 & -1 \\ 0 & 0 & 0 & 1 & 6 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\rightarrow \begin{cases} x_1 - x_3 - 7x_5 = 0 \\ x_2 + 2x_3 - x_5 = 0 \\ x_4 + 6x_5 = 0 \end{cases}$$

**1.** Set  $x_3 = 1, \quad x_5 = 0$

$$\rightarrow \begin{cases} x_1 - 1 = 0 & \underline{x_1 = 1} \\ x_2 + 2 = 0 & \underline{x_2 = -2} \\ \underline{x_4 = 0} \end{cases}$$

The special solution:  $\vec{s}_1 = (1, -2, 1, 0, 0)$

2. Set  $x_3 = 0, \quad x_5 = 1$

$$\rightarrow \begin{cases} x_1 - 7 = 0 & \underline{x_1 = 7} \\ x_2 - 1 = 0 & \underline{x_2 = 1} \\ x_4 + 6 = 0 & \underline{x_4 = -6} \end{cases}$$

The special solution:  $\vec{s}_2 = (7, 1, 0, -6, 1)$

The nullspace is the set  $\left\{ x_3 \begin{pmatrix} 1 \\ -2 \\ 1 \\ 0 \\ 0 \end{pmatrix} + x_5 \begin{pmatrix} 7 \\ 1 \\ 0 \\ -6 \\ 1 \end{pmatrix} \right\}$

f)  $\text{Rank}(A) = 3$

g)  $A\vec{x} = \vec{b}$ , where  $\vec{b} = A \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$

The complete solution = (the particular solution) + (special solution)

$$\vec{x} = \vec{x}_p + \vec{x}_n$$

$$\vec{x} = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} + x_3 \begin{pmatrix} 1 \\ -2 \\ 1 \\ 0 \\ 0 \end{pmatrix} + x_5 \begin{pmatrix} 7 \\ 1 \\ 0 \\ -6 \\ 1 \end{pmatrix}$$

### Exercise

$$\text{Let } A = \begin{pmatrix} 1 & 2 & 0 & 3 & 0 \\ 1 & 2 & -1 & -1 & 0 \\ 0 & 0 & 1 & 4 & 0 \\ 2 & 4 & 1 & 10 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

- a) Reduce  $A$  to row-reduced echelon form.
- b) What is the rank of  $A$ ?
- c) What the pivots variables?
- d) What are the free variables?
- e) Find the special solutions.
- f) What is the nullspace  $N(A)$ ?

### Solution

$$a) \begin{pmatrix} 1 & 2 & 0 & 3 & 0 \\ 1 & 2 & -1 & -1 & 0 \\ 0 & 0 & 1 & 4 & 0 \\ 2 & 4 & 1 & 10 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} \begin{array}{l} \\ R_2 - R_1 \\ \\ R_3 - 2R_1 \\ \end{array}$$

$$\begin{pmatrix} 1 & 2 & 0 & 3 & 0 \\ 0 & 0 & -1 & -4 & 0 \\ 0 & 0 & 1 & 4 & 0 \\ 0 & 0 & 1 & 4 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} \begin{array}{l} \\ -R_2 \\ R_3 - R_2 \\ R_4 - 2R_2 \\ \end{array}$$

$$\begin{pmatrix} 1 & 2 & 0 & 3 & 0 \\ 0 & 0 & 1 & 4 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} R_5 - R_4$$

$$R = \begin{pmatrix} 1 & 2 & 0 & 3 & 0 \\ 0 & 0 & 1 & 4 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \rightarrow \begin{array}{l} x_1 = -2x_2 - 3x_4 \\ x_3 = -4x_4 \\ x_5 = 0 \end{array}$$

- b)  $\text{Rank}(A) = 3$
- c) The pivots variables are:  $x_1, x_3, x_5$

d) The free variables are:  $x_2, x_4$

e) Let  $x = x_2 s_1 + x_4 s_2$

$$\begin{cases} x_1 = -2x_2 - 3x_4 \\ x_3 = -4x_4 \end{cases}$$

Set  $x_2 = 1, x_4 = 0$

The special solution:  $\vec{s}_1 = (-2, 1, 0, 0, 0)$

Set  $x_2 = 0, x_4 = 1$  ;

The special solution:  $\vec{s}_2 = (-3, 0, -4, 1, 0)$

f) The nullspace is the set  $\left\{ x_2 \begin{pmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -3 \\ 0 \\ -4 \\ 1 \\ 0 \end{pmatrix} \right\}$

$$N(A) = \begin{pmatrix} -2 & -3 \\ 1 & 0 \\ 0 & -4 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$$

### Exercise

$$\text{Let } A = \begin{pmatrix} 3 & 21 & 0 & 9 & 0 \\ 1 & 7 & -1 & -2 & -1 \\ 2 & 14 & 0 & 6 & 1 \\ 6 & 42 & -1 & 13 & 0 \end{pmatrix}$$

- Reduce  $A$  to row-reduced echelon form.
- What is the rank of  $A$ ?
- What the pivots?
- What are the free variables?
- Find the special solutions.
- What is the nullspace  $N(A)$ ?

### Solution

$$a) \begin{pmatrix} 3 & 21 & 0 & 9 & 0 \\ 1 & 7 & -1 & -2 & -1 \\ 2 & 14 & 0 & 6 & 1 \\ 6 & 42 & -1 & 13 & 0 \end{pmatrix} \begin{array}{l} \\ 3R_2 - R_1 \\ 3R_3 - 2R_1 \\ R_4 - 2R_1 \end{array}$$

$$\begin{pmatrix} 3 & 21 & 0 & 9 & 0 \\ 0 & 0 & -3 & -15 & -3 \\ 0 & 0 & 0 & 0 & 3 \\ 0 & 0 & -1 & -5 & 0 \end{pmatrix} \begin{array}{l} \\ -R_2 \\ \\ 3R_4 + R_2 \end{array}$$

$$\begin{pmatrix} 3 & 21 & 0 & 9 & 0 \\ 0 & 0 & 3 & 15 & 3 \\ 0 & 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 & 3 \end{pmatrix} \begin{array}{l} \frac{1}{3}R_1 \\ \frac{1}{3}R_2 \\ \frac{1}{3}R_3 \\ R_4 - R_3 \end{array}$$

$$\begin{pmatrix} 1 & 7 & 0 & 3 & 0 \\ 0 & 0 & 1 & 5 & 1 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \begin{array}{l} \\ R_2 - R_3 \\ \\ \end{array}$$

$$\begin{pmatrix} 1 & 7 & 0 & 3 & 0 \\ 0 & 0 & 1 & 5 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \begin{array}{l} x_1 = -7x_2 - 3x_4 \\ x_3 = -5x_4 \\ x_5 = 0 \end{array}$$

**b)** Rank( $A$ ) = 3

**c)** The pivots variables are:  $x_1, x_3, x_5$

**d)** The free variables are:  $x_2, x_4$

**e)** Let  $x = x_2 s_1 + x_4 s_2$

$$\begin{cases} x_1 = -7x_2 - 3x_4 \\ x_3 = -5x_4 \end{cases}$$

Set  $x_2 = 1, x_4 = 0$

The special solution:  $\vec{s}_1 = (-7, 1, 0, 0, 0)$

Set  $x_2 = 0, x_4 = 1$  ;

The special solution:  $\vec{s}_2 = (-3, 0, -5, 1, 0)$

f) The nullspace is the set  $\left\{ x_2 \begin{pmatrix} -7 \\ 1 \\ 0 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -3 \\ 0 \\ -5 \\ 1 \\ 0 \end{pmatrix} \right\}$

$$N(A) = \begin{pmatrix} -7 & -3 \\ 1 & 0 \\ 0 & -5 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$$

### Exercise

The 3 by 3 matrix  $A$  has rank 2.

$$A\vec{x} = \vec{b} \quad \text{is} \quad \begin{aligned} x_1 + 2x_2 + 3x_3 + 5x_4 &= b_1 \\ 2x_1 + 4x_2 + 8x_3 + 12x_4 &= b_2 \\ 3x_1 + 6x_2 + 7x_3 + 13x_4 &= b_3 \end{aligned}$$

- Reduce  $\begin{bmatrix} A & \vec{b} \end{bmatrix}$  to  $\begin{bmatrix} U & \vec{c} \end{bmatrix}$ , so that  $A\vec{x} = \vec{b}$  becomes triangular system  $U\vec{x} = \vec{c}$ .
- Find the condition on  $(b_1, b_2, b_3)$  for  $A\vec{x} = \vec{b}$  to have a solution
- Describe the column space of  $A$ . Which plane in  $\mathbb{R}^3$ ?
- Describe the nullspace of  $A$ . Which special solutions in  $\mathbb{R}^4$ ?
- Find a particular solution to  $A\vec{x} = (0, 6, -6)$  and then complete solution.

### Solution

$$a) \left[ \begin{array}{cccc|c} 1 & 2 & 3 & 5 & b_1 \\ 2 & 4 & 8 & 12 & b_2 \\ 3 & 6 & 7 & 13 & b_3 \end{array} \right] \quad \begin{array}{l} R_2 - 2R_1 \\ R_3 - 3R_1 \end{array}$$

$$\left[ \begin{array}{cccc|c} 1 & 2 & 3 & 5 & b_1 \\ 0 & 0 & 2 & 2 & b_2 - 2b_1 \\ 0 & 0 & -2 & -2 & b_3 - 3b_1 \end{array} \right] \quad R_3 + R_2$$

$$\left[ \begin{array}{cccc|c} 1 & 2 & 3 & 5 & b_1 \\ 0 & 0 & 2 & 2 & b_2 - 2b_1 \\ 0 & 0 & 0 & 0 & b_3 + b_2 - 5b_1 \end{array} \right]$$

**b)** The last equation  $b_3 + b_2 - 5b_1 = 0$  shows the solvability condition.

**c) (i)** The column space is the plane containing all combinations of the pivot columns: 1st (1, 2, 3) and 3<sup>rd</sup> (3, 8, 7).

**(ii)** The column space contains all vectors with  $b_3 + b_2 - 5b_1 = 0$ . That makes  $A\vec{x} = \vec{b}$

solvable, so  $\vec{b}$  is in the column space. All columns of  $A$  pass this test  $b_3 + b_2 - 5b_1 = 0$ . This is the equation for the plane in **(i)**.

**d)** The special solutions have free variables:

$$\begin{cases} x_1 + 2x_2 + 3x_3 + 5x_4 = 0 \\ 2x_3 + 2x_4 = 0 \end{cases}$$

$$\Rightarrow \begin{cases} x_1 = -2x_2 - 2x_4 \\ x_3 = -x_4 \end{cases}$$

Let  $x_2 = 1, x_4 = 0$

$$\Rightarrow \begin{cases} x_1 = -2 \\ x_3 = 0 \end{cases}$$

$$\vec{s}_1 = \begin{pmatrix} -2 \\ 1 \\ 0 \\ 0 \end{pmatrix}$$

Let  $x_2 = 0, x_4 = 1$

$$\Rightarrow \begin{cases} x_1 = -2 \\ x_3 = -1 \end{cases}$$

$$\vec{s}_2 = \begin{pmatrix} -2 \\ 0 \\ -1 \\ 1 \end{pmatrix}$$

The nullspace  $N(A)$  in  $\mathbb{R}^4$  contains all

$$\vec{x}_n = x_2 \begin{pmatrix} -2 \\ 1 \\ 0 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -2 \\ 0 \\ -1 \\ 1 \end{pmatrix}$$

**e)** One particular solution  $x_p$  has free variables = zero.

$$\begin{cases} x_1 + 2x_2 + 3x_3 + 5x_4 = 0 \\ 2x_3 + 2x_4 = 6 \end{cases}$$

$$\begin{cases} x_1 = -2x_2 - 3x_3 - 5x_4 \\ x_3 = 3 - x_4 \end{cases}$$

$$\begin{cases} x_1 = -2x_2 - 9 - 2x_4 \\ x_3 = 3 - x_4 \end{cases}$$

Let  $x_2 = x_4 = 0$

$$\begin{cases} x_1 = -9 \\ x_3 = 3 \end{cases}$$

$$\vec{x}_p = \begin{pmatrix} -9 \\ 0 \\ -3 \\ 0 \end{pmatrix}$$

The complete solution to  $A\vec{x} = (0, 6, -6)$  is  $\vec{x} = \vec{x}_p + \text{all } \vec{x}_n$

$$\vec{x} = \begin{pmatrix} -9 \\ 0 \\ -3 \\ 0 \end{pmatrix} + x_2 \begin{pmatrix} -2 \\ 1 \\ 0 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -2 \\ 0 \\ -1 \\ 1 \end{pmatrix}$$

### Exercise

Find the special solutions and describe the complete solution to  $A\vec{x} = \vec{0}$  for

$$A_1 = 3 \text{ by } 4 \text{ zero matrix} \quad A_2 = \begin{pmatrix} 3 & 6 \\ 1 & 2 \end{pmatrix} \quad A_3 = \begin{bmatrix} A_1 & A_2 \end{bmatrix}$$

Which are the pivot columns? Which are the free variables? What is the  $R$  (Reduced Row Echelon matrix) in each case?

### Solution

$A_1 \vec{x} = \vec{0}$  has 4 solutions. They are the columns  $\vec{s}_1, \vec{s}_2, \vec{s}_3, \vec{s}_4$  of the identity matrix (4 by 4).

The Nullspace is of  $\mathbb{R}^4$ .

The complete solution:  $\vec{x} = c_1 \vec{s}_1 + c_2 \vec{s}_2 + c_3 \vec{s}_3 + c_4 \vec{s}_4$  in  $\mathbb{R}^4$ .



There are no pivot columns; all variables are free; the reduced  $R$  is the same zero matrix as  $A_1$ .

$$A_2 \vec{x} = \vec{0}$$

$$A_2 \vec{x} = \begin{pmatrix} 3 & 6 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \rightarrow x_1 + 2x_2 = 0$$

The vector solution:  $\vec{s} = (-2, 1)$ , The first column of  $A_2$  is its pivot column, and  $x_2$  is the free variable.

$$\begin{pmatrix} 3 & 6 \\ 1 & 2 \end{pmatrix} \quad \frac{1}{3}R_1$$

$$\begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix} \quad R_2 - R_1$$

$$\begin{pmatrix} 1 & 2 \\ 0 & 0 \end{pmatrix}$$

$$R_3 = \begin{bmatrix} 1 & 2 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

All variables are free. There are three special solutions to  $A_3 \vec{x} = 0$

$$\vec{s}_1 = \begin{pmatrix} -2 \\ 1 \\ 0 \\ 0 \end{pmatrix} \quad \vec{s}_2 = \begin{pmatrix} -1 \\ 0 \\ 1 \\ 0 \end{pmatrix} \quad \vec{s}_3 = \begin{pmatrix} -2 \\ 0 \\ 0 \\ 1 \end{pmatrix}$$

The complete solution:

$$\vec{x} = c_1 \vec{s}_1 + c_2 \vec{s}_2 + c_3 \vec{s}_3$$

$$= c_1 \begin{pmatrix} -2 \\ 1 \\ 0 \\ 0 \end{pmatrix} + c_2 \begin{pmatrix} -1 \\ 0 \\ 1 \\ 0 \end{pmatrix} + c_3 \begin{pmatrix} -2 \\ 0 \\ 0 \\ 1 \end{pmatrix}$$

### Exercise

Create a 3 by 4 matrix whose special solutions to  $A\vec{x} = \vec{0}$  are  $\vec{s}_1$  and  $\vec{s}_2$  :

$$\vec{s}_1 = \begin{pmatrix} -3 \\ 2 \\ 0 \\ 0 \end{pmatrix} \quad \text{and} \quad \vec{s}_2 = \begin{pmatrix} -2 \\ 0 \\ -6 \\ 1 \end{pmatrix}$$

You could create the matrix  $A$  in row reduced form  $R$ . Then describe all possible matrices  $A$  with the required Nullspace  $N(A) = \text{all combinations of } \vec{s}_1 \text{ and } \vec{s}_2$ .

### Solution

We can write the solution:

$$\vec{x} = x_2 \vec{s}_1 + x_4 \vec{s}_2$$

$$x_2 \begin{pmatrix} -3 \\ 2 \\ 0 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -2 \\ 0 \\ -6 \\ 1 \end{pmatrix} = \begin{pmatrix} -3x_2 - 2x_4 \\ 2x_2 \\ -6x_4 \\ x_4 \end{pmatrix}$$

$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} -3x_2 - 2x_4 \\ 2x_2 \\ -6x_4 \\ x_4 \end{pmatrix} \rightarrow \begin{cases} x_1 = -3x_2 - 2x_4 \\ x_3 = -6x_4 \end{cases}$$

$$\rightarrow \begin{cases} x_1 + 3x_2 + 2x_4 = 0 \\ x_3 + 6x_4 = 0 \end{cases}$$

$$\Rightarrow \begin{pmatrix} 1 & 3 & 0 & 2 \\ 0 & 0 & 1 & 6 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

The entries 3, 2, 6 are the negatives of  $-3$ ,  $-2$ ,  $-6$  in the special solutions.

Every 3 by 4 matrix has at least one special solution. These  $A$ 's have two.

### Exercise

The plane  $x - 3y - z = 12$  is parallel to the plane  $x - 3y - z = 0$ . One particular point on this plane is  $(12, 0, 0)$ . All points on the plane have the form (fill the first components)

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} + y \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} + z \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

### Solution

$$x - 3y - z = 12$$

$$x = 3y + z + 12$$

$$\begin{aligned} \begin{bmatrix} x \\ y \\ z \end{bmatrix} &= \begin{pmatrix} 12 + 3y + z \\ y \\ z \end{pmatrix} \\ &= \begin{bmatrix} 12 \\ 0 \\ 0 \end{bmatrix} + y \begin{bmatrix} 3 \\ 1 \\ 0 \end{bmatrix} + z \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \end{aligned}$$

### Exercise

Construct a matrix whose column space contains  $(1, 1, 5)$  and  $(0, 3, 1)$  and whose Nullspace contains  $(1, 1, 2)$ .

### Solution

$$A = \begin{pmatrix} 1 & 0 & a \\ 1 & 3 & b \\ 5 & 1 & c \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 & a \\ 1 & 3 & b \\ 5 & 1 & c \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 1 + 2a \\ 1 + 3 + 2b \\ 5 + 1 + 2c \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\begin{cases} 1 + 2a = 0 & \rightarrow 2a = -1 & \Rightarrow a = -\frac{1}{2} \\ 4 + 2b = 0 & \rightarrow 2b = -4 & \Rightarrow b = -2 \\ 6 + 2c = 0 & \rightarrow 2c = -6 & \Rightarrow c = -3 \end{cases}$$

$$A = \begin{pmatrix} 1 & 0 & -\frac{1}{2} \\ 1 & 3 & -2 \\ 5 & 1 & -3 \end{pmatrix}$$

### Exercise

Construct a matrix whose column space contains  $(1, 1, 0)$  and  $(0, 1, 1)$  and whose Nullspace contains  $(1, 0, 1)$  and  $(0, 0, 1)$ .

### Solution

It is impossible. Matrix  $A$  must be 3 by 3.

Since the nullspace is supposed to contain two independent vectors,  $A$  can have at most  $3 - 2 = 1$  pivots.

Since the column space supposes to contain two independent vectors,  $A$  must have at least 2 pivots.

These conditions can't both be met.

### Exercise

Construct a matrix whose column space contains  $(1, 1, 1)$  and whose Nullspace contains  $(1, 1, 1, 1)$ .

### Solution

The matrix needs to be 3 by 4 matrix.

$$\begin{pmatrix} 1 & a & b & c \\ 1 & d & e & f \\ 1 & g & h & i \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\begin{cases} 1 + a + b + c = 0 \\ 1 + d + e + f = 0 \\ 1 + g + h + i = 0 \end{cases}$$

$$\begin{cases} a + b + c = -1 \\ d + e + f = -1 \\ g + h + i = -1 \end{cases}$$

$$\begin{cases} \text{if } b = c = 0 & a = -1 \\ \text{if } d = f = 0 & e = -1 \\ \text{if } g = h = 0 & i = -1 \end{cases}$$

$$\begin{pmatrix} 1 & -1 & 0 & 0 \\ 1 & 0 & -1 & 0 \\ 1 & 0 & 0 & -1 \end{pmatrix}$$

### Exercise

How is the Nullspace  $N(C)$  related to the spaces  $N(A)$  and  $N(B)$ , if  $C = \begin{bmatrix} A \\ B \end{bmatrix}$ ?

### Solution

$$Cx = \begin{bmatrix} Ax \\ Bx \end{bmatrix} = 0 = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

If and only if  $A\vec{x} = \vec{0}$  and  $B\vec{x} = \vec{0}$

$$N(C) = N(A) \cap N(B)$$

### Exercise

Why does no 3 by 3 matrix have a nullspace that equals its column space?

### Solution

If nullspace = column space, then  $n - r = r$  (there are  $r$  pivots).

For  $n = 3 \Rightarrow 3 = 2r$  is impossible.

### Exercise

If  $AB = 0$  then the column space  $B$  is contained in the \_\_\_\_\_ of  $A$ . Give an example of  $A$  and  $B$ .

### Solution

If  $AB = 0$  then the column space  $B$  is contained in the **nullspace** of  $A$ .

$$\text{Example: } A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \quad B = \begin{bmatrix} 1 & 1 \\ -1 & -1 \end{bmatrix}$$

### Exercise

True or false (with reason if true or example to show it is false)

- a) A square matrix has no free variables.
- b) An invertible matrix has no free variables.
- c) An  $m$  by  $n$  matrix has no more than  $n$  pivot variables.
- d) An  $m$  by  $n$  matrix has no more than  $m$  pivot variables.

### Solution

- a) False. Any matrix with fewer than full number of pivots will.  $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$

- b) True. Since it is invertible, we will get the full number of pivots. The nullspace has dimension, so we have 0 free variables.
- c) True, the number of pivot variables is the dimension of the nullspace, which is at most the number of columns. The nullspace dimension + column space dimension = number of columns.
- d) True, in reduced echelon matrix the pivot columns are all 0 except for a single 1, and there are only up to  $m$  vectors of this type.

### Exercise

Suppose an  $m$  by  $n$  matrix has  $r$  pivots. The number of special solutions is \_\_\_\_\_.

The Nullspace contains only  $x = 0$  when  $r =$  \_\_\_\_\_.

The column space is all of  $\mathbf{R}^m$  when  $r =$  \_\_\_\_\_.

### Solution

Suppose an  $m$  by  $n$  matrix has  $r$  pivots. The number of special solutions is  $\mathbf{n - r}$ .

The Nullspace contains only  $x = 0$  when  $r = \mathbf{n}$ .

The column space is all of  $\mathbf{R}^m$  when  $r = \mathbf{m}$ .

### Exercise

Find the complete solution in the form  $x_p + x_n$  to these full rank system:

$$\begin{array}{ll} a) & x + y + z = 4 \\ b) & \begin{array}{l} x + y + z = 4 \\ x - y + z = 4 \end{array} \end{array}$$

### Solution

a)  $x + y + z = 4$

The equivalent matrix is given by:  $\begin{cases} Ax = 4 \\ A = \begin{pmatrix} 1 & 1 & 1 \end{pmatrix} \end{cases}$

The complete solution in the form  $\vec{x} = \vec{x}_p + \vec{x}_n$

$\vec{x}_n$  is the homogeneous solution to  $A\vec{x}_n = \vec{0}$

Size of  $A$  is  $m = 1$  and  $n = 3$ ,  $\text{rank}(A) = r = 1$

$$A\vec{x}_n = \vec{0}$$

$$\begin{pmatrix} 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$x_1 + x_2 + x_3 = 0 \Rightarrow \underline{x_1 = -x_2 - x_3}$$

Set  $x_2 = 1, \quad x_3 = 0$

The special solution:  $\vec{s}_1 = (-1, 1, 0)$

Set  $x_2 = 0, \quad x_3 = 1$

The special solution:  $\vec{s}_2 = (-1, 0, 1)$

The nullspace is the set  $\left\{ x_2 \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} + x_3 \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} \right\}$

$$x = 4 - y - z$$

$$\Rightarrow x_1 = 4 - x_2 - x_3$$

Set  $x_2 = 0, \quad x_3 = 0$  that implies to the particular solution:  $\vec{x}_p = \begin{pmatrix} 4 \\ 0 \\ 0 \end{pmatrix}$

The complete solution in the form  $\vec{x} = \begin{pmatrix} 4 \\ 0 \\ 0 \end{pmatrix} + x_2 \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} + x_3 \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$

Note: that the null space of A is spanned by the two linearly independent vectors

$$(-1, 1, 0)^T \quad \text{and} \quad (-1, 0, 1)^T$$

**b)** 
$$\begin{cases} x + y + z = 4 \\ x - y + z = 4 \end{cases}$$

The equivalent matrix is given by:

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 1 & -1 & 1 \end{pmatrix} \text{ and } A \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

$$\left[ \begin{array}{ccc|c} 1 & 1 & 1 & 4 \\ 1 & -1 & 1 & 4 \end{array} \right] \quad R_2 - R_1$$

$$\left[ \begin{array}{ccc|c} 1 & 1 & 1 & 4 \\ 0 & -2 & 0 & 0 \end{array} \right] \quad -\frac{1}{2}R_2$$

$$\left[ \begin{array}{ccc|c} 1 & 1 & 1 & 4 \\ 0 & 1 & 0 & 0 \end{array} \right] \quad R_1 - R_2$$

$$\left[ \begin{array}{ccc|c} 1 & 0 & 1 & 4 \\ 0 & 1 & 0 & 0 \end{array} \right]$$

The pivots are  $x_1, x_2$  ; The free variable is  $x_3$

Rank  $r = 2$ ,  $n = 2$ ,  $m = 3$ .

The nullspace has dimension  $m - r = 1$ .

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\Rightarrow \begin{cases} x_1 + x_3 = 0 \rightarrow x_1 = -x_3 \\ x_2 = 0 \end{cases}$$

$$\text{If } x_3 = 1 \Rightarrow x_1 = -1$$

The special solution:  $\vec{s}_1 = (-1, 0, 1)$

$$\text{The nullspace is the set } \left\{ x_3 \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} \right\}$$

Set  $x_3 = 0$  that implies

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 4 \\ 0 \end{pmatrix}$$

$$\Rightarrow \begin{cases} x_1 = 4 \\ x_2 = 0 \end{cases}$$

$$\text{Then the particular solution: } \vec{x}_p = \begin{pmatrix} 4 \\ 0 \\ 0 \end{pmatrix}$$

The complete solution in the form:

$$\vec{x} = \begin{pmatrix} 4 \\ 0 \\ 0 \end{pmatrix} + x_3 \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$$



### Exercise

Find the complete solution in the form  $\vec{x}_p + \vec{x}_n$  to the system:

$$\begin{pmatrix} 1 & 3 & 1 & 2 \\ 2 & 6 & 4 & 8 \\ 0 & 0 & 2 & 4 \end{pmatrix} \vec{x} = \begin{pmatrix} 1 \\ 3 \\ 1 \end{pmatrix}$$

### Solution

$$\left( \begin{array}{cccc|c} 1 & 3 & 1 & 2 & 1 \\ 2 & 6 & 4 & 8 & 3 \\ 0 & 0 & 2 & 4 & 1 \end{array} \right) \quad R_2 - 2R_1$$

$$\left( \begin{array}{cccc|c} 1 & 3 & 1 & 2 & 1 \\ 0 & 0 & 2 & 4 & 1 \\ 0 & 0 & 2 & 4 & 1 \end{array} \right) \quad \begin{array}{l} 2R_1 - R_2 \\ R_3 - R_2 \end{array}$$

$$\left( \begin{array}{cccc|c} 2 & 6 & 0 & 0 & 1 \\ 0 & 0 & 2 & 4 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right) \quad \begin{array}{l} \frac{1}{2}R_1 \\ \frac{1}{2}R_2 \end{array}$$

$$\left( \begin{array}{cccc|c} 1 & 3 & 0 & 0 & \frac{1}{2} \\ 0 & 0 & 1 & 2 & \frac{1}{2} \\ 0 & 0 & 0 & 0 & 0 \end{array} \right)$$

The pivots are  $x_1, x_3$ ; The free variables are  $x_2, x_4$

$$\begin{pmatrix} 1 & 3 & 0 & 0 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\begin{cases} x_1 + 3x_2 = 0 & \underline{x_1 = -3x_2} \\ x_3 + 2x_4 = 0 & \underline{x_3 = -2x_4} \end{cases}$$

1. Set  $x_2 = 1, x_4 = 0$

The special solution:  $\vec{s}_1 = (-3, 1, 0, 0)$

2. Set  $x_2 = 0, x_4 = 1$

The special solution:  $\vec{s}_2 = (0, 0, -2, 1)$

The special solution:  $\vec{x}_n = x_2 \begin{pmatrix} -3 \\ 1 \\ 0 \\ 0 \end{pmatrix} + x_3 \begin{pmatrix} 0 \\ 0 \\ -2 \\ 1 \end{pmatrix}$

$$\begin{pmatrix} 1 & 3 & 0 & 0 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} = \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \\ 0 \end{pmatrix}$$

$$\begin{cases} x_1 + 3(0) = \frac{1}{2} & \underline{x_1 = \frac{1}{2}} \\ x_3 + 2(0) = \frac{1}{2} & \underline{x_3 = \frac{1}{2}} \end{cases}$$

Then the particular solution:

$$\vec{x}_p = \begin{pmatrix} \frac{1}{2} \\ 0 \\ \frac{1}{2} \\ 0 \end{pmatrix}$$

The complete solution in the form:

$$\vec{x} = \begin{pmatrix} \frac{1}{2} \\ 0 \\ \frac{1}{2} \\ 0 \end{pmatrix} + x_2 \begin{pmatrix} -3 \\ 1 \\ 0 \\ 0 \end{pmatrix} + x_3 \begin{pmatrix} 0 \\ 0 \\ -2 \\ 1 \end{pmatrix}$$

### Exercise

If  $A$  is  $3 \times 7$  matrix, its largest possible rank is \_\_\_\_\_. In this case, there is a pivot in every \_\_\_\_\_ of  $U$  and  $R$ , the solution to  $Ax = b$  \_\_\_\_\_ (always exists or is unique), and the column space of  $A$  is \_\_\_\_\_. Construct an example of such a matrix  $A$ .

### Solution

If  $A$  is  $3 \times 7$  matrix, its largest possible rank is **3**. In this case, there is a pivot in every **row** of  $U$  and  $R$ , the solution to  $Ax = b$  **always exists**, and the column space of  $A$  is  $\mathbb{R}^3$ .

$$A = \begin{pmatrix} * & * & * & * & * & * & * \\ * & * & * & * & * & * & * \\ * & * & * & * & * & * & * \end{pmatrix}$$

$\text{rank}(A) \leq 3$ , that implies that you have 3 pivots (1 each row)

$$A = \begin{pmatrix} 1 & 0 & 0 & * & * & * & * \\ 0 & 1 & 0 & * & * & * & * \\ 0 & 0 & 1 & * & * & * & * \end{pmatrix}$$
$$= \begin{pmatrix} 1 & 0 & 0 & 1 & 2 & 3 & 4 \\ 0 & 1 & 0 & 5 & 6 & 7 & 8 \\ 0 & 0 & 1 & 9 & 10 & 11 & 12 \end{pmatrix}$$

### Exercise

If  $A$  is  $6 \times 3$  matrix, its largest possible rank is \_\_\_\_\_. In this case, there is a pivot in every \_\_\_\_\_ of  $U$  and  $R$ , the solution to  $A\vec{x} = \vec{b}$  \_\_\_\_\_ (always exists or is unique), and the nullspace of  $A$  is \_\_\_\_\_. Construct an example of such a matrix  $A$ .

### Solution

If  $A$  is  $6 \times 3$  matrix, its largest possible rank is **3**. In this case, there is a pivot in every **column** of  $U$  and  $R$ , the solution to  $A\vec{x} = \vec{b}$  **is unique**, and the column space of  $A$  is  $\{\vec{0}\}$ .

$$A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}$$

### Exercise

Find the rank of  $A$ ,  $A^T A$  and  $AA^T$  for  $A = \begin{pmatrix} 1 & 1 \\ 0 & 2 \\ -1 & 2 \end{pmatrix}$

### Solution

$$A = \begin{pmatrix} 1 & 1 \\ 0 & 2 \\ -1 & 2 \end{pmatrix} \quad R_3 + R_1$$

$$\begin{pmatrix} 1 & 1 \\ 0 & 2 \\ 0 & 3 \end{pmatrix} \quad \frac{1}{2}R_2$$

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 0 & 3 \end{pmatrix} \quad R_3 - 3R_2$$

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$$

$$\underline{\text{rank}(A) = 2}$$

$$\begin{aligned} A^T A &= \begin{pmatrix} 1 & 0 & -1 \\ 1 & 2 & 2 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 2 \\ -1 & 2 \end{pmatrix} \\ &= \begin{pmatrix} 2 & -1 \\ -1 & 9 \end{pmatrix} \end{aligned}$$

$$\begin{pmatrix} 2 & -1 \\ -1 & 9 \end{pmatrix} \quad 2R_2 + R_1$$

$$\begin{pmatrix} 2 & -1 \\ 0 & 17 \end{pmatrix}$$

$$\underline{\text{rank}(A^T A) = 2}$$

$$AA^T = \begin{pmatrix} 1 & 1 \\ 0 & 2 \\ -1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 0 & -1 \\ 1 & 2 & 2 \end{pmatrix}$$

$$= \begin{pmatrix} 2 & 2 & 1 \\ 2 & 4 & 4 \\ 1 & 4 & 5 \end{pmatrix}$$

$$\begin{pmatrix} 2 & 2 & 1 \\ 2 & 4 & 4 \\ 1 & 4 & 5 \end{pmatrix} \begin{array}{l} \\ R_2 - R_1 \\ 2R_3 - R_1 \end{array}$$

$$\begin{pmatrix} 2 & 2 & 1 \\ 0 & 2 & 3 \\ 0 & 6 & 9 \end{pmatrix} \begin{array}{l} \\ \\ R_3 - 3R_2 \end{array}$$

$$\begin{pmatrix} 2 & 2 & 1 \\ 0 & 2 & 3 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\underline{\text{rank}(A^T A) = 2}$$

$\therefore \text{rank}(A) = \text{rank}(A^T A) = \text{rank}(AA^T)$  for any matrix,  $A$ .

### Exercise

Explain why these are all false:

- The complete solution is any linear combination of  $\vec{x}_p$  and  $\vec{x}_n$ .
- A system  $A\vec{x} = \vec{b}$  has at most one particular solution.
- The solution  $\vec{x}_p$  with all free variables zero is the shortest solution (minimum length  $\|\vec{x}\|$ ). Find a 2 by 2 counterexample.
- If  $A$  is invertible there is no solution  $\vec{x}_n$  in the null space.

### Solution

- The coefficient of  $\vec{x}_p$  must be one.
- If  $\vec{x}_n \in N(A)$  is the nullspace of  $A$  and  $\vec{x}_p$  is one particular solution, then  $\vec{x}_p$  and  $\vec{x}_n$  is also a particular solution.
- If  $A$  is a 2 by 2 matrix of rank 1, then the solution to  $A\vec{x} = \vec{b}$  form a line parallel to the line that the nullspace. The line  $x + y = 1$  gives such an example.

$$A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \quad \vec{b} = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad \vec{x}_p = \begin{pmatrix} \frac{1}{2} & \frac{1}{2} \end{pmatrix}$$

$$\begin{aligned}
 \text{Then } \|\vec{x}_p\| &= \sqrt{\left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right)^2} \\
 &= \sqrt{2 \cdot \frac{1}{4}} \\
 &= \frac{1}{\sqrt{2}} < 1
 \end{aligned}$$

while the particular solutions having some coordinate equal to zero are (1, 0) and (0, 1) and they both have  $\|\vec{x}_p\| = 1$

*d)* There is always  $\vec{x}_n = 0$

### Exercise

Write down all known relation between  $r$  and  $m$  and  $n$  if  $A\vec{x} = \vec{b}$  has

- a)* No solution for some  $\vec{b}$ .
- b)* Infinitely many solutions for every  $\vec{b}$ .
- c)* Exactly one solution for some  $\vec{b}$ , no solution for another  $\vec{b}$ .
- d)* Exactly one solution for every  $\vec{b}$ .

### Solution

- a)* The system has less than full row rank:  $r < m$ .
- b)* The system has full row rank and less than full column rank:  $m = r < n$ .
- c)* The system has full column rank and less than full row rank:  $n = r < m$ .
- d)* The system has full row and column rank (it is invertible):  $m = r = n$ .

### Exercise

Find a basis for its row space, find a basis for its column space, and determine its rank

$$\begin{array}{ll}
 a) \begin{bmatrix} 0 & 2 & -3 & 4 & 1 & 2 & 1 & 7 \\ 0 & 0 & 3 & -2 & 0 & 4 & -5 & 3 \\ 0 & 0 & 0 & 0 & 0 & 6 & -2 & 2 \\ 0 & 0 & 0 & 0 & 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} & b) \begin{bmatrix} 3 & 2 & -1 \\ 6 & 3 & 5 \\ -3 & -1 & -6 \\ 0 & -1 & 7 \end{bmatrix}
 \end{array}$$

### Solution

- a) Row Space:* every row

$$\text{Column Space: } \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -3 \\ 3 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 4 \\ 6 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -5 \\ -2 \\ 2 \\ 0 \end{bmatrix}$$

$$\text{Rank} = 4$$

$$b) \begin{pmatrix} 3 & 2 & -1 \\ 6 & 3 & 5 \\ -3 & -1 & -6 \\ 0 & -1 & 7 \end{pmatrix} \begin{array}{l} \\ R_2 - 2R_1 \\ R_3 + R_1 \\ \end{array}$$

$$\begin{pmatrix} 3 & 2 & -1 \\ 0 & -1 & 7 \\ 0 & 1 & -7 \\ 0 & -1 & 7 \end{pmatrix} \begin{array}{l} R_1 + 2R_2 \\ \\ R_3 + R_2 \\ R_4 - R_2 \end{array}$$

$$\begin{pmatrix} 3 & 0 & 13 \\ 0 & -1 & 7 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{array}{l} \frac{1}{3}R_1 \\ -R_2 \\ \\ \end{array}$$

$$\begin{pmatrix} 1 & 0 & \frac{13}{3} \\ 0 & 1 & -7 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\text{Row Space: } [3 \ 2 \ -1], [6 \ 3 \ 5]$$

$$\text{Column Space: } \begin{bmatrix} 3 \\ 6 \\ -3 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ -1 \\ -1 \end{bmatrix}$$

$$\text{Rank} = 2$$

### Exercise

Find a basis for the row space, find a basis for the null space, find  $\dim RS$ , find  $\dim NS$ , and verify  $\dim RS + \dim NS = n$

$$\begin{bmatrix} 1 & -2 & 4 & 1 \\ 3 & 1 & -3 & -1 \\ 5 & -3 & 5 & 1 \end{bmatrix}$$

### Solution

$$\begin{pmatrix} 1 & -2 & 4 & 1 \\ 3 & 1 & -3 & -1 \\ 5 & -3 & 5 & 1 \end{pmatrix} \quad \begin{array}{l} R_2 - 3R_1 \\ R_3 - 5R_1 \end{array}$$

$$\begin{pmatrix} 1 & -2 & 4 & 1 \\ 0 & 7 & -15 & -4 \\ 0 & 7 & -15 & -4 \end{pmatrix} \quad \begin{array}{l} 7R_1 + 2R_2 \\ R_3 - R_2 \end{array}$$

$$\begin{pmatrix} 7 & 0 & -2 & -1 \\ 0 & 7 & -15 & -4 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad \begin{array}{l} \frac{1}{7}R_1 \\ -\frac{1}{7}R_2 \end{array}$$

$$\begin{pmatrix} 1 & 0 & -\frac{2}{7} & -\frac{1}{7} \\ 0 & 1 & -\frac{15}{7} & -\frac{4}{7} \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

Row Space:  $\begin{bmatrix} 1 & -2 & 4 & 1 \end{bmatrix}, \begin{bmatrix} 3 & 1 & -3 & -1 \end{bmatrix}$

$$\text{Column Space: } \begin{pmatrix} 1 \\ 3 \\ 5 \end{pmatrix}, \begin{pmatrix} -2 \\ 1 \\ -3 \end{pmatrix}$$

$$\dim RS = 2$$

$$\dim NS = 2$$

$$2 + 2 = 4 \Rightarrow \dim RS + \dim NS = n$$



### Exercise

Determine if  $\vec{b}$  lies in the column space of the given matrix. If it does, express  $\vec{b}$  as linear combination of the column.

$$\begin{bmatrix} 2 & -3 \\ -4 & 6 \end{bmatrix}, \quad \vec{b} = \begin{bmatrix} 4 \\ -6 \end{bmatrix}$$

### Solution

$$\left[ \begin{array}{cc|c} 2 & -3 & 4 \\ -4 & 6 & -6 \end{array} \right] \quad R_2 + 2R_1$$

$$\left[ \begin{array}{cc|c} 2 & -3 & 4 \\ 0 & 0 & 2 \end{array} \right] \quad \begin{array}{l} \frac{1}{2}R_1 \\ \frac{1}{2}R_2 \end{array}$$

$$\left[ \begin{array}{cc|c} 1 & -\frac{3}{2} & 2 \\ 0 & 0 & 1 \end{array} \right] \quad R_1 - 2R_2$$

$$\left[ \begin{array}{cc|c} 1 & -\frac{3}{2} & 0 \\ 0 & 0 & 1 \end{array} \right]$$

$\vec{b}$  does not lie in the column space

### Exercise

Find the transition matrix from  $B$  to  $C$  and find  $[\vec{x}]_C$

$$a) \quad B = \{(3, 1), (-1, -2)\}, \quad C = \{(1, -3), (5, 0)\}, \quad [\vec{x}]_B = [-1 \quad -2]^T$$

$$b) \quad B = \{(1, 1, 1), (-2, -1, 0), (2, 1, 2)\}, \quad C = \{(-6, -2, 1), (-1, 1, 5), (-1, -1, 1)\},$$

$$[\vec{x}]_B = [-3 \quad 2 \quad 4]^T$$

### Solution

$$a) \quad \left[ \begin{array}{cc|cc} 1 & 5 & 3 & -1 \\ -3 & 0 & 1 & -2 \end{array} \right] \quad R_2 + 3R_1$$

$$\left[ \begin{array}{cc|cc} 1 & 5 & 3 & -1 \\ 0 & 15 & 10 & -5 \end{array} \right] \quad 3R_1 - R_2$$

$$\left[ \begin{array}{cc|cc} 3 & 0 & -1 & 2 \\ 0 & 15 & 10 & -5 \end{array} \right] \quad \begin{array}{l} \frac{1}{3}R_1 \\ \frac{1}{15}R_2 \end{array}$$

$$\left[ \begin{array}{cc|cc} 1 & 0 & -\frac{1}{3} & \frac{2}{3} \\ 0 & 1 & \frac{2}{3} & -\frac{1}{3} \end{array} \right]$$

$$\begin{aligned}
[\vec{x}]_c &= \begin{bmatrix} -\frac{1}{3} & \frac{2}{3} \\ \frac{2}{3} & -\frac{1}{3} \end{bmatrix} \begin{bmatrix} -1 \\ -2 \end{bmatrix} \\
&= \begin{bmatrix} -1 \\ 0 \end{bmatrix}
\end{aligned}$$

$$b) \left[ \begin{array}{ccc|ccc} -6 & -1 & -1 & 1 & -2 & 2 \\ -2 & 1 & -1 & 1 & -1 & 1 \\ 1 & 5 & 1 & 1 & 0 & 2 \end{array} \right] \begin{array}{l} 3R_2 - R_1 \\ 6R_3 + R_1 \end{array}$$

$$\left[ \begin{array}{ccc|ccc} -6 & -1 & -1 & 1 & -2 & 2 \\ 0 & 4 & -2 & 2 & -1 & 1 \\ 0 & 29 & 5 & 7 & -2 & 14 \end{array} \right] \begin{array}{l} 4R_1 + R_2 \\ 4R_3 - 29R_2 \end{array}$$

$$\left[ \begin{array}{ccc|ccc} -24 & 0 & -6 & 6 & -9 & 9 \\ 0 & 4 & -2 & 2 & -1 & 1 \\ 0 & 0 & 78 & -30 & 21 & 27 \end{array} \right] \begin{array}{l} 13R_1 + R_3 \\ 39R_2 + R_3 \end{array}$$

$$\left[ \begin{array}{ccc|ccc} -312 & 0 & 0 & 48 & -96 & 144 \\ 0 & 156 & 0 & 48 & -18 & 66 \\ 0 & 0 & 78 & -30 & 21 & 27 \end{array} \right] \begin{array}{l} -\frac{1}{312}R_1 \\ \frac{1}{156}R_2 \\ \frac{1}{78}R_3 \end{array}$$

$$\left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & -\frac{2}{13} & \frac{4}{13} & -\frac{6}{13} \\ 0 & 1 & 0 & \frac{4}{13} & -\frac{3}{26} & \frac{11}{26} \\ 0 & 0 & 1 & -\frac{5}{13} & \frac{7}{26} & \frac{9}{26} \end{array} \right]$$

$$\begin{aligned}
[\vec{x}]_c &= \begin{bmatrix} -\frac{2}{13} & \frac{4}{13} & -\frac{6}{13} \\ \frac{4}{13} & -\frac{3}{26} & \frac{11}{26} \\ -\frac{5}{13} & \frac{7}{26} & \frac{9}{26} \end{bmatrix} \begin{bmatrix} -3 \\ 2 \\ 4 \end{bmatrix} \\
&= \begin{bmatrix} -\frac{10}{13} \\ \frac{17}{13} \\ \frac{35}{13} \end{bmatrix}
\end{aligned}$$

### Exercise

Does  $A$  and  $A^T$  have the same number of pivots.

#### Solution

True

The number of pivots of  $A$  is its column rank,  $r$ .

We know that the column rank of  $A$  equals the row rank of  $A$ , which is the column rank of  $A^T$ .

Hence,  $A^T$  must have the same number of pivots as  $A$ .

### Exercise

Let  $A = \begin{pmatrix} 1 & 3 & 0 & 2 \\ 0 & 0 & 1 & 4 \\ 0 & 0 & 0 & 0 \end{pmatrix}$  where  $\vec{b} = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$

- a) What is the rank of  $A$ ?
- b) What is the dimension of  $A$ ?
- c) What are the pivots variables?
- d) What are the free variables?
- e) Find the special (homogeneous) solutions.
- f) What is the nullspace  $N(A)$ ?
- g) Find the particular solution to  $A\vec{x} = \vec{b}$
- h) Give the complete solution.

#### Solution

- a)  $\text{Rank}(A) = 2$
- b) Dimension of  $A = 2$
- c) The pivots variables are:  $x_1, x_3$
- d) The free variables are:  $x_2, x_4$
- e) Let  $x = x_2 s_1 + x_4 s_2$

$$A = \begin{pmatrix} 1 & 3 & 0 & 2 \\ 0 & 0 & 1 & 4 \\ 0 & 0 & 0 & 0 \end{pmatrix} \rightarrow \begin{cases} x_1 = -3x_2 - 2x_4 \\ x_3 = -4x_4 \end{cases}$$

Set  $x_2 = 1, x_4 = 0$

The special solution:  $\vec{s}_1 = (-3, 1, 0, 0)$

Set  $x_2 = 0, x_4 = 1$  ;

The special solution:  $\vec{s}_2 = (-2, 0, -4, 1)$

*f)* The nullspace is the set  $\left\{ x_2 \begin{pmatrix} -3 \\ 1 \\ 0 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -2 \\ 0 \\ -4 \\ 1 \end{pmatrix} \right\}$

$$N(A) = \begin{pmatrix} -3 & -2 \\ 1 & 0 \\ 0 & -4 \\ 0 & 1 \end{pmatrix}$$

*g)*  $\vec{x}_p = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$

*h)*  $\vec{x} = x_2 \begin{pmatrix} -3 \\ 1 \\ 0 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} -2 \\ 0 \\ -4 \\ 1 \end{pmatrix} + \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$

### Exercise

Let  $A = \begin{pmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$  where  $\vec{b} = A \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$

- What is the rank of  $A$ ?
- What is the dimension of  $A$ ?
- What are the pivots variables?
- What are the free variables?
- Find the special (homogeneous) solutions.
- What is the nullspace  $N(A)$ ?
- Find the particular solution to  $A\vec{x} = \vec{b}$
- Give the complete solution.

### Solution

- $\text{Rank}(A) = 3$
- Dimension of  $A = 1$
- The pivots variables are:  $x_1, x_2, x_4$

d) The free variables are:  $x_3$

e) Let  $x = x_3 s_1$

$$A = \begin{pmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \rightarrow \begin{cases} x_1 = -2x_3 \\ x_2 = 0 \\ x_3 = 0 \end{cases}$$

The special solution:  $\vec{s}_1 = (-2, 0, 1, 0)$

$$f) N(A) = \begin{pmatrix} -2 \\ 0 \\ 1 \\ 0 \end{pmatrix}$$

$$g) \vec{x}_p = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$$

$$h) \vec{x} = x_3 \begin{pmatrix} -2 \\ 0 \\ 1 \\ 0 \end{pmatrix} + \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$$

### Exercise

Let  $A = \begin{pmatrix} 1 & 2 & 0 & 4 \\ 0 & -3 & 1 & -12 \\ 0 & 0 & 0 & 0 \end{pmatrix}$  where  $\vec{b} = A \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$

- a) What is the rank of  $A$ ?
- b) What is the dimension of  $A$ ?
- c) What are the pivots variables?
- d) What are the free variables?
- e) Find the special (homogeneous) solutions.
- f) What is the nullspace  $N(A)$ ?
- g) Find the particular solution to  $A\vec{x} = \vec{b}$
- h) Give the complete solution.

### Solution

$$\begin{pmatrix} 1 & 2 & 0 & 4 \\ 0 & -3 & 1 & -12 \\ 0 & 0 & 0 & 0 \end{pmatrix} \xrightarrow{-\frac{1}{3}R_2}$$

$$\begin{pmatrix} 1 & 2 & 0 & 4 \\ 0 & 1 & -\frac{1}{3} & 4 \\ 0 & 0 & 0 & 0 \end{pmatrix} \xrightarrow{R_1 - 2R_2}$$

$$\begin{pmatrix} 1 & 0 & \frac{2}{3} & -4 \\ 0 & 1 & -\frac{1}{3} & 4 \\ 0 & 0 & 0 & 0 \end{pmatrix} \rightarrow \begin{cases} x_1 = -\frac{2}{3}x_3 + 4x_4 \\ x_2 = \frac{1}{3}x_3 - 4x_4 \end{cases}$$

**a)**  $\text{Rank}(A) = 2$

**b)** Dimension of  $A = 2$

**c)** The pivots variables are:  $x_1, x_2$

**d)** The free variables are:  $x_3, x_4$

**e)** Let  $x = x_3 \vec{s}_1 + x_4 \vec{s}_2$

Set  $x_3 = 1, x_4 = 0$

The special solution:  $\vec{s}_1 = \left(-\frac{2}{3}, \frac{1}{3}, 1, 0\right)$

Set  $x_3 = 0, x_4 = 1$  ;

The special solution:  $\vec{s}_2 = (4, -4, 0, 1)$

**f)** The nullspace is the set  $\left\{ x_3 \begin{pmatrix} -\frac{2}{3} \\ \frac{1}{3} \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} 4 \\ -4 \\ 0 \\ 1 \end{pmatrix} \right\}$

$$N(A) = \begin{pmatrix} -\frac{2}{3} & 4 \\ \frac{1}{3} & -4 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$g) \quad \vec{x}_p = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$$

$$h) \quad \vec{x} = x_3 \begin{pmatrix} -\frac{2}{3} \\ \frac{1}{3} \\ 1 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} 4 \\ -4 \\ 0 \\ 1 \end{pmatrix} + \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$$

### ***Exercise***

Let  $A = \begin{pmatrix} 1 & 0 & 0 & \frac{13}{11} \\ 0 & 1 & 0 & -\frac{17}{11} \\ 0 & 0 & 1 & \frac{6}{11} \\ 0 & 0 & 0 & 0 \end{pmatrix}$  where  $\vec{b} = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$

- What is the rank of  $A$ ?
- What is the dimension of  $A$ ?
- What are the pivots variables?
- What are the free variables?
- Find the special (homogeneous) solutions.
- What is the nullspace  $N(A)$ ?
- Find the particular solution to  $A\vec{x} = \vec{b}$
- Give the complete solution.

### **Solution**

- $\text{Rank}(A) = 3$
- Dimension of  $A = 1$
- The pivots variables are:  $x_1, x_2, x_3$
- The free variables are:  $x_4$
- Let  $x = x_4 \begin{pmatrix} s \\ 1 \end{pmatrix}$

$$A = \begin{pmatrix} 1 & 0 & 0 & \frac{13}{11} \\ 0 & 1 & 0 & -\frac{17}{11} \\ 0 & 0 & 1 & \frac{6}{11} \\ 0 & 0 & 0 & 0 \end{pmatrix} \rightarrow \begin{cases} x_1 = -\frac{13}{11} \\ x_2 = \frac{17}{11} \\ x_3 = -\frac{6}{11} \end{cases}$$

Set  $x_4 = 1$

The special solution:  $\vec{s}_1 = \left(-\frac{13}{11}, \frac{17}{11}, -\frac{6}{11}, 1\right)$

$$f) \quad N(A) = \begin{pmatrix} -\frac{13}{11} \\ \frac{17}{11} \\ -\frac{6}{11} \\ 1 \end{pmatrix}$$

$$g) \quad \vec{x}_p = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$$

$$h) \quad \vec{x} = x_4 \begin{pmatrix} -\frac{13}{11} \\ \frac{17}{11} \\ -\frac{6}{11} \\ 1 \end{pmatrix} + \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$$



### Exercise

Let  $A = \begin{pmatrix} 1 & 0 & 0 & -\frac{1}{3} \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -\frac{1}{3} \\ 0 & 0 & 0 & 0 \end{pmatrix}$  where  $\vec{b} = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$

- a) What is the rank of  $A$ ?
- b) What is the dimension of  $A$ ?
- c) What are the pivots variables?
- d) What are the free variables?
- e) Find the special (homogeneous) solutions.
- f) What is the nullspace  $N(A)$ ?
- g) Find the particular solution to  $A\vec{x} = \vec{b}$
- h) Give the complete solution.

### Solution

- a)  $\text{Rank}(A) = 3$
- b) Dimension of  $A = 1$
- c) The pivots variables are:  $x_1, x_2, x_3$
- d) The free variables are:  $x_4$
- e) Let  $x = x_4 s_1$

$$A = \begin{pmatrix} 1 & 0 & 0 & -\frac{1}{3} \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -\frac{1}{3} \\ 0 & 0 & 0 & 0 \end{pmatrix} \rightarrow \begin{cases} x_1 = \frac{1}{3} \\ x_2 = 1 \\ x_3 = \frac{1}{3} \end{cases}$$

Set  $x_4 = 1$

The special solution:  $\vec{s}_1 = \left(\frac{1}{3}, 1, \frac{1}{3}, 1\right)$

f)  $N(A) = \begin{pmatrix} \frac{1}{3} \\ 1 \\ \frac{1}{3} \\ 1 \end{pmatrix}$

$$g) \quad \vec{x}_p = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$$

$$h) \quad \vec{x} = x_4 \begin{pmatrix} \frac{1}{3} \\ 1 \\ \frac{1}{3} \\ 1 \end{pmatrix} + \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$$

### Exercise

$$\text{Let } A = \begin{pmatrix} 1 & 2 & 0 & -1 & 0 \\ 0 & 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 0 & 5 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \text{ where } \vec{b} = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$$

- What is the rank of  $A$ ?
- What is the dimension of  $A$ ?
- What are the pivots variables?
- What are the free variables?
- Find the special (homogeneous) solutions.
- What is the nullspace  $N(A)$ ?
- Find the particular solution to  $A\vec{x} = \vec{b}$
- Give the complete solution.

### Solution

- Rank( $A$ ) = 3
- Dimension of  $A = 2$
- The pivots variables are:  $x_1, x_3, x_5$
- The free variables are:  $x_2, x_4$

$$e) \text{ Let } x = x_2 s_1 + x_4 s_2$$

$$A = \begin{pmatrix} 1 & 2 & 0 & -1 & 0 \\ 0 & 0 & 1 & 3 & 0 \\ 0 & 0 & 0 & 0 & 5 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \rightarrow \begin{cases} x_1 = -2x_2 + x_4 \\ x_3 = -x_4 \\ x_5 = 0 \end{cases}$$

$$\text{Set } x_2 = 1 \quad x_4 = 0$$

The special solution:  $\vec{s}_1 = (-2, 1, 0, 0, 0)$

Set  $x_2 = 0$   $x_4 = 1$

The special solution:  $\vec{s}_2 = (1, 0, -1, 1, 0)$

$$f) \quad N(A) = \begin{pmatrix} -2 & 1 \\ 1 & 0 \\ 0 & -1 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$$

$$g) \quad \vec{x}_p = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$$

$$h) \quad \vec{x} = x_2 \begin{pmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} + x_4 \begin{pmatrix} 1 \\ 0 \\ -1 \\ 1 \\ 0 \end{pmatrix} + \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$$

### ***Exercise***

Find a basis for each of the four subspaces associated with the given matrix

$$A = \begin{pmatrix} 1 & 2 & 4 \\ 2 & 5 & 8 \end{pmatrix}$$

### **Solution**

$$\begin{pmatrix} 1 & 2 & 4 \\ 2 & 5 & 8 \end{pmatrix} \quad R_2 - 2R_1$$

$$\begin{pmatrix} 1 & 2 & 4 \\ 0 & 1 & 0 \end{pmatrix} \quad R_1 - 2R_2$$

$$\begin{pmatrix} 1 & 0 & 4 \\ 0 & 1 & 0 \end{pmatrix} \quad x_1 = -4x_3 \quad \leftarrow \text{Row space}$$

$$\text{Rank}(A) = 1$$

$$\text{Dimension of } A = 1$$

1. Basis for **row space**:  $\begin{pmatrix} 1 \\ 0 \\ 4 \end{pmatrix}$

The pivots variables are:  $x_1, x_2$

2. Basis of the **column spaces**:  $\begin{pmatrix} 1 \\ 2 \end{pmatrix} \begin{pmatrix} 2 \\ 5 \end{pmatrix}$

The free variable is:  $x_3$

Set  $x_3 = 1 \Rightarrow s_1 = (-4, 0, 1)$

3. Basis of the **Nullspace**:  $\begin{pmatrix} -4 \\ 0 \\ 1 \end{pmatrix}$

$$A^T = \begin{pmatrix} 1 & 2 \\ 2 & 5 \\ 4 & 8 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 2 \\ 2 & 5 \\ 4 & 8 \end{pmatrix} \begin{array}{l} \\ R_2 - 2R_1 \\ R_3 - 4R_1 \end{array}$$

$$\begin{pmatrix} 1 & 2 \\ 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{array}{l} R_1 - 2R_2 \\ \\ \end{array}$$

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix}$$

4. Basis of the **Left Nullspace**:  $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$

### ***Exercise***

Find a basis for each of the four subspaces associated with the given matrix

$$B = \begin{pmatrix} 1 & 3 & 0 & 5 \\ 2 & 6 & 1 & 16 \\ 5 & 15 & 0 & 25 \end{pmatrix}$$

### **Solution**

$$\begin{pmatrix} 1 & 3 & 0 & 5 \\ 2 & 6 & 1 & 16 \\ 5 & 15 & 0 & 25 \end{pmatrix} \begin{array}{l} \\ R_2 - 2R_1 \\ R_3 - 5R_1 \end{array}$$

$$\begin{pmatrix} 1 & 3 & 0 & 5 \\ 0 & 0 & 1 & 6 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{array}{l} x_1 = -3x_2 - 5x_4 \\ x_3 = -6x_4 \\ \\ \end{array} \begin{array}{l} \leftarrow \text{Row space} \\ \leftarrow \text{Row space} \\ \\ \end{array}$$

$$\text{Rank}(A) = 2$$

$$\text{Dimension of } A = 2$$

1. Basis for **row space**:

$$\begin{pmatrix} 1 \\ 3 \\ 0 \\ 5 \end{pmatrix} \begin{pmatrix} 0 \\ 0 \\ 1 \\ 6 \end{pmatrix}$$

$$\text{The pivots variables are: } x_1, x_3$$

2. Basis of the **column spaces**:

$$\begin{pmatrix} 1 \\ 2 \\ 5 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$$

$$\text{The free variable is: } x_2, x_4$$

$$\begin{cases} x_1 = -3x_2 - 5x_4 \\ x_3 = -6x_4 \end{cases}$$

$$\text{Set } x_2 = 1 \quad x_4 = 0$$

$$\text{The special solution: } s_1 = (-3, 1, 0, 0)$$

$$\text{Set } x_2 = 0 \quad x_4 = 1$$

$$\text{The special solution: } s_2 = (-5, 0, -6, 1)$$

3. Basis of the **Nullspace**:

$$\begin{pmatrix} -3 \\ 1 \\ 0 \\ 0 \end{pmatrix} \begin{pmatrix} -5 \\ 0 \\ -6 \\ 1 \end{pmatrix}$$

$$B^T = \begin{pmatrix} 1 & 2 & 5 \\ 3 & 6 & 15 \\ 0 & 1 & 0 \\ 5 & 16 & 25 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 2 & 5 \\ 3 & 6 & 15 \\ 0 & 1 & 0 \\ 5 & 16 & 25 \end{pmatrix} \quad \begin{array}{l} R_2 - 3R_1 \\ R_4 - 5R_1 \end{array}$$

$$\begin{pmatrix} 1 & 2 & 5 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 6 & 0 \end{pmatrix} \quad \begin{array}{l} R_4 - 2R_3 \\ R_4 - 6R_3 \end{array}$$

$$\begin{pmatrix} 1 & 0 & 5 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad y_1 = -5y_3$$

Let  $y_3 = 1$

4. Basis of the **Left Nullspace**:  $\begin{pmatrix} -5 \\ 0 \\ 1 \end{pmatrix}$

### Exercise

Find a basis for each of the four subspaces associated with the given matrix

$$C = \begin{pmatrix} 0 & 1 & 2 & 3 & 4 \\ 0 & 1 & 2 & 4 & 6 \\ 0 & 0 & 0 & 1 & 2 \end{pmatrix}$$

### Solution

$$\begin{pmatrix} 0 & 1 & 2 & 3 & 4 \\ 0 & 1 & 2 & 4 & 6 \\ 0 & 0 & 0 & 1 & 2 \end{pmatrix} \quad R_2 - R_1$$

$$\begin{pmatrix} 0 & 1 & 2 & 3 & 4 \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 1 & 2 \end{pmatrix} \quad \begin{array}{l} R_1 - 3R_2 \\ R_3 - R_2 \end{array}$$

$$\begin{pmatrix} 0 & 1 & 2 & 0 & -2 \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad \begin{array}{ll} x_2 = -2x_3 + 2x_5 & \leftarrow \text{Row space} \\ x_4 = -2x_5 & \leftarrow \text{Row space} \end{array}$$

Rank ( $A$ ) = 2

Dimension of  $A = 2$

1. Basis for **row space**:

$$\begin{pmatrix} 0 \\ 1 \\ 2 \\ 0 \\ -2 \end{pmatrix} \quad \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \\ 2 \end{pmatrix}$$

The pivots variables are:  $x_2, x_4$

2. Basis of the **column spaces**:

$$\begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \quad \begin{pmatrix} 3 \\ 4 \\ 1 \end{pmatrix}$$

The free variable is:  $x_3, x_5$

$$\begin{cases} x_2 = -2x_3 + 2x_5 \\ x_4 = -2x_5 \end{cases}$$

Set  $x_3 = 1 \quad x_5 = 0$

The special solution:  $s_1 = (0, -2, 1, 0, 2)$

Set  $x_3 = 0 \quad x_5 = 1$

The special solution:  $s_2 = (0, 2, 0, -2, 1)$

3. Basis of the **Nullspace**:

$$\begin{pmatrix} 0 \\ -2 \\ 1 \\ 0 \\ 2 \end{pmatrix} \quad \begin{pmatrix} 0 \\ 2 \\ 0 \\ -2 \\ 1 \end{pmatrix}$$

$$C^T = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 1 & 0 \\ 2 & 2 & 0 \\ 3 & 4 & 1 \\ 4 & 6 & 2 \end{pmatrix}$$

$$\begin{pmatrix} 0 & 0 & 0 \\ 1 & 1 & 0 \\ 2 & 2 & 0 \\ 3 & 4 & 1 \\ 4 & 6 & 2 \end{pmatrix} \quad \begin{matrix} \\ R_3 - 2R_2 \\ R_4 - 3R_2 \\ R_5 - 4R_2 \end{matrix}$$

$$\begin{pmatrix} 0 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 3 & 1 \\ 0 & 2 & 2 \end{pmatrix} \begin{array}{l} \\ 3R_2 - R_4 \\ \\ 3R_5 - 2R_4 \\ \end{array}$$

$$\begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & -1 \\ 0 & 0 & 0 \\ 0 & 3 & 1 \\ 0 & 0 & 4 \end{pmatrix} \begin{array}{l} \\ \frac{1}{3}R_4 \\ \frac{1}{4}R_5 \\ \\ \end{array}$$

$$\begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & -1 \\ 0 & 0 & 0 \\ 0 & 1 & \frac{1}{3} \\ 0 & 0 & 1 \end{pmatrix} \begin{array}{l} \\ R_2 + R_5 \\ \\ R_4 - \frac{1}{3}R_5 \\ \end{array}$$

$$\begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

4. Basis of the **Left Nullspace**:

$$\begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$



### Exercise

Find a basis for each of the four subspaces associated with the given matrix

$$D = \begin{pmatrix} 1 & 2 & 3 & 1 \\ 1 & 1 & 2 & 1 \\ 1 & 2 & 3 & 1 \end{pmatrix}$$

### Solution

$$\begin{pmatrix} 1 & 2 & 3 & 1 \\ 1 & 1 & 2 & 1 \\ 1 & 2 & 3 & 1 \end{pmatrix} \begin{array}{l} \\ R_2 - R_1 \\ R_3 - R_1 \end{array}$$

$$\begin{pmatrix} 1 & 2 & 3 & 1 \\ 0 & -1 & -1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{array}{l} R_2 + 2R_1 \\ -R_2 \\ \end{array}$$

$$\begin{pmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{array}{ll} x_1 = -x_3 - x_4 & \leftarrow \text{Row space} \\ x_2 = -x_3 & \leftarrow \text{Row space} \end{array}$$

$$\text{Rank}(A) = 2$$

$$\text{Dimension of } A = 2$$

1. Basis for **row space**:  $\begin{pmatrix} 1 \\ 0 \\ 1 \\ 1 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \\ 1 \\ 0 \end{pmatrix}$

The pivots variables are:  $x_1, x_2$

2. Basis of the **column spaces**:  $\begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \begin{pmatrix} 2 \\ 1 \\ 2 \end{pmatrix}$

The free variable is:  $x_3, x_4$

$$\begin{cases} x_1 = -x_3 - x_4 \\ x_2 = -x_3 \end{cases}$$

$$\text{Set } x_3 = 1 \quad x_4 = 0$$

$$\text{The special solution: } s_1 = (-1, 0, 1, 0)$$

$$\text{Set } x_3 = 0 \quad x_4 = 1$$

$$\text{The special solution: } s_2 = (-1, -1, 0, 1)$$

3. Basis of the **Nullspace**:

$$\begin{pmatrix} -1 \\ 0 \\ 1 \\ 0 \end{pmatrix} \quad \begin{pmatrix} -1 \\ -1 \\ 0 \\ 1 \end{pmatrix}$$

$$D^T = \begin{pmatrix} 1 & 1 & 1 \\ 2 & 1 & 2 \\ 3 & 2 & 3 \\ 1 & 1 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 1 & 1 \\ 2 & 1 & 2 \\ 3 & 2 & 3 \\ 1 & 1 & 1 \end{pmatrix} \begin{array}{l} \\ R_2 - 2R_1 \\ R_3 - 3R_1 \\ R_4 - R_1 \end{array}$$

$$\begin{pmatrix} 1 & 1 & 1 \\ 0 & -1 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{array}{l} R_1 + R_2 \\ \\ R_3 - 2R_2 \\ \end{array}$$

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{array}{l} \\ -R_2 \\ \\ \end{array}$$

$$\begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad y_1 = -y_3$$

Let  $y_3 = 1$

4. Basis of the **Left Nullspace**:

$$\begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$$

### Exercise

Find a basis for each of the four subspaces associated with the given matrix

$$M = \begin{pmatrix} 1 & -2 & 4 & 1 \\ 3 & 1 & -3 & -1 \\ 5 & -3 & 5 & 1 \end{pmatrix}$$

### Solution

$$\begin{pmatrix} 1 & -2 & 4 & 1 \\ 3 & 1 & -3 & -1 \\ 5 & -3 & 5 & 1 \end{pmatrix} \begin{array}{l} \\ R_2 - 3R_1 \\ R_3 - 5R_1 \end{array}$$

$$\begin{pmatrix} 1 & -2 & 4 & 1 \\ 0 & 7 & -15 & -4 \\ 0 & 7 & -15 & -4 \end{pmatrix} \begin{array}{l} 7R_1 + 2R_2 \\ \\ R_3 - R_2 \end{array}$$

$$\begin{pmatrix} 7 & 0 & -2 & -1 \\ 0 & 7 & -15 & -4 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{array}{l} \frac{1}{7}R_1 \\ \frac{1}{7}R_2 \\ \end{array}$$

$$\begin{pmatrix} 1 & 0 & -\frac{2}{7} & -\frac{1}{7} \\ 0 & 1 & -\frac{15}{7} & -\frac{4}{7} \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{array}{l} x_1 = \frac{2}{7}x_3 + \frac{1}{7}x_4 \\ x_2 = \frac{15}{7}x_3 + \frac{4}{7}x_4 \\ \end{array} \begin{array}{l} \leftarrow \text{Row space} \\ \leftarrow \text{Row space} \end{array}$$

$$\text{Rank}(A) = 2$$

$$\text{Dimension of } A = 2$$

1. Basis for **row space**:

$$\begin{pmatrix} 1 \\ 0 \\ -\frac{2}{7} \\ -\frac{1}{7} \end{pmatrix} \quad \begin{pmatrix} 0 \\ 1 \\ -\frac{15}{7} \\ -\frac{4}{7} \end{pmatrix}$$

The pivots variables are:  $x_1, x_2$

2. Basis of the **column spaces**:

$$\begin{pmatrix} 1 \\ 3 \\ 5 \end{pmatrix} \quad \begin{pmatrix} -2 \\ 1 \\ -3 \end{pmatrix}$$

The free variable is:  $x_3, x_4$

$$\begin{cases} x_1 = \frac{2}{7}x_3 + \frac{1}{7}x_4 \\ x_2 = \frac{15}{7}x_3 + \frac{4}{7}x_4 \end{cases}$$

$$\text{Set } x_3 = 1 \quad x_4 = 0$$

$$\text{The special solution: } s_1 = \left( \frac{2}{7}, \frac{15}{7}, 1, 0 \right)$$

$$\text{Set } x_3 = 0 \quad x_4 = 1$$

$$\text{The special solution: } s_2 = \left( \frac{1}{7}, \frac{4}{7}, 0, 1 \right)$$

3. Basis of the **Nullspace**:

$$\begin{pmatrix} \frac{2}{7} \\ \frac{15}{7} \\ 1 \\ 0 \end{pmatrix} \quad \begin{pmatrix} \frac{1}{7} \\ \frac{4}{7} \\ 0 \\ 1 \end{pmatrix}$$

$$M^T = \begin{pmatrix} 1 & 3 & 5 \\ -2 & 1 & -3 \\ 4 & -3 & 5 \\ 1 & -1 & 1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 3 & 5 \\ -2 & 1 & -3 \\ 4 & -3 & 5 \\ 1 & -1 & 1 \end{pmatrix} \begin{array}{l} \\ R_2 + 2R_1 \\ R_3 - 4R_1 \\ R_4 - R_1 \end{array}$$

$$\begin{pmatrix} 1 & 3 & 5 \\ 0 & 7 & 7 \\ 0 & -15 & -15 \\ 0 & -4 & -4 \end{pmatrix} \begin{array}{l} \\ \frac{1}{7}R_2 \\ \frac{1}{15}R_3 \\ \frac{1}{4}R_4 \end{array}$$

$$\begin{pmatrix} 1 & 3 & 5 \\ 0 & 1 & 1 \\ 0 & -1 & -1 \\ 0 & -1 & -1 \end{pmatrix} \begin{array}{l} R_1 - 3R_2 \\ \\ R_3 + R_2 \\ R_4 + R_2 \end{array}$$

$$\begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{array}{l} y_1 = -2y_3 \\ y_1 = -y_3 \\ \\ \end{array}$$

Let  $y_3 = 1$

4. Basis of the **Left Nullspace**:  $\begin{pmatrix} -2 \\ -1 \\ 1 \end{pmatrix}$

### ***Exercise***

Find a basis for each of the four subspaces associated with the given matrix

$$N = \begin{pmatrix} 3 & 2 & -1 \\ 6 & 3 & 5 \\ -3 & -1 & -6 \\ 0 & -1 & 7 \end{pmatrix}$$

### **Solution**

$$\begin{pmatrix} 3 & 2 & -1 \\ 6 & 3 & 5 \\ -3 & -1 & -6 \\ 0 & -1 & 7 \end{pmatrix} \begin{array}{l} \\ R_2 - 2R_1 \\ R_3 + R_1 \\ \end{array}$$

$$\begin{pmatrix} 3 & 2 & -1 \\ 0 & -1 & 7 \\ 0 & 1 & -7 \\ 0 & -1 & 7 \end{pmatrix} \begin{array}{l} R_1 + 2R_2 \\ \\ R_3 + R_2 \\ R_4 - R_2 \end{array}$$

$$\begin{pmatrix} 3 & 0 & 13 \\ 0 & -1 & 7 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{array}{l} \frac{1}{3}R_1 \\ -R_2 \\ \\ \end{array}$$

$$\begin{pmatrix} 1 & 0 & \frac{13}{3} \\ 0 & 1 & -7 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{array}{l} x_1 = -\frac{13}{3}x_3 \\ x_2 = 7x_3 \\ \\ \end{array} \begin{array}{l} \leftarrow \text{Row space} \\ \leftarrow \text{Row space} \\ \\ \end{array}$$

$$\text{Rank}(A) = 2$$

$$\text{Dimension of } A = 2$$

1. Basis for **row space**:  $\begin{pmatrix} 1 \\ 0 \\ \frac{13}{3} \\ 3 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \\ -7 \end{pmatrix}$

The pivots variables are:  $x_1, x_2$

2. Basis of the **column spaces**:  $\begin{pmatrix} 1 \\ 3 \\ 5 \end{pmatrix} \begin{pmatrix} -2 \\ 1 \\ -3 \end{pmatrix}$

The free variable is:  $x_3, x_4$

$$\begin{cases} x_1 = -\frac{13}{7}x_3 \\ x_2 = 7x_3 \end{cases}$$

Set  $x_3 = 1$ . The special solution:  $s_1 = \left(-\frac{13}{3}, 7, 1\right)$

3. Basis of the **Nullspace**:  $\begin{pmatrix} -\frac{13}{3} \\ 7 \\ 1 \end{pmatrix}$

$$N^T = \begin{pmatrix} 3 & 6 & -3 & 0 \\ 2 & 3 & -1 & -1 \\ -1 & 5 & -6 & 7 \end{pmatrix}$$

$$\begin{pmatrix} 3 & 6 & -3 & 0 \\ 2 & 3 & -1 & -1 \\ -1 & 5 & -6 & 7 \end{pmatrix} \begin{array}{l} \\ 3R_2 - 3R_1 \\ 3R_3 + R_1 \end{array}$$

$$\begin{pmatrix} 3 & 6 & -3 & 0 \\ 0 & -9 & 6 & -3 \\ 0 & 21 & -21 & 21 \end{pmatrix} \begin{array}{l} \frac{1}{3}R_1 \\ -\frac{1}{9}R_2 \\ \frac{1}{21}R_3 \end{array}$$

$$\begin{pmatrix} 1 & 2 & -1 & 0 \\ 0 & 1 & -\frac{2}{3} & \frac{1}{3} \\ 0 & 1 & -1 & 1 \end{pmatrix} \begin{array}{l} R_1 - 2R_2 \\ \\ R_3 - R_2 \end{array}$$

$$\begin{pmatrix} 1 & 0 & \frac{1}{3} & -\frac{2}{3} \\ 0 & 1 & -\frac{2}{3} & \frac{1}{3} \\ 0 & 0 & -\frac{1}{3} & -\frac{2}{3} \end{pmatrix} \begin{array}{l} R_1 + R_3 \\ R_1 - 2R_3 \end{array}$$

$$\begin{pmatrix} 1 & 0 & 0 & -\frac{4}{3} \\ 0 & 1 & 0 & \frac{5}{3} \\ 0 & 0 & -\frac{1}{3} & -\frac{2}{3} \end{pmatrix} -3R_3$$

$$\begin{pmatrix} 1 & 0 & 0 & -\frac{4}{3} \\ 0 & 1 & 0 & \frac{5}{3} \\ 0 & 0 & 1 & 2 \end{pmatrix} \begin{array}{l} y_1 = \frac{4}{3}y_4 \\ y_2 = -\frac{5}{3}y_4 \\ y_3 = -2y_4 \end{array}$$

Let  $y_4 = 1$

4. Basis of the **Left Nullspace**:  $\begin{pmatrix} \frac{4}{3} \\ -\frac{5}{3} \\ -2 \\ 1 \end{pmatrix}$