DYADIC BI-PARAMETER REPEATED COMMUTATOR AND DYADIC PRODUCT BMO

IRINA HOLMES, SERGEI TREIL, AND ALEXANDER VOLBERG

ABSTRACT. Let \mathbf{T} is a certain tensor product of simple dyadic shifts defined below. We prove here that for dyadic bi-parameter repeated commutator its norm can be estimated from below by Chang-Fefferman BMO norm pertinent to its symbol. See Theorems 9.1, 9.2 at the end of this article. But this is done below under an extra assumption on the Haar-Fourier side of the symbol. In Section 7 we carefully analyze what goes wrong in the absence of this extra assumption. At the end of this note we also list a counterexample to the existing proof of characterization of bi-parameter repeated commutator with the Hilbert transforms. This is a counterexample to the proof, and it is not a counterexample to the statement of factorization result in bi-disc, or to Nehari's theorem in bi-disc.

1. Introduction

In [HTV] we considered the characterization of the symbols of bounded commutator of multiplication and simple dyadic bi-parameter singular integral. In this note we are considering a dyadic model of *repeated* bi-parameter commutator. It has been noticed that repeated commutators are much harder to treat than the simple ones. The former still await the understanding, see below. The latter are well understood in many situations, see [FS], [DLWY], [DKPS], [HTV].

Bi-parameter theory is notoriously more difficult than the more classical one parameter theory of singular integrals. The good place to get acquainted with multi-parameter specifics are the papers of J.-L. Journé [JLJ], [JLJ2] and Muscalu-Pipher-Tao-Thiele [MPTT1], [MPTT2]. The applications to analysis in polydisc can be found in [Ch], [ChF], [ChF2].

It is well known [ChF, ChF2, JLJ, JLJ2] that in the multi-parameter setting all concepts of Carleson measure, BMO, John–Nirenberg inequality, Calderón–Zygmund decomposition (used in classical theory) are much more delicate. Paper [MPTT1] develops a completely new approach to prove natural tri-linear bi-parameter estimates on bi-parameter paraproducts, especially outside of Banach range. In [MPTT1] Journé's lemma [JLJ2] was used, but the approach did not generalize to multi-parameter paraproduct forms. This issue was resolved in [MPTT2], where a simplified method was used to address the multi-parameter paraproducts. But repeated bi-parameter commutators involve several bi-parameter paraproducts at once. Hence, the estimate of repeated bi-parameter commutators from below becomes tough: one needs to isolate those paraproducts one from another. And they do not like to be isolated.

One of the new feature of the multi-parameter theory is captured by several different definitions of BMO, see [FS], [Ch]. The necessity of those new effects was discovered first by Carleson [Car], see also [Tao].

The difficulties with multi-parameter theory was highlighted recently by two very different series of papers. One concerns itself with the Carleson measure and Carleson embedding on polydisc and on multi-tree, see, e.g. [AMPS18], [AHMV], [MPV], [MPVZ]. These papers, roughly speaking, are devoted to harmonic analysis (Carleson embedding in particular) on graphs with cycles. This theory is drastically different from the usual one, and it is much more difficult.

¹⁰⁰¹⁰ Mathematics Subject Classification. 42B100, 42B35, 47A100.

Key words and phrases. Bi-parameter Carleson embedding, bi-parameter Hankel operators, dyadic model.

IH is partially supported by the NSF an NSF Postdoc under Award No.1606270.

ST is partially supported by the NSF DMS 1856719.

AV is partially supported by the NSF grant DMS 1900268 and by Alexander von Humboldt foundation.

Another particularity of the multi-parameter theory is highlighted by the repeated commutator story. Lacey and Ferguson [FL] found the characterization of the symbols that give us bounded "small Hankel operators". It was a breakthrough article that gave a multi-parameter Nehari theorem and a long searched after factorization of bi-parameter Hardy space H^1 . It was exactly equivalent to a bi-parameter "repeated commutator characterization". Several papers followed where "bi-" was upgraded to "multi-", and where repeated commutation was performed with different classical singular integrals: [DP], [FL], [L], [LPPW], [LT].

There are many new and beautiful ideas in the above mentioned papers devoted to this subject. We also want to mention [OPS], where the authors use an argument inspired by Toeplitz operators to show the lower bounds. It assumes the lower norm for the Hilbert transform claimed in [FL], [LPPW] as a black box. But there is a problem with [FL] and with what followed.

That was a big breakthrough in multi-parameter theory. Unfortunately, [V] indicated a hole in the proofs of [FL], [L], and this circle of problems is still unsettled. See also Section 10 for the more detailed explanation.

We decided to simplify the problem to the "bare bones", and to consider repeated commutators with the simplest dyadic singular integral, namely, with the dyadic shift of order 1. The repeated commutator problem presented the same type of difficulty as in [FL] and subsequent papers. But we analyze these difficulties below. They become very clear (but still largely unsurmountable) in the dyadic case we consider below.

However, with an extra assumption on the symbol of repeated commutator we managed to get the expected answer, see Theorems 9.1, 9.2 at the end of this article. The expected answer is of course that the repeated commutator is bounded if and only if the symbol belongs to dyadic Chang-Fefferman BMO. If symbol does belong to dyadic Chang-Fefferman BMO then the repeated commutator is indeed bounded, see e.g. [LM].

In the opposite direction the only fully proved result so far is in [BP], but this paper requires that infinitely many commutators be bounded to conclude that the symbol is bounded.

As we already mentioned with an extra assumption on the symbol we can prove the necessity of dyadic Chang-Fefferman BMO (see Theorems 9.1, 9.2 at the end of this article). We list carefully what are the difficulties to be surmounted in the general case (without an extra assumption on the symbol), and what kind of combinatorial morass should be taken care of. This is left to a possibly more skillful reader.

2. Repeated commutator

Let \mathcal{D} be the usual dyadic lattice on the line and $\mathcal{D} \times \mathcal{D}$ be the family of dyadic rectangles on the plane. Haar functions are denote $\{h_I\}_{I \in \mathcal{D}}$. We consider $R = I \times J$ and $h_R = h_I \otimes h_J$. Let T_1 be the dyadic shift defined by on the functions of variable x:

$$T_1h_I = h_{I_{\perp}} - h_{I_{\perp}}$$
, if I is even; $T_1h_I = 0$, otherwise.

Let T_2 be the copy of T_1 , but acting on an independent variable y:

$$T_2h_J = h_{J_+} - h_{J_-}$$
, if J is even; $T_2h_J = 0$, otherwise.

Test functions are functions f = f(x, y), and $\mathbf{T} := T_1 \otimes T_2$.

We are interested in the commutator

$$\mathcal{T}^b := [T_2, [T_1, b]] = T_2(T_1b - bT_1) - (T_1b - bT_1)T_2 = T_2T_1b - T_2bT_1 - T_1bT_2 + bT_1T_2$$

and in characterization of its boundedness. The ultimate goal is to prove that its boundedness is equivalent to

$$b \in BMO^d$$
.

where BMO is understood in Chang–Fefferman definition. We always drop the superscript d as everything is dyadic below.

Simple commutator $[T_1, b]$ can be naturally split to 3 parts: D, π , Z parts. They correspond to different parts of the product operator $f \to b \cdot f$.

First D part follows.

$$(b,f) \to D(b,f) := \sum_{I} \sum_{I': I \subsetneq I'} \Delta_{I'} b \cdot \Delta_{I} f =$$
$$\sum_{I} (b, \mathbf{1}_{I}/|I|)(f, h_{I}) h_{I}.$$

Now compute commutator:

$$TD(b,f) - D(b,Tf) = \sum_{I} (b,\mathbf{1}_{I}/|I|)(f,h_{I})Th_{I} - \sum_{I} (b,\mathbf{1}_{I}/|I|)(Tf,h_{I})h_{I}.$$

Now compute commutator again:

$$TD(b,f) - D(b,Tf) = \sum_{I} \langle b \rangle_{I} (T(\Delta_{I}f) - \Delta_{I}(Tf)).$$

So let us consider

$$T(\Delta_I f) - \Delta_I (Tf)$$

for $f = h_K$. Let K = I and let I be even (meaning $|I| = 2^{-2k}$). Then $T(\Delta_I h_K) = Th_I = h_{I_+} - h_{I_-}$, and $\Delta_I(Th_K) = \Delta_I(Th_I) = 0$.

Let now K=I and let I be odd. Then $T(\Delta_I h_K)=Th_I=0$, and $\Delta_I(Th_K)=\Delta_I(Th_I)=\Delta_I(h_{I_+}-h_{I_-})=0$.

We conclude that TD(b, f) - D(b, Tf) will have terms:

$$\sum_{I \, even} \langle b \rangle_I(f, h_I)(h_{I_+} - h_{I_-}) = \sum_{\hat{I} \, even} \langle b \rangle_{\hat{I}}(f, h_{\hat{I}}) s(I, \hat{I}) h_I = \sum_{I \, odd} \langle b \rangle_{\hat{I}}(f, h_{\hat{I}}) s(I, \hat{I}) h_I,$$

where

(1)
$$s(I,\hat{I}) = \begin{cases} 1 & \text{if } I = (\hat{I})_{+} \\ -1 & \text{otherwise.} \end{cases}$$

There will be more terms. Denote by s(I) the dyadic sibling of I. Denote by \hat{I} the immediate predecessor of I. Let $K = \hat{I}$ and let \hat{I} be even. Then $\Delta_I h_K = 0$, so $T(\Delta_I h_K) = 0$, but $Th_K = Th_{\hat{I}} = h_{(\hat{I})_+} - h_{(\hat{I})_-}$ and so $Th_K = \pm (h_I - h_{s(I)})$, and, thus, $\Delta_I Th_K = \pm h_I$. So for $f = h_K$ with $K = \hat{I}$ and \hat{I} being even we have

On
$$I: T(\Delta_I f) - \Delta_I(T f) = -h_I$$
, if $I = (\hat{I})_+$; $T(\Delta_I f) - \Delta_I(T f) = +h_I$ if $I = (\hat{I})_-$.

We conclude that TD(b, f) - D(b, Tf) will have also the following terms:

$$-\sum_{\hat{I} \, even} \langle b \rangle_I(f, h_{\hat{I}}) s(I, \hat{I}) h_I = -\sum_{I \, odd} \langle b \rangle_I(f, h_{\hat{I}}) s(I, \hat{I}) h_I.$$

All other h_K give zero contribution.

We get finally that

(2)

$$(T_1D_b - D_bT_1)(f) = T_1D(b, f) - D(b, T_1f) = \sum_{I \text{ odd}} (\langle b \rangle_{\hat{I}} - \langle b \rangle_I)(f, h_{\hat{I}})s(I, \hat{I})h_I = \sum_{I \text{ odd}} \frac{-1}{|\hat{I}|^{1/2}}(b, h_{\hat{I}})(f, h_{\hat{I}})h_I.$$

Now we need to commute with T_2 :

$$[T_2, [T_1, D_b]](f) = \sum_{I \in \mathcal{A}} \frac{h_I}{|I|^{1/2}} \Big(T_2 \{ (b, h_{\hat{I}})(f, h_{\hat{I}}) \} - (b, h_{\hat{I}})(T_2 f, h_{\hat{I}}) \Big) .$$

We use now notations: $b_I := (b, h_I)$. Then we rewrite the previous display as follows:

$$[T_2, [T_1, D_b]](f) = \sum_{I \text{ odd}} \frac{h_I}{|I|^{1/2}} (T_2 b_{\hat{I}} - b_{\hat{I}} T_2)(f_{\hat{I}}) .$$

Now multiplication in the second variable $f_{\hat{I}} \to b_{\hat{I}} \cdot f_{\hat{I}}$ can be decomposed to D, π, Z parts.

2.1. DD part of repeated commutator.

$$DD = D_{2}([T_{2}, [T_{1}, D_{b}]](f)) = \sum_{Iodd} \frac{h_{I}}{|I|^{1/2}} D_{part}(T_{2}b_{\hat{I}} - b_{\hat{I}}T_{2})(f_{\hat{I}}) = \sum_{Iodd} \frac{h_{I}}{|I|^{1/2}} \sum_{Jodd} \frac{h_{J}}{|J|^{1/2}} (b_{\hat{I}}, h_{\hat{J}})(f_{\hat{I}}, h_{\hat{I}}) = \sum_{I,Jodd,odd} (b, h_{\hat{I}} \otimes h_{\hat{J}})(f, h_{\hat{I}} \otimes h_{\hat{J}}) \frac{h_{I}(x) \cdot h_{J}(y)}{|I|^{1/2}|J|^{1/2}} = \sum_{Rodd,odd} (b, h_{\hat{R}})(f, h_{\hat{R}}) \frac{h_{R}}{|R|^{1/2}}.$$

2.2. $D\pi$ part of repeated commutator.

$$D\pi = \pi_2([T_2, [T_1, D_b]](f)) = \sum_{Iodd} \frac{h_I}{|I|^{1/2}} \pi_{part}(T_2 b_{\hat{I}} - b_{\hat{I}} T_2)(f_{\hat{I}}) = \sum_{Iodd} \frac{h_I}{|I|^{1/2}} (T_2 \pi_{b_{\hat{I}}} - \pi_{b_{\hat{I}}} T_2)(f_{\hat{I}})$$

Let us recall what is π_b .

$$(b_{\hat{I}}, f_{\hat{I}}) \to \pi_{b_{\hat{I}}} f_{\hat{I}} := \sum_{J} \sum_{J': J \subsetneq J'} \Delta_{J} b_{\hat{I}} \cdot \Delta_{J'} f_{\hat{I}} = \sum_{J} \langle f_{\hat{I}} \rangle_{J} (b_{\hat{I}}, h_{J}) h_{J}.$$

Now compute commutator (below $\tilde{\mathbf{1}}_J := \mathbf{1}_J/|J|$).

(3)
$$D\pi = \sum_{Iodd} \frac{h_I}{|I|^{1/2}} \Big(T_2 \pi_{b_{\hat{I}}} f_{\hat{I}} - \pi_{b_{\hat{I}}} (T_2 f_{\hat{I}}) \Big) =$$

$$\sum_{Iodd} \sum_{Jeven} (b, h_{\hat{I}} \otimes h_J) (f, h_{\hat{I}} \otimes \tilde{\mathbf{1}}_J) \frac{h_I}{|I|^{1/2}} \otimes T_2 h_J -$$

$$\sum_{Iodd} \sum_{J} (b, h_{\hat{I}} \otimes h_J) (T_2 f, h_{\hat{I}} \otimes \tilde{\mathbf{1}}_J) \frac{h_I}{|I|^{1/2}} \otimes h_J.$$

Let us use the following notations for given I, J:

(4)
$$R = I \times J, \hat{R} = \hat{I} \times \hat{J}, \hat{R}_1 = \hat{I} \times J, \hat{R}_2 = I \times \hat{J}, h_R = h_I \otimes h_J$$

(5)
$$\tilde{\mathbf{1}}_R = \tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J, h\mathbf{1}_R = h_I \otimes \tilde{\mathbf{1}}_J, \mathbf{1}h_R = \tilde{\mathbf{1}}_I \otimes h_J$$

2.3. DZ part of repeated commutator.

$$DZ = Z_2([T_2, [T_1, D_b]](f)) = \sum_{Iodd} \frac{h_I}{|I|^{1/2}} Z_{part}(T_2 b_{\hat{I}} - b_{\hat{I}} T_2)(f_{\hat{I}}) = \sum_{Iodd} \frac{h_I}{|I|^{1/2}} (T_2 Z_{b_{\hat{I}}} - Z_{b_{\hat{I}}} T_2)(f_{\hat{I}})$$

Let us recall what is Z_b .

$$(b_{\hat{I}}, f_{\hat{I}}) \to Z_{b_{\hat{I}}} f_{\hat{I}} := \sum_{J} (b_{\hat{I}}, h_{J}) (f_{\hat{I}}, h_{J}) \tilde{\mathbf{1}}_{J} \,.$$

Therefore,

(6)
$$DZ = \sum_{I \text{ odd}} \frac{h_I}{|I|^{1/2}} \Big(T_2 Z_{b_{\hat{I}}}(f_{\hat{I}}) - Z_{b_{\hat{I}}}(T_2 f_{\hat{I}}) \Big) =$$

$$\sum_{I \text{ odd}} \sum_{J \text{ even}} (b, h_{\hat{I}} \otimes h_J) (f, h_{\hat{I}} \otimes h_J) \frac{h_I}{|I|^{1/2}} \otimes T_2 \tilde{\mathbf{1}}_J -$$

$$\sum_{I \text{ odd}} \sum_{J} (b, h_{\hat{I}} \otimes h_J) (T_2 f, h_{\hat{I}} \otimes h_J) \frac{h_I}{|I|^{1/2}} \otimes \tilde{\mathbf{1}}_J.$$

This finishes $DD, D\pi, DZ$ parts of repeated commutator.

Now we need to write $\pi D, Z\pi$ parts—they are just symmetric with respect to exchange the first and the second variables. But also we need ZZ part, $\pi Z, Z\pi$ parts, and the most dangerous $\pi\pi$ part.

2.4. π -parts. Here it is.

$$T_2(T_1\pi_b f - \pi_b(T_1 f)) - (T_1\pi_b(T_2 f) - \pi_b(T_1 T_2 f)) =$$

$$= (T_2T_1\pi_b f - T_1\pi_b(T_2 f)) - (T_2\pi_b(T_1 f) - \pi_b(T_1 T_2 f)) =$$

$$\sum_I (T_2 b_I - b_I T_2) \langle f \rangle_I T_1 h_I - \sum_I (T_2 b_I - b_I T_2) \langle T_1 f \rangle_I h_I.$$

Its πD part is obtained by calculating D parts of commutator $(T_2b_I - b_IT_2)$. So as before

(7)
$$\pi D = \sum_{Ieven} \sum_{Jodd} (b, h_I \otimes h_{\hat{J}}) (f, \tilde{\mathbf{1}}_I \otimes h_{\hat{J}}) T_1 h_I \otimes \frac{h_J}{|J|^{1/2}} - \sum_{I} \sum_{Jodd} (b, h_I \otimes h_{\hat{J}}) (T_1 f, \tilde{\mathbf{1}}_I \otimes h_J \otimes) h_I \otimes \frac{h_J}{|J|^{1/2}}.$$

We see that (7) is symmetric to (3), as it should be.

(8)
$$\pi Z = \sum_{Ieven} \sum_{J} (b, h_I \otimes h_J) (f, \tilde{\mathbf{1}}_I \otimes h_J) T_1 h_I \otimes T_2 \tilde{\mathbf{1}}_J - \sum_{Ieven} \sum_{J} (b, h_I \otimes h_J) (T_2 f, \tilde{\mathbf{1}}_I \otimes h_J) T_1 h_I \otimes \tilde{\mathbf{1}}_J - \sum_{Ieven} \sum_{J} (b, h_I \otimes h_J) (T_1 f, \tilde{\mathbf{1}}_I \otimes h_J) h_I \otimes T_2 \tilde{\mathbf{1}}_J + \sum_{Ieven} \sum_{J} (b, h_I \otimes h_J) (T_1 T_2 f, \tilde{\mathbf{1}}_I \otimes h_J) h_I \otimes \tilde{\mathbf{1}}_J.$$

Now the most dangerous $\pi\pi$ part of repeated commutator follows:

(9)
$$\pi\pi = \sum_{I,Jeven,even} (b, h_I \otimes h_J)(f, \tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J) T_1 h_I \otimes T_2 h_J - \sum_{Iany,Jeven} (b, h_I \otimes h_J)(T_1 f, \tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J) h_I \otimes T_2 h_J - \sum_{Ieven,Jany} (b, h_I \otimes h_J)(T_2 f, \tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J) T_1 h_I \otimes h_J + \sum_{Iany,Jany} (b, h_I \otimes h_J)(T_1 T_2 f, \tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J) h_I \otimes h_J.$$

2.5. **Z-parts.**

(10)
$$ZD = \sum_{I} \sum_{Jodd} (b, h_I \otimes h_{\hat{J}}) (f, h_I \otimes h_{\hat{J}}) T_1 \tilde{\mathbf{1}}_I \otimes h_J / |J|^{1/2} - \sum_{I} \sum_{Jodd} (b, h_I \otimes h_{\hat{J}}) (T_1 f, h_I \otimes h_{\hat{J}}) \tilde{\mathbf{1}}_I \otimes h_J / J|^{1/2}.$$

(11)
$$Z\pi = \sum_{Ieven} \sum_{J} (b, h_{I} \otimes h_{J})(f, h_{I} \otimes \tilde{\mathbf{1}}_{J})T_{1}\tilde{\mathbf{1}}_{I} \otimes T_{2}h_{J} - \sum_{Ieven} \sum_{J} (b, h_{I} \otimes h_{J})(T_{1}f, h_{I} \otimes \tilde{\mathbf{1}}_{J})\tilde{\mathbf{1}}_{I} \otimes T_{2}h_{J} - \sum_{Ieven} \sum_{J} (b, h_{I} \otimes h_{J})(T_{2}f, h_{I} \otimes \tilde{\mathbf{1}}_{J})T_{1}\tilde{\mathbf{1}}_{I} \otimes h_{J} + \sum_{Ieven} \sum_{J} (b, h_{I} \otimes h_{J})(T_{1}T_{2}f, h_{I} \otimes \tilde{\mathbf{1}}_{J})\tilde{\mathbf{1}}_{I} \otimes h_{J}.$$

$$ZZ = \sum_{Ieven} \sum_{J} (b, h_{I} \otimes h_{J})(f, h_{I} \otimes h_{J})T_{1}\tilde{\mathbf{1}}_{I} \otimes T_{2}\tilde{\mathbf{1}}_{J} - \sum_{Ieven} \sum_{J} (b, h_{I} \otimes h_{J})(T_{1}f, h_{I} \otimes h_{J})\tilde{\mathbf{1}}_{I} \otimes T_{2}\tilde{\mathbf{1}}_{J} - \sum_{Ieven} \sum_{J} (b, h_{I} \otimes h_{J})(T_{2}f, h_{I} \otimes h_{J})T_{1}\tilde{\mathbf{1}}_{J} \otimes \tilde{\mathbf{1}}_{J} + \sum_{Ieven} \sum_{J} (b, h_{I} \otimes h_{J})(T_{1}T_{2}f, h_{I} \otimes h_{J})\tilde{\mathbf{1}}_{J} \otimes \tilde{\mathbf{1}}_{J}.$$

3. Necessity of BMO_r

In this section we show that if \mathcal{T}^b is bounded, then $b \in BMO_r$. To do this, we test \mathcal{T}^b on a Haar function $h_{I''} \otimes h_{J''}$. This yields 25 operators, which we list below. Most paraproducts in Section 2 (like $\pi\pi$ for example) yield 2-4 terms, so names such as $\pi\pi 1$ below simply indicate that the term comes from $\pi\pi$. The operators below also have various restrictions on the parity of I'' and J'', i.e. most only appear for a certain combination of parities. This is indicated in parentheses next to each operator.

The operators are grouped by simplicity: Group I are the simplest operators containing finitely many terms, Groups IIa and IIb are product operators, and lastly the four operators in Group III are the most difficult. They are also color-coded...

Group I: Simple operators

$$\begin{array}{c} \boxed{\mathbf{DZ2}} \left(I'' \text{ even, } J'' \text{ even} \right) \\ (b, h_{I''} \otimes h_{J''_+}) \frac{1}{\sqrt{|I''|}} (h_{I''_+} + h_{I''_-}) \tilde{\mathbf{1}}_{J''_+} \\ - (b, h_{I''} \otimes h_{J''_-}) \frac{1}{\sqrt{|I''|}} (h_{I''_+} + h_{I''_-}) \tilde{\mathbf{1}}_{J''_-} \end{array}$$

$$\boxed{\mathbf{ZD2}} (I'' \text{ even, } J'' \text{ even}) \\
(b, h_{I''_{+}} \otimes h_{J''}) \frac{1}{\sqrt{|J''|}} \tilde{1}_{I''_{+}} (h_{J''_{+}} + h_{J''_{-}}) \\
-(b, h_{I''_{-}} \otimes h_{J''}) \frac{1}{\sqrt{|J''|}} \tilde{1}_{I''_{-}} (h_{J''_{+}} + h_{J''_{-}})$$

$$\begin{split} & \mathbf{ZZ4} \ (I'' \text{ even}, \, J'' \text{ even}) \\ & (b, h_{I''_+} \otimes h_{J''_+}) \tilde{\mathbf{1}}_{I''_+} \tilde{\mathbf{1}}_{J''_+} - (b, h_{I''_+} \otimes h_{J''_-}) \tilde{\mathbf{1}}_{I''_+} \tilde{\mathbf{1}}_{J''_-} \\ & - (b, h_{I''_-} \otimes h_{J''_+}) \tilde{\mathbf{1}}_{I''_-} \tilde{\mathbf{1}}_{J''_+} + (b, h_{I''_-} \otimes h_{J''_-}) \tilde{\mathbf{1}}_{I''_-} \tilde{\mathbf{1}}_{J''_-} \end{split}$$

ZD1 (
$$J''$$
 even)
$$-(b, h_{I''} \otimes h_{J''}) \frac{1}{\sqrt{|J''|}} (T_1 \tilde{\mathbf{1}}_{I''}) (h_{J''_+} + h_{J''_-})$$

$$\boxed{ \mathbf{DZ1} \ (I'' \ \mathbf{even}) } \\ - (b, h_{I''} \otimes h_{J''}) \frac{1}{\sqrt{|I''|}} (h_{I''_+} + h_{I''_-}) (T_2 \tilde{\mathbf{1}}_{J''})$$

$$\begin{split} & \boxed{\mathbf{ZZ2}} (J'' \text{ even}) \\ & - (b, h_{I''} \otimes h_{J''_+}) (T_1 \tilde{\mathbf{1}}_{I''}) \tilde{\mathbf{1}}_{J''_+} \\ & + (b, h_{I''} \otimes h_{J''_-}) (T_1 \tilde{\mathbf{1}}_{I''}) \tilde{\mathbf{1}}_{J''_-} \end{split}$$

ZZ1 (no restrictions) $(b, h_{I''} \otimes h_{J''})(T_1\tilde{\mathbf{1}}_{I''})(T_2\tilde{\mathbf{1}}_{J''})$

Group IIa: Product operators, summation over I

$$\boxed{\pi \mathbf{D2}} (I'' \text{ even, } J'' \text{ even)}$$

$$\boxed{\sum_{I \subsetneq I''_{+}} (b, h_{I} \otimes h_{J''}) h_{I''_{+}}(I) h_{I}} \frac{h_{J''_{+}} + h_{J''_{-}}}{\sqrt{|J''|}} - \left[\sum_{I \subsetneq I''_{-}} (b, h_{I} \otimes h_{J''}) h_{I''_{-}}(I) h_{I} \right] \frac{h_{J''_{+}} + h_{J''_{-}}}{\sqrt{|J''|}}$$

$$egin{aligned} \overline{\pi \mathbf{Z2}} & (I'' ext{ even}, \, J'' ext{ even}) \ & \left[\sum_{I \subsetneq I''_+} (b, h_I \otimes h_{J''_+}) h_{I''_+}(I) h_I
ight] ilde{\mathbf{1}}_{J''_+} - \left[\sum_{I \subsetneq I''_+} (b, h_I \otimes h_{J''_-}) h_{I''_+}(I) h_I
ight] ilde{\mathbf{1}}_{J''_-} \ & - \left[\sum_{I \subsetneq I''_-} (b, h_I \otimes h_{J''_+}) h_{I''_-}(I) h_I
ight] ilde{\mathbf{1}}_{J''_-} + \left[\sum_{I \subsetneq I''_-} (b, h_I \otimes h_{J''_-}) h_{I''_-}(I) h_I
ight] ilde{\mathbf{1}}_{J''_-} \end{aligned}$$

$$egin{aligned} \pi \mathbf{D} \mathbf{1} & (J'' ext{ even}) \ - & \left[\sum_{\substack{I \subsetneq I'' \ I ext{ even}}} (b, h_I \otimes h_{J''}) h_{I''}(I) (T_1 h_I)
ight] rac{h_{J''_+} + h_{J''_-}}{\sqrt{|J''|_-}} \end{aligned}$$

$$\begin{bmatrix} & & & \\ & & & \\ & & & \\ & & - \left[\sum_{I \subsetneq I''_+} (b, h_I \otimes h_{J''}) h_{I''_+}(I) h_I \right] (T_2 \tilde{\mathbf{1}}_{J''}) \\ & & + \left[\sum_{I \subsetneq I''_-} (b, h_I \otimes h_{J''}) h_{I''_-}(I) h_I \right] (T_2 \tilde{\mathbf{1}}_{J''}) \end{bmatrix}$$

$$egin{aligned} \pi \mathbf{Z4} & (J'' ext{ even}) \ & - \left[\sum_{\substack{I \subseteq I'' \ I ext{ even}}} (b, h_I \otimes h_{J''_+}) h_{I''}(I)(T_1 h_I)
ight] ilde{\mathbf{1}}_{J''_+} \ & + \left[\sum_{\substack{I \subseteq I'' \ I ext{ even}}} (b, h_I \otimes h_{J''_-}) h_{I''}(I)(T_1 h_I)
ight] ilde{\mathbf{1}}_{J''_-} \end{aligned}$$

$$egin{aligned} \pi \mathbf{Z3} & ext{ (no restrictions)} \ & \left[\sum_{\substack{I \subseteq I'' \ I ext{ even}}} (b, h_I \otimes h_{J''}) h_{I''}(I)(T_1 h_I)
ight] (T_2 ilde{\mathbf{1}}_{J''}). \end{aligned}$$

Group IIb: Product operators, summation over J

$$\boxed{ \begin{array}{c} \mathbf{D}\pi\mathbf{2} \left(I'' \text{ even, } J'' \text{ even} \right) \\ \\ \frac{h_{I''_+} + h_{I''_-}}{\sqrt{|I''|}} \left[\sum_{J \subsetneq J''_+} (b, h_{I''} \otimes h_J) h_{J''_+}(J) h_J \right] - \frac{h_{I''_+} + h_{I''_-}}{\sqrt{|I''|}} \left[\sum_{J \subsetneq J''_-} (b, h_{I''} \otimes h_J) h_{J''_-}(J) h_J \right] \end{array} }$$

$$\begin{split} \mathbf{\overline{Z}\pi\mathbf{4}} & \left(I'' \text{ even, } J'' \text{ even}\right) \\ & \tilde{\mathbf{1}}_{I_{+}''} \left[\sum_{J \subsetneq J_{+}''} (b, h_{I_{+}''} \otimes h_{J}) h_{J_{+}''}(J) h_{J} \right] - \tilde{\mathbf{1}}_{I_{-}''} \left[\sum_{J \subsetneq J_{+}''} (b, h_{I_{-}''} \otimes h_{J}) h_{J_{+}''}(J) h_{J} \right] \\ & - \tilde{\mathbf{1}}_{I_{+}''} \left[\sum_{J \subsetneq J_{-}''} (b, h_{I_{+}''} \otimes h_{J}) h_{J_{-}''}(J) h_{J} \right] + \tilde{\mathbf{1}}_{I_{-}''} \left[\sum_{J \subsetneq J_{-}''} (b, h_{I_{-}''} \otimes h_{J}) h_{J_{-}''}(J) h_{J} \right] \end{split}$$

$$-\frac{h_{I_+''}+h_{I_-''}}{\sqrt{|I''|}}\left[\sum_{\substack{J\subseteq J''\\J\text{ even}}}(b,h_{I''}\otimes h_J)h_{J''}(J)(T_2h_J)\right]$$

$$-(T_1\tilde{\mathbf{1}}_{I''})\left[\sum_{\substack{J\subseteq J''\\J\subseteq J_+''}}(b,h_{I''}\otimes h_J)h_{J_+''}(J)h_J\right]$$

$$+(T_1\tilde{\mathbf{1}}_{I''})\left[\sum_{\substack{J\subseteq J''\\J\subseteq J''}}(b,h_{I''}\otimes h_J)h_{J_-''}(J)h_J\right]$$

 $\mathbf{D}\pi\mathbf{1} \mid (I'' \text{ even})$

Group III: The difficult operators

$$\boxed{\pi\pi\mathbf{4} \left(I'' \text{ even, } J'' \text{ even}\right)}$$

$$\sum_{\substack{I\subsetneq I''_+\\ J\subsetneq J''_+\\ J\subsetneq J''_-\\ J\subsetneq J''_-}} (b,h_I\otimes h_J)h_{I''_+}(I)h_{J''_+}(J)h_I\otimes h_J - \sum_{\substack{I\subsetneq I''_+\\ J\subsetneq J''_-\\ J\subsetneq J''_-\\ J\subsetneq J''_-}} (b,h_I\otimes h_J)h_{I''_-}(I)h_{J''_+}(J)h_I\otimes h_J + \sum_{\substack{I\subsetneq I''_-\\ J\subsetneq J''_-\\ J\subsetneq J''_-\\ J\subsetneq J''_-}} (b,h_I\otimes h_J)h_{I''_-}(I)h_{J''_-}(J)h_I\otimes h_J + \sum_{\substack{I\subsetneq I''_-\\ J\subsetneq J''_-\\ J\subsetneq J''_-\\ J\subsetneq J''_-}} (b,h_I\otimes h_J)h_{I''_-}(I)h_{J''_-}(J)h_I\otimes h_J$$

$$\boxed{\pi\pi\mathbf{3}} (I'' \text{ even})$$

$$-\sum_{\substack{I \subsetneq I''_+\\ J \text{ even}, J \subsetneq J''}} (b, h_I \otimes h_J) h_{I''_+}(I) h_{J''}(J) h_I \otimes T_2 h_J + \sum_{\substack{I \subsetneq I''_-\\ J \text{ even}, J \subsetneq J''}} (b, h_I \otimes h_J) h_{I''_-}(I) h_{J''}(J) h_I \otimes T_2 h_J$$

$$\begin{array}{l} \boxed{\pi\pi\mathbf{2}} (J'' \text{ even}) \\ -\sum_{\substack{I \text{ even}, I \subseteq I'' \\ J \subseteq J''_+}} (b, h_I \otimes h_J) h_{I''}(I) h_{J''_+}(J) T_1 h_I \otimes h_J + \sum_{\substack{I \text{ even}, I \subseteq I'' \\ J \subseteq J''_-}} (b, h_I \otimes h_J) h_{I''}(I) h_{J''_-}(J) T_1 h_I \otimes h_J \end{array}$$

$$[\pi\pi\mathbf{1}]$$
 (no restrictions)
$$-\sum_{\substack{I \text{ even}, I \subseteq I'' \\ J \text{ even}, J \subseteq J''}} (b, h_I \otimes h_J) h_{I''}(I) h_{J''}(J) T_1 h_I \otimes T_2 h_J$$

The lemma below will bound most of these operators:

Lemma 1 If \mathcal{T}^b is L^2 -bounded, then the (one-parameter) commutators

$$\frac{1}{\sqrt{|J|}}[T_1, b_J] \quad and \quad \frac{1}{\sqrt{|I|}}[T_2, b_I]$$

are uniformly bounded, for any $I, J \in \mathcal{D}$. Consequently,

(13)
$$\sum_{I' \subset I} \left| (b, h_{I'} \otimes h_J) \right|^2 \lesssim |I||J|,$$

and

(14)
$$\sum_{J'\subset J} |(b, h_I \otimes h_{J'})|^2 \lesssim |I||J|,$$

for all $I, J \in \mathcal{D}$.

Proof. We prove the result in the first parameter x, the other being symmetric. We test \mathcal{T}^b on a function $f(x,y) = p(x) \otimes h_{J''}(y)$, where $p \in L^2(\mathbb{R})$ and $J'' \in \mathcal{D}$. Note that if \mathcal{T}^b is bounded, $\|\mathcal{T}^b f\|_2^2 \lesssim \|f\|_2^2 = \|p\|_2^2$. Computing $\mathcal{T}^b(p \otimes h_{J''})$ yields five operators: two with no restrictions on J'':

$$\begin{array}{lll} \mathtt{P_1} & = & \sum_{\substack{J \text{ even,} \\ J \subsetneq J''}} \left([T_1, b_J] p \right) h_{J''}(J) (T_2 h_J), \\ \\ \mathtt{P_2} & = & \left([T_1, b_{J''}] p \right) (T_2 \tilde{\mathbf{1}}_{J''}), \end{array}$$

and three which are only non-zero if J'' is even:

$$\begin{array}{lll} \mathtt{P_3} & = & -\left([T_1,b_{J''}]p\right)\frac{1}{\sqrt{|J''|}}(h_{J''_-}+h_{J''_+}), \\ \\ \mathtt{P_4} & = & -\sum_{J\subsetneq J''_+}\left([T_1,b_J]p\right)h_{J''_+}(J)h_J + \sum_{J\subsetneq J''_-}\left([T_1,b_J]p\right)h_{J''_-}(J)h_J, \\ \\ \mathtt{P_5} & = & -\left([T_1,b_{J''_-}]p\right)\tilde{\mathbf{1}}_{J''_-} + \left([T_1,b_{J''_-}]p\right)\tilde{\mathbf{1}}_{J''_-}. \end{array}$$

Suppose first that J'' is *even*. In this case, we have all five operators above. We look at the Haar coefficients of each:

$$(\mathtt{P_1}, h_{I_0} \otimes h_{J_0}) = \left\{ \begin{array}{ll} ([T_1, b_{J_0}] p, h_{I_0}) \, s(J_0, \hat{J}_0) h_{J''}(\hat{J}_0), & \text{if J_0 is odd, and $\hat{J}_0 \subsetneq J''$} \\ 0, & \text{otherwise.} \end{array} \right.$$

$$(\mathtt{P_2}, h_{I_0} \otimes h_{J_0}) = \left\{ \begin{array}{ll} ([T_1, b_{J''}] p, h_{I_0}) \, s(J_0, \hat{J}_0) h_{\hat{J}_0}(J''), & \text{if J_0 is odd, and $\hat{J}_0 \supsetneq J''$} \\ 0, & \text{otherwise.} \end{array} \right.$$

Since J'' is even, the inclusion conditions on J_0 are the same as $J_0 \subsetneq J''_+$ for P_1 , and $J_0 \supsetneq J''$ for P_2 .

$$(P_3, h_{I_0} \otimes h_{J_0}) = \begin{cases} -([T_1, b_{J''}]p, h_{I_0}) \frac{1}{\sqrt{|J''|}}, & \text{if } J_0 = J''_- \text{ or } J_0 = J''_+\\ 0, & \text{otherwise.} \end{cases}$$

$$(\mathtt{P_4}, h_{I_0} \otimes h_{J_0}) = \left\{ \begin{array}{ll} - \left([T_1, b_{J_0}] p, h_{I_0} \right) h_{J_+''}(J_0), & \text{if } J_0 \subsetneq J_+'' \\ + \left([T_1, b_{J_0}] p, h_{I_0} \right) h_{J_-''}(J_0), & \text{if } J_0 \subsetneq J_-'' \\ 0, & \text{otherwise.} \end{array} \right.$$

$$(\mathsf{P_5}, h_{I_0} \otimes h_{J_0}) = \left\{ \begin{array}{l} -\left([T_1, b_{J''_+}]p, h_{I_0}\right) h_{J_0}(J''_+) + \left([T_1, b_{J''_-}]p, h_{I_0}\right) h_{J_0}(J''_-), & \text{if } J_0 \supseteq J'' \\ 0, & \text{otherwise.} \end{array} \right.$$

Note that for $P_1 + P_2 + P_3$, the second component Haar functions run through all the *odd* dyadic intervals intersecting J''. For $P_4 + P_5$, the second component Haar functions run through *all* dyadic intervals intersecting J'', except J''_+ . Pick the components of P_4 and P_5 with even J on the second coordinate:

$$\begin{split} \mathbf{P}_{\text{even}} & := & -\sum_{\substack{J \subseteq J''_+ \\ J \text{ even}}} \left([T_1, b_J] p \right) h_{J''_+}(J) h_J + \sum_{\substack{J \subseteq J''_- \\ J \text{ even}}} \left([T_1, b_J] p \right) h_{J''_-}(J) h_J \\ & - \left([T_1, b_{J''_+}] p \right) \sum_{\substack{J \supseteq J''_- \\ L \text{ even}}} h_J(J''_+) h_J + \left([T_1, b_{J''_-}] p \right) \sum_{\substack{J \supseteq J''_- \\ L \text{ even}}} h_J(J''_-) h_J. \end{split}$$

Then $\mathcal{T}^b(p \otimes h_{J''}) - P_{\text{even}}$ and P_{even} are *orthogonal*, so if \mathcal{T}^b is bounded then these components must be individually bounded. In particular, P_{even} is bounded. Clearly

$$\begin{aligned} \|\mathbf{P}_{\text{even}}\|_{2}^{2} &\geq \sum_{\substack{J \subseteq J''_{+} \\ J \text{ even}}} \sum_{I_{0}} \left([T_{1}, b_{J}]p, h_{I_{0}} \right)^{2} \frac{1}{|J''_{+}|} + \sum_{\substack{J \subseteq J''_{-} \\ J \text{ even}}} \sum_{I_{0}} \left([T_{1}, b_{J}]p, h_{I_{0}} \right)^{2} \frac{1}{|J''_{-}|} \\ &= \frac{2}{|J''|} \sum_{\substack{J \subseteq J''_{-} \\ J \text{ even}}} \|[T_{1}, b_{J}]p\|_{2}^{2}, \end{aligned}$$

giving us that

(15)
$$\sum_{\substack{J \subsetneq J'' \\ J}} \|[T_1, b_J]p\|_2^2 \lesssim |J''| \|p\|_2^2, \text{ for all } J'' \text{ even.}$$

Now suppose J'' is odd, so we only have the operators P_1 and P_2 . As before,

$$(\mathtt{P_1}, h_{I_0} \otimes h_{J_0}) = \left\{ \begin{array}{cc} ([T_1, b_{J_0}] p, h_{I_0}) \, s(J_0, \hat{J}_0) h_{J''}(\hat{J}_0), & \text{if } J_0 \text{ is odd, and } J_0 \subsetneq J'' \\ 0, & \text{otherwise.} \end{array} \right.$$

$$(\mathtt{P}_2, h_{I_0} \otimes h_{J_0}) = \left\{ \begin{array}{cc} ([T_1, b_{J''}]p, h_{I_0}) \, s(J_0, \hat{J_0}) h_{\hat{J_0}}(J''), & \text{if } J_0 \text{ is odd, and } J_0 \supsetneq J'' \\ 0, & \text{otherwise.} \end{array} \right.$$

Obviously P_1 and P_2 are orthogonal, so $\|\mathcal{T}^b(p \otimes h_{J''})\|_2^2 = \|P_1\|^2 + \|P_2\|_2^2$ for odd J''. From $\|P_1\|_2^2 \leq \|\mathcal{T}^b\|_2^2$, it is easy to see that

(16)
$$\sum_{J \subseteq J'' \atop J \subseteq J''} \|[T_1, b_J]p\|_2^2 \lesssim |J''| \|p\|_2^2, \text{ for all } J'' \text{ odd.}$$

From (15) and (16), it is obvious that, if \mathcal{T}^b is bounded, then

(17)
$$\sum_{J \subset J''} \|[T_1, b_J]\|_2^2 \lesssim |J''|, \text{ for all } J'' \in \mathcal{D}.$$

This is a much stronger assertion than the Lemma needs, and immediately implies (13) using the one-parameter implication:

$$\frac{1}{\sqrt{|J|}}[T_1, b_J] \text{ is } L^2\text{-bounded} \Rightarrow \frac{1}{\sqrt{|J|}}b_J \in BMO(\mathbb{R}) \Rightarrow \sum_{I' \subset I} |(b, h_{I'} \otimes h_J)|^2 \lesssim |I||J|.$$

Theorem 3.1 If \mathcal{T}^b is bounded, then $b \in BMO_r$.

Proof. The results in Lemma 1 immediately show that all operators in Groups I and II are a priori bounded and can be disregarded. What remains are the four operators in Group III.

Case 1: I'' and J'' are both odd. In this case, we only have the operator $\pi\pi 1$, which we rewrite as

$$\pi\pi 1 = -\sum_{\substack{I \text{ odd}, I \subseteq I'' \\ J \text{ odd}, J \subseteq J''}} (b, h_{\hat{I}} \otimes h_{\hat{J}}) h_{I''}(\hat{I}) h_{J''}(\hat{J}) s(I, \hat{I}) s(J, \hat{J}) h_{I} \otimes h_{J}.$$

So boundedness of \mathcal{T}^b in this case reduces to just this $\pi\pi 1$. Computing its norm gives us

$$\|\pi\pi 1\|_{2}^{2} = \frac{1}{|I''||J''|} \sum_{\substack{I \text{ odd}, I \subseteq I'' \\ J \text{ odd}, J \subseteq J''}} |(b, h_{\hat{I}} \otimes h_{\hat{J}})|^{2} = \frac{1}{|I''||J''|} \sum_{\substack{I \text{ even}, I \subseteq I'' \\ J \text{ even}, J \subseteq J''}} |(b, h_{I} \otimes h_{J})|^{2} \lesssim C,$$

and from here immediately

(18)
$$\sum_{\substack{I \text{ even}, I \subseteq I_0 \\ J \text{ even}, J \subseteq J_0}} |(b, h_I \otimes h_J)|^2 \lesssim |I_0||J_0|, \text{ for all } I_0, J_0 \in \mathcal{D}.$$

Case 2: I'' is odd and J'' is even. In this case we have two operators, $\pi\pi 1$ and $\pi\pi 2$. But now we know from (18) that $\pi\pi 1$ is already bounded, so we only need to look at $\pi\pi 2$:

$$\pi\pi2 = -\sum_{\substack{I \text{ odd}, I \subseteq I'' \\ J \subseteq J_{\perp}^{J''}}} (b, h_{\hat{I}} \otimes h_J) h_{I''}(\hat{I}) h_{J_{+}^{\prime\prime}}(J) s(I, \hat{I}) h_I \otimes h_J + \sum_{\substack{I \text{ odd}, I \subseteq I'' \\ J \subseteq J_{\perp}^{J''}}} (b, h_{\hat{I}} \otimes h_J) h_{I''}(\hat{I}) h_{J_{-}^{\prime\prime}}(J) s(I, \hat{I}) h_I \otimes h_J.$$

Separate $\pi\pi 2$ into $\pi\pi 2_{\text{odd}}$, where we only sum over $odd\ J \subsetneq J''_{\pm}$, and $\pi\pi 2_{\text{even}}$, where we only sum over $even\ J \subsetneq J''_{\pm}$. These are orthogonal, so they are individually bounded. Now

$$\|\pi\pi 2_{\text{odd}}\|_{2}^{2} = \frac{2}{|I''||J''|} \sum_{\substack{I \text{ even}, I \subseteq I'' \\ J \text{ odd}, J \subseteq J''_{\perp}}} |(b, h_{I} \otimes h_{J})|^{2} \lesssim C.$$

From here, it follows easily that

(19)
$$\sum_{\substack{I \text{ even}, I \subseteq I_0 \\ J \text{ odd}, J \subseteq J_0}} |(b, h_I \otimes h_J)|^2 \lesssim |I_0||J_0|, \text{ for all } I_0, J_0 \in \mathcal{D}.$$

Case 3: I'' is even and J'' is odd. This case is symmetrical to Case 2, and yields

(20)
$$\sum_{\substack{I \text{ odd}, I \subseteq I_0 \\ J \text{ even}, J \subseteq J_0}} |(b, h_I \otimes h_J)|^2 \lesssim |I_0||J_0|, \text{ for all } I_0, J_0 \in \mathcal{D}.$$

Case 4: I'' and J'' are both even. In this case we have all four operators in Group III, but we already know from (18), (19) and (20) that all but $\pi\pi 4$ are already bounded. Now

$$\|\pi\pi 4\|_{2}^{2} = \frac{4}{|I''||J''|} \sum_{\substack{I \subsetneq I''_{\pm} \\ J \subsetneq J''_{\pm}}} |(b, h_{I} \otimes h_{J})|^{2},$$

which gives us exactly that

$$\sum_{\substack{I \subseteq I_0 \\ J \subseteq J_0}} |(b, h_I \otimes h_J)|^2 \lesssim |I_0||J_0|, \text{ for all } I_0, J_0 \in \mathcal{D},$$

so $b \in BMO_r$.

4. Analyzing what is left to estimate if BMO_r of the symbol is small

In the previous section we proved

We can normalize b to have

$$||b||_{BMO_{ChF}} = 1$$
.

Hence, to get an estimate of \mathcal{T}^b from below we can assume

$$||b||_{BMO_r} \lesssim \varepsilon_0,$$

where ε_0 is a very small positive absolute number.

We recall that everything is dyadic.

Definition 4.1. The Haar support of function α is the collection of dyadic rectangles such that Haar coefficient does not vanish: $\sigma_H(\alpha) := \{R : (\alpha, h_R) \neq 0\}.$

Definition 4.2. Let
$$R = I \times J$$
, $p(R) := \hat{R} = \hat{I} \times \hat{J}$, $p_m(R) = \hat{p}_{m-1}(R)$.

Definition 4.3. Let U_0 be a finite union of dyadic rectangles. We call such U_0 a dyadic open set. For every such U_0 we denote by \mathcal{U}_0 the collection of all dyadic rectangles lying inside U_0 .

Definition 4.4. We say that function α is deep inside a dyadic open set U if $p_m(\sigma_H(\alpha)) \subset \mathcal{U}$. We can call such α m-deep inside U.

Definition 4.5. $\mathcal{T}_*^b = [T_2^*, [T_1^*, b]].$

- 4.1. **Plan.** We will try to estimate \mathcal{T}^b from below under assumptions:
 - (1) $\sigma_H(b)$ is finite;
 - (2) $||b||_{BMO_{ChF}} = 1;$
 - (3) $||b||_{BMO_r} \leq \varepsilon_0$;
 - (4) U_0 is a dyadic open set such that $||b||_{BMO_{ChF}} = \frac{1}{|U_0|} \sum_{R \in \mathcal{U}_0} (b, h_R)^2$.

By rescaling we can assume that $|U_0| = 1$.

We will stay soon only with $\pi\pi$ part of repeated commutator.

Lemma 2 (Symbol is "smaller" than test function) $\mathcal{T}_*^{h_{I'} \otimes h_{J'}}(h_{I''} \otimes h_{J''}) \neq 0 \Rightarrow I' \leq \hat{I}'', J' \leq \hat{J}''$.

Lemma 3 (Symbol is "smaller" than test function) $\mathcal{T}^{h_{I'} \otimes h_{J'}}(h_{I''} \otimes h_{J''}) \neq 0 \Rightarrow I' \leq I'', J' \leq J''$.

Proof. Direct calculation.
$$\Box$$

Definition 4.6. Given U_0 , we use a standard notation U for the set $=\{M_s^d \mathbf{1}_{U_0} \geq \frac{1}{16}\}$, where M_s^d denotes the strong dyadic maximal function.

Lemma 4 Let U_0 be a finite union of dyadic rectangles. Let U be as in the Definition 4.6. Let ρ be such that $\sigma_H(\rho) \subset \mathcal{U}_0$. Then $\mathcal{T}_*^{\beta}(\rho) = 0$ for any β such that $\sigma_H(\beta) \cap \mathcal{U} = \emptyset$.

Proof. Follows from Lemma 2.
$$\Box$$

Lemma 5 Let U_0 be a finite union of dyadic rectangles. Let U be as in the Definition 4.6. Let ρ be such that $\sigma_H(\rho) \subset \mathcal{U}$. Then $\mathcal{T}^{\beta}(\rho) = 0$ for any β such that $\sigma_H(\beta) \cap \mathcal{U} = \emptyset$.

Proof. Follows from Lemma 3.

Definition 4.7. $Edge_m(U_0) = \{R : R \in \mathcal{U}_0, p_{m-1}(R) \in \mathcal{U}_0, p_m(R) \cap U_0^c \neq \emptyset\}$. $Edge(U_0) := Edge_1(U_0)$. The same for U from Definition 4.6. $Edge(U) = \{R : R \subset U, \hat{R} \cap U^c \neq \emptyset\}$.

So $Edge_m(U_0)$ describes rectangles in U_0 that are "deep, but not so deep" in U_0 , they are on the edge.

Given b and U_0 , we consider U built as in Definition 4.6 and we consider the decomposition

$$(23) b = \alpha_0 + \beta,$$

where

- (1) $\alpha_0 := \sum_{R \subset U} (b, h_R) h_R;$
- (2) β is the rest, meaning that rectangles giving non-zero β coefficient have to be like this: $R \cap U^c \neq \emptyset$.

By Lemma 41 that says that symbol should be "smaller" than the test function, we get that if ρ is such that $\sigma_H(\rho) \subset \mathcal{U}_0$, then

$$\mathcal{T}_*^{\beta}(\rho) = 0.$$

So-in estimating $\mathcal{T}_*^b(\rho)$ from below we can forget about $\|\mathcal{T}_*^\beta(\rho)\|_2$ and look only at $\mathcal{T}_*^{\alpha_0}(\rho)$. And the only thing we used for claiming that is that Haar spectrum of ρ is in \mathcal{U}_0 and Definition 4.6 of U.

By Definition 4.6 and weak boundedness of M_s^d we know that

$$(25) |U| \le C_0 |U_0|$$

Obviously, if we normalize $|U_0| = 1$, $||\rho||_2 = 1$, we have

$$C_0^{1/2} \| \mathcal{T}^b \| \ge \| \mathcal{T}^b \| |U|^{1/2} \ge \| \mathcal{T}^b \| \| \rho \| |U|^{1/2} \ge |(\mathbf{1}_U, \mathcal{T}^b_*(\rho))| = |(\mathbf{1}_U, \mathcal{T}^{\alpha_0}_*(\rho))|.$$

So we are left to estimate $|(\mathbf{1}_U, \mathcal{T}_*^{\alpha_0}(\rho))|$ from below, where $\sigma_H(\rho) \subset \mathcal{U}_0$.

5. The estimate of
$$|(\mathcal{T}^{\alpha_0}(\mathbf{1}_U), \rho|)$$
 from Below, where $\sigma_H(\rho) \subset \mathcal{U}_0$

To estimate $|(\mathbf{1}_U, \mathcal{T}_{\star}^{\alpha_0}(\rho))|$ from below let us look at our formulas for $\pi\pi, \pi Z, \ldots, DD$ for

$$b = \alpha_0, f = \mathbf{1}_U$$
.

- **Remark 5.1.** If needed we can always replace α_0 that was such that $\sigma_H(\alpha_0) \subset \mathcal{U}$, by α_{m-1} , that is $\sigma_H(\alpha_{m-1}) \subset \mathcal{U}$. This is only if we need this. The result one obtains after such change is weaker, but it still would give the information on some part of Chang–Fefferman BMO norm of b.
- 5.1. Signature and parts of commutator that vanish. We use the notion of signature. We look at the functions employed in the definition of each part. There are 3 functions in I position and 3 functions in J position. We write signature as a matrix, where the first row is assigned to 3 functions in I position and the second row is assigned to 3 functions in J position. The element of the matrix is 0 if the integral of corresponding function is 0 (e.g. h_I), and the element of the matrix is 1 if the integral of corresponding function is not 0 (e.g. $\mathbf{1}_I$). For example,

$$\begin{bmatrix} 0,0,0\\0,0,0 \end{bmatrix}$$

is the signature of DD part of repeated commutator, and

$$\begin{bmatrix} 0,0,0\\0,1,0 \end{bmatrix}.$$

is the signature of $D\pi$ part. DZ part has signature

$$\begin{bmatrix} 0,0,0\\0,0,1 \end{bmatrix} .$$

Lemma 6 All parts having only one 1 part in the signature can be estimates by $C\|\alpha_0\|_{BMO_r} \leq C\varepsilon_0$.

Proof. Direct calculation.

Thus those parts can be disregarded. These are all D..., ...D parts. Consider now parts with Z..., ...Z. These are terms with signatures

 $\begin{bmatrix} 0,1,0\\0,0,1 \end{bmatrix},$

and

 $\begin{bmatrix} 0,0,1\\0,1,0 \end{bmatrix}$

Also ZZ has signature

$$\begin{bmatrix} 0,0,1\\0,0,1 \end{bmatrix} \; .$$

They will necessarily consist of the terms having the following factors:

- (1) $(\mathbf{1}_U, h_I \otimes h_J);$
- (2) $(\mathbf{1}_U, h_I \otimes \tilde{\mathbf{1}}_J);$
- (3) $(\mathbf{1}_U, T_1^* h_I \otimes \tilde{\mathbf{1}}_J);$
- (4) $(\mathbf{1}_U, \tilde{\mathbf{1}}_I \otimes T_1^* h_J);$
- (5) $(\mathbf{1}_U, h_I \otimes T_2^* \tilde{\mathbf{1}}_J);$
- (6) $(\mathbf{1}_U, T_1^*h_I \otimes T_2^*h_J);$
- (7) et cetera....

It is important to remember that here we can count only those I, J such that $I \times J$ are inside U_0 . This follows from our choice of ρ : $\sigma_H(\rho) \subset U_0$. But as $I \times J$ are inside U_0 almost all those scalar products above are equal to zero. Unfortunately not all, number (5) does not vanish as $T_2^* \tilde{\mathbf{1}}_J$ is a very non-localized function. But let us even assume for a while that all such terms disappear (which is false as we just explained, some terms stay).

Let ρ be an arbitrary test function with Haar spectrum inside \mathcal{U}_0 (we can actually use functions with Haar spectrum m-deep in U_0 if needed). Build U as in Definition 4.6. We have (25). We denoted by α_0 the part of b having Haar spectrum in \mathcal{U} .

After the bookkeeping we see that in $|(\mathbf{1}_U, \mathcal{T}_*^{\alpha_0} \rho)|$ only $\pi\pi$ part rests because all other parts either vanish or are smaller than $C\|b\|_{BMO_r} \leq C\varepsilon_0$.

Let us write down all four terms of $\pi\pi$ part of $(\mathbf{1}_U, \mathcal{T}_*^{\alpha_0}\rho), \sigma_H(\rho) \subset U_0$,

(26)
$$\pi\pi(\alpha_{0}, \mathbf{1}_{U}; \mathbf{T}) = \sum_{I} (\alpha_{0}, h_{I} \otimes h_{J}) (\mathbf{1}_{U}, \tilde{\mathbf{1}}_{I} \otimes \tilde{\mathbf{1}}_{J}) T_{1} h_{I} \otimes T_{2} h_{J} - \sum_{I} (\alpha_{0}, h_{I} \otimes h_{J}) (\mathbf{1}_{U}, T_{1}^{*} \tilde{\mathbf{1}}_{I} \otimes \tilde{\mathbf{1}}_{J}) h_{I} \otimes T_{2} h_{J} - \sum_{I} (\alpha_{0}, h_{I} \otimes h_{J}) (\mathbf{1}_{U}, \tilde{\mathbf{1}}_{I} \otimes T_{2}^{*} \tilde{\mathbf{1}}_{J}) T_{1} h_{I} \otimes h_{J} + \sum_{I} (\alpha_{0}, h_{I} \otimes h_{J}) (\mathbf{1}_{U}, T_{1}^{*} \tilde{\mathbf{1}}_{I} \otimes T_{2}^{*} \tilde{\mathbf{1}}_{J}) h_{I} \otimes h_{J} = I - II - III + IV.$$

The range of I, J can be naturally associated with the definition of T_1, T_2 , and with the nature of ρ : $I \times J \subset U_0$.

5.2. Conjugate operator. It seems to be obvious that we can replace both T_i by T_i^* , and get the similar expression. Now we want to estimate $|(\mathbf{1}_U, \mathcal{T}_*^{\alpha_0}(\rho))|$ from below. Let ρ be such that $\sigma_H(\rho) \subset \mathcal{U}$. We denoted by α_0 the part of b having Haar spectrum in \mathcal{U} . By Lemma 3

$$(\mathbf{1}_U, \mathcal{T}^{\alpha_0}(\rho)) = (\mathbf{1}_U, \mathcal{T}^b(\rho)).$$

In estimating $(\mathbf{1}_U, \mathcal{T}^{\alpha_0}(\rho))$ we the terms having the following factors:

- (1) $(\mathbf{1}_U, h_I \otimes h_J)$;
- (2) $(\mathbf{1}_U, h_I \otimes \tilde{\mathbf{1}}_J);$
- (3) $(\mathbf{1}_U, T_1 h_I \otimes \tilde{\mathbf{1}}_J);$
- $(4) (\mathbf{1}_U, \mathbf{1}_I \otimes T_1 h_J);$
- (5) $(\mathbf{1}_U, h_I \otimes T_2 \tilde{\mathbf{1}}_J);$
- (6) $(\mathbf{1}_U, T_1 h_I \otimes T_2 h_J);$
- (7) et cetera... .

By the choice of α_0 just mentioned above, $I \times J \subset U$ here. Therefore all these scalar products seem to vanish. But the attentive reader can notice that, yes, e.g. $(\mathbf{1}_U, T_1 h_I \otimes \tilde{\mathbf{1}}_J)$ does vanish, but $(\mathbf{1}_U, h_I \otimes T_2 \tilde{\mathbf{1}}_J)$ does not. This will create a big difficulty below.

However, let us assume for a while that in estimating $(\mathbf{1}_U, \mathcal{T}^{\alpha_0}(\rho))$ from below we will be left with sum of terms estimated by $C\|b\|_{BMO_r}$ and with the following $\pi\pi$ terms that we cannot cancel out:

(27)
$$\pi\pi(\alpha_{0}, \mathbf{1}_{U}; \mathbf{T}^{*}) = \sum_{I} (\alpha_{0}, h_{I} \otimes h_{J}) (\mathbf{1}_{U}, \tilde{\mathbf{1}}_{I} \otimes \tilde{\mathbf{1}}_{J}) T_{1}^{*} h_{I} \otimes T_{2}^{*} h_{J} - \sum_{I} (\alpha_{0}, h_{I} \otimes h_{J}) (\mathbf{1}_{U}, T_{1} \tilde{\mathbf{1}}_{I} \otimes \tilde{\mathbf{1}}_{J}) h_{I} \otimes T_{2}^{*} h_{J} - \sum_{I} (\alpha_{0}, h_{I} \otimes h_{J}) (\mathbf{1}_{U}, \tilde{\mathbf{1}}_{I} \otimes T_{2} \tilde{\mathbf{1}}_{J}) T_{1}^{*} h_{I} \otimes h_{J} + \sum_{I} (\alpha_{0}, h_{I} \otimes h_{J}) (\mathbf{1}_{U}, T_{1} \tilde{\mathbf{1}}_{I} \otimes T_{2} \tilde{\mathbf{1}}_{J}) h_{I} \otimes h_{J} = A - B - C + D.$$

We do not write the scalar product with $\rho, \sigma_H(\rho) \subset \mathcal{U}$, but we keep it in mind. The range of I, J can be naturally associated with the definition of T_1, T_2 , and with the nature of α_0 : definitely $I \times J \subset U$, but for example the term with $T_1^*h_I \otimes T_2^*h_J$ disappears if $I \times J \subset U$, but $\hat{I} \times \hat{J}$ is not inside U. This term will be "cut-out" by the scalar product with $\rho, \sigma_H(\rho) \subset \mathcal{U}$.

But notice that all terms such that $I \times J \subset U_0$ stay, they are not cut-out. This is because by Definition 4.6 we have $\hat{I} \times \hat{J} \subset U$ as soon as $I \times J \subset U_0$.

We like the first line of (27) (and we also like the first line of (26)). If the first line were alone in (27) then:

$$\|\pi\pi(\alpha_0, \mathbf{1}_U; \mathbf{T}^*)\|_2^2 = \|A\|_2^2 \ge \sum_{\substack{R \text{ odd odd} \\ R \subset U_0}} (b, h_R)^2.$$

And we would get

(28)
$$\|\mathcal{T}^b\| = \|\mathcal{T}^b_*\| \ge c \sum_{\substack{R \text{ odd odd} \\ R \subset U_0}} (b, h_R)^2,$$

which is "almost" what we want (in fact that would be 25 percent of what we want). Of course we want more, namely, the summation not restricted to just even-even R. So we have several problems:

- (1) how to get the summation over all R's inside U_0 ?
- (2) what to do with terms II, III, IV?

Of course we need summation over all R not just (odd, odd), but still, (28) would be a step in the direction of proving the ultimate estimate:

(29)
$$\|\mathcal{T}^b\| \ge c \sum_{R \subset U_0} (b, h_R)^2.$$

5.3. Coming back to $(\mathbf{1}_U, \mathcal{T}_*^{\alpha_0}(\rho))$ and trying other test functions.

Remark 5.2. We can try to use $T_1\mathbf{1}_U, T_2\mathbf{1}_U, T_1T_2\mathbf{1}_U$ as test functions instead of $\mathbf{1}_U$ in (26). Again it is easy to see that if we choose instead of α_0 a deeper function $\alpha_1 = \sum_{Ris\ 2-deep\ in\ U_0} (b, h_R) h_R$, then after bookkeeping we see that in $|(T_i\mathbf{1}_U, \mathcal{T}^{\alpha_1}(\alpha_1))|$ only the $\pi\pi$ part rests because all other parts either vanish or are smaller than $C||b||_{BMO_r} \leq C\varepsilon_0$.

Let us look at (we use that $(T_1^*)^2 = 0$)

(30)
$$\pi\pi(\alpha_1, T_1\mathbf{1}_U) = \sum_{I} (\alpha_1, h_I \otimes h_J) (\mathbf{1}_U, T_1^* \tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J) T_1 h_I \otimes T_2 h_J + \sum_{I} (\alpha_1, h_I \otimes h_J) (\mathbf{1}_U, T_1^* \tilde{\mathbf{1}}_I \otimes T_2^* \tilde{\mathbf{1}}_J) T_1 h_I \otimes h_J.$$

We immediately conclude that the second sum restricted to I and J even is orthogonal to what is rest, so

(31)
$$\sum_{I,J \, even, even} (\alpha_1, h_I \otimes h_J)^2 (\mathbf{1}_U, T_1^* \tilde{\mathbf{1}}_I \otimes T_2^* \tilde{\mathbf{1}}_J)^2 \leq C \|\mathcal{T}^b\|^2.$$

Let us look at (we use that $(T_2^*)^2 = 0$)

(32)
$$\pi\pi(\alpha_1, T_2\mathbf{1}_U) = \sum_{I} (\alpha_1, h_I \otimes h_J) (\mathbf{1}_U, \tilde{\mathbf{1}}_I \otimes T_2^* \tilde{\mathbf{1}}_J) T_1 h_I \otimes T_2 h_J + \sum_{I} (\alpha_1, h_I \otimes h_J) (\mathbf{1}_U, T_1^* \tilde{\mathbf{1}}_I \otimes T_2^* \tilde{\mathbf{1}}_J) h_I \otimes T_2 h_J.$$

We again conclude that the second sum restricted to I and J even is orthogonal to what is rest, so the same inequality follows:

(33)
$$\sum_{I,J \, even, even} (\alpha_1, h_I \otimes h_J)^2 (\mathbf{1}_U, T_1^* \tilde{\mathbf{1}}_I \otimes T_2^* \tilde{\mathbf{1}}_J)^2 \leq C \|\mathcal{T}^b\|^2.$$

Exactly the same inequality follows if we consider $\pi\pi(\alpha_1, T_1T_2\mathbf{1}_U)$.

But what would be needed is the "opposite" sum estimate:

(34)
$$\sum_{I,J \text{ odd,odd}} (\alpha_1, h_I \otimes h_J)^2 (\mathbf{1}_U, T_1^* \tilde{\mathbf{1}}_I \otimes T_2^* \tilde{\mathbf{1}}_J)^2 \leq C \|\mathcal{T}^b\|^2,$$

because exactly this sum entangles with the first line of (26): $\sum (\alpha_0, h_I \otimes h_J) (\mathbf{1}_U, \tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J) T_1 h_I \otimes T_2 h_J$.

Problem. So far we did not guess how to get (28) from estimating formula (27), not to mention that (29) rests even less clear. But we will proceed in the next two sections.

Let us take a look at the one parameter case. We obviously can assume that we have localization, and that eventually we just need to find out how to estimate the analog of, say, (27) from below.

Here is this analog. It has two rather than four sums because we are in one parameter situation. The "open" set U in one parameter case is just a dyadic interval I_0 . We can assume that

$$|I_0| = 1$$
.

(35)
$$\pi(\alpha_0, \mathbf{1}_{I_0}; T^*) = \sum_{I \in \mathcal{D}(I_0)} (b, h_I) \langle T^* \mathbf{1}_{I_0} \rangle_I h_I - \sum_{I \in \mathcal{D}(I_0)} (b, h_I) \langle \mathbf{1}_{I_0} \rangle_I T^* h_I = \sum_{I \in \mathcal{D}(I_0)} (b, h_I) \langle T^* \mathbf{1}_{I_0} \rangle_I h_I - \sum_{I \in \mathcal{D}(I_0)} (b, h_I) T^* h_I = A - B.$$

Before plunging into the consideration of (35), let us notice that the choice of $\mathbf{1}_U$ in two parameter case was initiated by the wish to stay only with $\pi\pi$ part and to bring many other parts of the commutator to zero.

Such a problem is not present in one parameter case. In this case we can notice that the estimate of the commutator can be reduced to the estimate from below of the following expression, where the test function $f \neq \mathbf{1}_{I_0}$. In fact we can choose test function differently. So consider the following more general expression

(36)
$$\pi(\alpha_0, f_{I_0}; T) = \sum_{I \in \mathcal{D}(I_0)} (b, h_I) \langle T^* f_{I_0} \rangle_I h_I - \sum_{I \in \mathcal{D}(I_0)} (b, h_I) \langle f_{I_0} \rangle_I T^* h_I = A_f - B_f.$$

Let us now choose a very simple $f_{I_0} \neq \mathbf{1}_{I_0}$. Here it is convenient to think that I_0 is odd (and normalized $|I_0| = 1/2$). In this case our choice of test function is the "uncle" function:

$$f_{I_0} = h_{sI_0}$$
.

Here sI_0 is a sibling of I_0 . By the definition of T^* ,

$$h_{sI_0}|I_0=0, \quad T^*h_{sI_0}|I_0=\operatorname{sgn}(I_0,\hat{I}_0)h_{\hat{I}_0}=\pm\frac{1}{\sqrt{2}}.$$

We plug into (36) above and B_f disappears. The estimate of A_f gives

$$\sum_{I \in \mathcal{D}(I_0)} (b, h_I)^2 \le C \|\pi(\alpha_0, f_{I_0}; T)\|_2^2 \le C \|\text{the commutator}\|_2^2.$$

In fact, for even I_0 the choice $f = h_{s(\hat{I}_0)}$ would work. We are done.

Remark 6.1. What serves as "the uncle" function $F = f_U$ of U in the two parameter case—this is not clear.

Now let us come back to (35), and let us try to obtain $\sum_{I \in \mathcal{D}(I_0)} (b, h_I)^2 \leq C$ the commutator $\|_2^2$ directly from (35).

Recall that $T^*h_I = \operatorname{sgn}(I, \hat{I})h_{\hat{I}}$ for odd I and 0 for even.

Let us consider $I \in \mathcal{D}(I_0)$ such that $I \in \mathcal{D}(I_0)$, and then we have a term $-\operatorname{sgn}(I,I)(b,h_I)h_{\hat{I}}$ in -B sum (this is when I is odd, when I is even this term vanishes), and we have another term $h_{\hat{I}} \cdot (b,h_{\hat{I}})$ with coefficient $c_I := \langle T^* \mathbf{1}_{I_0} \rangle_{\hat{I}}$. This term is always present, for even and odd I.

Let us understand, what is c_I , which is equal to the constant value of the constant function $T^*\mathbf{1}_{I_0}|I_0$ (recall that $\hat{I} \subset I_0$). Let I_1 denote the father of I_0 , I_2 denote the father of I_1 , et cetera.

$$\mathbf{1}_{I_0} = (\mathbf{1}_{I_0}, h_{I_1})h_{I_1} + (\mathbf{1}_{I_0}, h_{I_2})h_{I_2} + \dots$$

Hence

$$(T^*\mathbf{1}_{I_0})|I_0 = (\mathbf{1}_{I_0}, h_{I_1})(T^*h_{I_1})|I_0 + (\mathbf{1}_{I_0}, h_{I_2})(T^*h_{I_2})|I_0 + \dots$$

We recall that $|I_0|=1$. Definitely the first term of the series is at most $\frac{1}{2\sqrt{2}}$ by absolute value (it can be zero too). In fact, h_{I_1} gives the contribution of $\frac{1}{\sqrt{2}}$ and $T^*h_{I_1}$ gives the contribution $\frac{1}{2}$. The next term of the series is at most $\frac{1}{2\sqrt{2}}$: h_{I_2} gives the contribution of $\frac{1}{2}$ and $T^*h_{I_2}$ gives the contribution 0 or $\frac{1}{2\sqrt{2}}$. It is clear that

(37)
$$\frac{1}{8} \le |c_I| \le \frac{1}{\sqrt{2}} < 1.$$

We see that for I such that $\hat{I} \subset I_0$ the sum A - B of (35) has the term

$$h_{\hat{I}} \cdot (-\operatorname{sgn}(I, \hat{I}) \cdot [(b, h_I) - d_I(b, h_{\hat{I}})], \quad |d_I| \le \frac{1}{\sqrt{2}},$$

when I is odd, and the term

$$h_{\hat{I}} \cdot (\operatorname{sgn}(I, \hat{I}) \cdot d_I(b, h_{\hat{I}}), \quad |d_I| \ge \frac{1}{8},$$

when I is even.

All terms are orthogonal, and we get

(38)
$$\sum_{I:\hat{I} \in \mathcal{D}(I_0), I \, odd} |(b, h_I) - d_I(b, h_{\hat{I}})|^2 \le C \|\text{the commutator}\|_2^2,$$

and

(39)
$$\sum_{I: I \in \mathcal{D}(I_0), Iodd} |(b, h_I)|^2 \le \sum_{I: \hat{I} \in \mathcal{D}(I_0), Ieven} |(b, h_{\hat{I}})|^2 \le C \|\text{the commutator}\|_2^2,$$

From these two inequalities and from the estimates of d_I , we can immediately obtain

(40)
$$\sum_{I:I\in\mathcal{D}(I_0)} |(b,h_I)|^2 \le C \|\text{the commutator}\|_2^2,$$

Notice that (38) alone would give us an interesting information.

We want to get from (38) alone that

(41)
$$\sum_{I:\hat{I}\in\mathcal{D}(I_0),Iodd} (b,h_I)^2 \le C \|\pi(\alpha_0,f_{I_0};T)\|_2^2 \le C \|\text{the commutator}\|_2^2.$$

(This is stated in (39), but suppose that we do not have it, suppose that we have only (38).) If this is obtained, then it is easy to get

(42)
$$\sum_{I \in \mathcal{D}(I_0), Iodd} (b, h_I)^2 \le C \|\pi(\alpha_0, f_{I_0}; T)\|_2^2 \le C \|\text{the commutator}\|_2^2.$$

Now $(38) \Rightarrow (41)$ seems is easy by the following lemmas.

Lemma 7 Let $|d_n| \le q < 1$, let $b_n := a_n - d_n a_{n-1}$. Assume that $b := \{b_n\}_{n=0}^{\infty} \in \ell^2$, then $a := \{a_n\}_{n=0}^{\infty} \in \ell^2$. Proof. We can write

$$a_n = b_n + d_n a_{n-1} = b_n + d_n b_{n-1} + d_n d_{n-1} a_{n-2} = \dots = b_n + D_1 b_{n-1} + D_2 b_{n-2} + \dots + D_n b_0$$

where $D_1 := d_n, D_2 = d_n d_{n-1}, \dots, |D_k| \le q^k$. Therefore, a = Ab, where A is a triangular matrix (with 1's one the diagonal), which clearly satisfies Schur test. Hence, $A : \ell^2 \to \ell^2$ and lemma is proved.

In fact we need this type of lemma on the dyadic tree T of dyadic lattice $\mathcal{D}(I_0)$. But here it is.

Lemma 8 Let m(I,J) is a matrix on T such that m(I,J)=0 if $I\subseteq J$ and for c>0

$$m(I,J) = \left(\frac{|I|}{|J|}\right)^c$$

otherwise. Then this matrix is a bounded operator from $\ell^2(T)$ to itself.

Proof. This matrix acts boundedly from $\ell^{\infty}(T)$ to $\ell^{1}(T)$. In fact,

$$\sum_{I} m(I,J) \le C_0 \quad \forall I .$$

But its adjoint also acts boundedly from $\ell^{\infty}(T)$ to $\ell^{1}(T)$. In fact

$$\sum_{I} m(I,J) \le C_0 \quad \forall J .$$

By interpolation it maps $\ell^2(T)$ to itself boundedly.

7. Back to the estimate from below of (26), (27)

First we just loosely discuss what we are going to do. The "actual idea" follows in Subsection 7.1. Disclaimer: Subsection 7.1 contains three deliberate mistakes. We will indicate all of them at the end of Subsection 7.1. If the reader can circumvent those problems, then the main result will be proved: repeated dyadic paraproduct is bounded if and only if its symbol is in Chang–Fefferman dyadic BMO.

Remark 7.1. We actually prove this result in Theorem 9.2 below, but only under the extra conditions on the symbol. We cannot get rid of those extra assumptions at this time.

Let us repeat: we just need to be able to estimate the sum in (26) from below by constant or by $c \sum_{I \times I \subset U} (b, h_I \otimes h_J)^2$. Even the estimate from below by

$$c \sum_{I \times J \subset U, I \times J \text{ is } m \text{ deep, } I, J \text{ even, even}} (b, h_I \otimes h_J)^2$$

would be a meaningful achievement. Similarly, even the estimate from below by

$$c \sum_{I \times J \subset U, \, I \times J \, is \, \, m \, deep, \, I, J \, odd, odd} (b, h_I \otimes h_J)^2$$

would be a meaningful achievement as well.

Recall

(43)
$$\pi\pi(\alpha_{0}, \mathbf{1}_{U}; \mathbf{T}) = \sum_{I} (\alpha_{0}, h_{I} \otimes h_{J}) (\mathbf{1}_{U}, \tilde{\mathbf{1}}_{I} \otimes \tilde{\mathbf{1}}_{J}) T_{1} h_{I} \otimes T_{2} h_{J} - \sum_{I} (\alpha_{0}, h_{I} \otimes h_{J}) (\mathbf{1}_{U}, T_{1}^{*} \tilde{\mathbf{1}}_{I} \otimes \tilde{\mathbf{1}}_{J}) h_{I} \otimes T_{2} h_{J} - \sum_{I} (\alpha_{0}, h_{I} \otimes h_{J}) (\mathbf{1}_{U}, \tilde{\mathbf{1}}_{I} \otimes T_{2}^{*} \tilde{\mathbf{1}}_{J}) T_{1} h_{I} \otimes h_{J} + \sum_{I} (\alpha_{0}, h_{I} \otimes h_{J}) (\mathbf{1}_{U}, T_{1}^{*} \tilde{\mathbf{1}}_{I} \otimes T_{2}^{*} \tilde{\mathbf{1}}_{J}) h_{I} \otimes h_{J} = I - III - IIII + IV.$$

Definition 7.2. Let $0 < \varepsilon_1 \le \frac{1}{100}$. The rectangle $I \times J \subset U$ is called ε_1 small if

$$|(\mathbf{1}_U, T_1^* \tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J)| \leq \varepsilon_1$$
.

Lemma 9 Let $I \times J$ be m-deep in U, $m \geq 2$. Let $R = I \times J$ be ε_1 -small. Then $R_1 = \hat{I} \times J$ and $R_2 = I \times \hat{J}$, $\hat{R} = \hat{I} \times \hat{J}$ are ε_1 -small.

Proof. Let us consider R_1 . We look at

$$(\mathbf{1}_U,T_1^*\tilde{\mathbf{1}}_I\otimes \tilde{\mathbf{1}}_J)-(\mathbf{1}_U,T_1^*\tilde{\mathbf{1}}_{\hat{I}}\otimes \tilde{\mathbf{1}}_J)$$

and notice that $\tilde{\mathbf{1}}_I - \tilde{\mathbf{1}}_{\hat{I}}$ is proportional to $h_{\hat{I}}$, so $\tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J - \tilde{\mathbf{1}}_{\hat{I}}I \otimes \tilde{\mathbf{1}}_J$ is proportional to $h_{\hat{I}} \otimes h_J$. Hence $T_1^*(\tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J - \tilde{\mathbf{1}}_{\hat{I}}I \otimes \tilde{\mathbf{1}}_J)$ is proportional to $h_{\mathcal{I}} \otimes h_J$, where \mathcal{I} is the parent of \hat{I} (grandparent of I). The scalar product with $\mathbf{1}_U$ returns 0 if R is at least 2 deep.

Definition 7.3. Let $\tilde{\mathbf{1}}_R$ temporarily denote $\tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J$ for $R = I \times J$.

We proved that

(44)
$$(\mathbf{1}_{U}, T_{1}^{*}\tilde{\mathbf{1}}_{R}) = (\mathbf{1}_{U}, T_{1}^{*}\tilde{\mathbf{1}}_{R_{1}}),$$

if R is at least 2-deep. Therefore, R_1 is also ε_1 small.

The same proof works for R_2 . And then going from R_2 to \hat{R} involves the same reasoning. In particular we get

$$(\mathbf{1}_{U}, T_{1}^{*}\tilde{\mathbf{1}}_{R}) = (\mathbf{1}_{U}, T_{1}^{*}\tilde{\mathbf{1}}_{R_{1}}) = (\mathbf{1}_{U}, T_{1}^{*}\tilde{\mathbf{1}}_{R_{2}}) = (\mathbf{1}_{U}, T_{1}^{*}\tilde{\mathbf{1}}_{\hat{R}}).$$

And we conclude that all R_1, R_2, \hat{R} are ε_1 small as soon as R is such.

Lemma 10 Let $R = I \times J$ be such that $R \subset U$ and moreover R is at least 1 deep. Let R be ε_1 -small. Let $I' \in ch(I)$, $J' \in ch(J)$. Then $R^1 = \hat{I}' \times J$ and $R^2 = I \times \hat{J}'$, $R^{12} = \hat{I}' \times \hat{J}'$ are ε_1 -small.

Proof. The proof is exactly the same, again the difference of two characteristic functions normalized as we did gives us a proportional to Haar function, T_1^* moves it only one step up, and so the result is still orthogonal to $\mathbf{1}_U$.

Now we can prove (only with I and J interchanged) that the same (50) will hold for the terms $(\mathbf{1}_U, \tilde{\mathbf{1}}_I \otimes T_2^* \tilde{\mathbf{1}}_J)$ instead of $(\mathbf{1}_U, T_1^* \tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J)$. And then, again we can repeat the same reasoning for the terms $(\mathbf{1}_U, T_1^* \tilde{\mathbf{1}}_I \otimes T_2^* \tilde{\mathbf{1}}_J)$.

Conclusion:

- 1) in (26) either all terms $(\mathbf{1}_U, T_1^*\tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J)$, $(\mathbf{1}_U, \tilde{\mathbf{1}}_I \otimes T_2^*\tilde{\mathbf{1}}_J)$, $(\mathbf{1}_U, T_1^*\tilde{\mathbf{1}}_I \otimes T_2^*\tilde{\mathbf{1}}_J)$ are $\frac{1}{100}$ small, or
- 2) two brand of terms, e.g. all $(\mathbf{1}_U, T_1^* \tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J)$, $(\mathbf{1}_U, \tilde{\mathbf{1}}_I \otimes T_2^* \tilde{\mathbf{1}}_J)$ are $\frac{1}{100}$ small and all $(\mathbf{1}_U, T_1^* \tilde{\mathbf{1}}_I \otimes T_2^* \tilde{\mathbf{1}}_J)$ are NOT $\frac{1}{100}$ small (there are three choices of such a situation), or
- 3) one brand of terms, e.g. all $(\mathbf{1}_U, T_1^* \tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J)$ are all $\frac{1}{100}$ small, and two other brands are NOT $\frac{1}{100}$ small, or
- 4) all terms of all three brands are NOT $\frac{1}{100}$ small.

Let us consider item 1) and item 4). The rest should follow suit.

So suppose all terms of all three brands are $\frac{1}{100}$ small. Then notice that $(b, h_I \otimes h_J)$ from the first line of (26) has coefficient $(\mathbf{1}_U \tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J) = 1$ —and of course all this stands in front of some $h_{I'} \otimes h_{J'}$ (children). If we collect the same $h_{I'} \otimes h_{J'}$ terms from all 3 other lines of (26) the b-Haar coefficient (maybe not $(b, h_I \otimes h_J)$, but some other Haar coefficient of a *some* rectangle "close" to $I \times J$) will appear but with the factor less or equal than $\frac{1}{100}$ by absolute value. So we will have that the sum in (26) is exactly equal to the sum of orthogonal terms

$$[1 \cdot (b, h_I \otimes h_J) + \varepsilon_1(b, \dots) + \varepsilon_2(b, \dots) + \varepsilon_3(b, \dots)]h_{I'} \otimes h_{J'}$$

where ... stand for some $h_{I_i'' \times J_i''}$, i = 1, 2, 3, where $I_i'' \times J_i''$ is a close relative of $I \times J$. Hence (26) has the norm squared bounded from below by the sum

$$|1 \cdot (b, h_I \otimes h_J) + \varepsilon_1(b, \dots) + \varepsilon_2(b, \dots) + \varepsilon_3(b, \dots)|^2$$

where the summation is over all those $I \times J$, that are, say, at least 2-deep in U and I, J are both even. And of course $|\varepsilon_i| \le 1/100$.

Then using the analog of Lemma 8 but for double tree, we conclude that the square of the norm of the sum in (26) is bounded from below by the sum

$$c |(b, h_I \otimes h_J)|^2$$
,

where the summation is over all $I \times J$, say, at least 2-deep in U and and I, J are both even, and we got what we wanted at least for even, even part of b.

Let us consider now item 4): all coefficients of all three brands are bigger than $\frac{1}{100}$ by absolute value. Let us start with the fourth line of (26). It contains I even, J even sum, and this sum is orthogonal to the rest. So immediately we get (using that $|(\mathbf{1}_U, T_1^* \tilde{\mathbf{1}}_I \otimes T_2^* \tilde{\mathbf{1}}_J)| \geq 1/100$ by assumption 4))

(46)
$$\sum_{Ieven, J even} (b, h_I \otimes h_J)^2 \le C \|\text{sum of } (26)\|_2^2.$$

So looks like we estimated $\|b^{even,even}\|_{BMO_{ChF}}$ from above by the norm of the repeated commutator.

7.1. "A proof" of the Chang-Fefferman BMO norm estimate of $b^{odd,odd}$. In this subsection we prefer to work with T_1^*, T_2^* rather than T_1, T_2 and we notice that this change is basically symmetric. So our goal will be to estimate Chang-Fefferman norm of $b^{even,even}$ by the norm of $[T_1^*, [T_2^*, b]]$.

Absolutely symmetrically to what has been done above we can reduce the above mentioned task to estimating the norm of the following sum from below by Chang–Fefferman norm of $b^{odd,odd}$.

(47)
$$\pi\pi(\alpha_{0}, \mathbf{1}_{U}; \mathbf{T}^{*}) = \sum_{I} (\alpha_{0}, h_{I} \otimes h_{J}) (\mathbf{1}_{U}, \tilde{\mathbf{1}}_{I} \otimes \tilde{\mathbf{1}}_{J}) T_{1}^{*} h_{I} \otimes T_{2}^{*} h_{J} - \sum_{I} (\alpha_{0}, h_{I} \otimes h_{J}) (\mathbf{1}_{U}, T_{1} \tilde{\mathbf{1}}_{I} \otimes \tilde{\mathbf{1}}_{J}) h_{I} \otimes T_{2}^{*} h_{J} - \sum_{I} (\alpha_{0}, h_{I} \otimes h_{J}) (\mathbf{1}_{U}, \tilde{\mathbf{1}}_{I} \otimes T_{2} \tilde{\mathbf{1}}_{J}) T_{1}^{*} h_{I} \otimes h_{J} + \sum_{I} (\alpha_{0}, h_{I} \otimes h_{J}) (\mathbf{1}_{U}, T_{1} \tilde{\mathbf{1}}_{I} \otimes T_{2} \tilde{\mathbf{1}}_{J}) h_{I} \otimes h_{J} = I - II - III + IV.$$

The summation is over $I, J, I \times J \subset U$.

Definition 7.4. Let $0 < \varepsilon_1 \le \frac{1}{100}$. The rectangle $I \times J \subset U$ is called ε_1 small if

$$|(\mathbf{1}_U, T_1\tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J)| \leq \varepsilon_1$$
.

Let us start with the fourth line of (47). It contains I, J (odd, odd) sum, and this sum is orthogonal to the rest. So immediately we get, using the assumption $|(\mathbf{1}_U, T_1\tilde{\mathbf{1}}_I \otimes T_2\tilde{\mathbf{1}}_J)| \geq \varepsilon_2$, that

(48)
$$\sum_{Iodd, Jodd} (b, h_I \otimes h_J)^2 \le C\varepsilon_2^{-2} \|\text{sum of } (26)\|_2^2.$$

The next lemma says that if $|(\mathbf{1}_U, T_1\tilde{\mathbf{1}}_I \otimes T_2\tilde{\mathbf{1}}_J)| \geq \varepsilon_2$ happens for one $I \times J$ inside U it happens for all such rectangles inside U. (If U is connected, which we can assume by considering connected components one by one.)

We just saw that the assumption $|(\mathbf{1}_U, T_1\tilde{\mathbf{1}}_I \otimes T_2\tilde{\mathbf{1}}_J)| \geq \varepsilon_2$ implies (48). Let us now assume that $|(\mathbf{1}_U, T_1\tilde{\mathbf{1}}_I \otimes T_2\tilde{\mathbf{1}}_J)| \leq \varepsilon_2$ for (odd, odd) I, J, then by lemmas below we can assume that for all $I \times J$ inside U we have

$$|(\mathbf{1}_U, T_1\tilde{\mathbf{1}}_I \otimes T_2\tilde{\mathbf{1}}_J)| \leq \varepsilon_2$$
.

Definition 7.5. Let $\tilde{\mathbf{1}}_R$ temporarily denote $\tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J$ for $R = I \times J$.

Lemma 11 Let $I \times J$ be m-deep in U, $m \geq 2$. Let $R = I \times J$ be ε_1 -small. Then $R_1 = \hat{I} \times J$ and $R_2 = I \times \hat{J}$, $\hat{R} = \hat{I} \times \hat{J}$ are ε_1 -small.

Proof. Let us consider R_1 . We look at

$$(\mathbf{1}_U, T_1 \tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J) - (\mathbf{1}_U, T_1 \tilde{\mathbf{1}}_{\hat{I}} \otimes \tilde{\mathbf{1}}_J)$$

and notice that $\tilde{\mathbf{1}}_I - \tilde{\mathbf{1}}_{\hat{I}}$ is proportional to $h_{\hat{I}}$, so $\tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J - \tilde{\mathbf{1}}_{\hat{I}}I \otimes \tilde{\mathbf{1}}_J$ is proportional to $h_{\hat{I}} \otimes h_J$. Hence $T_1(\tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J - \tilde{\mathbf{1}}_{\hat{I}}I \otimes \tilde{\mathbf{1}}_J)$ is proportional to $h_{\mathcal{I}} \otimes h_J$, where \mathcal{I} is the parent of \hat{I} (grandparent of I). The scalar product with $\mathbf{1}_U$ returns 0 if R is in U.

We proved that

$$(\mathbf{1}_U, T_1 \tilde{\mathbf{1}}_R) = (\mathbf{1}_U, T_1 \tilde{\mathbf{1}}_{R_1}),$$

if R is in U. Therefore, R_1 is also ε_1 small.

The same proof works for R_2 . And then going from R_2 to \hat{R} involves the same reasoning. In particular we get

(50)
$$(\mathbf{1}_{U}, T_{1}\tilde{\mathbf{1}}_{R}) = (\mathbf{1}_{U}, T_{1}\tilde{\mathbf{1}}_{R_{1}}) = (\mathbf{1}_{U}, T_{1}\tilde{\mathbf{1}}_{R_{2}}) = (\mathbf{1}_{U}, T_{1}\tilde{\mathbf{1}}_{\hat{R}}).$$

And we conclude that all R_1, R_2, \hat{R} are ε_1 small as soon as R is such.

Notice that there was nothing special about T_1 in this equality. So we can write the same thing for the action by T_2 and for the action of $T_1 \otimes T_2$:

(51)
$$(\mathbf{1}_U, T_2 \tilde{\mathbf{1}}_R) = (\mathbf{1}_U, T_2 \tilde{\mathbf{1}}_{R_1}) = (\mathbf{1}_U, T_2 \tilde{\mathbf{1}}_{R_2}) = (\mathbf{1}_U, T_2 \tilde{\mathbf{1}}_{\hat{R}}).$$

$$(\mathbf{1}_U, T_1 \otimes T_2 \tilde{\mathbf{1}}_R) = (\mathbf{1}_U, T_1 \otimes T_2 \tilde{\mathbf{1}}_{R_1}) = (\mathbf{1}_U, T_1 \otimes T_2 \tilde{\mathbf{1}}_{R_2}) = (\mathbf{1}_U, T_1 \otimes T_2 \tilde{\mathbf{1}}_{\hat{R}}).$$

Lemma 12 Let $R = I \times J$ be such that $R \subset U$ and moreover R is at least 1 deep. Let R be inside U, and let it be ε_1 -small. Let $I' \in ch(I)$, $J' \in ch(J)$. Then $R^1 = I' \times J$ and $R^2 = I \times J'$, $R^{12} = I' \times J'$ are ε_1 -small.

Proof. The proof is exactly the same, again the difference of two characteristic functions normalized as we did gives us a function proportional to Haar function, T_1 moves it only one step down, and so the result is still orthogonal to $\mathbf{1}_U$.

Let us fix two small constants $\varepsilon_2 \ll \varepsilon_1 \leq \frac{1}{100}$. We look at the terms $(\mathbf{1}_U, T_1 \tilde{\mathbf{1}}_I \otimes T_2 \tilde{\mathbf{1}}_J)$ in (47) and we know by Lemma 11 and Lemma 12 that all those terms are the same. So if they are for all I, J (even, even) are like this:

$$|(\mathbf{1}_U, T_1 \tilde{\mathbf{1}}_I \otimes T_2 \tilde{\mathbf{1}}_J)| \geq \varepsilon_2$$

then (46) is immediately true with $C = \varepsilon_2^{-2}$. Now suppose that all of those terms are like that:

$$|(\mathbf{1}_U, T_1\tilde{\mathbf{1}}_I \otimes T_2\tilde{\mathbf{1}}_J)| \leq \varepsilon_2$$
.

Notice that then the same is true for I, J (even, odd). In fact, this follows from (52). Those (even, odd) output terms also happens in this sum

$$\sum (\alpha_0, h_{I'} \otimes h_{J'}) (\mathbf{1}_U, \tilde{\mathbf{1}}_{I'} \otimes T_2 \tilde{\mathbf{1}}_{J'}) T_1^* h_{I'} \otimes h_{J'},$$

when I', J' are (odd, odd). Notice that the sum itself is in (even, odd) space. Therefore, when we consider the (even, odd) output it can come only from the fourth and the third line of (47). In fact, the first line of (47) is in (even, even) space and the second line is totally in (odd, even) space. Therefore, each term of this (even, odd) part of the output will have the form (I odd, J odd):

$$h_I \otimes h_J[(\alpha_0, h_{\tilde{I}} \otimes h_J)(\text{smaller by abs. value than } \varepsilon_2) + (\alpha_0, h_I \otimes h_J) \cdot (\mathbf{1}_U, \tilde{\mathbf{1}}_I \otimes T_2 \tilde{\mathbf{1}}_J)].$$

Hence, taking (even, odd) part of (47) we have

 $\sum_{Iodd, Jodd} |(\alpha_0, h_{\hat{I}} \otimes h_J)(\text{smaller by abs. value than } \varepsilon_2) + (\alpha_0, h_{\hat{I}} \otimes h_J)(\mathbf{1}_U, \tilde{\mathbf{1}}_I \otimes T_2 \tilde{\mathbf{1}}_J)|^2 \leq C \|\text{sum in } (47)\|_2^2.$

Now we remember that either 1) all $|(\mathbf{1}_U, \tilde{\mathbf{1}}_{\hat{I}} \otimes T_2 \tilde{\mathbf{1}}_J)| \geq \varepsilon_1 >> \varepsilon_2$; or 2) all $|(\mathbf{1}_U, \tilde{\mathbf{1}}_{\hat{I}} \otimes T_2 \tilde{\mathbf{1}}_J)| \leq \varepsilon_1$.

Let us formulate now an analog of Lemma 8, but for a dyadic bi-tree $\mathcal{T} \times \mathcal{T}$. Bi-tree is a graph of dyadic rectangles, where the edges connect rectangles iff one of them is inside the other and the ratio of their areas is 2. In the next lemma $R = I \times J$, $R' = I' \times J'$. In other words R, R' are two end-points of the same edge if either I is a parent of I', J' = J, or I' is a parent of I, J' = I, or J' is a parent of J, I' = I.

Lemma 13 Let us consider a matrix enumerated by the couples of vertices (R, R'), $R' \in \mathcal{T} \times \mathcal{T}$, $R' \in \mathcal{T} \times \mathcal{T}$. Let its matrix elements m(R, R') = 0 unless $R \subset R'$ or $R' \subset R$. In the latter cases, if $R \subset R'$, then $m(R, R') = \left(\frac{|R|}{|R'|}\right)^A$, and if $R' \subset R$, then $m(R, R') = \left(\frac{|R'|}{|R|}\right)^A$, where A is a sufficiently large constant. Then this matrix defines a bounded operator from $\ell^2(\mathcal{T} \times \mathcal{T})$ to itself.

Proof. This is easy to see by applying Schur's test as in the proof of Lemma 8.

If we have 1), then by Lemma 13 (it is an analog of Lemma 8 for bi-tree) we will get from the last display

(53)
$$\sum_{Iodd,Iodd} (\alpha_0, h_I \otimes h_J)^2 \leq C\varepsilon_1^{-2} \|\text{sum in } (47)\|_2^2.$$

This is because $\varepsilon_1 >> \varepsilon_2$. And (53) is exactly (48).

At this moment we established the following dichotomy: either we have proved (48), or, otherwise we have $|(\mathbf{1}_U, \tilde{\mathbf{1}}_I \otimes T_2 \tilde{\mathbf{1}}_J)| \leq \varepsilon_1$ (for all I, J!), and we also have $|(\mathbf{1}_U, T_1 \tilde{\mathbf{1}}_I \otimes T_2 \tilde{\mathbf{1}}_J)| \leq \varepsilon_2$ (again for all I, J!). When we say "all" we mean all I, J such that $I \times J \subset U$. (We think here that U is connected, this does not restrict the generality as we can consider the connected components of U one by one.)

Now we call the attention of the reader that third and second lines of (47) are totally symmetric under the interchange of I and J. So we consider again two cases: either 1) $|(\mathbf{1}_U, T_1 \tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J)| \leq \varepsilon_1$ (for all I, J); or, 2) $|(\mathbf{1}_U, T_1 \tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J)| \geq \varepsilon_1$ (for all I, J).

In the second case we obtain a form of (53) symmetric under I versus J exchange:

(54)
$$\sum_{Iodd, Jodd} (\alpha_0, h_I \otimes h_J)^2 \le C\varepsilon_1^{-2} \|\text{sum in } (47)\|_2^2.$$

But (54) is exactly (48).

Now we are left with case 1), namely, $|(\mathbf{1}_U, T_1\tilde{\mathbf{1}}_I \otimes \tilde{\mathbf{1}}_J)| \leq \varepsilon_1$ (for all I, J). We remind that we came to the following conclusion: either we already proved (48) or we have two other sets of inequalities: $|(\mathbf{1}_U, \tilde{\mathbf{1}}_I \otimes T_2\tilde{\mathbf{1}}_J)| \leq \varepsilon_1$ (for all I, J!), and we also have $|(\mathbf{1}_U, T_1\tilde{\mathbf{1}}_I \otimes T_2\tilde{\mathbf{1}}_J)| \leq \varepsilon_2$ (again for all I, J!). Again, when we say "all" we mean all I, J such that $I \times J \subset U$.

So all three terms $|(\mathbf{1}_U, T_1\tilde{\mathbf{1}}_I \otimes T_2\tilde{\mathbf{1}}_J)|$, $|(\mathbf{1}_U, \tilde{\mathbf{1}}_I \otimes T_2\tilde{\mathbf{1}}_J)|$, $|(\mathbf{1}_U, T_1\tilde{\mathbf{1}}_I \otimes T_2\tilde{\mathbf{1}}_J)|$ for all I, J are small.

At this moment we consider the first line of (47). Notice that its output is in the (even, even) space by the definition of T_1^*, T_2^* .

All other three lines also have some of their output in (even, even) space. So we collect the factor in front of the same $h_{I'} \otimes h_{J'}$, I', J' being (even, even) from all four lines of (47).

We will have that the (even, even) part of the sum in (47) is exactly equal to the sum of orthogonal terms $(\hat{I}, \hat{J} \text{ are (even, even)})$:

$$[1 \cdot (b, h_I \otimes h_J) + \tilde{\varepsilon}_1(b, \dots) + \tilde{\varepsilon}_2(b, \dots) + \tilde{\varepsilon}_3(b, \dots)]h_{\hat{\iota}} \otimes h_{\hat{\iota}}$$

where I, J are children of of \hat{I}, \hat{J} correspondingly, where dots ... stand for some $h_{I_i'' \times J_i''}$, i = 1, 2, 3, and where $I_i'' \times J_i''$ is a close relative of $I \times J$. These close relatives will be $h_{\hat{I}} \otimes h_J$, $h_I \otimes h_{\hat{J}}$ and $h_{\hat{I}} \otimes h_{\hat{J}}$. Also $|\tilde{\varepsilon}_1| \leq \varepsilon_1 << \varepsilon_2$, $|\tilde{\varepsilon}_2| \leq \varepsilon_1 << \varepsilon_2$, $|\tilde{\varepsilon}_3| \leq \varepsilon_2$.

Hence (47) has the norm squared bounded from below by the sum

$$\sum_{I,J \ are \ odd,odd...} |1 \cdot (b,h_I \otimes h_J) + \tilde{\varepsilon}_1(b,h_{\hat{I}} \otimes h_J) + \tilde{\varepsilon}_2(b,h_I \otimes h_{\hat{J}}) + \tilde{\varepsilon}_3(b,h_{\hat{I}} \otimes h_{\hat{J}})|^2,$$

where the summation is over all those $I \times J$, that are inside U and I, J are both odd. As we have seen $|\tilde{\varepsilon}_1| \leq \varepsilon_1 << \varepsilon_2, |\tilde{\varepsilon}_2| \leq \varepsilon_1 << \varepsilon_2, |\tilde{\varepsilon}_3| \leq \varepsilon_2$. Using this smallness and Lemma 13 for double tree, we conclude that the square of the norm of the sum in (47) is bounded from below by the sum

$$c |(b, h_I \otimes h_J)|^2$$
,

where the summation is over all $I \times J$, say, at least 1-deep in U and and I, J are both odd, and we got what we wanted at least for odd, odd part of b, namely we got (46).

We just proved the following "theorem".

Theorem 7.6 Let T_1, T_2 be dyadic shifts introduced at the beginning of this article. There is a universal constant $C < \infty$ such that

$$||b^{odd,odd}||_{BMO_{ChF}} \le C ||[T_1^*, [T_2^*, b]]||.$$

Now if we have repeated commutator with not just one but with 4 shifts of our type (our shift is even even to odd odd, but there are three analogous shifts obviously with different input/output).

If repeated commutator of b with all these 4 shifts are bounded we conclude as before that the full $||b||_{BMO_{ChF}}$ norm is the estimate from above for the maximum of norms of those repeated commutators with multiplication by b. To formulate the theorem, let us denote the shifts we used above by

$$T_1 =: T_1^{eo}, T_2 =: T_2^{eo}$$
.

Notice that those shifts have one drawback: they have huge kernels. So it is easy to understand that may be repeated commutator looses some information about the symbol b.

However we have three more couple T_1^{oe} , T_2^{oe} , T_1^{eo} , T_2^{oe} , T_1^{oe} , T_2^{eo} . Then we can repeat the argument above for each of those couples. Then we get the "theorem":

Theorem 7.7 There s a universal constant $C < \infty$ such that

$$\|b\|_{BMO_{ChF}} \leq C \, \max \left[\|[T_1^{eo}, [T_2^{eo}, b]]\|, \|[T_1^{oe}, [T_2^{oe}, b]]\|, \|[T_1^{eo}, [T_2^{oe}, b]]\|, \|[T_1^{oe}, [T_2^{eo}, b]]\| \right].$$

It is well known that the estimates from above in this theorem also holds, see e.g. [LM].

7.2. Discussion of the deliberate mistakes.

- (1) In Lemmas 9, 11, and equation (49) we checked the claim for $R_1 = \hat{I} \times J$, and it is correct. But it is incorrect for and $R_2 = I \times \hat{J}$. In other words it is not true that one small coefficients automatically means that all of them are small. We can "spread smallness" horizontally, but not vertically. Lemmas 9, 11 were about T_1 . The similar claims about T_2 would allow to "spread smallness" vertically, but not horizontally. So we cannot "spread smallness" throughout the whole of connected set U.
- (2) In estimating the repeated paraproduct from below we can disregard all parts with D (see Section 8), as they are bounded by BMO_r norm of the symbol. This is true. Also it is trues that substituting $f = \mathbf{1}_U$ eliminates all ZZ sums. But it is not true that substituting $f = \mathbf{1}_U$ eliminates all πZ , $Z\pi$ terms of repeated paraproduct and leaves us only with $\pi\pi$ part. In fact, if one take a look at all the 16 first sums comprising repeated paraproduct except for parts with D (see the beginning of Section 9), we can see that πZ unfortunately does not cancel out completely. The term $\pi Z3$ contains $(T_1^*, \tilde{\mathbf{1}}_I \otimes h_J)$, and this term is not obliged to be zero. (But $(T_2^*, \tilde{\mathbf{1}}_I \otimes h_J)$ is zero.) Again $\pi Z4$ contains $(T_1^*T_2^*, \tilde{\mathbf{1}}_I \otimes h_J)$, and this term is not obliged to be zero. So $\pi Z3, \pi Z4$ survive, which is very unfortunate, they join all for lines of $\pi\pi$ survivors. Moreover, by the same reason $Z\pi 2$ and $Z\pi 4$ also survive. The more terms survive (now we have 6 of them, the more difficult is to separate their action.
- (3) At least all $\pi\pi$ terms are almost orthogonal. Adding $\pi Z3, \pi Z4, Z\pi 2$ and $Z\pi 4$, makes the survival part not to be the sum of almost orthogonal terms.
- (4) Let us assume for a while that we are left only with almost orthogonal part, that is, with $\pi\pi$. This would happen if we consider the repeated commutator, but not with multiplication by b but with bi-parameter paraproduct π_b . This is also interesting, in principle. Let us take a look how we can use almost orthogonality. So $f = \mathbf{1}_U$, $b = \alpha_0$ and we collect the terms in front of $h_K \otimes h_L$ in $\pi\pi$ all four

lines below. Here is the coefficient in front of $h_K \otimes h_L$. Recall that K, L are such that $\hat{K} \times \hat{L} \subset U$.

$$\operatorname{sgn}(K, \hat{K}) \operatorname{sgn}(L, \hat{L})(b, h_{\hat{K}} \otimes h_{\hat{L}}) \cdot (\mathbf{1}_{U}, \tilde{\mathbf{1}}_{\hat{K}} \otimes \tilde{\mathbf{1}}_{\hat{L}}) -$$

$$- \operatorname{sgn}(K, \hat{K})(b, h_{\hat{K}} \otimes h_{L})(\mathbf{1}_{U}, \tilde{\mathbf{1}}_{\hat{K}} \otimes T_{2}\tilde{\mathbf{1}}_{L}) -$$

$$- \operatorname{sgn}(L, \hat{L})(b, h_{K} \otimes h_{\hat{L}})(\mathbf{1}_{U}, T_{1}\tilde{\mathbf{1}}_{K} \otimes \tilde{\mathbf{1}}_{\hat{L}}) +$$

$$(b, h_{K} \otimes h_{L})(\mathbf{1}_{U}, T_{1}\tilde{\mathbf{1}}_{K} \otimes T_{2}\tilde{\mathbf{1}}_{L}) =$$

$$\pm (b, h_{\hat{K}} \otimes h_{\hat{L}}) -$$

$$\pm (b, h_{\hat{K}} \otimes h_{L})(\mathbf{1}_{U}, \tilde{\mathbf{1}}_{\hat{K}} \otimes T_{2}\tilde{\mathbf{1}}_{L}) -$$

$$\pm (b, h_{K} \otimes h_{\hat{L}})(\mathbf{1}_{U}, T_{1}\tilde{\mathbf{1}}_{K} \otimes \tilde{\mathbf{1}}_{\hat{L}}) +$$

$$+ (b, h_{K} \otimes h_{L})(\mathbf{1}_{U}, T_{1}\tilde{\mathbf{1}}_{K} \otimes T_{2}\tilde{\mathbf{1}}_{L})$$

$$=: x_{K,L}.$$

So we are able to conclude immediately that the boundedness of the repeated commutator with bi-parameter π_b (not with multiplication by b!) implies that

(55)
$$\sum_{K,L:\hat{K}\times\hat{L}\subset U} x_{K,L}^2 \leq \|\text{repeated commutator with } \pi_b\|^2 |U|.$$

But the ultimate goal would be to get a different inequality:

(56)
$$\sum_{K,L:\hat{K}\times\hat{L}\subset U} (b,h_{\hat{K}}\otimes h_{\hat{L}})^2 \leq \|\text{repeated commutator with } \pi_b\|^2|U| .$$

Those inequalities are close to each other: just by the previous calculation of $x_{K,L}$ above. But they are not the same. How to get rid of extra terms in $x_{K,L}$? Lemma 13 should help, but to apply it we need the smallness of "parasite" terms $(\mathbf{1}_U, \tilde{\mathbf{1}}_{\hat{K}} \otimes T_2 \tilde{\mathbf{1}}_L), (\mathbf{1}_U, T_1 \tilde{\mathbf{1}}_K \otimes \tilde{\mathbf{1}}_{\hat{L}}), (\mathbf{1}_U, T_1 \tilde{\mathbf{1}}_K \otimes T_2 \tilde{\mathbf{1}}_L)$. But the size of those terms is probably approximately 1, and as we already noticed Lemma 11, that dealt with this size, is only partially true.

(5) And a final difficulty is that in $(b, h_{\hat{K}} \otimes h_{\hat{L}})$ above we have only \hat{K}, \hat{L} (even, even) by the definition of our dyadic shifts.

8. The lists of building blocks of repeated commutators with ${f T}$ and ${f T}^*$

Here are all the blocks of repeated commutator with $T_1^* \otimes T_2^*$. The blocks of repeated commutator with $T_1 \otimes T_2$ are exactly the same, only stars are deleted. All parts with D letter are estimated by $\lesssim ||b||_{BMO_r}$. Under the localization to U_0 we put $f = \mathbf{1}_U$, where U is the enlargement of U_0 .

Then ZZ part vanishes and $Z\pi 3$. The rest stays and should be estimated from below. In the next section we show that this is easy if b has some extra property. In this section we show how difficult is the problem when this extra property is not assumed. However, there is a hope because the main part that is left after localization is $\pi\pi$ and it has the property of almost orthogonality.

$$\pi\pi : \sum (b, h_I \otimes h_J) \langle f \rangle_{I \times J} (T_1^* h_I) \otimes (T_2^* h_J)$$

$$- \sum (b, h_I \otimes h_J) \langle T_2^* f \rangle_{I \times J} (T_1^* h_I) \otimes (h_J)$$

$$- \sum (b, h_I \otimes h_J) \langle T_1^* f \rangle_{I \times J} (h_I) \otimes (T_2^* h_J)$$

$$+ \sum (b, h_I \otimes h_J) \langle T_1^* T_2^* f \rangle_{I \times J} (h_I) \otimes (h_J)$$

$$\pi Z : \sum (b, h_I \otimes h_J)(f, \tilde{\mathbf{1}}_I \otimes h_J)(T_1^* h_I) \otimes (T_2^* \tilde{\mathbf{1}}_J)$$

$$- \sum (b, h_I \otimes h_J)(T_2^* f, \tilde{\mathbf{1}}_I \otimes h_J)(T_1^* h_I) \otimes (\tilde{\mathbf{1}}_J)$$

$$- \sum (b, h_I \otimes h_J)(f, T_1^* \tilde{\mathbf{1}}_I \otimes h_J)(h_I) \otimes (T_2^* \tilde{\mathbf{1}}_J)$$

$$+ \sum (b, h_I \otimes h_J)(T_1^* T_2^* f, \tilde{\mathbf{1}}_I \otimes h_J)(h_I) \otimes (\tilde{\mathbf{1}}_J)$$

$$Z\pi : \sum (b, h_I \otimes h_J)(f, h_I \otimes \tilde{\mathbf{1}}_J)(T_1^* \tilde{\mathbf{1}}_I) \otimes (T_2^* h_J)$$

$$- \sum (b, h_I \otimes h_J)(T_2^* f, h_I \otimes \tilde{\mathbf{1}}_J)(T_1^* \tilde{\mathbf{1}}_I) \otimes (h_J)$$

$$- \sum (b, h_I \otimes h_J)(T_1^* f, h_I \otimes \tilde{\mathbf{1}}_J)(\tilde{\mathbf{1}}_I) \otimes (T_2^* h_J)$$

$$+ \sum (b, h_I \otimes h_J)(T_1^* T_2^* f, h_I \otimes \tilde{\mathbf{1}}_J)(\tilde{\mathbf{1}}_I) \otimes (h_J)$$

$$ZZ : \sum (b, h_I \otimes h_J)(f, h_I \otimes h_J)(T_1^* \tilde{\mathbf{1}}_I) \otimes (T_2^* \tilde{\mathbf{1}}_J)$$

$$- \sum (b, h_I \otimes h_J)(f, T_2^* h_I \otimes h_J)(T_1^* \tilde{\mathbf{1}}_I) \otimes (\tilde{\mathbf{1}}_J)$$

$$- \sum (b, h_I \otimes h_J)(T_1^* f, h_I \otimes h_J)(\tilde{\mathbf{1}}_I) \otimes (T_2^* \tilde{\mathbf{1}}_J)$$

$$+ \sum (b, h_I \otimes h_J)(T_1^* T_2^* f, h_I \otimes h_J)(\tilde{\mathbf{1}}_I) \otimes (\tilde{\mathbf{1}}_J)$$

$$\pi D : -\sum (b, h_I \otimes h_{\hat{J}})(f, \tilde{\mathbf{1}}_I \otimes h_J) \frac{1}{\sqrt{|\hat{J}|}} (T_1^* h_I) \otimes h_{\hat{J}}$$
$$+ \sum (b, h_I \otimes h_{\hat{J}})(T_1^* f, \tilde{\mathbf{1}}_I \otimes h_J) \frac{1}{\sqrt{|\hat{J}|}} (h_I) \otimes h_{\hat{J}}$$

$$ZD: -\sum (b, h_I \otimes h_{\hat{J}})(f, h_I \otimes h_J) \frac{1}{\sqrt{|\hat{J}|}} (T_1^* \tilde{\mathbf{1}}_I) \otimes h_{\hat{J}}$$
$$+ \sum (b, h_I \otimes h_{\hat{J}}) (T_1^* f, h_I \otimes h_J) \frac{1}{\sqrt{|\hat{J}|}} (\tilde{\mathbf{1}}_I) \otimes h_{\hat{J}}$$

$$D\pi : -\sum (b, h_{\hat{I}} \otimes h_{J})(f, h_{I} \otimes \tilde{\mathbf{1}}_{J}) \frac{1}{\sqrt{|\hat{I}|}} (h_{\hat{I}}) \otimes T_{2}^{*} h_{J}$$
$$+ \sum (b, h_{\hat{I}} \otimes h_{J})(T_{2}^{*} f, h_{I} \otimes \tilde{\mathbf{1}}_{J}) \frac{1}{\sqrt{|\hat{I}|}} (h_{\hat{I}}) \otimes h_{J}$$

$$DZ : -\sum (b, h_{\hat{I}} \otimes h_J)(f, h_I \otimes h_J) \frac{1}{\sqrt{|\hat{I}|}} (h_{\hat{I}}) \otimes (T_2^* \tilde{\mathbf{1}}_J)$$
$$+ \sum (b, h_{\hat{I}} \otimes h_J)(T_2^* f, h_I \otimes h_J) \frac{1}{\sqrt{|\hat{I}|}} (h_{\hat{I}}) \otimes (\tilde{\mathbf{1}}_J)$$

$$DD: \sum (b, h_{\hat{I}} \otimes h_J)(f, h_I \otimes h_J)h_{\hat{I}} \otimes h_{\hat{J}}$$

9. Why everything works for b skipping scales?

As nothing works for general b, let us assume that our b skips scales. We mean here extra assumption that

(57)
$$(b, h_I \otimes h_J) \neq 0 \Rightarrow IJ$$
, are even, even.

In this special case we get the following

Theorem 9.1 Let b satisfy (57). Then for any dyadic open set U_0 we have

$$\sum_{I \times J \subset U_0} (b, h_I \otimes h_J)^2 \le C_1 \|\mathcal{T}^b\|^2 |U_0| + C_2 \|b\|_{BMO_r}^2 |U_0|.$$

Proof. Just in front of this section the reader can see the table of all parts of the commutator. It is easy to prove that all parts $D\pi$, πD , ZD, DZ are bounded by $||b||_{BMO_r}^2|U_0$. The DD part is bounded just be

$$\sup_{I,J} \frac{(b, h_I \otimes h_J)^2}{|I||J|} ||f||_2^2,$$

which is bounded by $||b||_{BMO_r}^2 ||f||_2^2$ again.

If b is reduced to α_0 —the part of b with Haar spectrum in U_0 . We can consider such reduced $b = \alpha_0$ by Lemma 2. At the same times this means that I, J in $\pi\pi$, $\pi Z, Z\pi$, ZZ parts are such that $I \times J \subset U_0$.

If $f = \mathbf{1}_U$, where U is chosen in (4.6), $|U| \leq C|U_0|$, then ZZ part vanishes, if b is reduced to α_0 —the part of b with Haar spectrum in U_0 .

If $f = \mathbf{1}_U$, in πnZ part the first and the second lines vanish, but $\pi Z3$, $\pi Z4$ do not vanish, in $Z\pi$ part the first and the third lines vanish but $Z\pi 2$, $Z\pi 4$ do not vanish.

Notice that by assumption (57) $\pi Z3$, $\pi Z4$ are in (even, any), and $Z\pi 2$, $Z\pi 4$ are in (any, even).

Let us take a look at the $\pi\pi$ 1-the first line of $\pi\pi$ part. It is is (odd, odd) part, and so it is orthogonal to $\pi Z3, \pi Z4, Z\pi 2, Z\pi 4$.

It is also orthogonal to $\pi\pi2$, $\pi\pi3$, $\pi\pi4$ -the second, third and fourth lines of $\pi\pi$ part. In fact, by assumption (57) $\pi\pi2$ is in (odd, even), $\pi\pi3$ is in (even, odd), $\pi\pi2$ is in (even, even) parts. Therefore,

(58)
$$\sum_{I \times J \subset U_0} (\alpha_0, h_I \otimes h_J)^2 \leq \|\pi \pi 1(\alpha_0, \mathbf{1}_U)\|_2^2 \leq \|\pi \pi(\alpha_0, \mathbf{1}_U) + \pi Z(\alpha_0, \mathbf{1}_U) + Z\pi(\alpha_0, \mathbf{1}_U)\|_2^2.$$

On the other hand,

$$\sup_{\|\rho\|_2 = 1, \sigma_H(\rho) \subset U_0} |(\mathbf{1}_U, \mathcal{T}_*^b(\rho))| = \sup_{\|\rho\|_2 = 1, \sigma_H(\rho) \subset U_0} |(\mathbf{1}_U, \mathcal{T}_*^{\alpha_0}(\rho))| =$$

$$\|\pi\pi(\alpha_0, \mathbf{1}_U) + \pi Z(\alpha_0, \mathbf{1}_U) + Z\pi(\alpha_0, \mathbf{1}_U) + ZZ(\alpha_0, \mathbf{1}_U) + \text{parts with } D(\alpha_0, \mathbf{1}_U)\|_2 \ge$$

$$\|\pi\pi(\alpha_0, \mathbf{1}_U) + \pi Z(\alpha_0, \mathbf{1}_U) + Z\pi(\alpha_0, \mathbf{1}_U)\|_2 - \|\text{parts with } D(\mathbf{1}_U, \alpha_0)\|_2.$$

Now we already noticed that all parts $D\pi$, πD , ZD, DZ, DD are bounded by $||b||_{BMO_r}^2 |U_0$, and $ZZ(\alpha_0, \mathbf{1}_U) = 0$. This and (58) prove theorem.

Theorem 9.2 Let b satisfy (57). Then for any dyadic open set U_0

$$\sum_{I \times J \subset U_0} (b, h_I \otimes h_J)^2 \le C_1 \|\mathcal{T}^b\|^2 |U_0|.$$

Proof. Let us consider two cases: 1) $||b||_{BMO_r} \le \frac{1}{4C_2} ||b||_{BMO_{CHF}}$, 2) $||b||_{BMO_r} \ge \frac{1}{4C_2} ||b||_{BMO_{CHF}}$. In the second case we use Section 3 and conclude the claim of the theorem.

In the first case we use Theorem 9.1. We choose the dyadic open set U_0 that gives us the Chang-Fefferman BMO norm of b, and we write down the conclusion of Theorem 9.1. The left hand side of

$$\sum_{I \times J \subset U_0} (b, h_I \otimes h_J)^2 \le C_1 \|\mathcal{T}^b\|^2 |U_0| + C_2 \|b\|_{BMO_r}^2 |U_0|$$

is $||b||_{BMO_{ChF}}^2 |U_0|$ because of extremal property of U_0 . Now we use $||b||_{BMO_r} \leq \frac{1}{4C_2} ||b||_{BMO_{CHF}}$ to absorb $C_2 ||b||_{BMO_r}^2 |U_0|$ into the left hand side. It is well known that the estimates from above in this theorem also holds, see e.g. [LM].

10. A COUNTEREXAMPLE

Many results suffer from the same gap as [FL], if they all rely on the claim of the form that follows or a similar claim:

(59)
$$\|\phi\|_2 \lesssim \|P\phi\|_2$$
,

where P is a projection given by a Fourier multiplier whose symbol is the indicator of a product of half-spaces (in the original case of [FL], simply $\mathbb{R}_+ \times \mathbb{R}_+ \subset \mathbb{R}^2$) and the Fourier transform of ϕ is "symmetric". In practice, ϕ takes the form $\phi = |\alpha|^2$ for some function α analytic in \mathbb{C}^2_+ . Obviously ϕ is non-negative and in particular real-valued, so some symmetry of Fourier transform should follow.

In the one parameter case the simple symmetry shows that the previous claim is obviously correct. But not so in two parameter case anymore, as an example below shows.

The statement (59) should be avoided (in this generality $\phi = |\alpha|^2$, α being nice analytic function in \mathbb{C}_+^2) in the future corrected proof. However, something like that should be in the proof, as it is the only estimate from below in the whole reasoning: all other estimates in [FL] are the estimates of operators from above.

Let α be an analytic function in $\mathbb{C}_+ \times \mathbb{C}_+$ and let it be just Fourier transform of $\psi(\xi), \xi = (\xi_1, \xi_2) \in \mathbb{R}^2_{++}, \psi \in L^2(d\xi_1 d\xi_2)$. Let the support of ψ be in $Q := [0, 2]^2$. In what follows $d\xi = d\xi_1 d\xi_2$.

$$|\alpha(x_1, x_2)|^2 = \int \int e^{ix_1(\xi_1 - \eta_1) + ix_2(\xi_2 - \eta_2)} \psi(\xi) \bar{\psi}(\eta) d\xi d\eta =$$

$$\int \int e^{ix_1 t_1 + ix_2 t_2} \psi(\xi_1, \xi_2) \overline{\psi(\xi_1 - t_1, \xi_2 - t_2)} d\xi_1 d\xi_2 dt_1 dt_2$$

Hence Fourier transform

$$(\mathcal{F}|\alpha|^2)(t_1, t_2) = \int_{Q_0} \psi(\xi_1, \xi_2) \overline{\psi(\xi_1 - t_1, \xi_2 - t_2)} d\xi_1 d\xi_2$$

If ψ is a characteristic function of ϵ neighborhood of $\xi_1 + \xi_2 = 1, \xi_i \geq 0$, then for positive vectors t this quantity is uniformly bounded by $C\epsilon$. Moreover, this quantity is zero if positive vector t has norm bigger than $C\epsilon$.

Hence

$$||P_{++}|\alpha|^2||_2 \simeq \sqrt{\epsilon^2 \cdot \epsilon^2} = \epsilon^2$$

On the other hand,

$$\|\alpha\|_2 \simeq \epsilon^{1/2}$$
.

Also let us look at $P_{+-}|\alpha|^2$. Now we look at

$$(\mathcal{F}|\alpha|^2)(t_1, -t_2) = \int_{Q_0} \psi(\xi_1, \xi_2) \overline{\psi(\xi_1 - t_1, \xi_2 + t_2)} d\xi_1 d\xi_2$$

for positive vector t. We can see that this is of order ϵ for positive vectors t such that they fill-in the domain inside angle of size ϵ and diameter 1 and of total measure of order ϵ . Thus

$$||P_{+-}|\alpha|^2||_2^2 \simeq \epsilon^3$$
.

So

$$\|\alpha\|_4^2 = \||\alpha|^2\|_2 \ge \epsilon^{3/2}$$
.

In fact,

$$\|\alpha\|_4^2 = \||\alpha|^2\|_2 \approx \epsilon^{3/2}$$
.

So both inequalities below are false:

$$||P_{++}|\alpha|^2||_2 \ge c||\alpha||_2^2; \quad ||P_{++}|\alpha|^2||_2 \ge c||\alpha||_4^2.$$

Normalization of α does not help. Let $\tilde{\alpha} = \epsilon^{-1/2} \alpha$. Then

$$\begin{split} \|\tilde{\alpha}\|_2 &\asymp 1, \\ \|\tilde{\alpha}\|_4^2 &= \||\tilde{\alpha}|^2\|_2 \asymp \epsilon^{1/2} \,. \end{split}$$

And

$$||P_{++}|\tilde{\alpha}|^2||_2 \asymp \epsilon$$
.

So the inequality below is false:

$$||P_{++}|\tilde{\alpha}|^2||_2 \ge c.$$

Even though $\|\tilde{\alpha}\|_2 \approx 1$.

Thus the inequality on page 208 line 4 top of [L] cannot be true as written. Similarly, inequalities in [FL] on page 149 line 3 at the end of the page cannot be true as written.

Symmetry works in estimating from below positive frequency part of the Fourier (of real function) via the whole Fourier transform of this function perfectly well for functions of one variable. But for functions of two variables this does not work. Even if the function is a) positive and b) the modulus squared of analytic function in $\mathbb{C}_+ \times \mathbb{C}_+$ as the example above shows.

One can guess what is missing is one more normalization (which we do not have in the example above):

$$(60) 1 \approx \|\tilde{\alpha}\|_2 \lesssim \|\tilde{\alpha}\|_4.$$

However, there is a modification of this counterexample suggested by T. Hytönen that has this property as well.

References

- [AHMV] N. Arcozzi; I. Holmes; P. Mozolyako; A. Volberg, Bi-parameter embedding and measures with restricted energy conditions. Math. Ann. 377 (2020), no. 1-2, 643-674.
- [AMPS18] N. Arcozzi, P. Mozolyako, K.-M. Perfekt, and G. Sarfatti, *Bi-parameter Potential theory and Carleson measures for the Dirichlet space on the bidisc.* Preprint. Nov. 10018. arXiv: 1811.04990.
- [BP] O. Blasco; S. Pott, Dyadic BMO on the bidisk. Rev. Mat. Iberoamericana 21 (2005), no. 2, 483–510.
- [Car] L. Carleson, A counterexample for measures bounded on H^p for the bi-disc. Mittag-Leffler Report 1974, no. 7.
- [Ch] Sun-Yung A. Chang Carleson measure on the bi-disc. Ann. of Math. (2) v. 109, no. 3, 1979, 613-6100.
- [ChF] Sun-Yung A. Chang, R. Fefferman, A continuous version of duality of H¹ with BMO on the bidisc. Ann. of Math. (2) v. 112, no. 1, 1980, 179–1001.
- [ChF2] Sun-Yung A. Chang, R. Fefferman, Some recent developments in Fourier analysis and Hp-theory on product domains. Bull. Amer. Math. Soc. (N.S.) 12, 1985, 1–43.
- [DP] L. Dalenc and S. Petermichl. A lower bound criterion for iterated commutators. J. Funct. Anal., 266(8):5300–5320, 2014.
- [DKPS] K. DOMELEVO, S. KAKOROUMPOS, S. PETERMICHL, O. SOLER I GIBERT, Dyadic little BMO estimates, Preprint. Dec. 2020.
- [FS] Ferguson, Sarah H.; Sadosky, Cora Characterizations of bounded mean oscillation on the polydisc in terms of Hankel operators and Carleson measures. J. Anal. Math. 81 (2000), 239–267.
- [DLWY] X. T. DUONG, J. LI, BRETT D. WICK, D. YANG, Commutators, Little BMO and weal factorization, arXiv1609.0784v2.
- [DP] L. DALENC AND S. PETERMICHL. A lower bound criterion for iterated commutators. J. Funct. Anal., 266(8):5300-5320, 2014.

- [FS] FERGUSON, SARAH H.; SADOSKY, CORA Characterizations of bounded mean oscillation on the polydisc in terms of Hankel operators and Carleson measures. J. Anal. Math. 81 (2000), 239–267.
- [FL] S. FERGUSON, M. LACEY, A characterization of product BMO by commutators, Acta Math. 189 (2002), 143–160.
- [HPW] I. HOLMES, S. PETERMICHL, B. WICK, Weighted little bmo and two-weight inequalities for Journé commutators, Analysis&PDE 11 No. 7, 2018.
- [HTV] I. Holmes, S. Treil, A. Volberg, Dyadic bi-parameter simple commutator and dyadic little bmo, arXiv:2012.05376, pp. 1–14.
- [JLJ] J.-L. JOURNÉ, Two problems of Calderón-Zygmund theory on product-spaces. Ann. Inst. Fourier (Grenoble), v. 38, 1988, no. 1, 111–132.
- [JLJ2] J.-L. JOURNÉ, Calderoón–Zygmund operators on product spaces. Rev. Mat. Iberoamericana 1, 1985, 55–91.
- [LM] M.LACEY, J. METCALFE, Paraproducts in one and several parameters, Forum Math. 19 (2007), 325–351.
- [L] M. LACEY, Lectures on Nehari's Theorem on the Polydisc, Contemporary Mathematics, Volume 444, 2007, pp. 185–213.
- [MPV] P. MOZOLYAKO; G. PSAROMILIGKOS; A. VOLBERG, Counterexamples for multi-parameter weighted paraproducts. C. R. Math. Acad. Sci. Paris 358 (2020), no. 5, 529–534.
- [MPVZ] P. MOZOLYAKO; G. PSAROMILIGKOS; A. VOLBERG; P. ZORIN-KRANICH, Combinatorial property of all positive measures in dimensions 2 and 3. C. R. Math. Acad. Sci. Paris 358 (2020), no. 6, 721–725.
- [MPTT1] C. Muscalu, J. Pipher, T. Tao, C. Thiele, Bi-parameter paraproducts, Acta Math., 193 (2004), 269–296.
- [MPTT2] C. Muscalu, J. Pipher, T. Tao, C. Thiele, Multi-parameter paraproducts, Rev. Mat. Iberoamericana 22 (2006), no. 3, 963–976
- [LPPW] M. T. LACEY, S. PETERMICHL, J. C. PIPHER, AND B. D. WICK. Multiparameter Riesz commutators. Amer. J. Math., 131(3):731–769, 2009.
- [LT] M. T. LACEY AND E. TERWILLEGER. Hankel operators in several complex variables and product BMO. Houston J. Math., 35(1):159–183, 2009.
- [OPS] Y. Ou, S. Petermichl, and E. Strouse. Higher order Journé commutators and characterizations of multi-parameter BMO. Adv. Math., 291:24–58, 2016.
- [Tao] T. Tao, Dyadic product H¹, BMO, and Carleson's counterexample, preprint, pp. 1-12.
- [V] A. Volberg, Non-symmetry for Fourier transform in two variables, preprint 2019, pp.1–2.

Department of Mathematics, Texas A&M University, College Station, TX 77843-3368

DEPARTMENT OF MATHEMATICS, BROWN UNIVERSITY, 151 THAYER STR./BOX 1917, PROVIDENCE, RI 02912, USA Email address: treil@math.brown.edu (S. Treil)

Department of Mathematics, Michigan Sate University, East Lansing, MI. 48823 Email address: volberg@math.msu.edu (A. Volberg)