


Latest results on invisibly decaying Higgs bosons

Patrick Dunne - Imperial College London
on behalf of the ATLAS and CMS Collaborations
DM@LHC 2016 - 31/03/2016



Outline

- ▶ How to search for invisibly decaying Higgs bosons:
 - direct and indirect searches
- ▶ Overview of Run 1 results from ATLAS and CMS
- ▶ New Run 2 results from CMS
- ▶ Projections of future sensitivity (NB all limits at 95% CL unless stated otherwise)


Why look for invisibly decaying Higgs bosons?

Theoretical Motivations

- ▶ All massive SM particles get their mass through Higgs boson couplings
- ▶ Why not dark matter?

Experimental motivation

- ▶ Measurements of the Higgs boson made so far are impressive:
 - Mass measured with 0.2% error
- ▶ A lot of parameters are still relatively unconstrained:
 - Indirect limit on width is $\sim 4\Gamma_{SM}$
- ▶ Plenty of room for Higgs boson couplings to dark matter



CMS-PAS-HIG-15-002
ATLAS-CONF-2015-044


How to search for invisibly decaying Higgs bosons

Indirect searches

- ▶ Compare visible width to total width:

$$- \text{BR}_{\text{BSM}} = \frac{\Gamma_H - \Gamma_{\text{vis}}}{\Gamma_H}$$

- ▶ No measurement of Γ_H , need to make an assumption
- Usually assume SM width
- ▶ ATLAS+CMS combination gives an observed (expected) limit on BR_{BSM} of 0.34 (0.35)



CMS-PAS-HIG-15-002
ATLAS-CONF-2015-044


How to search for invisibly decaying Higgs bosons

Direct searches

- ▶ Look for associated Higgs boson products plus E_T^{miss}


Production channels

- ▶ VBF mode is most sensitive
 - Second highest rate and distinctive topology
- ▶ Gluon fusion has no visible products, needs ISR
 - High rate, difficult final state
- ▶ VH has clean final states but low rate



Run 1 ATLAS direct searches - $Z(\ell\ell)H$


- Selects two leptons opposite large E_T^{miss}
- E_T^{miss} shape analysis with data driven backgrounds
- Observed (expected) limit on $\mathcal{B}(H \rightarrow inv.)$ for $m_H = 125.5$ GeV is 75 (62)%



PRL 112, 201802 (2014)

Run 1 ATLAS direct searches - $V(\text{had})H$

- Targets $W/Z \rightarrow qq$ final state
- E_T^{miss} and dijet p_T shape analysis with data driven backgrounds
- Observed (expected) limit on $\mathcal{B}(H \rightarrow \text{inv.})$ for $m_H = 125$ GeV is 78 (86)%



Eur. Phys. J. C (2015) 75:337

Run 1 ATLAS direct searches - VBF


- ▶ Select two jets with large $\Delta\eta$ opposite large E_T^{miss}
- ▶ Counting experiment with data driven background estimation
- $W \rightarrow e\nu$, $W \rightarrow \mu\nu$ and $W \rightarrow \tau\nu$ and $Z \rightarrow \nu\nu$ normalisation tied together
- ▶ Observed (expected) limit on $\mathcal{B}(H \rightarrow inv.)$ for $m_H = 125$ GeV is 28 (31)%

Signal region Process	SR1	SR2a	SR2b
ggF signal	20 ± 15	58 ± 22	19 ± 8
VBF signal	286 ± 57	182 ± 19	105 ± 15
$Z(\rightarrow \nu\nu)+jets$	339 ± 37	1580 ± 90	335 ± 23
$W(\rightarrow \ell\nu)+jets$	235 ± 42	1010 ± 50	225 ± 16
Multijet	2 ± 2	20 ± 20	4 ± 4
Other backgrounds	1 ± 0.4	64 ± 9	19 ± 6
Total background	577 ± 62	2680 ± 130	583 ± 34
Data	539	2654	636

JHEP 01 (2016) 172

Run 1 ATLAS direct searches - Combination


- ▶ Combining searches significantly improves limits
- ▶ Direct searches provide most sensitivity
 - Observed (expected) limit on $\mathcal{B}(H \rightarrow \text{inv.})$ for $m_H = 125$ GeV is 25 (27)%
- ▶ Adding indirect results adds assumption on Higgs total width
 - Observed (expected) limit on $\mathcal{B}(H \rightarrow \text{inv.})$ for $m_H = 125$ GeV is 23 (24)%



JHEP11(2015)206

Run 1 CMS direct searches - ZH


- ▶ Searches in $Z \rightarrow \ell\ell$ and $Z \rightarrow b\bar{b}$ channels
- ▶ $Z(\ell\ell)H$ search is a 2D shape analysis with data driven backgrounds
- ▶ $Z(b\bar{b})H$ search is a BDT shape analysis with data driven backgrounds
- ▶ Combined ZH searches observed (expected) limit on $\mathcal{B}(H \rightarrow \text{inv.})$ for $m_H = 125$ GeV is 81 (83)%



Eur. Phys. J. C 74 (2014) 2980

Run 1 CMS direct searches - Monojet+V(had)H


- Search has categories targeting $V(had)H$ and ggH production modes
- E_T^{miss} shape analysis with data driven background estimation
- Observed (expected) limit on $\mathcal{B}(H \rightarrow inv.)$ for $m_H = 125$ GeV is 53 (62)%



CMS-PAS-EXO-12-055

Run 1 CMS direct searches - VBF



- ▶ Select two jets with large $\Delta\eta$ separated from large E_T^{miss}
- ▶ Update of first search Eur. Phys. J. C 74 (2014) 2980
- ▶ Dedicated “parked data” trigger
- ▶ Counting experiment with data driven backgrounds
- ▶ V+jets backgrounds separately normalised
- ▶ Observed (expected) limit on $\mathcal{B}(H \rightarrow inv.)$ for $m_H = 125$ GeV is 57 (40)%
 - If all normalisations had same uncertainty as $W \rightarrow \mu\nu$ expected limit would be 33%



CMS-PAS-HIG-14-038

Run 1 CMS direct searches - Combination


- Combine by production mode as well as full combination
 - ggH-tagged is monojet, VH-tagged is $Z(\ell\ell)H+Z(bb)H+V(\text{had})H$, VBF-tagged is VBF
- Obs. (exp.) limit on $\mathcal{B}(H \rightarrow \text{inv.})$ at $m_H = 125$ GeV is 36 (30)%



CMS-PAS-HIG-15-012

Run 2 CMS direct searches - ZH



- ▶ Targets $Z \rightarrow \ell\ell$ final state
- ▶ 2D shape analysis with leading backgrounds estimated using MC
- ▶ Observed (expected) limit on $\mathcal{B}(H \rightarrow inv.)$ for $m_H = 125$ GeV is 124 (124)%



CMS-PAS-HIG-16-008

Run 2 CMS direct searches - VBF



- Dedicated trigger used again
- Counting experiment with data driven background estimation
- V+jets backgrounds all taken to have same normalisation
- Observed (expected) limit on $\mathcal{B}(H \rightarrow inv.)$ for $m_H = 125$ GeV is 69 (62)%



CMS-PAS-HIG-16-009

Run 2 CMS direct searches - Combination


- ▶ First CMS analysis combining 8 and 13 TeV results
- ▶ Limit calculated both by production mode and overall
- ▶ Combined observed (expected) limit on $\mathcal{B}(H \rightarrow inv.)$ for $m_H = 125$ GeV is 32 (26)%



CMS-PAS-HIG-16-009


Projections

- CMS VBF analysis projected to increased luminosity at 13 TeV
- If systematics scale as $\sqrt{\mathcal{L}}$ can exclude $\mathcal{B}(H \rightarrow inv.) = 5\%$ with full LHC dataset





arXiv:1603.07739

- ▶ Several models for interpreting invisible Higgs limits
- ▶ “Higgs portal” has been quite commonly used
 - Assume 125 GeV Higgs acts as mediator between visible and dark matter sectors
 - Assume scalar, fermion or vector dark matter
- ▶ Scalar dark matter would mix with Higgs boson
 - mixing angle must be small
- ▶ Vector dark matter width goes to infinity as mass decreases



Dark matter interpretations - Projections

- ▶ Models with electroweak couplings studied: favour VBF channel
- ▶ VBF topology allows looser E_T^{miss} selection making EFTs valid
- ▶ Also investigate simplified models with scalar/pseudoscalar mediator
- ▶ Projections of CMS VBF channel sensitivity at several luminosities



arXiv:1603.07739

Summary

- ▶ Both collaborations are sensitive to $\mathcal{B}(H \rightarrow inv.) \sim 25\%$ with current datasets
 - Current 95% CL upper observed (expected) limits from direct searches are CMS: 32 (26)%, ATLAS: 25 (27) %
 - Combination of channels allow sensitivity to be greatly improved
- ▶ Projected limit on $\mathcal{B}(H \rightarrow inv.) \sim 10\text{-}20\%$ from VBF alone by the end of LHC Run 2 and 5% by end of LHC running assuming systematics scale as $\sqrt{\mathcal{L}}$