

VBF Higgs to Invisible - Update


AN-14-243

P. Dunne on behalf of the VBF $H \rightarrow$ invisible analysis group

Why Higgs to Invisible?

Experimental motivation

- ▶ Current measurements of the 125 GeV Higgs boson are compatible with Standard Model (SM) expectations
- large uncertainties can still accommodate significant beyond the SM (BSM) properties
- ▶ Additional Higgs bosons with exotic decays are not excluded



Theoretical motivation



- ▶ Many BSM theories predict Higgs boson decays to invisible final states:
 - e.g. SUSY, extra dimensions, fourth-generation neutrinos
- ▶ These final state particles are often dark matter candidates

Talk outline

- ▶ For run 1 we had two sets of triggers
 - Prompt trigger used for current published result: HIG-13-30
 - Parked triggers analysis progress presented today


Overview

- ▶ Reminder of prompt analysis
- ▶ Why we chose our new analysis strategy
- ▶ Details of parked analysis
 - Emphasis on changes from established prompt analysis
- ▶ Brief look at some other analysis techniques we are investigating



Prompt Analysis

- ▶ Single bin counting experiment
 - Signal region chosen to eliminate QCD and be above trigger turn ons
- ▶ Major backgrounds use data driven estimates:
- ▶ $Z \rightarrow \nu\nu, W \rightarrow \ell\nu$, QCD
- ▶ Minor backgrounds taken from MC:
- ▶ $VV, W\gamma, t\bar{t}$, single top
- ▶ Expected limit 49% at $m_H = 125$ GeV



Data driven background estimation

$$W: N_S = N_S^{MC} \frac{N_C^{Data} - N_C^{Bkg}}{N_C^{MC}}$$

Z:

$$N_S^{Z \rightarrow \nu\nu} = (N_C^{Data} - N_C^{bkg}) \cdot \frac{\sigma(Z \rightarrow \nu\nu)}{\sigma(Z/\gamma^* \rightarrow \mu\mu)} \cdot \frac{\epsilon_S^{ZMC}}{\epsilon_C^{ZMC}}$$

Parked Triggers

- ▶ Use already analysed prompt trigger for run A
- ▶ One parked trigger for runs B and C, another for run D
- ▶ All parked and prompt triggers are seeded by L1ETM40
- ▶ Parked triggers have looser HLT thresholds
- ▶ This allows us to look at new regions of phase space and different analysis techniques

HLT

Run period	MET cut	dijet p_T cut	dijet mass cut
A	METnoMuons > 65 GeV	DiPFJet40	MJJ800
B&C	N/A	DiJet35	MJJ700
D	N/A	DiJet30	MJJ700

Software framework strategy

Prompt analysis

- ▶ Two frameworks: Analyses A and B
- ▶ independent ntuples and analysis code

Parked analysis

- ▶ Insufficient manpower to maintain and develop two frameworks
- ▶ Moved to one fully developed framework
 - New framework is development of analysis B and uses same ntuples
- ▶ Synchronised yields in signal and control regions between new framework and old analyses A and B
- ▶ Repeated expected limit calculation from HIG-13-030 analysis with the new framework and parked data
 - Agrees with HIG-13-030 to within 2%, which is good given rereco, and change of global tag and triggers

Analysis strategy

Initial investigations

- ▶ Planned to define a loose pre-selection and model QCD shape
- ▶ Several options for analysis strategy:
 - Rectangular cuts and counting experiment
 - Rectangular cuts and shape experiment
 - MVA and counting experiment
 - MVA and shape experiment
- ▶ Due to trigger conditions no appropriate QCD control region found - details later

Final plan for parked data

- ▶ During QCD studies found new variables which remove QCD
 - Signal efficiency significantly higher than prompt selection
- ▶ Remaining backgrounds very signal like in variables studied so far
 - Opted for cut and count analysis

QCD options tried

Several methods tried to model QCD

Standard MC

- doesn't have enough events

Private VBF+MET enriched QCD MC sample

- Can only enrich in events with real met
- Can't model met from mismeasurement

Data-driven shape using different jet pairs in the event

- Jet kinematics are very biased
- Ordering in p_T and angle have been tried
- Reweighting individual distributions to fix others has been tried

Changes since prompt analysis

Trigger

- ▶ Parked trigger efficiency has been measured including variable correlation
 - This allows the trigger turn on region to be used

Signal region


- ▶ The signal region has been reoptimised for the looser parked triggers
 - New region uses new variable has higher signal efficiency with much less QCD

Background estimation

- ▶ A top control region has been added
- ▶ Minor modifications made to $W \rightarrow \tau\nu$ background estimation method
- ▶ QCD background estimation method changed
- ▶ $W\gamma$ contribution found to be modelled already by our $W \rightarrow \ell\nu$ Monte Carlo




Trigger efficiency

- ▶ The variables used in prompt and parked triggers are highly correlated:
 - dijet mass, METnoMU, jet 2 p_T
- ▶ In the prompt analysis we neglected correlations and cut to ensure trigger was $> 95\%$ efficient
- ▶ For the parked analysis we use a 2D binning in dijet mass and jet 2 p_T
 - MJJ: 0,600,800,900,1000,5000
 - Jet 2 p_T : 30,40,50,60,1000
- ▶ In each bin we fit the METnoMU trigger turn on using an error function
- ▶ We then combine the turn ons from runs A, BC and D weighted by luminosity and apply this to MC events




Preselection choice

- ▶ Trigger turn ons and detector acceptance impose the following cuts:
 - $\eta_{j1} \cdot \eta_{j2} < 0$, $|\eta_{j1,2}| < 4.7$, jet 1 $p_T > 50$ GeV, $\Delta\eta_{jj} > 3.6$, jet 2 $p_T > 40$ GeV, $MET_{nomu} > 90$ GeV, $M_{jj} > 800$ GeV
- ▶ As in the prompt analysis we also veto events with 'veto' electrons or muons
- ▶ QCD in plots is VBF enriched MC doesn't model all QCD
- ▶ Poor data-MC agreement from QCD contamination motivates the following additional cuts:
- ▶ $\frac{MET_{nomu}}{\sigma_{MET_{nomu}}} > 3.0$, $\text{Min}\Delta\phi(\text{all jets } p_T > 30 \text{ GeV}, MET_{nomu}) > 1.0$, $M_{jj} > 1000$ GeV




Signal region selection

- ▶ Can't model QCD shape so alter cuts to remove most QCD
 - Can then tolerate a larger uncertainty on QCD estimation
- ▶ Remaining QCD in region $\frac{MET_{noMU}}{\sigma_{MET_{noMU}}} < 4$ and $Min\Delta\phi(all\ jets, MET_{noMU}) < 2.0$
- ▶ Signal contribution also large for some regions of parameter space
 - We blind these regions and use as a basis for signal region optimisation



Signal region selection



- ▶ We optimise by choosing the cut values with the best 95% C.L. expected limit
 - Limit calculation details later
- ▶ We scanned through jet 2 p_T , dijet mass and $\text{Min}\Delta\phi(\text{all jets}, \text{MET}_{\text{no mu}})$
- ▶ Best limit was found for:
 - No additional jet 2 p_T cut
 - $\frac{\text{MET}_{\text{no MU}}}{\sigma_{\text{MET}_{\text{no MU}}} > 4$
 - $\text{Min}\Delta\phi(\text{all jets}, \text{MET}_{\text{no mu}}) > 2.5$
- ▶ This was used as our “signal region”
- ▶ Discrepancy outside signal region is from QCD



Top control region



- Top contribution to V+jets control regions is non-negligible
 - up to 16% in $W \rightarrow \tau \nu$
- Use method used for W backgrounds in prompt analysis
- Region: signal region with lepton veto replaced with requirement for 1 tight muon and 1 tight electron
 - Very few events in $e\mu$ region so also removed $\text{Min}\Delta\phi(\text{all jets}, \text{MET}_{\text{nomu}})$ cut

N_C^{data}	$28 \pm 5.3(\text{stat.})$
N_C^{bkg}	$0.6 \pm 0.2(\text{MC stat.})$
$N_S^{\text{top MC}}$	$9.6 \pm 1.8(\text{MC stat.})$
$N_C^{\text{top MC}}$	$42.6 \pm 5.2(\text{MC stat.})$
N_S^{top}	$6.1 \pm 1.2(\text{stat.}) \pm 1.4(\text{MC stat.})$





$$W \rightarrow e\nu$$

- Data-MC agreement good
- Same method used as for prompt analysis





$$W \rightarrow \mu\nu$$

- ▶ Data-MC agreement good
- ▶ Same method used as for prompt analysis




$$Z \rightarrow \nu\nu$$

- Data-MC agreement good for limited statistics
- Same method used as for prompt analysis



$W \rightarrow \tau \nu$ control region


- For other W +jets backgrounds control region is:
 - signal region with lepton veto replaced with a requirement for a single lepton
- For $W \rightarrow \tau \nu$ there are not enough events in this region: 2 events for N_C^{Data}
 - In prompt analysis we removed the central jet veto (CJV)
- CJV no longer used, so we remove the $\text{Min}\Delta\phi(\text{all jets}, \text{MET}_{\text{nomu}})$ cut
 - This leads to QCD contamination so we require:
 - $\text{Min}\Delta\phi(\text{leading 2 jets}, \text{MET}_{\text{nomu}}) > 1.0$
 - We also add an $m_T > 20$ GeV cut on the lepton-MET system to remove QCD contamination



$W \rightarrow \tau \nu$ control region


- Estimate error from difference between control region and signal region cuts
- $W \rightarrow \mu \nu$ has enough events to see data driven weight variation with $\text{Min}\Delta\phi(\text{all jets}, \text{MET}_{\text{nomu}})$ cut
 - weight changes by 20% when loosening cut from 2.5 to 1.0.
 - We add a 20% systematic on the $W \rightarrow \tau \nu$ background

N_{C}^{data}	$88 \pm 9.4(\text{stat.})$
N_{C}^{bkg}	$15.2 \pm 4.8(\text{MCstat.})$
N_{S}^{MC}	$176.1 \pm 10.5(\text{MCstat.})$
N_{S}^{MC}	$133.9 \pm 8.0(\text{MCstat.})$
N_{S}^{W}	$95.7 \pm 12.3(\text{stat.}) \pm 10.2(\text{MCstat.})$




QCD background estimation: Shape region choice

- ▶ Try modelling QCD shape in preselection region
- ▶ QCD met must come from a jet, use region with low $\text{Min}\Delta\phi(\text{all jets}, \text{MET}_{\text{nomu}})$
- ▶ Try inverted region with:
 - $\text{Min}\Delta\phi(\text{all jets}, \text{MET}_{\text{nomu}}) < 1.0$
 - $\text{Min}\Delta\phi(\text{leading jets}, \text{MET}_{\text{nomu}}) > 1.0$
- ▶ Has good shape agreement with enriched QCD MC
- ▶ Use shape taken from requiring:
 - $\text{Min}\Delta\phi(\text{all jets}, \text{MET}_{\text{nomu}}) < 1.0$
- ▶ And replacing $\text{Min}\Delta\phi(\text{all jets}, \text{MET}_{\text{nomu}})$ with $\text{Min}\Delta\phi(\text{leading jets}, \text{MET}_{\text{nomu}})$




QCD background estimation: Scale factor

- ▶ Cannot use ABCD to normalise
- ▶ Scale factor shows strong dependence on cut variables
- ▶ Norm 2 and 3 have large signal contamination
 - Norm 3 also has low stats and odd because we forbid jets opposite the met in phi
- ▶ Fit scale factor variation in norm 1
- ▶ Check consistency in norm 2 and 3




Scale factor variation

Norm 1



Norm 1+2



QCD background estimation: Result and systematics

Region	Factor	Extrapolation $\text{mindphi} > 2.5$	Extrapolation $\text{metsig} > 4$
Norm 1	0.22 ± 0.03	0.016 ± 0.009	0.04 ± 0.03
Norm 1+2	0.15 ± 0.02	0.010 ± 0.003	0.03 ± 0.02
Norm 1+3	0.41 ± 0.03	0.036 ± 0.062	1.10 ± 0.10
Norm 2	0.08 ± 0.02	-	0.05 ± 0.04
Norm 3	1.22 ± 0.15	0.60 ± 0.25	-

- ▶ Good agreement in all $\text{Min}\Delta\phi(\text{all jets}, \text{MET}_{\text{nomu}})$ extrapolations
- ▶ Norm 3 agreement in metsig is poor
 - As norm 3 has low statistics and is an odd region: drop
- ▶ Use envelope of norm 1 scale factors
- ▶ Final prediction: $N_5^{\text{QCD}} = 14 \pm 10$
- ▶ Expected limit 0.5% better with no QCD, 0.5% worse with double error

Results

Process	Number of events
$Z \rightarrow \nu\nu$	$141.9 \pm 36.4(\text{stat.}) \pm 15.0(\text{MC stat.})$
$W \rightarrow e\nu$	$59.7 \pm 7.7(\text{stat.}) \pm 5.2(\text{MC stat.})$
$W \rightarrow \mu\nu$	$81.2 \pm 5.6(\text{stat.}) \pm 5.8(\text{MC stat.})$
$W \rightarrow \tau\nu$	$95.7 \pm 12.3(\text{stat.}) \pm 10.2(\text{MC stat.})$
QCD	14 ± 10
Top	$6.1 \pm 1.2(\text{stat.}) \pm 1.4(\text{MC stat.})$
VV	$6.0 \pm 0.6(\text{MC stat.})$
Total bkg.	$405 \pm 39.6(\text{stat.}) \pm 19.8(\text{MC stat.})$
VBF signal	$313.5 \pm 9.4(\text{MC stat.})$
ggH signal	$22.5 \pm 6.0(\text{MC stat.})$
Total signal	$336 \pm 11.1(\text{MC stat.})$

Expected limits

- ▶ Used Higgs combine package with Asymptotic CLs method
- ▶ Performed a single bin counting experiment
- ▶ Analysis blind so have expected limits only
- ▶ 95% C.L. Median limit on $B(H \rightarrow \text{inv.})$ for $m_H = 125$ GeV is: **31%**
 - 1σ band is 23-42%
 - 2σ band is 17-57%
- ▶ Prompt analysis expected limit was 49%
- ▶ We intend to run other mass points:
- ▶ 110, 150, 200, 300 and 400 GeV

Uncertainties by decreasing impact


Control region statistics
 $Z \rightarrow \nu\nu$ - $Z/\gamma^* \rightarrow \mu\mu$ extrapolation
nb above to be revisited
JES
 $W \rightarrow \tau\nu$ extrapolation
MC statistics
QCD systematics
lepton ID efficiency
JER
UES
luminosity
PU weighting
theory uncertainties

Uncertainty impact table - impacts larger than 0%

Median expected limit with:		
All Nuisances: 30.8%	No Nuisances: 12.2%	
Nuisance	Removal impact/%	Addition impact/%
CMS_VBFHinv_zvv_stat:	-12.7%	64.6%
CMS_VBFHinv_zvv_extrapfacunc:	-7.6%	47.4%
CMS_scale_j:	-6.3%	19.2%
CMS_VBFHinv_tau_extrapfacunc:	-4.7%	24.9%
CMS_VBFHinv_zvv_norm:	-1.9%	18.4%
CMS_VBFHinv_wtau_stat:	-1.9%	12.0%
CMS_VBFHinv_wtau_norm:	-1.3%	8.8%
CMS_VBFHinv_wel_stat:	-1.3%	4.8%
CMS_VBFHinv_qcd_norm:	-1.3%	4.0%
CMS_VBFHinv_tau_eff:	-0.6%	5.6%
CMS_VBFHinv_wmu_norm:	-0.6%	3.2%
CMS_eff_m:	-0.6%	2.4%
CMS_VBFHinv_wmu_stat:	-0.6%	2.4%
CMS_VBFHinv_wel_norm:	-0.6%	2.4%
CMS_res_j:	-0.6%	0.8%
CMS_scale_met:	0.0%	0.8%

BDT Study

- ▶ Had a quick look at MVA analysis
- ▶ Started from cut based signal region
 - Only region with negligible QCD
- ▶ Best expected limit obtained 30%
 - Does not take into account any increased systematic
 - Therefore unlikely to be worthwhile
- ▶ New variables could make MVA worthwhile
- ▶ Ability to model QCD would enable looser starting selection which may make MVA worthwhile



Summary

Parked analysis

- ▶ New cut based analysis found with negligible QCD and higher signal efficiency
 - Can then accept remaining QCD estimate with large error
 - Made possible by looser parked trigger thresholds
- ▶ MVA investigated
 - Not sufficient benefit without new variables or looser preselection
- ▶ Full cut based analysis presented
 - Expected limit **31%**
 - Improved from 49% for prompt analysis

Run 2 prospects

- ▶ Prescaled looser control triggers planned to enable better QCD control region
- ▶ Will reinvestigate shape analysis and MVA

Backup

VBF enriched QCD MC


MC Filter: Vectorial sum of neutrino E_T

$$\triangleright \sum E_{\perp}(\vec{\nu}) > 40 \text{ GeV}$$

MC Filter: Dijet Filter

- ▶ Select jets with:
 - ▶ $p_{\perp} > 20 \text{ GeV}$
 - ▶ $|\eta| < 5.0$
- ▶ From selected jets at least one pair with:
 - ▶ $m_{jj} > 700 \text{ GeV}$
 - ▶ $\Delta\eta > 3.2$


QCD Inc 80-600 GeV




There 2 distinct populations of events: real and fake met.

Sample	Ev. Gen.	Filter Eff.	Events	XS [pb]	Eq. Lumi. [fb^{-1}]
QCD-Pt-80to120	3937600000	0.000049	1614416	1033680	38.09
QCD-Pt-120to170	7000000000	0.000283	2051000	156293.3	44.79
QCD-Pt-170to300	1375000000	0.000987	1391500	34138.15	40.28
QCD-Pt-300to470	800000000	0.002659	207840	1759.549	45.47
QCD-Pt-470to600	250000000	0.004127	104675	113.8791	219.53



QCD Data-Driven: $j1j3$, $j2j3$ shape





MVA Details





MVA Details



MVA Details



MVA Details



No systematics limits

- ▶ Prompt limit with no systematics 16.6%