


## WORKED BY: mathematics teacher PJETËR NEÇAJ

TIRANA, ALBANIA on July 6, 2025

### TOPIC: A PROBLEM from PLANE GEOMETRY

Given any quadrilateral ABCD. Let there be any point P inside this quadrilateral. We join the point P with the midpoints of the sides of the quadrilateral and thus 4 quadrilaterals are formed. We denote by  $S_1, S_2, S_3, S_4$  the surfaces of these quadrilaterals that were formed, taking them in turn, in a sequence (according to the movement of the clock). The problem is: Can we find a connection between the surfaces  $S_1, S_2, S_3, S_4$ ?



• We are reasoning as follows:


First we are making auxiliary constructions, we join the point P with the vertices ABCD. In the triangle PAB we have the two triangles PAM and PMB where  $AM=MB$  which means that these two triangles have equal areas. We denote  $a=S_{PAM}=S_{PMB}$

In triangle PBC we have two triangles PBN and PNC where  $BN=NC$  means that these two triangles have equal areas.

We denote  $b=S_{PBN}=S_{PNC}$

We denote  $c=S_{PCK}=S_{PKD}$

We denote  $d=S_{PDL}=S_{PLA}$



We evaluate  $S_1+S_3=(a+d)+(b+c)=a+b+c+d$

We evaluate  $S_2+S_4=(c+d)+(a+b)=a+b+c+d$

It is clear that the amounts:  **$S_1+S_3=S_2+S_4$**


**So we have a True Statement, we have proven a THEOREM.**

**Now I am formulating the THEOREM that I PROVEN.**

**THEOREM:** If any point P inside any concave quadrilateral, is joined to the midpoints of the sides of the quadrilateral, 4 quadrilaterals with surfaces by  $S_1, S_2, S_3, S_4$  (marked in a direction of rotation) are formed, then the equation is true:  $S_1 + S_3 = S_2 + S_4$ .

ATTENTION: We are continuing the study of the PROBLEM, now in the general case, for any concave polygon with an even number of sides  $n=6, n=8, n=10, \dots$ . And we reason as follows:

• Let's have any concave hexagon:



Point P anywhere inside the hexagon, and M, N, K, L, Q, H are the midpoints of the sides.

We connect the point P with the midpoints of the sides and 6 quadrilaterals will be formed as in the figure. Auxiliary construction, we connect the point P with all the vertices of the hexagon ABCDEF.

In triangle PAB we have  $AM = MB$ , which means:

Surface  $S_{PAM} = S_{PMB} = a$  square unit.

We continue the reasoning as above and in triangle PFA we have  $FH = HA$ , leading to:

$S_{PFH} = S_{PHA} = f$  square unit.

We denote:  $S_1 = S_{PMAH}$ ,  $S_2 = S_{PHFQ}$ ,  $S_3 = S_{PQEL}$ ,  $S_4 = S_{PLDK}$ ,  $S_5 = S_{PKCN}$ ,  $S_6 = S_{PNBM}$

We evaluate:  $S_1 + S_3 + S_5 = (a+f) + (e+d) + (c+b) = a+b+c+d+e+f$

We also have  $S_2 + S_4 + S_6 = (f+e) + (d+c) + (b+a) = a+b+c+d+e+f$

• It is clear that the sums:  $S_1 + S_3 + S_5 = S_2 + S_4 + S_6$

It is clear that the Theorem we proved above for  $n=4$  and  $n=6$  is true when the number of sides  $n$  is even for  $n=4, 6, 8, 10, 12, \dots$

So it is true that:  $S_1 + S_3 + S_5 + S_7 + \dots + S_{n-1} = S_2 + S_4 + S_6 + S_8 + \dots + S_n$

• I believe that I DISCOVERED a truth in PLANE GEOMETRY, that is, I PROVEN a THEOREM, and I wish and believe that I have the right to give it a NAME, I am naming it:

**THEOREM PJETËR NEÇAJ**

Tirana, Albania on JULY 6, 2025