

The emergence of geometric order in proliferating metazoan epithelia.

by Gibson *et al.* *Nature*, 2006

Computational Biology and Bioinformatics Seminar

Philip Hartout

May 28, 2020

ETH Zurich

Introduction


'The organisation of cells into epithelial sheets is an essential feature of animal design.'

Gibson et al [?]

Predictable geometries arise when *minimising surface energy* or *maximising space filling* in biological and non-biological structures:

Examples


- *Drosophila Melanogaster*'s retinal cells
- Honeycombs
- Coins on tabletops



Predictable geometries arise when *minimising surface energy* or *maximising space filling* in biological and non-biological structures:

Examples

- *Drosophila Melanogaster*'s retinal cells
- Honeycombs
- Coins on tabletops



How is the geometry of *growing* epithelial cells organized and how can it be explained?

Emergent behaviour

→ requires understanding emergent behaviours

Emergent behaviour

→ requires understanding emergent behaviours

Emergent behaviour

It is the behaviour of a system that can *only* be explained by examination of a system's **parts** *and* their **relationships**.

Model Assumptions

Post-mitotic relationship between two daughter cells

Stochastically marked cell lineages with GFP revealed stable post-mitotic patterns.




Figure 1: Daughter-cell relationship.




Figure 2: Observed post-mitotic relationship between daughter cells.

Post-mitotic relationship between two daughter cells

Stochastically marked cell lineages with GFP revealed stable post-mitotic patterns.




Figure 1: Daughter-cell relationship.

→ required to model relationships between cells





Figure 2: Observed post-mitotic relationship between daughter cells.

Typical topology changes of a dividing cell

Figure 3: Topology changes during cell division



A Discrete Markov Model for Proliferating Epithelia

A Markov chain is a discrete stochastic process with the Markov property:

$$P(X_t|X_{t-1}, \dots, X_1) = P(X_t|X_{t-1})$$

where X_t is the time-dependent r.v. describing the state of the system at time t .

It is determined by a probability transition matrix P and an initial probability distribution.

Applying a discrete Markov model to growing epithelia

s state of the cell as its number of sides, where $s > 3$.

Applying a discrete Markov model to growing epithelia

s state of the cell as its number of sides, where $s > 3$.
 p_s the relative frequency of s -sided cells in the population

Applying a discrete Markov model to growing epithelia

s	state of the cell as its number of sides, where $s > 3$.
p_s	the relative frequency of s -sided cells in the population
$\mathbf{p}^{(t)}$	the infinite row vector $\mathbf{p}^{(t)} = [p_4, p_5, p_6, \dots]$ state of the population at generation t .

Applying a discrete Markov model to growing epithelia

s	state of the cell as its number of sides, where $s > 3$.
p_s	the relative frequency of s -sided cells in the population
$\mathbf{p}^{(t)}$	the infinite row vector $\mathbf{p}^{(t)} = [p_4, p_5, p_6, \dots]$ state of the population at generation t .
$\mathbf{p}^{(t+1)} = \mathbf{p}^{(t)}PS$	the state dynamics, where P and S are the probabilistic transition matrices.

Applying a discrete Markov model to growing epithelia

s	state of the cell as its number of sides, where $s > 3$.
p_s	the relative frequency of s -sided cells in the population
$\mathbf{p}^{(t)}$	the infinite row vector $\mathbf{p}^{(t)} = [p_4, p_5, p_6, \dots]$ state of the population at generation t .
$\mathbf{p}^{(t+1)} = \mathbf{p}^{(t)}PS$	the state dynamics, where P and S are the probabilistic transition matrices.
P_{ij}	the probability that an i sided cell divides to produce a j -sided daughter cell.

Applying a discrete Markov model to growing epithelia

s	state of the cell as its number of sides, where $s > 3$.
p_s	the relative frequency of s -sided cells in the population
$\mathbf{p}^{(t)}$	the infinite row vector $\mathbf{p}^{(t)} = [p_4, p_5, p_6, \dots]$ state of the population at generation t .
$\mathbf{p}^{(t+1)} = \mathbf{p}^{(t)}PS$	the state dynamics, where P and S are the probabilistic transition matrices.
P_{ij}	the probability that an i sided cell divides to produce a j -sided daughter cell.
S_{ij}	the probability of an i -sided cell will gain sides from dividing neighbour cells divisions to become j -sided.

Transition matrices

$$P = \begin{matrix} & \begin{matrix} 4 & 5 & 6 & 7 & 8 & 9 & \dots \end{matrix} \\ \begin{matrix} 4 \\ 5 \\ 6 \\ 7 \\ 8 \\ 9 \\ \vdots \end{matrix} & \begin{pmatrix} 1 & & & & & & \\ 1 & 1 & & & & & \\ 1 & 2 & 1 & & & & \\ 1 & 3 & 3 & 1 & & & \\ 1 & 4 & 6 & 4 & 1 & & \\ 1 & 5 & 10 & 10 & 5 & 1 & \\ & & & & & & \ddots \end{pmatrix} \end{matrix}$$

P_{ij} : the probability that an i sided cell divides to produce a j -sided daughter cell.

$$S = \begin{matrix} & \begin{matrix} 4 & 5 & 6 & 7 & 8 & 9 & \dots \end{matrix} \\ \begin{matrix} 4 \\ 5 \\ 6 \\ 7 \\ 8 \\ 9 \\ \vdots \end{matrix} & \begin{pmatrix} 0 & 1 & & & & & \\ & 0 & 1 & & & & \\ & & 0 & 1 & & & \\ & & & 0 & 1 & & \\ & & & & 0 & 1 & \\ & & & & & 0 & 1 \\ & & & & & & \ddots \end{pmatrix} \end{matrix}$$

S_{ij} the probability of an i -sided cell will gain sides from dividing neighbour cells divisions

Predicting the equilibrium E from $T = PS$

The Perron-Frobenius theorem guarantees that a Markov chain will converge to a unique equilibrium E which is the principal eigenvector of the transition matrix $T = PS$.




Figure 4: Distribution of the predicted and observed polygons (yellow: predicted equilibrium)

Robust equilibrium topology




Figure 5: Discrete state dynamics




Figure 6: Figure showing the same attained equilibrium, no matter the initial conditions

Additional insights on mitotic cells

As seen in S , a mitotic cell gains on average one side, and daughter cells have one less side. Experimental evidence confirms this:

Non-mitotic cells	(5.94 ± 0.06)
Mitotic cells	(6.99 ± 0.07)

→ cells accumulate sides until division.




Figure 7: Polygon distributions of mitotic cells vs. non-mitotic cells

Polygon distribution of mitotic cells in non-proliferating tissues

- The model does not apply to non-proliferating tissue
- Forcing mitosis using *string* (*stg*) in tissues does not replicate the proliferating tissue distribution.




Figure 8: Distribution of polygons in non-proliferating tissue

→ Differential proliferation influences cell shape and morphogenesis.

Subsequent works

Subsequent work

→ additional work done to factor in biophysical aspects [?].

Subsequent work

→ additional work done to factor in biophysical aspects [?].

Work on 3D structures




Figure 9: The scutoid has been discovered as a solution to 3D cell packing.[?]

Work on dynamics




Figure 10: Modelling and prediction of speed and direction of proliferating epithelia. [?]

Conclusions

The proposed **discrete Markov chain**:

1. shows how epithelial topology can be irregular, but **not random**.

The proposed **discrete Markov chain**:

1. shows how epithelial topology can be irregular, but **not random**.
2. **explains the predominantly hexagonal topology** of growing epithelia as well as the distribution of other polygons.

The proposed **discrete Markov chain**:

1. shows how epithelial topology can be irregular, but **not random**.
2. **explains the predominantly hexagonal topology** of growing epithelia as well as the distribution of other polygons.
3. shows an **emergent mechanism** by which epithelia accomodate **rapid proliferation** while maintaining **structural integrity**.

Questions?



https://github.com/pjhartout/CBB_Seminar

Geometry of *growing* epithelial cells

Time-lapse movies collected revealed the following findings:

- Initially polygonal prophase cells rounded up and divided
- Cell-neighbour relationships were stably maintained throughout the cell cycle
- The topology of the divided cells is invariant over time.




Figure 11: General hexagon-dominated topology of growing epithelial in *Drosophila Melanogaster*.

Derivation of Markov state dynamics - transition matrix P

- P_{ij} : the probability that an i sided cell divides to produce a j -sided daughter cell.
- K_t : the number of junctions distributed to one daughter cell on division at generation t . $s_{t-1} - K_t$ are left for the other.

Derivation of Markov state dynamics - transition matrix P

- P_{ij} : the probability that an i sided cell divides to produce a j -sided daughter cell.
- K_t : the number of junctions distributed to one daughter cell on division at generation t . $s_{t-1} - K_t$ are left for the other.
- Because no triangular cells are observed, each daughter receives at least two junctions, leaving $s_t - 4$ junctions to be distributed among the daughters with equal probability.
- So, the number of junctions received by the first daughter is $K_t - 2 \sim \mathcal{B}(n = s_{t-1}, p = \frac{1}{2})$.

Derivation of Markov state dynamics - transition matrix P

- P_{ij} : the probability that an i sided cell divides to produce a j -sided daughter cell.
- K_t : the number of junctions distributed to one daughter cell on division at generation t . $s_{t-1} - K_t$ are left for the other.
- Because no triangular cells are observed, each daughter receives at least two junctions, leaving $s_t - 4$ junctions to be distributed among the daughters with equal probability.
- So, the number of junctions received by the first daughter is $K_t - 2 \sim \mathcal{B}(n = s_{t-1}, p = \frac{1}{2})$.

- As a result

$$P_{ij} = P(K_t + 2 = j | s_{t-1} = i) = \binom{i-4}{j-4} \cdot \frac{1}{2^{i-4}}$$

Derivation of Markov state dynamics - transition matrix P

- P_{ij} : the probability that an i sided cell divides to produce a j -sided daughter cell.
- K_t : the number of junctions distributed to one daughter cell on division at generation t . $s_{t-1} - K_t$ are left for the other.
- Because no triangular cells are observed, each daughter receives at least two junctions, leaving $s_t - 4$ junctions to be distributed among the daughters with equal probability.
- So, the number of junctions received by the first daughter is $K_t - 2 \sim \mathcal{B}(n = s_{t-1}, p = \frac{1}{2})$.

- As a result

$$P_{ij} = P(K_t + 2 = j | s_{t-1} = i) = \binom{i-4}{j-4} \cdot \frac{1}{2^{i-4}}$$

- Therefore, the unnormalized entries of P are the coefficients of Pascal's triangle:

$$P = \begin{matrix} & 4 & 5 & 6 & 7 & 8 & 9 & \dots \\ \begin{matrix} 4 \\ 5 \\ 6 \\ 7 \\ 8 \\ 9 \\ \vdots \end{matrix} & \left(\begin{array}{cccccc} 1 & & & & & \\ 1 & 1 & & & & \\ 1 & 2 & 1 & & & \\ 1 & 3 & 3 & 1 & & \\ 1 & 4 & 6 & 4 & 1 & \\ 1 & 5 & 10 & 10 & 5 & 1 \\ & & & & & \ddots \end{array} \right) \end{matrix}$$

Derivation of Markov state dynamics - transition matrix S

- S_{ij} : the probability of an i -sided cell will gain sides from dividing neighbour cells divisions

Derivation of Markov state dynamics - transition matrix S

- S_{ij} : the probability of an i -sided cell will gain sides from dividing neighbour cells divisions
- On mitosis, a cell adds one side to each of two neighbouring cells.

Derivation of Markov state dynamics - transition matrix S

- S_{ij} : the probability of an i -sided cell will gain sides from dividing neighbour cells divisions
- On mitosis, a cell adds one side to each of two neighbouring cells.
- Assuming N cells in an epithelium, the number of cells after one round of divisions is $2N$

Derivation of Markov state dynamics - transition matrix S

- S_{ij} : the probability of an i -sided cell will gain sides from dividing neighbour cells divisions
- On mitosis, a cell adds one side to each of two neighbouring cells.
- Assuming N cells in an epithelium, the number of cells after one round of divisions is $2N$
- On average the number of sides gained per cell is $\frac{2N}{2N} = 1$. That is, a cell will gain 1 side every cycle on average.

Derivation of Markov state dynamics - transition matrix S

- S_{ij} : the probability of an i -sided cell will gain sides from dividing neighbour cells divisions
- On mitosis, a cell adds one side to each of two neighbouring cells.
- Assuming N cells in an epithelium, the number of cells after one round of divisions is $2N$
- On average the number of sides gained per cell is $\frac{2N}{2N} = 1$. That is, a cell will gain 1 side every cycle on average.
- Thus, $S_{ij} = 1$ if $j = i + 1$ and 0 otherwise.

$$S = \begin{matrix} & \begin{matrix} 4 & 5 & 6 & 7 & 8 & 9 & \dots \end{matrix} \\ \begin{matrix} 4 \\ 5 \\ 6 \\ 7 \\ 8 \\ 9 \\ \vdots \end{matrix} & \begin{pmatrix} 0 & 1 & & & & & \\ & 0 & 1 & & & & \\ & & 0 & 1 & & & \\ & & & 0 & 1 & & \\ & & & & 0 & 1 & \\ & & & & & 0 & 1 \\ & & & & & & 0 & 1 \\ & & & & & & & \ddots \end{pmatrix} \end{matrix}$$

Backup slides - Assumptions of the recurrence system

From the aforementioned observations we make the following assumptions about the mathematical modelling of cell epithelia:

- Cells are polygons with a minimum of four sides
- Cells do not resort
- Mitotic siblings retain a common common junctional interface
- Cells have asynchronous but uniform cell cycle times
- Cleavage places cut a side rather than a vertex of the mother polygon
- Mitotic cleavage orientation randomly distributes existing tricellular junctions to both daughter cells.

Recurrence system

Table 1: Cell features, topological equivalence and evolution at division t

Cell feature	Graph equivalent	Evolution at division t
Tricellular junction	Vertex, v_t	$v_t = v_{t-1} + 2f_t$
Cell side	Edge, e_t	$e_t = e_{t-1} + 3f_{t-1}$
Apical cell surface	Face, f_t	$e_t = e_{t-1} + 3f_{t-1}$

So we can construct the following system:

$$s_t = \frac{2(e_t + 3f_{t-1})}{2f_{t-1}} = \frac{s_{t-1}}{2} + 3$$

which exponentially converges to 6 provided the initial condition:

$$s_t = 6 + 2^{-t}(s_0 - 6)$$

