

Computational Biology

Lecturers:


Tanja Stadler, Carsten Magnus & Tim Vaughan

Teaching Assistants:

Jūlija Pečerska, Jérémie Sciré,
Sarah Nadeau & Marc Manceau

Computational Evolution
Department of Biosystems Science and Engineering

HS 2019



The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

The easiest way to study something is by observation.

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

The easiest way to study something is by observation.

► Wetlab

- Very realistic;
- Time-consuming and expensive;
- Impossible (sometimes).

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

The easiest way to study something is by observation.

► Wetlab

- Very realistic;
- Time-consuming and expensive;
- Impossible (sometimes).

► Simulation

- A virtual experiment in which we mimic a (biological) process on a computer to study its properties
- Not necessarily realistic
- Allows us to:
 - * generate data with given assumptions;
 - * test predictive properties of models.

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

The easiest way to study something is by observation.

► Wetlab

- Very realistic;
- Time-consuming and expensive;
- Impossible (sometimes).

► Simulation

- A virtual experiment in which we mimic a (biological) process on a computer to study its properties
- Not necessarily realistic
- Allows us to:
 - * generate data with given assumptions;
 - * test predictive properties of models.

Today we will simulate evolution!

The tree of great apes

CB




Figure adapted from [?]

The Simulation Game

Studying evolution


Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm



- Format for tree representation
- To record a tree in Newick format:
 - Assign a label to each tip
 - Choose two tips that are a cherry (e.g. C and D)
 - Replace selected tips with a new tip of the form (tip1:branch1,tip2:branch2) (e.g. (C : 1, D : 1))
 - Branch length to the new tip is the branch length to the cherry
 - Repeat until the full tree is rewritten
- What is the Newick format for the rooted tree above?

The Simulation Game

Studying evolution


Simulating evolution

Initializing the starting sequence

Simulating the substitutions

Pen and paper exercise

Algorithm



- Format for tree representation
- To record a tree in Newick format:
 - Assign a label to each tip
 - Choose two tips that are a cherry (e.g. C and D)
 - Replace selected tips with a new tip of the form (tip1:branch1,tip2:branch2) (e.g. (C : 1, D : 1))
 - Branch length to the new tip is the branch length to the cherry
 - Repeat until the full tree is rewritten
- What is the Newick format for the rooted tree above?
(((C : 1, D : 1) : 1, A : 2) : 1, B : 3);

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting sequence

Simulating the substitutions

Pen and paper exercise

Algorithm

Steps:

1. Initialization of the starting sequence:

- ▶ Sample a starting nucleotide for each position in the sequence

2. Iterative simulation of sequence evolution, along all branches of the tree

- ▶ Compute the transition probability matrix $P(t_b)$.
- ▶ Sample a new nucleotide for each position in the sequence.

Step 1: Initialization of the starting sequence

CB

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

1a. Sample a starting nucleotide n

Step 1: Initialization of the starting sequence

CB

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

1a. Sample a starting nucleotide n

From the vector of equilibrium frequencies of nucleotides

	T	C	A	G
Π	0.22	0.26	0.33	0.19

Step 1: Initialization of the starting sequence

CB

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

1a. Sample a starting nucleotide n

From the vector of equilibrium frequencies of nucleotides


	T	C	A	G
Π	0.22	0.26	0.33	0.19

Knowing Π , how do we sample a nucleotide?

Inverse transform method

CB

- Sample u from $U(0, 1)$;



The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions


Pen and paper exercise

Algorithm

Inverse transform method

CB

- ▶ Sample u from $U(0, 1)$;
- ▶ Transform u into a sample from the desired distribution using the **CDF** == **C**umulative **D**istribution **F**unction $F_X(x) = P(X \leq x)$.



The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting sequence

Simulating the substitutions


Pen and paper exercise

Algorithm

Inverse transform method

CB

- ▶ Sample u from $U(0, 1)$;
- ▶ Transform u into a sample from the desired distribution using the **CDF** == **C**umulative **D**istribution **F**unction $F_X(x) = P(X \leq x)$.



The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting sequence


Simulating the substitutions

Pen and paper exercise

Algorithm

Inverse transform method

- ▶ Sample u from $U(0, 1)$;
- ▶ Transform u into a sample from the desired distribution using the **CDF** == **C**umulative **D**istribution **F**unction
 $F_X(x) = P(X \leq x)$.



The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting sequence


Simulating the substitutions

Pen and paper exercise

Algorithm

Inverse transform method

- ▶ Sample u from $U(0, 1)$;
- ▶ Transform u into a sample from the desired distribution using the **CDF** == **C**umulative **D**istribution **F**unction
 $F_X(x) = P(X \leq x)$.



The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting sequence

Simulating the substitutions


Pen and paper exercise

Algorithm

Sampling discrete random variables

CB

	T	C	A	G
Π	0.22	0.26	0.33	0.19
CDF	0.22	0.48	0.81	1.00



The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting sequence

Simulating the substitutions


Pen and paper exercise

Algorithm

Sampling discrete random variables

CB

	T	C	A	G
Π	0.22	0.26	0.33	0.19
CDF	0.22	0.48	0.81	1.00



Sample u from $U(0, 1)$.

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequenceSimulating the
substitutions


Pen and paper exercise

Algorithm

Sampling discrete random variables

CB

	T	C	A	G
Π	0.22	0.26	0.33	0.19
CDF	0.22	0.48	0.81	1.00



Sample u from $U(0, 1)$.

E.g. $u = 0.62$.

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequenceSimulating the
substitutions


Pen and paper exercise

Algorithm

Sampling discrete random variables

CB

	T	C	A	G
Π	0.22	0.26	0.33	0.19
CDF	0.22	0.48	0.81	1.00



Sample u from $U(0, 1)$.

E.g. $u = 0.62$.

Select nucleotide **A**.

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequenceSimulating the
substitutions


Pen and paper exercise

Algorithm

Sampling discrete random variables

CB

	T	C	A	G
Π	0.22	0.26	0.33	0.19
CDF	0.22	0.48	0.81	1.00



Sample u from $U(0, 1)$.

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequenceSimulating the
substitutions


Pen and paper exercise

Algorithm

Sampling discrete random variables

CB

	T	C	A	G
Π	0.22	0.26	0.33	0.19
CDF	0.22	0.48	0.81	1.00



Sample u from $U(0, 1)$.

E.g. $u = 0.18$.

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequenceSimulating the
substitutions

Pen and paper exercise

Algorithm

Sampling discrete random variables

CB

The Simulation Game

Studying evolution


Simulating evolution

Initializing the starting
sequenceSimulating the
substitutions

Pen and paper exercise

Algorithm

	T	C	A	G
Π	0.22	0.26	0.33	0.19
CDF	0.22	0.48	0.81	1.00



Sample u from $U(0, 1)$.


E.g. $u = 0.18$.

Select nucleotide **T**.

Step 1: Initializing the starting sequence

CB

1b. Place n on the root node;



The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions


Pen and paper exercise

Algorithm

Step 1: Initializing the starting sequence

CB

1b. Place n on the root node;



The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions


Pen and paper exercise

Algorithm

Step 2a: Choose the next branch for simulation

CB

Get a branch b with a nucleotide at the start;
 $t_b = \text{length}(b)$;
 n = nucleotide at start of branch b ;



The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions


Pen and paper exercise

Algorithm

Step 2a: Choose the next branch for simulation

CB

Get a branch b with a nucleotide at the start;
 $t_b = \text{length}(b)$;
 n = nucleotide at start of branch b ;



The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

Step 2b-d: Sample the new nucleotide

CB

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

$$P(t_b) = e^{Qt_b};$$

Sample new nucleotide n_{new} from row n in $P(t_b)$;

Place n_{new} at the end of branch b ;

Step 2b-d: Sample the new nucleotide

CB

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

$$P(t_b) = e^{Qt_b};$$

Sample new nucleotide n_{new} from row n in $P(t_b)$;

Place n_{new} at the end of branch b ;

To sample new nucleotide n_{new} we will need the substitution rate matrix Q , and transition probability matrix P .

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequenceSimulating the
substitutions

Pen and paper exercise

Algorithm

$\Pi = (\pi_T, \pi_C, \pi_A, \pi_G)$ - equilibrium frequencies.

α_1, α_2 - transition ratios.

β - transversion ratios.

$$Q_{\text{TN93}} = \begin{array}{c} \begin{array}{cc} & \begin{array}{cccc} & \text{T} & \text{C} & \text{A} & \text{G} \end{array} \\ \begin{array}{c} \text{T} \\ \text{C} \\ \text{A} \\ \text{G} \end{array} & \left(\begin{array}{cccc} \cdot & \alpha_1 \pi_C & \beta \pi_A & \beta \pi_G \\ \alpha_1 \pi_T & \cdot & \beta \pi_A & \beta \pi_G \\ \beta \pi_T & \beta \pi_C & \cdot & \alpha_2 \pi_G \\ \beta \pi_T & \beta \pi_C & \alpha_2 \pi_A & \cdot \end{array} \right) \end{array} \end{array}$$

The diagonals are set such that each row sums up to zero, e.g.

$$q_{\text{TT}} = -(\alpha_1 \pi_C + \beta \pi_A + \beta \pi_G).$$

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

$$\Pi = (0.22, 0.26, 0.33, 0.19)$$

$$\alpha_1 = 44.229, \alpha_2 = 21.781$$

$$\beta = 1$$

$$Q_{\text{TN93}} = \begin{matrix} & \begin{matrix} \text{T} & \text{C} & \text{A} & \text{G} \end{matrix} \\ \begin{matrix} \text{T} \\ \text{C} \\ \text{A} \\ \text{G} \end{matrix} & \begin{pmatrix} -0.01957 & 0.01873 & 0.00054 & 0.00031 \\ 0.01584 & -0.01669 & 0.00054 & 0.00031 \\ 0.00036 & 0.00042 & -0.00752 & 0.00674 \\ 0.00036 & 0.00042 & 0.01170 & -0.01249 \end{pmatrix} \end{matrix}$$

Note: the matrix is scaled to 0.0135 substitutions per mya so that we get reasonable sequences.

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

$\Pi = (\pi_T, \pi_C, \pi_A, \pi_G)$ - equilibrium frequencies.

α_1, α_2 - transition ratios.

β - transversion ratios.

t_b - branch length

$$P(t_b) = e^{t_b Q_{TN93}(\alpha_1, \alpha_2, \beta, \Pi)}$$

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

$$\Pi = (0.22, 0.26, 0.33, 0.19)$$

$$\alpha_1 = 44.229, \alpha_2 = 21.781$$

$$\beta = 1$$

$$t_b = 13 \text{ mya}$$

$$P_{\text{TN93}}(13\text{mya}) = \begin{matrix} & \begin{matrix} \text{T} & \text{C} & \text{A} & \text{G} \end{matrix} \\ \begin{matrix} \text{T} \\ \text{C} \\ \text{A} \\ \text{G} \end{matrix} & \begin{pmatrix} 0.795 & 0.194 & 0.007 & 0.004 \\ 0.164 & 0.824 & 0.007 & 0.004 \\ 0.005 & 0.005 & 0.913 & 0.077 \\ 0.005 & 0.005 & 0.134 & 0.856 \end{pmatrix} \end{matrix}$$

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

We start with nucleotide **T**, so we are interested in row T:

$$P_{\text{TN}_{93}}(13\text{mya}) = \begin{matrix} & \text{T} & \text{C} & \text{A} & \text{G} \\ \begin{matrix} \text{T} \\ \text{C} \\ \text{A} \\ \text{G} \end{matrix} & \begin{pmatrix} 0.795 & 0.194 & 0.007 & 0.004 \\ \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot \end{pmatrix} \end{matrix}$$

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

We start with nucleotide **T**, so we are interested in row T:

$$P_{\text{TN}_{93}}(13\text{mya}) = \begin{matrix} & \begin{matrix} \text{T} & \text{C} & \text{A} & \text{G} \end{matrix} \\ \begin{matrix} \text{T} \\ \text{C} \\ \text{A} \\ \text{G} \end{matrix} & \begin{pmatrix} 0.795 & 0.194 & 0.007 & 0.004 \\ . & . & . & . \\ . & . & . & . \\ . & . & . & . \end{pmatrix} \end{matrix}$$

Sample new nucleotide n_{new} with the weights

$[p_{\text{T} \rightarrow \text{T}}, p_{\text{T} \rightarrow \text{C}}, p_{\text{T} \rightarrow \text{A}}, p_{\text{T} \rightarrow \text{G}}]$

Sample u from $U(0, 1)$.

E.g. $u = 0.81$.

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise


Algorithm

Getting the substitution

CB

Sample u from $U(0, 1)$.

E.g. $u = 0.81$.



Selected substitution is $T \rightarrow C$.

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm


Step 2b-d: Sample the new nucleotide

CB

$$P(t_b) = e^{Qt_b};$$

Sample new nucleotide n_{new} from row n in $P(t_b)$;

Place n_{new} at the end of branch b ;



The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm


Step 2b-d: Sample the new nucleotide

CB

$$P(t_b) = e^{Qt_b};$$

Sample new nucleotide n_{new} from row n in $P(t_b)$;

Place n_{new} at the end of branch b ;



The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

Repeat step 2

CB

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

while *not all branches are used* **do**

 Get a branch **b** with a nucleotide at the start;

$t_b = \text{length}(\mathbf{b})$;

\mathbf{n} = nucleotide at start of branch **b**;

$P(t_b) = e^{Q t_b}$;

 Sample new nucleotide \mathbf{n}_{new} from row \mathbf{n} in $P(t_b)$;

 Place \mathbf{n}_{new} at the start of the daughter branches of **b**;

end

The Simulation Game

Studying evolution


Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm



The Simulation Game

Studying evolution


Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm



The Simulation Game

Studying evolution


Simulating evolution


Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm





The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting sequence

Simulating the substitutions

Pen and paper exercise

Algorithm

The Simulation Game

Studying evolution


Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm



The Simulation Game

Studying evolution


Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm



The Simulation Game

Studying evolution


Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm



The Simulation Game

Studying evolution


Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm



The Simulation Game

Studying evolution


Simulating evolution

Initializing the starting sequence

Simulating the substitutions

Pen and paper exercise

Algorithm



The Simulation Game

Studying evolution


Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm



The Simulation Game

Studying evolution


Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm



Studying evolution

Simulating evolution

Initializing the starting sequence

Simulating the substitutions

Pen and paper exercise

Algorithm



The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

1. Split into pairs;
2. Get the materials;
3. Evolve a character along the tree;

All of the characters together will produce an alignment.

Using dice to generate random numbers

CB

We will be using 10-sided dice for our random number generation.

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm

We will be using 10-sided dice for our random number generation.

Samples from $\text{Unif}(0,1)$ with 2 decimal point precision:

1. Take 2 dice of different colours;
2. Assign a fixed decimal position to each of the dice (and keep it for the whole simulation);
 - ▶ E.g. red is 1st position, blue is 2nd;
3. Roll the dice to get 2 numbers;
 - ▶ E.g. red = 5, blue = 8;
4. Combine the numbers to get a sample;
 - ▶ E.g. 0.58.


```
N = number of sites in the alignment;  
Q = substitution rate matrix;  
for i = 1 to N do  
  | Sample a nucleotide  $n$  from the initial distribution;  
  | Add  $n$  to the sequence of the root node;  
end  
while not all branches are visited do  
  | Get a branch  $b$  with a sequence at the start;  
  |  $t_b = \text{length}(b)$ ;  
  |  $P(t_b) = e^{Qt_b}$ ;  
  | for i = 1 to N do  
    |  $n =$  nucleotide at position i at the start of branch  $b$ ;  
    | Sample new nucleotide  $n_{\text{new}}$  from row  $n$  in  $P(t_b)$ ;  
    | Place  $n_{\text{new}}$  at the end of sequences in the daughter  
    |   branches of  $b$ ;  
  | end  
end
```

The Simulation Game

Studying evolution

Simulating evolution

Initializing the starting
sequence

Simulating the
substitutions

Pen and paper exercise

Algorithm