Computational Biology

Lecturers:

Tanja Stadler, Carsten Magnus & Tim Vaughan

Teaching Assistants: Jūlija Pečerska, Jérémie Sciré, Sarah Nadeau & Marc Manceau

Computational Evolution
Department of Biosystems Science and Engineering

HS 2019



Independent Contrast Method

A 4 - 17 - 17 -

Intuition

Notation

Illustration

Pen and Paper exercise


Solution

Modelling traits on a tree

Each of j traits:

- starts at a single root value,
- evolves independently as a Brownian Motion on distinct branches.
- finishes evolving upon reaching the tips.

The traits are correlated because they evolved along the same tree.



Independent Contrast Method

Motivation

Intuition

Illustration

Pen and Paper exercise

Exercis

How can we do regression after accounting for this tree-driven correlation ?

Intuition


We look for alternative quantities (*contrasts*) that would be:

1. independent:

- independent Brownian trajectories in subtrees.
- use the tree structure to find independent variables.

2. identically distributed:

- traits are normally distributed with variance that scales with branch lengths,
- use branch lengths to define contrasts with equal variance.



Independent Contrast Method

Motivation Intuition

Notation


Illustration

Pen and Paper exercise

Exercis Solutio

Notation

- $\triangleright X_i^j$: value of trait j at node i
- ▶ t_i: length of the branch leading to node i
- ▶ t_i: corrected branch length leading to node i
- Z_k^j: value of contrast corresponding to trait j on node k.



Independent Contrast Method

//otivation

Intuition Notation

Algorithm

Pen and Paper exercise

Exercise


Algorithm - Overview

If k is a tip:

 X_k is the observed value $t_{
u}'=t_k$

Else if k is the parent of i, l:

$$\begin{split} t_k' &= \frac{t_i' \cdot t_l'}{t_i' + t_l'} + t_k \\ X_k^j &= X_i^j \frac{t_l'}{t_i' + t_l'} + X_l^j \frac{t_l'}{t_i' + t_l'} \\ Z_k^j &= \frac{X_i^j - X_l^j}{\sqrt{t_i' + t_l'}} \end{split}$$



Always visit the parent node k after the two child nodes i, l.

Independent Contrast Method

Motivation

Intuition

Algorithm

Pen and Paper exercise

Solutio

Algorithm - Pseudo-code

else

Compute t_i', X_l^j, Z_l^j where i is the first child of k; Compute t_l', X_l^j, Z_l^j where l is the second child of k; $t_k' \leftarrow \frac{t_l't_l'}{t_l'+t_l'} + t_k;$ $X_k^j \leftarrow X_l^j \frac{t_l'}{t_l'+t_l'} + X_l^j \frac{t_l'}{t_l'+t_l'};$ $Z_k^j \leftarrow \frac{X_l^j-X_l^j}{\sqrt{t_l'+t_l'}};$

end

Independent Contrast Method


Motivatio

Intuition

Algorithm

Pen and Paper

Exercise



Independent Contrast Method

Motivation

Intuition

Notation Algorithm

Illustration

Pen and Paper exercise

Exercise Solution

Pen and Paper exercise

Calculate the normalised independent contrasts for both traits on this tree.

► Corrected branch lengths


$$t_k' = \frac{t_i' \cdot t_l'}{t_i' + t_l'} + t_k$$

► Trait values of trait j

$$X_k^j = X_i^j \frac{t_l'}{t_l'+t_l'} + X_l^j \frac{t_l'}{t_l'+t_l'} \label{eq:Xk}$$

Normalised contrasts of trait j

$$Z_k^j = \frac{X_i^j - X_l^j}{\sqrt{t_i' + t_l'}}$$



Independent Contrast

Motivation

Intuition

Algorithm

Pen and Paper exercise

Exercise

Pen and Paper exercise - solution


Calculate the normalised independent contrasts for both traits on this tree.

Trait 1:

$$Z_6^1 = 0.1933545$$
 $Z_7^1 = -0.7244694$
 $Z_8^1 = 1.5428162$
 $Z_9^1 = -0.9045340$

Trait 2:

$$Z_6^2 = -0.4540037$$
 $Z_7^2 = -1.0717747$
 $Z_8^2 = 2.1362070$
 $Z_9^2 = -1.8090681$



ndependent Contrast Method

Motivation

Algorithm

Pen and Paper

Exercise Solution