ILO-PTB CXR form - study

Who is the reviewer?

- ◯ Prof Rees
- ◯ Dr Iddy
- ◯ Dr Patrick
- ◯ Other

Study ID of participant

_____

What is the participant's age?

_____

What is the technical quality of the Xray?

- ◯ Good - free of technical imperfections or artifacts
- ◯ Acceptable - with no technical defects or artifacts likely to impair classification of the radiograph for pneumoconiosis
- ◯ Acceptable - with technical defects or artifacts but still adequate for classification purposes
- ◯ Unreadable - unreadable or unacceptable for classification purposes

Reason for reduced image quality

- ☐ Overexposed (dark)
- ☐ Underexposed (light
- ☐ Artifacts
- ☐ Improper position
- ☐ Poor contrast
- ☐ Poor processing
- ☐ Underinflation
- ☐ Mottle
- ☐ Excessive Edge Enhancement

Is the Xray normal?

- ◯ Yes
- ◯ No

Are there parenchymal opacities?

- ◯ Yes
- ◯ No

What is the shape of the opacities? (first choice)

- p - rounded diameter up to ~1.5 mm
- q - rounded diameter 1.5 to ~3 mm
- r - rounded diameter 3 to ~10 mm
- s - irregular width up to ~1.5 mm
- t - irregular width 1.5 to ~3 mm
- u - irregular width 3 to ~10 mm
- None

What is the shape of the opacities? (second choice)

- p - rounded diameter up to ~1.5 mm
- q - rounded diameter 1.5 to ~3 mm
- r - rounded diameter 3 to ~10 mm
- s - irregular width up to ~1.5 mm
- t - irregular width 1.5 to ~3 mm
- u - irregular width 3 to ~10 mm
- None

In which zones are these opacities distributed?

- [ ] Right upper
- [ ] Left upper
- [ ] Right middle
- [ ] Left middle
- [ ] Right lower
- [ ] Left lower

What is the (average) profusion of these opacities?

- 0/0 - None
- 0/-
- 0/1
- 1/0
- 1/1
- 1/2
- 2/1
- 2/2
- 2/3
- 3/2
- 3/3
- 3/+

Are there any large opacities?

- A - One or more large opacity with a diameter of <50mm but >10mm or several opacities >10mm but with a SUM of their dimensions <50mm
- B - One large opacity having the longest dimension exceeding 50 mm but not exceeding the equivalent area of the right upper zone, or several large opacities with the sum of their longest dimensions exceeding 50 mm but not exceeding the equivalent area of the right upper zone.
- C - One large opacity which exceeds the equivalent area of the right upper zone, or several large opacities which, when combined, exceed the equivalent area of the right upper zone.
- No

Are there any classifiable pleural abnormalities?

- Yes
- No

Are there any pleural plaques?

- Yes
- No

Where are the pleural plaques?

- [ ] Chest wall - right
- [ ] Chest wall - left
- [ ] Face on - right
- [ ] Face on - left
- [ ] Diaphragm - right
- [ ] Diaphragm - left
- [ ] Other - right
- [ ] Other - left

Is there calcification of the plaques?

- [ ] Chest wall - right
- [ ] Chest wall - left
- [ ] Face on - right
- [ ] Face on - left
- [ ] Diaphragm - right
- [ ] Diaphragm - left
- [ ] Other - right
- [ ] Other - left

What is the extent of the RIGHT sided plaques?

- ( ) None
- ( ) < 1/4 of the chest wall
- ( ) >1/4 but <1/2 of the chest wall
- ( ) >1/2 of the chest wall

What is the extent of the LEFT sided plaques?

- ( ) None
- ( ) < 1/4 of the chest wall
- ( ) >1/4 but <1/2 of the chest wall
- ( ) >1/2 of the chest wall

What is the width of the RIGHT sided plaques?

- ( ) None
- ( ) >3 mm to <5 mm
- ( ) > 5mm to <10 mm
- ( ) > 10 mm

What is the width of the LEFT sided plaques?

- None
- >3 mm to <5 mm
- > 5mm to <10 mm
- > 10 mm

Is there costophrenic angle obliteration?

- Yes
- No

Where is the diffuse pleural plaque thickening?

- In profile - right
- In profile - left
- Face on - right
- Face on - left
- Other - right
- Other - left
- None

Is there calcification?

- In profile - right
- In profile - left
- Face on - right
- Face on - left
- Other - right
- Other - left
- None

What is the extent of the RIGHT sided thickening?

- None
- < 1/4 of the chest wall
- >1/4 but <1/2 of the chest wall
- >1/2 of the chest wall

What is the extent of the LEFT sided thickening?

- None
- < 1/4 of the chest wall
- >1/4 but <1/2 of the chest wall
- >1/2 of the chest wall

What is the width of the RIGHT sided thickening?

- None
- >3 mm to <5 mm
- > 5mm to <10 mm
- > 10 mm

What is the width of the LEFT sided thickening?

- None
- >3 mm to <5 mm
- > 5mm to <10 mm
- > 10 mm

Are there signs of active tuberculosis?

- Yes - probable
- Yes - possible
- No

Does there appear to be miliary TB disease?

- Yes
- No

On a scale of 0 (not at all likely) to 50 (equally likely to be TB or not) to 100 (certain this is TB), how likely is the diagnosis of active tuberculosis?



Are there other findings?

- Yes
- No

9. Are there any other findings?

- [ ] aa - atherosclerotic aorta
- [ ] at - significant apical pleural thickening
- [ ] ax - coalescence of small opacities
- [ ] bu - bulla(e)
- [ ] ca - cancer, thoracic malignancies excluding mesothelioma
- [ ] cg - calcified non-pneumoconiotic nodules (e.g. granuloma) or nodes
- [ ] cn - calcification in small pneumoconiotic opacities
- [ ] co - abnormality of cardiac size or shape
- [ ] cp - cor pulmonale
- [ ] di - marked distorition of intrathoracic cavity
- [ ] ef - pleural effusion
- [ ] em - emphysema
- [ ] es - eggshell calcification of hilar or mediastinal lymph nodes
- [ ] fr - fractured rib(s) (acute or healed)
- [ ] hi - enlargement of non-calcified hilar or mediastinal lymph nodes
- [ ] ho - honeycomb lung
- [ ] id - ill-defined diaphragm border
- [ ] ih - ill-defined heart border
- [ ] kl - septal (Kerley) lines
- [ ] me – mesothelioma
- [ ] pa - plate atelectasis
- [ ] pi - pleural thickening of an interlobar fissure
- [ ] px - pneumothorax
- [ ] ra - rounded atelectasis
- [ ] rp - rheumatoid pneumoconiosis
- [ ] od - other disease or significant abnormality

Are there changes consistent with post TB lung disease?

- ( ) Yes - probable
- ( ) Yes - possible
- ( ) No

Is there fibrocystic or fibrotic (including parenchymal bands) disease?

- ( ) Yes
- ( ) No

In which zones is the fibrocystic disease or fibrotic disease distributed?

- [ ] Right upper
- [ ] Left upper
- [ ] Right middle
- [ ] Left middle
- [ ] Right lower
- [ ] Left lower

What overall proportion of lung do the fibrocystic or fibrotic disease fill?



Are there inflitrates (non-nodular) consisent with TB disease?

- ( ) Yes
- ( ) No

In which zones is/are the infiltrate(s) distributed?

- [ ] Right upper
- [ ] Left upper
- [ ] Right middle
- [ ] Left middle
- [ ] Right lower
- [ ] Left lower

What overall proportion of the lung is covered by the infiltrates?



Are there nodules that are consisent with TB?

- Yes
- No

In which zones are the nodules distributed?

- Right upper
- Left upper
- Right middle
- Left middle
- Right lower
- Left lower

What proportion of the lung do the nodules cover



Are there cavities?

- Yes
- No

In which zones are the cavities distributed?

- Right upper
- Left upper
- Right middle
- Left middle
- Right lower
- Left lower

What overall proportion of the lung is covered by cavities?



Is there bronchiectasis?

- Yes
- No

In which zones is the bronchiectasis distributed?

- Right upper
- Left upper
- Right middle
- Left middle
- Right lower
- Left lower

What overall proportion of the lung is bronchiectatic?



What proportion of the lung is destroyed due to post TB lung disease (not active disease)?



Is there consolidation (not inflitrates)?

- ○ Yes
- ○ No

In which zones is the consolidation distributed?

- ☐ Right upper
- ☐ Left upper
- ☐ Right middle
- ☐ Left middle
- ☐ Right lower
- ☐ Left lower

What overall proportion of the lung is consolidated?



10. Comments and verbal report