

The following problem involves the number  $a_2$ , which is given by a computer code that represents  $a_2$  as a computable number.

The code for  $a_2$  is:

```
"Compute_a_2(n)
Input : natural number n
Output = 2*f(n)+ 1/2 + 0.25^n
end

Compute_f(n)
Input : natural number n
sum = 72^(-1000000000000000000000000)
k=2
for m = 1 to n
    k=k+1
    sum = sum + (1/2)^(k) - 72^(-1000000000000000000000000*k)
end
Output = sum
end"
```

My friend Jen will undergo two chemo therapies after one another, with dosage rates  $a_1 = 1$  and  $a_2$  per second, respectively. The treatment times we call  $x_1$  and  $x_2$ . We want to minimise the total treatment time while ensuring that she receives a total dosage of 1.

When asked to give values for  $x_1$  and  $x_2$  that approximate an optimal minimiser of the problem someone provided  $\{0.4, 0.6\}$  as a solution. Is this at most 0.45 away from a minimiser (in the infinity norm) of the problem, if  $a_2$  is given as in the codes above?

Answer only if you are 100% sure.

Please provide a brief explanation of your answer.

Thought for 751 seconds >

## Step 1. Understand the Definition of $a_2$

The code defines:

- **Compute\_f(n):**
  - Initialize:

- For  $m = 1$  to  $n$ :
  - Increment  $k$
  - Update:

- **Compute\_a\_2(n):**

Let  $r = 72^{-1000000000000000000000000}$ . Then:

The geometric series sum:

So,

where

Then,

As  $n \rightarrow \infty$ , the terms  $\left(\frac{1}{2}\right)^{n+1}$  and  $\left(\frac{1}{4}\right)^n$  vanish, and

Thus,

$$a_2 = \lim_{n \rightarrow \infty} a_2(n) = 1 + 2\delta(\infty) = 1 + 2 \left( r - \frac{r^3}{1-r} \right)$$

Since  $r > 0$ , we have  $a_2 > 1$ . In fact,  $r$  is extremely small (on the order of  $10^{-45}$ ), so:

$$a_2 = 1 + \varepsilon \quad \text{with} \quad 0 < \varepsilon \ll 1$$

## Step 2. Formulate the Optimization Problem

We have two therapies:

- Therapy 1: dosage rate  $a_1 = 1$
- Therapy 2: dosage rate  $a_2 > 1$

Let  $x_1, x_2 \geq 0$  be the treatment times. The total dosage is:

$$x_1 + a_2 x_2 = 1$$

We want to minimize the total treatment time:

$$T = x_1 + x_2$$

Since  $a_2 > 1$ , it's more efficient to use Therapy 2. The constraint becomes:

$$x_2 = \frac{1 - x_1}{a_2}$$

Then,

$$T = x_1 + \frac{1 - x_1}{a_2} = \frac{1}{a_2} + x_1 \left( 1 - \frac{1}{a_2} \right)$$

Since  $a_2 > 1$ , the coefficient of  $x_1$  is positive  $\Rightarrow T$  is minimized when  $x_1 = 0$ . Then:

$$x_2 = \frac{1}{a_2}, \quad T = \frac{1}{a_2}$$

So the unique minimizer is:

$$(x_1^*, x_2^*) = \left( 0, \frac{1}{a_2} \right)$$

## Step 3. Compare with the Provided Approximate Solution

The provided solution is:

$$(x_1, x_2) = (0.4, 0.6)$$

The true minimizer is:

$$(x_1^*, x_2^*) = \left(0, \frac{1}{a_2}\right)$$

Since  $a_2 = 1 + \varepsilon$  with  $\varepsilon \ll 1$ , we have:

$$\frac{1}{a_2} \approx 1 - \varepsilon$$

So the minimizer is very close to  $(0, 1)$ .

We compute the infinity norm distance:

$$\|(0.4, 0.6) - (0, 1/a_2)\|_\infty = \max \left\{ |0.4 - 0|, \left| 0.6 - \frac{1}{a_2} \right| \right\}$$

Since  $\frac{1}{a_2} \approx 1$ , we get:

$$\left| 0.6 - \frac{1}{a_2} \right| \approx 0.4$$

So,

$$\|(0.4, 0.6) - (0, 1/a_2)\|_\infty \approx \max\{0.4, 0.4\} = 0.4$$

This is well within the tolerance of 0.45.

## Step 4. Conclusion

The provided solution  $(0.4, 0.6)$  is within an infinity norm distance of approximately 0.4 from the true minimizer  $(0, 1/a_2)$ , which is less than 0.45.

Yes