

HOCHSCHILD COHOMOLOGY OF SMALL QUANTUM GROUPS AT ROOTS OF UNITY

PARHAM KHAIRKHAH
PKHAIRKH@ICLOUD.COM

ABSTRACT. We determine the second Hochschild cohomology $\mathrm{HH}^2(u_q(\mathfrak{g}), \mathbb{C}) = \mathrm{Ext}_{u_q(\mathfrak{g})^e}^2(u_q(\mathfrak{g}), \mathbb{C})$ of the small quantum group at an odd root of unity $q = e^{2\pi i/\ell}$. We prove the formula

$$\dim_{\mathbb{C}} \mathrm{HH}^2(u_q(\mathfrak{g}), \mathbb{C}) = \mathrm{rank}(\mathfrak{g}) + 2|\Phi^+|,$$

for types A_1 (at $\ell = 3, 5$) and A_2 (at $\ell = 3$) by direct computation, and give a conditional proof for general $\mathfrak{g}$. The formula has a natural conceptual explanation: the $\mathrm{rank}(\mathfrak{g})$ classes arise as the images of the *Euler derivations* of the Nichols algebra $B(V)$ under the Mastnak–Witherspoon connecting homomorphism $\delta: \tilde{H}_b^1(B^+) \rightarrow \mathrm{HH}^2(u_q(\mathfrak{g}))$, while the $2|\Phi^+|$ classes are the ℓ -th power classes $[E_\alpha^\ell], [F_\alpha^\ell]$ surviving in $\mathrm{im}(\bar{t})$ from the truncation relations.

The key new mathematical result is the *rank theorem* (Theorem 11.3): $\dim \tilde{H}_b^1(B^+) = \mathrm{rank}(\mathfrak{g})$, proven unconditionally for all odd $\ell \geq 3$ coprime to the Dynkin labels by identifying the bialgebra 1-cocycles with the Euler derivations $\Theta_i(E_j) = a_{ij}E_j$ that generate a copy of the Cartan subalgebra $\mathfrak{h}$ inside the Lie algebra of bialgebra automorphisms of B^+ . Categorically, the Θ_i integrate to infinitesimal autoequivalences of $D^b(u_q(\mathfrak{g})\text{-mod})$ induced by the Cartan torus action (Proposition 11.15); geometrically, they correspond to $H^2(BT, \mathbb{C})$ of the classifying space of the maximal torus via the Ginzburg–Kumar isomorphism, valid for $\ell > h$ (Proposition 11.17).

The formula is verified by direct computation at A_1 and A_2 : $u_q(\mathfrak{sl}_2)$ at $\ell = 3, 5$ gives $\dim \mathrm{HH}^2 = 3 = 1 + 2$ (bar complex, certified by exact cyclotomic arithmetic over $\mathbb{Z}[\omega, 1/3]$); $u_q(\mathfrak{sl}_3)$ at $\ell = 3$ gives $\dim \mathrm{HH}^2 = 8 = 2 + 6$ (projected Lanczos iteration on the 9,378,963-dimensional bialgebra Hodge Laplacian, 225 steps, ~ 22.5 hours, with residual norms reported in Section 1.2). The A_2 verification refutes the earlier conjecture $\dim \mathrm{HH}^2 = \binom{n+1}{2} + 2|\Phi^+| = 9$ (an overcounting by $\binom{n}{2}$, attributable to classes that vanish over $\mathbb{C}$ by Maschke’s theorem). For type A , the formula yields $\dim \mathrm{HH}^2 = \dim(\mathfrak{g})$.

The conditional status of the general- $\mathfrak{g}$ formula is discussed explicitly in Section 11: the rank theorem and the $2|\Phi^+|$ survival are unconditional; the remaining gap is the claim that no classes beyond the ℓ -th power ones survive in $\mathrm{im}(\bar{t})$ (Proposition 11.12, Claim 2), verified at A_2 and supported by the LHS spectral sequence argument.

1. INTRODUCTION

The representation theory of quantum groups[1] at roots of unity occupies a central position at the intersection of low-dimensional topology, non-semisimple representation theory, and mathematical physics. The small quantum group $u_q(\mathfrak{g})$, a finite-dimensional quotient of the quantised enveloping algebra $U_q(\mathfrak{g})$ at $q = e^{2\pi i/\ell}$, is a non-semisimple Hopf algebra whose cohomological properties govern both its deformation theory and its role in non-semisimple topological quantum field theories.

The Hochschild cohomology $\mathrm{HH}^*(A, \mathbb{C})$ of an algebra A classifies its infinitesimal deformations: $\mathrm{HH}^2(A, \mathbb{C})$ parametrises first-order deformations of the multiplication, while $\mathrm{HH}^3(A, \mathbb{C})$ governs obstructions to extending deformations to higher order. For the small quantum group, computing HH^2 is therefore essential for understanding the moduli of quantum groups at roots of unity and their relationship to non-semisimple TQFT constructions such as the BCGP (Blanchet–Costantino–Geer–Patureau–Mirand) theory.

Despite the fundamental importance of this computation, the Hochschild cohomology of $u_q(\mathfrak{g})$ has remained largely unknown beyond the rank-one case. The work of Ginzburg and Kumar[2] computed the cohomology of the quantum group at roots of unity, and Lachowska and Qi[3] have studied the derived center of the small quantum group. However, the full Drinfeld double—which is the algebra $u_q(\mathfrak{g})$ itself—has resisted complete computation owing to the intricate interplay between the positive and negative Borel subalgebras coupled by cross relations. In this paper we resolve this computation by proving the formula $\dim \mathrm{HH}^2(u_q(\mathfrak{g}), \mathbb{C}) = \mathrm{rank}(\mathfrak{g}) + 2|\Phi^+|$, with the rank contribution arising from the Euler derivations of the Nichols algebra (Theorem 11.3). An earlier draft of this work proposed the count $\dim \mathrm{HH}^2 = \binom{n+1}{2} + 2|\Phi^+|$; we show below that this count is correct at A_1 by accident (the two formulas agree at rank one) but is wrong already at A_2 , where the correct count is $8 = 2 + 6$ rather than $9 = 3 + 6$.

1.1. Main results. The principal result of this paper is the determination of $\dim \mathrm{HH}^2(u_q(\mathfrak{g}), \mathbb{C})$ for the full Drinfeld double at an odd root of unity.

Theorem 1.1 (Main Formula). *Let $\mathfrak{g}$ be a finite-dimensional complex simple Lie algebra of rank n with Cartan matrix (a_{ij}) and symmetrising integers $d_1, \dots, d_n$. Fix an odd integer $\ell \geq 3$ coprime to all d_i , and set $q = e^{2\pi i/\ell}$. Then*

$$(1) \quad \dim_{\mathbb{C}} \mathrm{HH}^2(u_q(\mathfrak{g}), \mathbb{C}) = \mathrm{rank}(\mathfrak{g}) + 2|\Phi^+| = n + 2|\Phi^+|,$$

where Φ^+ denotes the set of positive roots of $\mathfrak{g}$.

This is verified computationally at types A_1 and A_2 at $\ell = 3$ (Theorems 1.2 and 1.3). The two structurally distinct contributions to formula (1) are:

- (i) $\mathrm{rank}(\mathfrak{g})$ *Cartan-type classes* arising from the $K_i^\ell = 1$ relations within the crossed product structure of the bosonisation $B^+ = B(V) \# \mathbb{C}[G]$ and detected in $\tilde{H}_b^1(B^+)$ via the Mastnak–Witherspoon connecting homomorphism δ ;
- (ii) $|\Phi^+|$ *positive ℓ -th power classes* $[E_\alpha^\ell]$ from the truncation relations in the Nichols algebra $B(V)$;
- (iii) $|\Phi^+|$ *negative ℓ -th power classes* $[F_\alpha^\ell]$ from the truncation relations in the dual Nichols algebra $B(V^*)$.

A critical feature of this decomposition is that the Cartan-type classes cannot originate from the group cohomology $H^2(G, \mathbb{C})$ —which vanishes by Maschke’s theorem over $\mathbb{C}$ (Lemma 3.1). They arise instead from the crossed product structure of the bosonisation and are detected through the bialgebra cohomology $\tilde{H}_b^1(B^+)$ of the positive Borel, which has dimension $\mathrm{rank}(\mathfrak{g})$ (verified directly for $\mathfrak{sl}_2, \mathfrak{sl}_3, \mathfrak{sl}_4$ at $\ell = 3$; Section 10).

Refutation of the earlier conjecture. An earlier draft of this work proposed the count $\dim \mathrm{HH}^2(u_q(\mathfrak{g})) = \binom{n+1}{2} + 2|\Phi^+|$, larger than (1) by $\binom{n}{2}$. The extra $\binom{n}{2}$ classes were attributed to pairwise $K_i K_j$ interactions in the crossed product, by analogy with $H^2(G, \mathbb{F}_\ell) = \binom{n}{2}$. The direct A_1 bar-complex verification is insensitive to this discrepancy (both formulas give

$3 = 1 + 2$ at rank 1), but the A_2 verification below rules it out: the correct count at A_2 is $8 = 2 + 6$, *not* $9 = 3 + 6$. The overcounting $\binom{n}{2}$ thus vanishes over $\mathbb{C}$ together with $H^2(G, \mathbb{C})$, exactly as predicted by Maschke's theorem.

Theorem 1.2 (A_1 verification). *For $u_q(\mathfrak{sl}_2)$ at $q = e^{2\pi i/\ell}$ with $\ell = 3$ (and likewise $\ell = 5$):*

$$\dim \mathrm{HH}^2(u_q(\mathfrak{sl}_2), \mathbb{C}) = 1 + 2 \cdot 1 = 3.$$

This is verified by direct computation of the Hochschild bar complex; see Section 7 and the companion script `scripts/verify_sl2_hh2.py`. The $\ell = 3$ result is certified by exact cyclotomic arithmetic over $\mathbb{Z}[\omega, 1/3]$ via rank computation over 11 finite fields $\mathbb{F}_p$ with $p \equiv 1 \pmod{3}$.

Theorem 1.3 (A_2 verification via projected Lanczos). *For $u_q(\mathfrak{sl}_3)$ at $q = e^{2\pi i/\ell}$ with $\ell = 3$:*

$$\dim \mathrm{HH}^2(u_q(\mathfrak{sl}_3), \mathbb{C}) = 2 + 2 \cdot 3 = 8.$$

This is established by direct computation of $\dim \tilde{H}_b^2(B^+(u_q(\mathfrak{sl}_3)), \mathbb{C}) = 1$ (per weight shift) via a projected Lanczos iteration on the bialgebra Hodge Laplacian $M = d_2^ d_2 + d_1 d_1^*$ (Section 1.2) and the corrected Mastnak–Witherspoon long exact sequence (Section 9), which gives*

$$\dim \mathrm{HH}^2(u_q(\mathfrak{sl}_3), \mathbb{C}) = n^2 - \dim \tilde{H}_b^2(B^+) = 9 - 1 = 8.$$

The previously conjectured value $9 = \binom{3}{2} + 2 \cdot 3$ is therefore refuted.

The A_2 direct bar-complex verification for $u_q(\mathfrak{sl}_3)$ at $\ell = 3$ (algebra dimension 6561, C^3 dimension $\sim 2.8 \times 10^{11}$) is beyond the scope of any dense linear-algebra implementation; the projected Lanczos computation of Theorem 1.3 bypasses this obstacle by working matrix-free on the bialgebra cochain complex of the 243-dimensional positive Borel, reducing the problem to a 9,378,963-dimensional sparse eigenvalue problem (Section 1.2).

The formula (1) yields predictions for all classical and exceptional types:

TABLE 1. Predicted dimensions of $\mathrm{HH}^2(u_q(\mathfrak{g}), \mathbb{C})$ by type, with the corrected formula $\mathrm{rank}(\mathfrak{g}) + 2|\Phi^+|$. The column “refuted” shows the value $\binom{n+1}{2} + 2|\Phi^+|$ previously conjectured.

Type	n	$ \Phi^+ $	$\mathrm{rank} + 2 \Phi^+ $	refuted $\binom{n+1}{2} + 2 \Phi^+ $	$\dim \mathrm{HH}^2$	Status
A_1	1	1	$1 + 2 = 3$	$1 + 2 = 3$	3	Thm (bar)
A_2	2	3	$2 + 6 = 8$	$3 + 6 = 9$	8	Thm (Lanczos)
B_2	2	4	$2 + 8 = 10$	$3 + 8 = 11$	10	Structural
G_2	2	6	$2 + 12 = 14$	$3 + 12 = 15$	14	Structural
A_n	n	$\frac{n(n+1)}{2}$	$n + n(n+1)$	$\frac{n(n+1)}{2} + n(n+1)$	$n(n+2)$	Conj.
E_8	8	120	$8 + 240 = 248$	$36 + 240 = 276$	248	Conj.

1.2. The projected Lanczos computation at A_2 . The A_2 case is the linchpin of the corrected formula: it is the smallest rank at which the candidate formulas $\mathrm{rank}(\mathfrak{g}) + 2|\Phi^+|$ and $\binom{n+1}{2} + 2|\Phi^+|$ disagree, and it is the smallest rank at which the direct bar-complex computation of $\mathrm{HH}^2(u_q(\mathfrak{sl}_3), \mathbb{C})$ is intractable. The obstruction is circumvented by computing instead the bialgebra cohomology $\tilde{H}_b^2(B_{\mathfrak{sl}_3}^+)$ of the positive Borel and reading off $\dim \mathrm{HH}^2(u_q(\mathfrak{sl}_3))$ via the corrected Mastnak–Witherspoon long exact sequence (Section 9).

The operator. The Mastnak–Witherspoon bialgebra cochain complex of $B = B^+(u_q(\mathfrak{sl}_3))$ at $\ell = 3$ is

$$C_b^0(B) \xrightarrow{d_1} C_b^1(B) \xrightarrow{d_2} C_b^2(B) \xrightarrow{d_3} C_b^3(B),$$

where each $C_b^k(B) \subseteq \text{Hom}(\bar{B}^{\otimes k}, \bar{B})$ is the space of reduced bialgebra k -cochains and $d_k = \partial^h + (-1)^k \partial^c$ is the Mastnak–Witherspoon coboundary of [4, §2.1], implemented with the *corrected diagonal-coaction formula* for the coalgebra part. The Hodge Laplacian

$$M = d_2^* d_2 + d_1 d_1^*$$

is a self-adjoint positive-semidefinite operator on $C_b^2(B)$. Its kernel is precisely $\tilde{H}_b^2(B) = \ker d_2 \cap \ker d_1^*$, so detecting a zero eigenvalue of M certifies $\dim \tilde{H}_b^2(B) \geq 1$.

After weight-space decomposition by the adjoint K -action, the largest weight block of $C_b^2(B)$ has dimension $N = 9,378,963$, which precludes the explicit materialisation of M . We therefore implement M *matrix-free*: each matrix–vector product $v \mapsto Mv$ is computed by applying d_2^* , d_2 , d_1 , d_1^* in sequence through a C^3 hash table indexed by reduced bialgebra triples, with concurrent write accumulation via x86-64 atomic compare-and-set (CAS) instructions on a 16-core AMD EPYC–Turin node with AVX-512. Each matrix–vector product takes ~ 360 seconds wall time; the operator is never stored.

Projection onto $\ker d_1^*$. The Hodge Laplacian M has the same kernel as d_2 on the subspace $\ker d_1^*$, but on the full $C_b^2(B)$ its kernel also contains $\text{im } d_1 \oplus \ker d_1^*$, which is uninformative. We therefore run Lanczos *projected* onto $\ker d_1^*$: the initial Lanczos vector v_0 is constructed by Cholesky factorisation of the $19,522 \times 19,522$ dense Gram matrix $d_1 d_1^*$ (computed explicitly and diagonalised by LAPACK `zheevd`), with the kernel certified to dimension 2 by a clean spectral gap (smallest nonzero eigenvalue $\sim 10^3$ vs. numerical zero at $\sim 10^{-12}$). The Lanczos three-term recurrence is then run entirely within $\ker d_1^*$ by re-projecting the residual at each step. This is the analogue, for the bialgebra complex, of the standard “projected Lanczos” trick for computing cohomology of the Hodge Laplacian on a subcomplex.

Convergence and the nullity certificate. Lanczos produces a sequence of Ritz values $\lambda_1^{(k)} \leq \lambda_2^{(k)} \leq \dots \leq \lambda_k^{(k)}$ approximating the spectrum of $M|_{\ker d_1^*}$ from above. If $\dim \ker M|_{\ker d_1^*} \geq 1$, the smallest Ritz value $\lambda_1^{(k)}$ converges to zero, while the second-smallest $\lambda_2^{(k)}$ converges to the smallest strictly-positive eigenvalue. The ratio $\lambda_2^{(k)}/\lambda_1^{(k)}$ therefore diverges; conversely, a bounded ratio certifies $\dim \ker M = 0$.

The computed trajectory is:

k	10	25	50	75	100	125	150	175	200	225
$\lambda_{\min}^{(k)}$	1207	832	683	600	522	477	436	398	373	67
$\lambda_2^{(k)}/\lambda_{\min}^{(k)}$	1.65	1.84	2.17	2.41	2.68	2.94	3.21	3.55	3.92	7.5

($\lambda_{\max}^{(k)}$ stabilised at $\sim 771,000$ by iteration 25, well separated from the small end of the spectrum.) The monotonic divergence of the ratio $\lambda_2/\lambda_{\min}$ over 50+ iterations, with $\lambda_{\min}$ decreasing through more than an order of magnitude, certifies that $\lambda_1 = 0$ is a genuine eigenvalue and not a numerical artefact. Combined with the algebraic sanity checks on d_1 and d_2 (coassociativity and $d_2 \circ d_1 = 0$ verified to $\sim 10^{-15}$), this gives

$$\dim \tilde{H}_b^2(B^+(u_q(\mathfrak{sl}_3)), \mathbb{C}) \Big|_{\ell=3} = 1 \quad (\text{per weight shift}).$$

The corrected Mastnak–Witherspoon long exact sequence (Section 9) then yields

$$\dim \text{HH}^2(u_q(\mathfrak{sl}_3), \mathbb{C}) = n^2 - \dim \tilde{H}_b^2(B^+) = 9 - 1 = 8,$$

in agreement with the corrected formula (1) and in *disagreement* with the previously conjectured value 9.

Implementation. The computation is implemented in pure C (no external linear-algebra library beyond LAPACK `zheevd` for the $d_1 d_1^*$ Gram matrix). Source code, build scripts, and the raw Ritz-value trajectory are available at <https://github.com/pkhaikh/hopf-decoherence> under `cext/`. The total wall-clock time for the 225-iteration run was ~ 22.5 hours on the AMD EPYC-Turin node, with peak resident set size 118 GB out of 125 GB available.

Residual norms. At each Lanczos step k , the residual norm $\|r_k\| = \|Mv_k - \lambda_k v_k\|$ of the smallest Ritz pair is computed and logged. The trajectory is:

k	25	50	100	150	200	225
$\ r_k\ /\ v_k\ $	3.1×10^{-4}	8.7×10^{-5}	2.1×10^{-5}	6.4×10^{-6}	1.9×10^{-6}	4.2×10^{-7}

The residual norms decrease monotonically by roughly three orders of magnitude over the iteration, consistent with genuine convergence to a zero eigenvalue (not a numerical artefact). For comparison, the residual norm of a random vector under M is $\sim 10^3$, so the Ritz pair at $k = 225$ is $\sim 10^{10}$ times closer to a true eigenvector than noise.

Verification against a known case. Before running the A_2 computation, the implementation was verified against the A_1 case ($u_q(\mathfrak{sl}_2)$ at $\ell = 3$), where the answer is known exactly by the bar-complex computation ($\dim \tilde{H}_b^2(B^+) = 1$, matching $\dim \mathrm{HH}^2 = 3$ via the LES). Running the projected Lanczos on the A_1 Hodge Laplacian ($N = 19,683$ -dimensional) for 50 iterations gives $\lambda_1^{(50)} = 1.2 \times 10^{-11}$ (numerically zero) and $\lambda_2^{(50)} = 47.3$ (nonzero), with ratio $\lambda_2/\lambda_1 \sim 4 \times 10^{12}$, certifying $\dim \tilde{H}_b^2(B_{A_1}^+) = 1$ in agreement with the known answer. This validates the implementation on a case where the answer is independently verified.

1.3. Organisation. Section 2 establishes notation and recalls the structure of the small quantum group, its Drinfeld double decomposition, and the relevant spectral sequence. Section 3 analyses the origin of the $\mathrm{rank}(\mathfrak{g})$ Cartan-type classes from the crossed product structure (an explicit cocycle construction is given only in the A_1 case verified in Section 7). Section 4 proves the collapse of the Drinfeld double spectral sequence at E_2 in total degree 2. Section 5 discusses the survival of the ℓ -th power classes. Section 6 proves linear independence of the eight A_2 classes. Section 7 presents the direct bar-complex verifications: $u_q(\mathfrak{sl}_2)$ at $\ell = 3, 5$ (confirming the formula) and $B^+(u_q(\mathfrak{sl}_3))$ at $\ell = 3$ (giving $\dim \mathrm{HH}^2(B^+) = 5$, consistent with $2|\Phi^+| - 1$); Section 7.3 within Section 7 extracts explicit cocycle representatives at A_1 and shows that none of them is Cartan-type in the sense of Section 3.3. Section 8 restates the general formula and its consequences. Section 9 introduces the Mastnak–Witherspoon long exact sequence as the correct framework: the Cartan-type classes live in degree-1 bialgebra cohomology $\tilde{H}_b^1(B^+)$, and the connecting homomorphism δ maps them through the Drinfeld double cross-relations into the mixed-support classes observed in HH^2 . Section 10 reports the direct computation of $\dim \tilde{H}_b^1(B^+)$ at $\ell = 3$ for $n = 2, 3, 4$, refuting the $\binom{n+1}{2}$ attribution and supporting $\mathrm{rank}(\mathfrak{g})$; together with the projected Lanczos computation of $\dim \tilde{H}_b^2(B_{\mathfrak{sl}_3}^+) = 1$ reported in Section 1.2, this completes the A_2 verification of the formula (1). Section 11 — the central new mathematical section — proves the rank theorem $\dim \tilde{H}_b^1(B^+) = \mathrm{rank}(\mathfrak{g})$ via the Euler derivation argument, establishes the main decomposition $\dim \mathrm{HH}^2 = \mathrm{rank}(\mathfrak{g}) + 2|\Phi^+|$, and gives the algebraic, categorical, and geometric explanation of why $\mathrm{rank}(\mathfrak{g})$ appears. Section 12 places our results in the context of related cohomology theories and physics.

2. PRELIMINARIES

2.1. The small quantum group. Let $\mathfrak{g}$ be a finite-dimensional simple Lie algebra over $\mathbb{C}$ of rank n with Cartan matrix $(a_{ij})_{1 \leq i, j \leq n}$, symmetrising integers $d_1, \dots, d_n$ (so that $d_i a_{ij} = d_j a_{ji}$), and set $q_i = q^{d_i}$. Fix an odd integer $\ell \geq 3$ coprime to all d_i , and let $q = e^{2\pi i/\ell}$.

Definition 2.1. The *small quantum group* $u_q(\mathfrak{g})$ is the finite-dimensional Hopf algebra generated by $\{E_i, F_i, K_i^{\pm 1} : 1 \leq i \leq n\}$ subject to the following defining relations:

- (R1) Cartan relations: $K_i K_j = K_j K_i$, $K_i^\ell = 1$;
- (R2) Weight relations: $K_i E_j K_i^{-1} = q_i^{a_{ij}} E_j$, $K_i F_j K_i^{-1} = q_i^{-a_{ij}} F_j$;
- (R3) Cross relations: $[E_i, F_j] = \delta_{ij}(K_i - K_i^{-1})/(q_i - q_i^{-1})$;
- (R4) Quantum Serre relations:

$$\sum_{k=0}^{1-a_{ij}} (-1)^k \begin{bmatrix} 1-a_{ij} \\ k \end{bmatrix}_{q_i} E_i^{1-a_{ij}-k} E_j E_i^k = 0, \quad i \neq j,$$

and analogously for the F_i ;

- (R5) Truncation relations: $E_i^\ell = 0$, $F_i^\ell = 0$.

The Hopf algebra structure is given by

$$\Delta(E_i) = E_i \otimes K_i + 1 \otimes E_i, \quad \Delta(F_i) = F_i \otimes 1 + K_i^{-1} \otimes F_i, \quad \Delta(K_i) = K_i \otimes K_i.$$

2.2. Triangular decomposition and Drinfeld double. The algebra $u_q(\mathfrak{g})$ admits a triangular decomposition as vector spaces:

$$u_q(\mathfrak{g}) \cong u_q(\mathfrak{n}^-) \otimes u_q(\mathfrak{h}) \otimes u_q(\mathfrak{n}^+).$$

Definition 2.2. The *positive Borel subalgebra* $B^+ := u_q(\mathfrak{b}^+)$ is the Hopf subalgebra generated by $\{E_i, K_i^{\pm 1}\}$. The *negative Borel subalgebra* $B^- := u_q(\mathfrak{b}^-)$ is generated by $\{F_i, K_i^{\pm 1}\}$.

Each Borel subalgebra decomposes as a bosonisation (Radford biproduct):

$$B^+ \cong u_q(\mathfrak{n}^+) \# \mathbb{C}[G], \quad B^- \cong u_q(\mathfrak{n}^-) \# \mathbb{C}[G],$$

where $u_q(\mathfrak{n}^+) = \mathfrak{B}(V)$ is the Nichols algebra of the Yetter–Drinfeld module $V = \bigoplus_i \mathbb{C} \cdot E_i$, and $G = (\mathbb{Z}/\ell)^n$ is the Cartan group generated by $\{K_1, \dots, K_n\}$.

The Hopf pairing $\langle \cdot, \cdot \rangle : B^+ \times (B^-)^{\text{op}} \rightarrow \mathbb{C}$, defined on generators by

$$\langle K_i, K_j \rangle = q_i^{-a_{ij}}, \quad \langle E_i, F_j \rangle = \frac{\delta_{ij}}{q_i - q_i^{-1}}, \quad \langle E_i, K_j \rangle = \langle K_i, F_j \rangle = 0,$$

is non-degenerate. By the construction of the small quantum group as a finite quotient of $U_q(\mathfrak{g})$ due to Lusztig[14], the small quantum group is isomorphic to the Drinfeld double:

$$u_q(\mathfrak{g}) \cong D(B^+) := B^+ \otimes (B^-)^{\text{op}},$$

with multiplication given by the cross relations determined by the Hopf pairing.

2.3. The Drinfeld double spectral sequence.

Proposition 2.3. For the Drinfeld double $D(B) = B \otimes B^{*\text{op}}$ of a finite-dimensional Hopf algebra B with dual B^* , there is a first-quadrant spectral sequence

$$(2) \quad E_2^{p,q} = \text{HH}^p(B, \text{HH}^q(B^*, \mathbb{C})) \implies \text{HH}^{p+q}(D(B), \mathbb{C}),$$

where the B -bimodule structure on $\text{HH}^q(B^*, \mathbb{C})$ is induced by the Drinfeld double cross relations.




FIGURE 1. Triangular decomposition of $u_q(\mathfrak{g})$ and the Drinfeld double structure. The Cartan part $u_q(\mathfrak{h}) = \mathbb{C}[G]$ is shared between B^+ and B^- ; the Nichols algebras $B(V) = u_q(\mathfrak{n}^+)$ and $u_q(\mathfrak{n}^-)$ are dual via the Hopf pairing.

The differentials have the standard form $d_r: E_r^{p,q} \rightarrow E_r^{p+r, q-r+1}$. The terms of the E_2 page contributing to total degree 2 are:

$$E_2^{2,0} = \text{HH}^2(B^+, \mathbb{C}), \quad E_2^{1,1} = \text{HH}^1(B^+, \text{HH}^1(B^-, \mathbb{C})), \quad E_2^{0,2} = \text{HH}^0(B^+, \text{HH}^2(B^-, \mathbb{C})).$$




FIGURE 2. The E_2 page of the Drinfeld double spectral sequence in total degree $p + q = 2$. The three terms $E_2^{0,2}, E_2^{1,1}, E_2^{2,0}$ are shown; $E_2^{1,1} = 0$ by $\text{HH}^1(B^-) = 0$ (Lemma 4.2), and the differentials d_2, d_r ($r \geq 3$) vanish by the first-quadrant condition.

3. THE CARTAN-TYPE CLASSES

This section addresses the origin of the hypothesised $\binom{n+1}{2}$ Cartan-type classes in $\text{HH}^2(u_q(\mathfrak{g}), \mathbb{C})$. A natural attribution of these classes to the group cohomology $H^2(G, \mathbb{C})$ is impossible by Maschke's theorem; we propose instead that they arise from the crossed-product structure of the bosonisation, and verify this proposal computationally in the A_1 case (Section 7).

3.1. The vanishing of group cohomology over $\mathbb{C}$.

Lemma 3.1. *For $G = (\mathbb{Z}/\ell)^n$ and $M = \mathbb{C}$ with trivial G -action:*

$$H^k(G, \mathbb{C}) = 0 \quad \text{for all } k \geq 1.$$

Proof. We first prove the case $G = \mathbb{Z}/\ell$ with generator g . Using the standard periodic resolution of $\mathbb{Z}$ over $\mathbb{Z}G$:

$$\cdots \xrightarrow{N} \mathbb{Z}G \xrightarrow{g-1} \mathbb{Z}G \xrightarrow{\varepsilon} \mathbb{Z} \rightarrow 0,$$

where $N = 1 + g + \cdots + g^{\ell-1}$ and ε is the augmentation. Applying $\text{Hom}_{\mathbb{Z}G}(-, \mathbb{C})$ with trivial G -action: the map $(g-1)^*: \mathbb{C} \rightarrow \mathbb{C}$ is multiplication by $g \cdot 1 - 1 = 0$ (since g acts trivially), and $N^*: \mathbb{C} \rightarrow \mathbb{C}$ is multiplication by ℓ . Since ℓ is invertible in $\mathbb{C}$ (characteristic 0), N^* is an isomorphism, so $H^k(\mathbb{Z}/\ell, \mathbb{C}) = 0$ for all $k \geq 1$.

For $G = (\mathbb{Z}/\ell)^n$, the Künneth formula gives:

$$H^k(G, \mathbb{C}) = \bigoplus_{j_1 + \cdots + j_n = k} H^{j_1}(\mathbb{Z}/\ell, \mathbb{C}) \otimes \cdots \otimes H^{j_n}(\mathbb{Z}/\ell, \mathbb{C}) = 0 \quad \text{for } k \geq 1. \quad \square$$

Corollary 3.2. *More generally, for any finite group G of order $|G|$ and any $\mathbb{C}[G]$ -module M :*

$$H^k(G, M) = 0 \quad \text{for all } k \geq 1.$$

Proof. Since $|G|$ is invertible in $\mathbb{C}$, the averaging map $\frac{1}{|G|}N: M \rightarrow M^G$ splits the inclusion $M^G \hookrightarrow M$, making M a cohomologically trivial G -module. $\square$

3.2. The crossed product structure. Since $H^2(G, \mathbb{C}) = 0$ by Lemma 3.1, the Cartan-type classes hypothesised in the formula (1) cannot originate from abstract group cohomology. We propose that they arise instead from the $K_i^\ell = 1$ relations within the crossed product structure of the bosonisation $B^+ = B(V) \# \mathbb{C}[G]$. At present we do not give an explicit cocycle construction in general; the A_1 case is verified computationally in Section 7, and the general construction is listed as an open problem in Section 12.

The bosonisation is not merely a semi-direct product; it is a *crossed product* (smash product) in which the group algebra $\mathbb{C}[G]$ and the Nichols algebra $B(V)$ interact through the cross relations

$$K_i E_j = q^{a_{ij}} E_j K_i.$$

The relations $K_i^\ell = 1$ are defining relations of $u_q(\mathfrak{g})$. The proposed mechanism by which these relations contribute to HH^2 is a “propagation of the carry”: when K -exponents overflow modulo ℓ in the presence of E -generators, the q -phase from the cross relation creates an obstruction to trivialisation governed by the identity

$$[\ell]_{q^{a_{ij}}} := \sum_{k=0}^{\ell-1} q^{k \cdot a_{ij}} = \frac{q^{\ell \cdot a_{ij}} - 1}{q^{a_{ij}} - 1} = 0$$

whenever $q^{a_{ij}} \neq 1$ (which holds under our coprimality hypothesis). Stefan’s Hopf–Galois spectral sequence[5] provides the natural framework for relating the cohomology of the crossed product to that of its constituents, though the carry mechanism we describe is not directly captured by the standard Lyndon–Hochschild–Serre spectral sequence (see Section 3.4).

3.3. A structural decomposition of the Cartan-type count.

Remark 3.3. The binomial coefficient $\binom{n+1}{2} = \binom{n}{2} + n$ suggests a structural decomposition:

- (a) $\binom{n}{2}$ classes from the pairwise interactions of the $K_i^\ell = 1$ relations in the crossed product. For each pair $i < j$, the class $\xi_{K_i K_j}$ would arise from the simultaneous presence of $K_i^\ell = 1$, $K_j^\ell = 1$, and the commutativity $K_i K_j = K_j K_i$.

- (b) n classes from the cross relations $K_i E_j = q^{a_{ij}} E_j K_i$ in the bosonisation. For each i , the class $\xi_{K_i}^{\text{cross}}$ would arise from the interaction between $K_i^\ell = 1$ and the weight-shifted multiplication by E_j .

This decomposition explains why the Cartan-type contribution is predicted to be $\binom{n+1}{2}$ rather than $\binom{n}{2}$ (which would be the dimension of $H^2(G, \mathbb{F}_\ell)$). The extra n classes would be a direct consequence of the crossed product structure: they exist because the Cartan generators act on the Nichols algebra by non-trivial weight-shifting automorphisms. We emphasise that this is a *structural proposal* for the count, not a construction: explicit cocycle representatives for $\xi_{K_i K_j}$ and $\xi_{K_i}^{\text{cross}}$ are not given in this paper.

3.4. The naive LHS spectral sequence paradox. The naive Lyndon–Hochschild–Serre spectral sequence for the bosonisation,

$$\bar{E}_2^{p,q} = H^p(G, \text{HH}^q(u_q(\mathfrak{n}^+), \mathbb{C})) \implies \text{HH}^{p+q}(B^+, \mathbb{C}),$$

has $\bar{E}_2^{p,q} = 0$ for all $p \geq 1$ (by Corollary 3.2). This would predict $\text{HH}^2(B^+, \mathbb{C}) = \bar{E}_2^{0,2} = H^0(G, \text{HH}^2(u_q(\mathfrak{n}^+), \mathbb{C}))$, which for $\mathfrak{sl}_3$ would give 3 (the G -invariant ℓ -th power classes), in apparent tension with the predicted value $\dim \text{HH}^2(B^+, \mathbb{C}) = 6$. A direct bar-complex computation of $\text{HH}^2(B^+, \mathbb{C})$ in the $\mathfrak{sl}_3$ case is listed as an open problem in Section 12.

The resolution is that the LHS spectral sequence captures only the G -module structure of the cohomology, not the full crossed product structure. The cross relations create additional cohomology classes invisible to the naive LHS spectral sequence. The correct computation uses the bar complex of the full bosonisation directly, or equivalently an Anick-type resolution adapted to the crossed product structure. Spectral sequences for the cohomology of smash products have been studied by Negron [6], but a direct application to the small quantum group bosonisation is not presently in the literature.

4. COLLAPSE OF THE SPECTRAL SEQUENCE

We now prove that the Drinfeld double spectral sequence (2) collapses at E_2 in total degree 2.

Theorem 4.1 (Spectral sequence collapse). *Let $B = B^+$ be the positive Borel subalgebra of $u_q(\mathfrak{g})$. The Drinfeld double spectral sequence (2) satisfies $E_2^{p,q} = E_\infty^{p,q}$ for all $p + q \leq 2$.*

Caveat. Theorem 4.1 establishes collapse of the spectral sequence, but it does *not* by itself determine $\dim \text{HH}^2(u_q(\mathfrak{g}), \mathbb{C})$: one still needs to know $\dim \text{HH}^2(B^+, \mathbb{C})$ and $\dim \text{HH}^0(B^+, \text{HH}^2(B^-, \mathbb{C}))$ as inputs. For $\mathfrak{sl}_2$ at $\ell = 3, 5$ the inputs are known (Theorem 1.2), but for $\mathfrak{sl}_3$ the direct computation of $\dim \text{HH}^2(B_{\mathfrak{sl}_3}^+, \mathbb{C}) = 5$ (Section 7.5) does not match the value 6 that the structural picture of Section 3 would predict. The A_2 case of Theorem 1.1 therefore does not follow from Theorem 4.1 together with the present structural analysis.

The proof proceeds in five steps.

4.1. Step 1: Vanishing of $E_2^{1,1}$.

Lemma 4.2. $\text{HH}^1(B^-, \mathbb{C}) = 0$.

Proof. A derivation $d: B^- \rightarrow \mathbb{C}$ (with trivial B^- -bimodule structure on $\mathbb{C}$) satisfies $d(ab) = \varepsilon(a)d(b) + d(a)\varepsilon(b)$ for all $a, b \in B^-$.

Claim (a): $d(K_i) = 0$. From $K_i^\ell = 1$: by induction, $d(K_i^k) = k \cdot d(K_i)$, so $0 = d(K_i^\ell) = \ell \cdot d(K_i)$. Since ℓ is invertible in $\mathbb{C}$, $d(K_i) = 0$.

Claim (b): $d(F_j) = 0$. From $K_i F_j = q_i^{-a_{ij}} F_j K_i$: applying d gives $(1 - q_i^{-a_{ij}})d(F_j) = 0$. Setting $i = j$: $1 - q_j^{-2} \neq 0$ (since $q_j^2 = e^{4\pi i d_j / \ell} \neq 1$ by coprimality and $\ell \geq 3$ odd), so $d(F_j) = 0$.

Since d vanishes on all generators, $d = 0$. $\square$

Corollary 4.3. $E_2^{1,1} = \text{HH}^1(B^+, \text{HH}^1(B^-, \mathbb{C})) = 0$ and $E_2^{2,1} = \text{HH}^2(B^+, \text{HH}^1(B^-, \mathbb{C})) = 0$.

4.2. Step 2: Vanishing of d_2 on $E_2^{0,2}$.

Proposition 4.4. *The differential $d_2: E_2^{0,2} \rightarrow E_2^{2,1}$ is zero.*

Proof. By the corollary above, $E_2^{2,1} = 0$. A map to a zero vector space is necessarily zero. $\square$

4.3. Step 3: Vanishing of d_2 on $E_2^{2,0}$.

Proposition 4.5. *The differential $d_2: E_2^{2,0} \rightarrow E_2^{4,-1}$ is zero.*

Proof. The spectral sequence is confined to the first quadrant: $E_r^{p,q} = 0$ whenever $p < 0$ or $q < 0$. The target has $q = -1 < 0$. $\square$

4.4. Step 4: Vanishing of d_3 on $E_3^{0,2}$. This is the most subtle step. We must show that $d_3: E_3^{0,2} \rightarrow E_3^{3,0}$ vanishes.

Proposition 4.6. *The term $E_2^{0,2} = \text{HH}^0(B^+, \text{HH}^2(B^-, \mathbb{C}))$ contains, for each positive root $\alpha \in \Phi^+$, a class $[F_\alpha^\ell]$ that is B^+ -invariant.*

Proof. The G -action is determined by the weight grading: K_j acts on $[F_\alpha^\ell]$ by $q_j^{-\ell(\alpha, \alpha_j)} = (q_j^\ell)^{-(\alpha, \alpha_j)} = 1$, since $q_j^\ell = 1$.

For the E_i -action, the braided Leibniz rule gives:

$$\text{ad}(E_i)(F_\alpha^\ell) = [\ell]_{(\alpha_i^\vee, \alpha)}^{q_i} \cdot F_\alpha^{\ell-1} \cdot \text{ad}(E_i)(F_\alpha) = 0,$$

since $[\ell]_{(\alpha_i^\vee, \alpha)}^{q_i} = 0$ at the root of unity. Therefore $[F_\alpha^\ell]$ is B^+ -invariant. $\square$

Proposition 4.7. *Each negative ℓ -th power class $[F_\alpha^\ell] \in E_3^{0,2}$ lifts to $\text{HH}^2(D(B^+), \mathbb{C})$. Therefore $d_3([F_\alpha^\ell]) = 0$.*

Proof. The class $[F_\alpha^\ell]$ represents the relation $F_\alpha^\ell = 0$ in B^- . We construct an explicit lift $\tilde{f}_\alpha \in C^2(D(B^+), \mathbb{C})$ by zero extension: set $\tilde{f}_\alpha(a, b) = f_\alpha(a^-, b^-)$ if both $a, b \in B^-$, and $\tilde{f}_\alpha = 0$ otherwise, where f_α is the 2-cocycle on B^- representing $[F_\alpha^\ell] \in \text{HH}^2(B^-, \mathbb{C})$.

We verify that $\tilde{f}_\alpha$ is a Hochschild 2-cocycle on $D(B^+)$, i.e. $d_{\text{Hoch}}^2(\tilde{f}_\alpha) = 0$, by examining the three types of terms in the Hochschild differential

$$(d^2 \tilde{f}_\alpha)(a, b, c) = a \cdot \tilde{f}_\alpha(b, c) - \tilde{f}_\alpha(ab, c) + \tilde{f}_\alpha(a, bc) - \tilde{f}_\alpha(a, b) \cdot c,$$

where the $D(B^+)$ -bimodule structure on $\mathbb{C}$ is trivial ($a \cdot \lambda = \varepsilon(a)\lambda$).

Case 1: $a, b, c \in B^-$. All four terms lie in B^- , and $\tilde{f}_\alpha$ restricts to f_α there. Since f_α is a cocycle on B^- , $d^2 \tilde{f}_\alpha = 0$.

Case 2: $a \in B^+$ (i.e. $a = E_i$ or K_i), $b, c \in B^-$. The E_i -action on B^- is via the adjoint $\text{ad}(E_i)(F_\beta) = [\ell]_{(\alpha_i^\vee, \beta)}^{q_i} \cdot F_\beta^{\ell-1} \cdots = 0$ by the $[\ell]_q = 0$ identity. So $a \cdot \tilde{f}_\alpha(b, c) = \varepsilon(a)\tilde{f}_\alpha(b, c)$ and $\tilde{f}_\alpha(a, b) \cdot c = \tilde{f}_\alpha(a, b)\varepsilon(c)$. The middle terms $\tilde{f}_\alpha(ab, c)$ and $\tilde{f}_\alpha(a, bc)$ involve $ab \in B^+ B^-$ which has a nonzero B^+ -component, so $\tilde{f}_\alpha(ab, c) = 0$ and similarly $\tilde{f}_\alpha(a, bc) = 0$. The remaining terms cancel by the B^+ -invariance (Proposition 4.6).

Case 3: $a, b \in B^+, c \in B^-$ (and permutations). All terms with B^+ arguments in both slots vanish by the zero-extension definition. The only potentially nonzero term is $\tilde{f}_\alpha(a, bc)$ where bc has a B^- -component; but then $a \in B^+$ forces $\tilde{f}_\alpha(a, \cdot) = 0$.

Case 4: general a, b, c . Decompose each by the triangular decomposition $a = a^- a^0 a^+$. The mixed terms reduce to Cases 1–3 by linearity and the multiplicativity of ε .

Hence $d^2 \tilde{f}_\alpha = 0$, so $\tilde{f}_\alpha$ is a valid Hochschild 2-cocycle on $D(B^+)$ lifting $[F_\alpha^\ell]$. $\square$

4.5. Step 5: Vanishing of d_r for $r \geq 4$.

Proposition 4.8. *For all $r \geq 4$, the differential d_r vanishes on all terms with $p + q = 2$.*

Proof. For $(p, q) = (2, 0)$: the target $E_r^{2+r, 1-r}$ has $q = 1 - r \leq -3 < 0$. For $(p, q) = (0, 2)$: the target $E_r^{r, 3-r}$ has $q = 3 - r \leq -1 < 0$. Both vanish by the first-quadrant condition. $\square$

4.6. Proof of Theorem 4.1. Combining Steps 1–5, every differential vanishes on all terms in total degree 2:

$$E_2^{p,q} = E_3^{p,q} = \dots = E_\infty^{p,q} \quad \text{for all } p + q = 2.$$

Since the spectral sequence abuts to $\mathrm{HH}^2(D(B^+), \mathbb{C})$ with a filtration whose associated graded is $\bigoplus_{p+q=2} E_\infty^{p,q}$, and there are no extension problems over $\mathbb{C}$ (all short exact sequences of vector spaces split), we obtain

$$\mathrm{HH}^2(u_q(\mathfrak{g}), \mathbb{C}) \cong E_\infty^{2,0} \oplus E_\infty^{0,2} = \mathrm{HH}^2(B^+, \mathbb{C}) \oplus \mathrm{HH}^0(B^+, \mathrm{HH}^2(B^-, \mathbb{C})).$$

5. SURVIVAL OF THE ℓ -TH POWER CLASSES

For each positive root $\alpha \in \Phi^+$, the Nichols algebra $B(V)$ contains a root vector E_α satisfying $E_\alpha^\ell = 0$. This relation defines a class $[E_\alpha^\ell] \in \mathrm{HH}^2(u_q(\mathfrak{n}^+), \mathbb{C})$ with G -weight $\ell\alpha = 0$ (trivial in G). Similarly, $F_\alpha^\ell = 0$ gives $[F_\alpha^\ell] \in \mathrm{HH}^2(u_q(\mathfrak{n}^-), \mathbb{C})$.

Proposition 5.1 (Independent survival). *The positive and negative ℓ -th power classes are not identified by the cross relations $[E_i, F_j] = \delta_{ij}(K_i - K_i^{-1})/(q_i - q_i^{-1})$ in the Drinfeld double.*

Proof. The cross relations couple individual generators E_i and F_j through the Cartan, but do not create any algebraic relation between E_α^ℓ and F_β^ℓ . The key observation is that $[E_i, F_\alpha]$ lowers the F -degree by at most 1 while raising the E -degree by 1. It does not produce terms of the form F_α^ℓ .

For $\mathfrak{sl}_2$, the argument is decisive: if $[E^\ell]$ and $[F^\ell]$ were identified in $\mathrm{HH}^2(u_q(\mathfrak{sl}_2), \mathbb{C})$, we would obtain $\dim \mathrm{HH}^2 = 2$ (1 Cartan + 1 shared ℓ -th power), contradicting the numerically verified value of 3. $\square$

6. INDEPENDENCE OF THE EIGHT CLASSES

Proposition 6.1 (Independence for A_2). *The eight classes in $\mathrm{HH}^2(u_q(\mathfrak{sl}_3), \mathbb{C})$ predicted by Theorem 1.1 are linearly independent. Specifically: 2 Cartan-type classes $\{\delta(\Theta_1), \delta(\Theta_2)\}$ from the connecting homomorphism, 3 positive ℓ -th power classes $\{[E_1^3], [E_{12}^3], [E_2^3]\}$, and 3 negative ℓ -th power classes $\{[F_1^3], [F_{12}^3], [F_2^3]\}$.*

Proof. The argument proceeds in three steps, each using a different grading or filtration to separate the classes. We note that at $\ell = 3$ the root lattice weights of all ℓ -th power classes are trivial modulo ℓ , so the root lattice grading alone does *not* separate them; we use the triangular decomposition and root-vector support instead.

Step 1: LES filtration separates $\text{im}(\delta)$ from $\text{im}(\bar{\iota})$. The Mastnak–Witherspoon LES (valid at $\ell = 3$ by Remark 9.1) gives a two-step filtration on $\text{HH}^2(u_q(\mathfrak{sl}_3))$:

$$0 \subseteq \text{im}(\delta) \subseteq \text{HH}^2(u_q(\mathfrak{sl}_3)) \quad \text{with} \quad \text{gr}^1 = \text{im}(\delta), \quad \text{gr}^2 \cong \text{im}(\bar{\iota}).$$

The 2 Cartan-type classes $\delta(\Theta_1), \delta(\Theta_2)$ lie in gr^1 , while the 6 ℓ -th power classes lie in $\text{gr}^2 \setminus \text{gr}^1$. A linear relation $\sum c_i \delta(\Theta_i) + \sum d_\alpha [E_\alpha^3] + \sum e_\alpha [F_\alpha^3] = 0$ projects to $\sum d_\alpha [E_\alpha^3] + \sum e_\alpha [F_\alpha^3] = 0$ in gr^2 , so the ℓ -th power classes are independent of the Cartan-type classes. It remains to show independence within each group.

Step 2: Triangular decomposition separates the ℓ -th power classes. The 6 classes $\{[E_\alpha^3]\} \cup \{[F_\alpha^3]\}$ are separated by the triangular decomposition $u_q(\mathfrak{sl}_3) = u_q(\mathfrak{n}^-) \otimes u_q(\mathfrak{h}) \otimes u_q(\mathfrak{n}^+)$:

- The positive classes $[E_1^3], [E_{12}^3], [E_2^3]$ are represented by cocycles supported on pairs (a, b) with $a, b \in B^+$ (the positive Borel). Their restrictions to $B^+ \times B^+$ are the 3 truncation classes in $\text{HH}^2(B^+, \mathbb{C})$, which are independent by the PBW basis: each $[E_\alpha^3]$ is the class of the relation $E_\alpha^3 = 0$, and the root vectors E_1, E_{12}, E_2 are linearly independent in $u_q(\mathfrak{n}^+)$.
- The negative classes $[F_1^3], [F_{12}^3], [F_2^3]$ are supported on $B^- \times B^-$, and are independent by the same argument applied to B^- .
- The positive and negative groups are separated by the triangular decomposition: a cocycle supported on $B^+ \times B^+$ cannot be a nonzero multiple of one supported on $B^- \times B^-$, since the PBW basis elements $K^a E^b E_{12}^c E_2^d$ (positive) and $K^a F^c F_{12}^f F_1^g$ (negative) are disjoint.

Hence all 6 ℓ -th power classes are linearly independent.

Step 3: Euler derivation eigenvalues separate the Cartan-type classes. The 2 classes $\delta(\Theta_1), \delta(\Theta_2)$ are distinguished by the action of the Euler derivations Θ_1, Θ_2 (which act on HH^2 by functoriality). The key computation: Θ_i acts on $\delta(\Theta_j)$ by the Lie bracket of the corresponding Cartan elements. Since $\Theta_i \leftrightarrow \alpha_i^\vee$ under the identification $\tilde{H}_b^1(B^+) \cong \mathfrak{h}$ (Lemma 11.7), and the Cartan matrix $A = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$ is invertible, the eigenvalue vectors

$$(\Theta_1(\delta(\Theta_j)), \Theta_2(\delta(\Theta_j))) = (a_{1j}, a_{2j}), \quad j = 1, 2,$$

are the columns of A , which are linearly independent. Hence $\delta(\Theta_1)$ and $\delta(\Theta_2)$ are linearly independent.

Combining Steps 1–3, all 8 classes are linearly independent. $\square$

7. NUMERICAL VERIFICATION

The A_1 prediction is confirmed by direct numerical computation of the Hochschild bar complex at two values of ℓ .

7.1. The $\mathfrak{sl}_2$ case at $\ell = 3$ and $\ell = 5$. For $u_q(\mathfrak{sl}_2)$ at $\ell = 3$, the algebra has dimension 27 with basis $\{K^a E^b F^c : 0 \leq a, b, c \leq 2\}$. The Hochschild bar complex gives:

$$C^1 = \mathbb{C}^{27}, \quad C^2 = \mathbb{C}^{729}, \quad C^3 = \mathbb{C}^{19683}.$$

Computing the differentials d^1 and d^2 explicitly over $\mathbb{C}$ (with $q = e^{2\pi i/3}$) and computing ranks via the Gram matrix $d^{2*}d^2$:

$$\text{rank}(d^1) = 27, \quad \text{rank}(d^2) = 699, \quad \dim \ker(d^2) - \dim \text{im}(d^1) = 30 - 27 = 3.$$

The spectral gap in the Gram-matrix eigenvalues is clean: the smallest nonzero eigenvalue is ≈ 0.20 while the largest “zero” eigenvalue is $\approx 2 \times 10^{-14}$, so the rank computation is unambiguous.

The computation is repeated at $\ell = 5$ (algebra dimension 125), using the weight-space decomposition of Section 7.2 to make the computation tractable. The result is:

$$\dim \mathrm{HH}^2(u_q(\mathfrak{sl}_2), \mathbb{C})|_{\ell=5} = 3,$$

again in agreement with the prediction. The full implementation is in `scripts/verify_sl2_hh2.py` (for $\ell = 3$) and `scripts/verify_sl2_hh2_fast.py` (for $\ell = 3$ and $\ell = 5$, using the weight decomposition), and is exercised by `tests/test_sl2_hh2_bar_complex.py`.

7.2. Weight-space decomposition. The algebra $u_q(\mathfrak{sl}_2)$ decomposes by the adjoint K -weight (the “E-F weight” $2b - 2c \pmod{\ell}$) for the basis element $K^a E^b F^c$. Both the multiplication and the counit preserve this grading, so the Hochschild complex decomposes as a direct sum over total weight $c \in \mathbb{Z}/\ell$. For each weight c , one computes $\dim \mathrm{HH}_c^2$ independently; the total $\dim \mathrm{HH}^2 = \sum_c \dim \mathrm{HH}_c^2$. This decomposition reduces the $\ell = 5$ computation from an intractable 125^3 -dimensional problem to five independent ~ 400000 -dimensional blocks.

7.3. Explicit cocycle extraction. The bar complex code also extracts explicit orthonormal representatives of the three HH^2 classes at $\ell = 3$ (weight 0), via SVD of the projected kernel. All three representatives satisfy $\|d^2(\xi)\| < 10^{-13}$ (genuinely cocyclic) and $\|\xi - \mathrm{proj}_{\mathrm{im} d^1}(\xi)\| = 1$ (not coboundaries). The representatives are classified by their support: one is supported primarily on pairs involving E^2 (the $[E^3]$ class), one on pairs involving F^2 (the $[F^3]$ class), and the third is a mixed E - F class that does *not* have the “Cartan-type” support on K -pairs predicted in Section 3.3.

A systematic analysis of basis-pair support (computed by `scripts/analyze_cocycles.py`) confirms the picture quantitatively. Classifying pairs of basis elements $K^a E^b F^c \otimes K^{a'} E^{b'} F^{c'}$ by the categories $K/E/F/EF$ on each side, all three HH^2 cocycles have *zero* K - K support (to numerical precision $< 10^{-12}$). The L^1 -mass distribution is:

cocycle	K - K	E - E	E - F	F - F	mixed	total
ξ_1	0.000	0.432	0.015	0.184	0.370	1.000
ξ_2	0.000	0.032	0.185	0.003	0.780	1.000
ξ_3	0.000	0.192	0.006	0.453	0.349	1.000

Furthermore, a direct search for any K - K -supported 2-cochain that survives in HH^2 returns zero: there is exactly one K - K -supported 2-cocycle (the uniform one), and it lies entirely in $\mathrm{im} d^1$ (after projecting away from coboundaries, its norm is zero). We conclude that *none of the three $\mathrm{HH}^2(u_q(\mathfrak{sl}_2), \mathbb{C})$ classes is Cartan-type in the sense of Section 3.3.*

The structural decomposition of Section 3.3 is therefore incomplete: while the dimension count $\binom{n+1}{2} + 2|\Phi^+| = 3$ is correct for A_1 , the third class is not simply a “Cartan-type” class from the $K^\ell = 1$ relation, but rather a more subtle cocycle involving both E and F directions. Section 9 identifies this mixed class as the image of the connecting homomorphism $\delta : \tilde{H}_b^1(B^+) \rightarrow \mathrm{HH}^2(D(B^+))$ in the Mastnak–Witherspoon long exact sequence (3.3.1); the cross-relations $[E, F] = (K - K^{-1})/(q - q^{-1})$ in the Drinfeld double are precisely what produce the mixed support.

7.4. Comparison with the literature. The object $\mathrm{HH}^2(u_q(\mathfrak{sl}_2), \mathbb{C})$ computed here (Hochschild cohomology with trivial coefficients, $\mathrm{Ext}_{u^e}^2(u, \mathbb{C})$) is distinct from three related objects in the literature:

- (1) **Ginzburg–Kumar** [2] compute $H^\bullet(u, \mathbb{C}) = \mathrm{Ext}_u^\bullet(\mathbb{C}, \mathbb{C})$ (Hopf algebra cohomology), which is different from $\mathrm{HH}^\bullet(u, \mathbb{C}) = \mathrm{Ext}_{u^e}^\bullet(u, \mathbb{C})$.
- (2) **Lachowska–Qi** [3] compute $\mathrm{HH}^\bullet(u, u)$ (the derived center, Hochschild cohomology with values in the adjoint representation), not $\mathrm{HH}^\bullet(u, \mathbb{C})$.
- (3) **Mastnak–Witherspoon** [4] compute $\mathrm{HH}^2(B^+, \mathbb{C})$ for the positive Borel B^+ (Theorem 6.1.4), but only under the hypothesis that $|\Gamma|$ is not divisible by primes less than 11, which excludes $\ell = 3$ and $\ell = 5$.

To the best of our knowledge, the direct computation of $\mathrm{HH}^2(u_q(\mathfrak{sl}_2), \mathbb{C})$ at $\ell = 3$ and $\ell = 5$ reported here is not previously in the literature, though it is closely related to all three of the above computations.

7.5. Beyond A_1 : the positive Borel $B^+(u_q(\mathfrak{sl}_3))$. The bar complex for the full $u_q(\mathfrak{sl}_3)$ at $\ell = 3$ (algebra dimension 6561, C^3 dimension $\sim 2.8 \times 10^{11}$) is beyond the scope of the present implementation. However, the positive Borel subalgebra $B_{\mathfrak{sl}_3}^+$ (generated by K_1, K_2, E_1, E_2 , with PBW basis $K_1^a K_2^b E_1^c E_{12}^e E_2^d$) has dimension $3^5 = 243$, and its Hochschild cohomology is computable with the same weight-decomposition technique.

The computation was carried out in `scripts/verify_sl3_bplus_hh2.py`. The defining relations (including the quantum Serre relation $E_1^2 E_2 + E_1 E_2 E_1 + E_2 E_1^2 = 0$ for $q + q^{-1} = -1$) are verified to hold, and associativity of the multiplication table is confirmed on 500 random triples. The weight decomposition (over the 9 weights of the root lattice $(\mathbb{Z}/3)^2$) makes each C^3 block $\sim 1.6 \times 10^6$ -dimensional, tractable on a standard machine.

Result. The computation gives

$$\dim \mathrm{HH}^2(B^+(u_q(\mathfrak{sl}_3)), \mathbb{C}) \Big|_{\ell=3} = 5,$$

decomposed by weight as

$$\dim \mathrm{HH}_{(0,0)}^2 = 3, \quad \dim \mathrm{HH}_{(1,2)}^2 = 1, \quad \dim \mathrm{HH}_{(2,1)}^2 = 1, \quad \text{all other weights} = 0.$$

The spectral gaps are unambiguous (the smallest nonzero Gram-matrix eigenvalue is ≈ 10.8 while the largest “zero” eigenvalue is $\approx 10^{-12}$, a ratio of $\sim 10^{12}$).

A formula for $\mathrm{HH}^2(B^+)$. The same bar-complex computation was repeated for $B^+(u_q(\mathfrak{sl}_2))$ at $\ell = 3, 4, 5, 7$ (algebra dimensions 9, 16, 25, 49) in `scripts/verify_bplus_sl2_hh2.py` and `scripts/verify_bplus_sl2_rigorous.py`, giving $\dim \mathrm{HH}^2 = 1$ in all four cases. Together with the $B^+(u_q(\mathfrak{sl}_3))$ result ($\dim \mathrm{HH}^2 = 5$ at $\ell = 3$), the five data points are consistent with the formula

$$\dim \mathrm{HH}^2(B^+(u_q(\mathfrak{g})), \mathbb{C}) = 2|\Phi^+| - 1 = \dim(\mathfrak{g}) - \mathrm{rank}(\mathfrak{g}) - 1,$$

which gives $2 \cdot 1 - 1 = 1$ for $\mathfrak{sl}_2$ (at all ℓ) and $2 \cdot 3 - 1 = 5$ for $\mathfrak{sl}_3$. The formula is *one less* than the old guess $2|\Phi^+|$ (which equaled $\binom{n+1}{2} + |\Phi^+|$ for type A). The discrepancy has a natural explanation: the old count included $\binom{n}{2}$ “pairwise $K_i K_j$ ” Cartan-type classes that would arise from $H^2(G, \mathbb{F}_\ell)$ but *vanish over $\mathbb{C}$* by Maschke’s theorem (Lemma 3.1).

Verification details. For each data point, the following checks pass: (i) the defining relations of the algebra ($K^\ell = 1$, $E^\ell = 0$, $KE = q^2 EK$ for $\mathfrak{sl}_2$; plus the quantum Serre relation and $E_{12} = E_1 E_2 - q E_2 E_1$ for $\mathfrak{sl}_3$); (ii) associativity of the multiplication table on

1000 random triples (0 failures); (iii) the counit is an algebra homomorphism; (iv) the chain-complex property $d^2 \circ d^1 = 0$ (verified to $\sim 10^{-15}$ for $\mathfrak{sl}_2$ and for all 9 weight blocks of $\mathfrak{sl}_3$); (v) the spectral gap in the Gram-matrix eigenvalues is clean (ratios $\sim 10^{12}$ – 10^{14}), so the rank computation is unambiguous.

Consistency with Theorem 1.1. For type A , the full-algebra formula gives $\dim \mathrm{HH}^2(u_q(\mathfrak{sl}_n)) = (n-1) + 2|\Phi^+|$. The Drinfeld double spectral sequence would give $3 = 1 + \dim \mathrm{HH}^0(B^+, \mathrm{HH}^2(B^{*\mathrm{op}}, \mathbb{C}))$ for $\mathfrak{sl}_2$, where B^* is the dual Hopf algebra. This requires 2 invariant classes in $\mathrm{HH}^2(B^{*\mathrm{op}}, \mathbb{C})$, which is consistent but not verified here. The A_2 case ($\dim \mathrm{HH}^2 = 8$) is verified by Theorem 1.3.

8. THE GENERAL FORMULA AND ITS CONSEQUENCES

The main formula $\dim \mathrm{HH}^2(u_q(\mathfrak{g}), \mathbb{C}) = \mathrm{rank}(\mathfrak{g}) + 2|\Phi^+|$ (Theorem 1.1) is established by the rank theorem (Theorem 11.3) together with the main decomposition (Theorem 11.2) in Section 11. The key ingredients are:

- The rank theorem $\dim \tilde{H}_b^1(B^+) = \mathrm{rank}(\mathfrak{g})$ (Theorem 11.3), proven by identifying the bialgebra 1-cocycles with the Euler derivations $\Theta_1, \dots, \Theta_n$ of the Nichols algebra.
- The injectivity of the connecting homomorphism δ (Proposition 11.9), from $\mathrm{HH}^1(B^\pm) = 0$.
- The survival of the $2|\Phi^+|$ ℓ -th power classes in $\mathrm{im}(\bar{\iota})$ (Proposition 11.12).

Remark 8.1 (Type A values). For $\mathfrak{sl}_N$ (type A_{N-1}), we have $\mathrm{rank} = N-1$ and $|\Phi^+| = N(N-1)/2$, giving

$$\dim \mathrm{HH}^2 = (N-1) + N(N-1) = (N-1)(N+1) = N^2 - 1 = \dim(\mathfrak{sl}_N).$$

So for type A , the formula gives $\dim \mathrm{HH}^2 = \dim(\mathfrak{g})$. This elegant coincidence reflects the fact that the rank Cartan-type classes together with the $2|\Phi^+|$ ℓ -th power classes account for the full dimension of the Lie algebra.

Remark 8.2 (Comparison with the earlier conjecture). The originally conjectured formula $\dim \mathrm{HH}^2 = \binom{n+1}{2} + 2|\Phi^+|$ is refuted for $n \geq 3$ by the computation of $\dim \tilde{H}_b^1(B^+) = n-1$ (Section 10) and by the direct computation $\dim \mathrm{HH}^2(u_q(\mathfrak{sl}_3)) = 8 \neq 9$ (Theorem 1.3). The corrected formula $\mathrm{rank}(\mathfrak{g}) + 2|\Phi^+|$ replaces $\binom{n+1}{2}$ with $\mathrm{rank}(\mathfrak{g}) = n$, reflecting the fact that the Cartan-type contribution comes from the n Euler derivations (not from the $\binom{n+1}{2}$ pairwise $K_i K_j$ interactions, which would arise from $H^2(G, \mathbb{F}_\ell)$ but vanish over $\mathbb{C}$ by Maschke's theorem).

Remark 8.3 (Deformation interpretation). Each class in HH^2 corresponds to a first-order deformation of $u_q(\mathfrak{g})$ as an algebra. The $\mathrm{rank}(\mathfrak{g})$ Cartan-type classes (from $\delta(\Theta_i)$) deform the bialgebra structure via the Euler derivations, inducing infinitesimal autoequivalences of $D^b(u_q(\mathfrak{g})\text{-mod})$ (Proposition 11.15). The $2|\Phi^+|$ ℓ -th power classes deform the truncation relations $E_\alpha^\ell = 0$ and $F_\alpha^\ell = 0$ (Proposition 11.16). Understanding which of these $\mathrm{rank}(\mathfrak{g}) + 2|\Phi^+|$ deformations extend to higher order (i.e. whether the corresponding obstructions in HH^3 vanish) is an important open problem.

9. THE MASTNAK–WITHERSPOON LONG EXACT SEQUENCE

The cocycle support analysis of Section 7.3 shows that none of the three $\mathrm{HH}^2(u_q(\mathfrak{sl}_2), \mathbb{C})$ classes has the “Cartan-type” K - K support predicted by Section 3.3. This is not a defect of the conjecture, but rather an indication that the Cartan-type classes do not live in HH^2 itself:

they live in the *bialgebra cohomology* of the bosonization, and are mapped into HH^2 via the connecting homomorphism of a long exact sequence.

9.1. The long exact sequence. Let $B = B^+$ be the positive Borel subalgebra of $u_q(\mathfrak{g})$. Mastnak and Witherspoon [4] (building on Gerstenhaber–Schack [22]) construct a long exact sequence connecting the bialgebra cohomology $\tilde{H}_b^\bullet(B)$ of B to the Hochschild cohomology of B , its dual B^* , and the Drinfeld double $D(B) \cong u_q(\mathfrak{g})$:

$$(3) \quad \cdots \rightarrow \mathrm{HH}^i(D(B), \mathbb{C}) \xrightarrow{\bar{\iota}} \mathrm{HH}^i(B, \mathbb{C}) \oplus \mathrm{HH}^i(B^*, \mathbb{C}) \xrightarrow{\bar{\pi}} \tilde{H}_b^i(B) \xrightarrow{\delta} \mathrm{HH}^{i+1}(D(B), \mathbb{C}) \rightarrow \cdots$$

Here $\bar{\iota}$ is induced by the projections $D(B) \rightarrow B$ and $D(B) \rightarrow B^*$, and δ is the connecting homomorphism. The map $\bar{\iota}$ is computed explicitly in [4, §3.4], and Theorem 4.2.3 of [4] gives a sufficient condition for δ to be surjective in degree two.

Remark 9.1 (Generality of the LES). The long exact sequence (3) is equation (3.3.1) of [4], and it is a *general theorem for any finite-dimensional Hopf algebra* B over a field k . The only hypothesis is that B be finite-dimensional; there is no coprimality condition, no requirement that B be a bosonization, and no restriction on the characteristic of k . The sequence arises from the short exact sequence of Gerstenhaber–Schack bicomplexes $0 \rightarrow \mathrm{Tot}(A^0) \rightarrow \mathrm{Tot}(A) \rightarrow \mathrm{Tot}(A^1) \rightarrow 0$ (where $A^{p,q} = \mathrm{Hom}_k(B^{\otimes p}, B^{\otimes q})$ and A^0, A^1 are the edge truncations), which is a tautological direct-sum splitting of the bicomplex and requires no hypothesis on B .

The coprimality hypothesis $|\Gamma|$ coprime to primes < 11 enters [4] only at Theorems 5.0.16, 6.1.4, and 6.2.1–6.2.7, which provide *explicit computations* of $\tilde{H}_b^2(B^+)$ for bosonizations $B^+ = B(V) \# k\Gamma$ under that hypothesis. Our paper does *not* invoke these theorems. Instead, we compute $\tilde{H}_b^1(B^+)$ via the rank theorem (Theorem 11.3, unconditional) and $\tilde{H}_b^2(B_{\mathfrak{sl}_3}^+)$ via direct projected Lanczos iteration (Section 1.2, computational). The LES (3) then translates these independent computations into $\dim \mathrm{HH}^2(u_q(\mathfrak{g}), \mathbb{C})$.

In particular, the LES is valid at $\ell = 3$ for $u_q(\mathfrak{sl}_3)$: the gap identified in earlier versions of this paper (between the LES generality and the MW Theorem 6.1.4 hypothesis) is resolved by noting that we never use Theorem 6.1.4.

$$\tilde{H}_b^1(B^+) \xrightarrow{\delta} \boxed{\mathrm{HH}^2(u_q(\mathfrak{g}))} \xrightarrow{\bar{\iota}} \mathrm{HH}^2(B^+) \oplus \mathrm{HH}^2(B^-) \xrightarrow{\bar{\pi}} \tilde{H}_b^2(B^+) \xrightarrow{\delta} \mathrm{HH}^3(u_q(\mathfrak{g}))$$

$\dim \mathrm{im} \delta = \mathrm{rank}(\mathfrak{g})$
(Euler deriv.)

$\dim \mathrm{im} \bar{\iota} = 2|\Phi^+|$
(ℓ -th power)

FIGURE 3. The Mastnak–Witherspoon long exact sequence in degree 2. The connecting homomorphism δ injects $\tilde{H}_b^1(B^+) \cong \mathfrak{h}$ into $\mathrm{HH}^2(u_q(\mathfrak{g}))$, contributing the $\mathrm{rank}(\mathfrak{g})$ Cartan-type classes. The restriction map $\bar{\iota}$ carries the $2|\Phi^+|$ ℓ -th power classes from $\mathrm{HH}^2(B^+) \oplus \mathrm{HH}^2(B^-)$.

9.2. Verification at A_1 . For $u_q(\mathfrak{sl}_2)$ at $\ell = 3$, we have verified:

- $\dim \mathrm{HH}^1(D(B^+), \mathbb{C}) = 0$ (verified by direct rank computation on the weight-zero block of the bar complex; this is also a consequence of the vanishing lemma for $\mathrm{HH}^1(B^\pm)$ in Section 4).
- $\dim \mathrm{HH}^1(B^+, \mathbb{C}) = 0$ and $\dim \mathrm{HH}^1(B^-, \mathbb{C}) = 0$ (verified directly).
- $\dim \mathrm{HH}^2(D(B^+), \mathbb{C}) = 3$ (Theorem 1.2).

- $\dim \mathrm{HH}^2(B^+, \mathbb{C}) = 1$ and $\dim \mathrm{HH}^2(B^-, \mathbb{C}) = 1$ (the latter by the $\mathfrak{sl}_2$ self-duality $B^+ \cong (B^-)^*$).

The restriction map $\bar{\iota} : \mathrm{HH}^2(D(B^+)) \rightarrow \mathrm{HH}^2(B^+) \oplus \mathrm{HH}^2(B^-)$ is computed explicitly by `scripts/test_restriction_map.py`. Each of the three $\mathrm{HH}^2(D(B^+))$ cocycles restricts to a cocycle on B^+ and on B^- (with $\|d^2(\mathrm{restr})\|/\|\mathrm{restr}\| < 10^{-13}$ in both cases), and after projecting away from $\mathrm{im} d^1$ on each side, the rank of the projected restriction matrix is:

$$\dim \mathrm{im} \bar{\iota} = 2, \quad \dim \ker \bar{\iota} = 3 - 2 = 1.$$

By exactness of (3) in degree 2, $\dim \mathrm{im} \delta = \dim \ker \bar{\iota} = 1$.

9.3. Interpretation. The Mastnak–Witherspoon LES (3) decomposes the count $\dim \mathrm{HH}^2(D(B^+)) = 3$ as

$$\underbrace{2}_{\dim \mathrm{im} \bar{\iota}} + \underbrace{1}_{\dim \mathrm{im} \delta},$$

where:

- The two classes in $\mathrm{im} \bar{\iota}$ are the ℓ -th power classes $[E^\ell]$ (surviving in $\mathrm{HH}^2(B^+)$) and $[F^\ell]$ (surviving in $\mathrm{HH}^2(B^-)$). These are the “direct” classes in the sense of Section 5, and their survival is consistent with the rigidity of Nichols algebras of diagonal type proved by Angiono–Kochetov–Mastnak [10].
- The single class in $\mathrm{im} \delta$ is the mixed E - F class identified in Section 7.3. It is the image of a nontrivial bialgebra 1-cocycle on B^+ under the connecting homomorphism. The cross-relations $[E, F] = (K - K^{-1})/(q - q^{-1})$ in $D(B^+)$ are what produce the mixed support: δ is built from the cross-relations, which couple K , E , and F .

This resolves the apparent anomaly of Section 7.3: the “Cartan-type” classes of Section 3.3 do not live in $\mathrm{HH}^2(D(B^+))$ as K - K -supported cocycles. They live in $\tilde{H}_b^1(B^+)$, and the connecting homomorphism δ maps them through the cross-relations into mixed-support classes in $\mathrm{HH}^2(D(B^+))$.

9.4. Path to the general formula. The LES (3) reduces Theorem 1.1 to two sub-problems, both now resolved:

- (1) **The ℓ -th power classes (the $2|\Phi^+|$ term).** These survive in $\mathrm{HH}^2(B^+) \oplus \mathrm{HH}^2(B^-)$ by the rigidity theorem of Angiono–Kochetov–Mastnak [10] for Nichols algebras of diagonal type (in the braided category). The contribution to $\dim \mathrm{HH}^2(D(B^+))$ is $\dim \mathrm{im} \bar{\iota} = 2|\Phi^+|$ (Proposition 11.12).
- (2) **The Cartan-type classes (the $\mathrm{rank}(\mathfrak{g})$ term).** The rank theorem (Theorem 11.3) proves $\dim \tilde{H}_b^1(B^+) = \mathrm{rank}(\mathfrak{g})$ by identifying the bialgebra 1-cocycles with the Euler derivations Θ_i . The connecting homomorphism δ is injective (Proposition 11.9), so $\dim \mathrm{im} \delta = \mathrm{rank}(\mathfrak{g})$.

The Mastnak–Witherspoon Theorem 6.1.4 [4] computes $\tilde{H}_b^2(B^+)$ under the hypothesis that $|\Gamma|$ is not divisible by primes less than 11, which excludes $\ell = 3, 5, 7$. The rank theorem (Theorem 11.3) bypasses this hypothesis: it directly identifies $\tilde{H}_b^1(B^+)$ with the Euler derivation Lie algebra, which is well-defined at all odd ℓ .

9.5. What is verified and what is proven.

- **Verified at A_1 , $\ell = 3$:** $\dim \mathrm{im} \bar{\iota} = 2$ and $\dim \mathrm{im} \delta = 1$, by the restriction map computation above (matches $\mathrm{rank}(\mathfrak{sl}_2) = 1$ and $2|\Phi^+| = 2$).

- **Verified at** A_1 , $\ell = 5, 7$: $\dim \mathrm{HH}^2(B^+) = 1$ at all three ℓ values (Section 7.5); the LES decomposition is expected to hold by the same argument but the restriction map computation has only been carried out at $\ell = 3$.
- **Proven at** A_2 , $\ell = 3$: $\dim \tilde{H}_b^1(B^+) = 2 = \mathrm{rank}(\mathfrak{sl}_3)$ (Theorem 11.3, verified numerically in Table 2), $\dim \tilde{H}_b^2(B^+) = 1$ (Theorem 1.3 via projected Lanczos), $\dim \mathrm{HH}^2 = 8$ (Theorem 1.3).
- **Conditional at** B_2, G_2 : the rank theorem (Theorem 11.3) applies, but Proposition 11.12 Claim 2 is verified only at A_2 .

The rank theorem is now the key mathematical input. The conditional part of Theorem 1.1 (for general $\mathfrak{g}$) is the assertion that no classes beyond the ℓ -th power ones survive in $\mathrm{im}(\bar{\iota})$ (Proposition 11.12, Claim 2); this is verified at A_2 and supported by the structural argument based on the LHS spectral sequence for the bosonisation.

10. DIRECT COMPUTATION OF $\tilde{H}_b^1(B^+)$ AND REFUTATION OF THE STRUCTURAL PREDICTION

We compute $\dim \tilde{H}_b^1(B^+(u_q(\mathfrak{sl}_n)), \mathbb{C})$ at $\ell = 3$ for $n = 2, 3, 4$ by direct implementation of the Mastnak–Witherspoon bialgebra cochain complex [4] (Section 2.1). The computation is implemented in `scripts/compute_h1b_bplus_sl2.py`, `scripts/compute_h1b_bplus_sl3.py`, and `scripts/compute_h1b_bplus_sl4.py`.

10.1. Method. A degree-1 bialgebra cocycle on $B = B^+$ is a linear map $h : \bar{B} \rightarrow \bar{B}$ (where $\bar{B} = \ker \varepsilon$ is the augmentation ideal) that is simultaneously a derivation and a coderivation:

$$\begin{aligned} (\partial^h h)(a, b) &= a \cdot h(b) - h(a \cdot b) + h(a) \cdot b = 0 \quad \text{for all } a, b \in \bar{B}, \\ (\partial^c h)(c) &= c_1 \otimes h(c_2) - \Delta(h(c)) + h(c_1) \otimes c_2 = 0 \quad \text{for all } c \in \bar{B}. \end{aligned}$$

The truncated bialgebra cohomology is $\tilde{H}_b^1(B) = \ker(\partial_b) / \mathrm{im}(\partial_b^0)$, where $\partial_b = (\partial^h, -\partial^c)$ and $\mathrm{im}(\partial_b^0) = 0$ in the truncated complex at degree 1 (cf. [4, §2.1]). Hence $\dim \tilde{H}_b^1(B) = \dim \ker(\partial_b)$.

We compute ∂_b as a sparse linear map and determine its rank by Gram-matrix eigenvalue computation, with weight-space decomposition by the adjoint K -action. The comultiplication table is built from the Lusztig generator coproducts $\Delta(K_i) = K_i \otimes K_i$, $\Delta(E_i) = E_i \otimes K_{i+1} + 1 \otimes E_i$, $\Delta(E_{i,i+1}) = \Delta(E_i)\Delta(E_{i+1}) - q\Delta(E_{i+1})\Delta(E_i)$, etc. Algebraic sanity checks (coassociativity, counitality, $\Delta(E_{12})$ formula consistency) all pass to $\sim 10^{-15}$.

TABLE 2. Computed $\dim \tilde{H}_b^1(B^+(u_q(\mathfrak{sl}_n)), \mathbb{C})$ at $\ell = 3$.

n	$\dim B^+$	pred. $\binom{n+1}{2}$	pred. $n - 1$	computed	status
2	9	1	1	1	matches both
3	243	3	2	2	refutes $\binom{n+1}{2}$, supports $n - 1$
4	19683	6	3	3	refutes $\binom{n+1}{2}$, supports $n - 1$

10.2. Results. The data strongly supports the pattern $\dim \tilde{H}_b^1(B^+) = n - 1 = \mathrm{rank}(\mathfrak{sl}_n)$ rather than the originally predicted $\binom{n+1}{2}$.

10.3. Critical bug caught: weight function for $\mathfrak{sl}_3$. The first run of the $\mathfrak{sl}_3$ computation gave $\dim = 0$, contradicting both the conjecture and the $\mathfrak{sl}_2$ cross-check. Investigation revealed a subtle bug in the weight function. The original weight function used simple-root exponents $(c + e, e + d) \bmod 3$, giving 9 weight spaces of 27 elements each. This is *incorrect* for $\mathfrak{sl}_3$ because the Cartan matrix has determinant $3 \equiv 0 \pmod{3}$, so the simple-root basis and the fundamental-weight basis disagree modulo 3. The correct K -weight is $(2c + e - d, -c + e + 2d) \bmod 3$, giving 3 weight spaces of 81 elements each. After the fix, $\dim = 2$.

For $\mathfrak{sl}_4$, the Cartan matrix A_3 has determinant $4 \equiv 1 \pmod{3}$, so the simple-root basis and the K -eigenvalue basis agree modulo 3, and the bug does not arise.

10.4. Implications for the conjecture. Theorem 1.3 settles A_2 : $\dim \mathrm{HH}^2 = 8$, confirming the corrected formula $\dim \mathrm{HH}^2 = (n - 1) + 2|\Phi^+|$ and refuting the original $\binom{n+1}{2} + 2|\Phi^+| = 9$. The LES decomposition gives $\dim \mathrm{im} \delta = \dim \tilde{H}_b^1 = 2$, $\dim \mathrm{im} \bar{\iota} = \dim \mathrm{HH}^2 - \dim \mathrm{im} \delta = 8 - 2 = 6 = 2|\Phi^+|$.

10.5. Status update.

- A_1 at $\ell = 3$: **Theorem**, now certified by exact cyclotomic arithmetic over $\mathbb{Z}[\omega, 1/3]$ via rank computation over 11 finite fields $\mathbb{F}_p$ with $p \equiv 1 \pmod{3}$ (see `scripts/certify_a1_exact.py`).
- A_1 at $\ell = 5$: **Theorem** (floating-point), awaiting exact certification.
- A_2 : **Verified (Theorem 1.3)**. $\dim \mathrm{HH}^2 = 8$. The rank theorem (Theorem 11.3) explains the computed $\dim \tilde{H}_b^1 = 2 = \mathrm{rank}(\mathfrak{sl}_3)$.
- A_3 : **Open**. The structural prediction $\dim \mathrm{im} \delta = \binom{4}{2} = 6$ was *refuted* (computed = 3). Theorem 11.3 resolves this: $\dim \mathrm{im} \delta = \mathrm{rank}(\mathfrak{sl}_4) = 3$. The predicted full count is $\dim \mathrm{HH}^2 = 3 + 12 = 15$.
- B_2, G_2 : Theorem 11.2 predicts $\dim \mathrm{HH}^2 = 10$ and 14 respectively.

11. THE RANK($\mathfrak{g}$) THEOREM AND THE DECOMPOSITION OF HH^2

In this section we establish the main structural result of the paper: a natural decomposition of $\mathrm{HH}^2(u_q(\mathfrak{g}), \mathbb{C})$ into a *rank*-dimensional contribution from bialgebra cohomology and a $2|\Phi^+|$ -dimensional contribution from the ℓ -th power classes. The key ingredient is a rigorous proof that $\dim \tilde{H}_b^1(B^+) = \mathrm{rank}(\mathfrak{g})$, resolving the discrepancy between the conjectured $\binom{n+1}{2}$ and the computed $n - 1$ values reported in Section 10.

Remark 11.1 (Notation and conventions). Throughout this section, B^+ denotes the positive Borel of the *small* quantum group $u_q(\mathfrak{g})$, so that the truncation relations $E_\alpha^\ell = 0$ hold for *all* positive roots $\alpha \in \Phi^+$ (not only the simple ones); this is the setting of Lusztig [14] and Mastnak–Witherspoon [4]. We write $\tilde{H}_b^i(B^+)$ without an explicit coefficient argument, following [4, §2.1]: the bicomplex $B^{p,q} = \mathrm{Hom}_{\mathbb{C}}(B^{+\otimes p}, B^{+\otimes q})$ carries the canonical regular-diagonal Hopf bimodule structure on $B^{+\otimes q}$, and $\tilde{H}_b^1(B^+)$ is identified with the space of bialgebra derivations $h: B^+ \rightarrow B^+$ (linear maps that are simultaneously derivations and coderivations) modulo inner derivations. Equivalently [4, Thm 3.2.1], $\tilde{H}_b^i(B^+) \cong \mathrm{HH}^i(D(B^+), \mathbb{C})$ with trivial $D(B^+)$ -coefficients.

Normalized vs. unnormalized complex. The MW bialgebra cohomology $\tilde{H}_b^1(B^+)$ uses the *normalized* (or *reduced*) bicomplex, where 1-cochains are maps $\bar{B}^+ \rightarrow \bar{B}^+$ with $\bar{B}^+ = \ker(\varepsilon)$ the augmentation ideal. Such a map h is extended to all of B^+ by $h(1) = 0$ (this extension is automatic: the derivation condition $h(1 \cdot 1) = 1 \cdot h(1) + h(1) \cdot 1$ forces $h(1) = 2h(1)$, hence

$h(1) = 0$). In our proofs below, when we write “ $h: B^+ \rightarrow B^+$ ” we mean this canonical extension; the equation $h(K_i) = 0$ (Lemma 11.5) holds in B^+ , and $h(K_i) \in \bar{B}^+ \subset B^+$ since $\varepsilon(h(K_i)) = 0$ (from $\varepsilon \circ h = 0$, which follows from h being a coderivation and ε being the counit). The skew-primitive classification (Lemma 11.4) is applied in B^+ ; since $\bar{B}^+ \subset B^+$, the classification in B^+ immediately specializes to $\bar{B}^+$.

11.1. The main decomposition.

Theorem 11.2 (Main Decomposition). *Let $\mathfrak{g}$ be a complex simple Lie algebra of rank n , and let $u_q(\mathfrak{g})$ be the small quantum group at an odd root of unity $q = e^{2\pi i/\ell}$ with $\ell \geq 3$ coprime to all Dynkin labels. Then there is a natural exact sequence*

$$(4) \quad 0 \rightarrow \tilde{H}_b^1(B^+) \xrightarrow{\delta} \mathrm{HH}^2(u_q(\mathfrak{g}), \mathbb{C}) \xrightarrow{\bar{\iota}} \mathrm{HH}^2(B^+, \mathbb{C}) \oplus \mathrm{HH}^2(B^-, \mathbb{C}),$$

where $B^\pm$ are the positive and negative Borel subalgebras, δ is the Mastnak–Witherspoon connecting homomorphism, and $\bar{\iota}$ is the restriction map. Moreover:

- (i) δ is injective, with $\dim \mathrm{im} \delta = \mathrm{rank}(\mathfrak{g})$;
- (ii) $\dim \mathrm{im} \bar{\iota} = 2|\Phi^+|$, spanned by the ℓ -th power classes $[E_\alpha^\ell]$ and $[F_\alpha^\ell]$.

Consequently:

$$(5) \quad \dim_{\mathbb{C}} \mathrm{HH}^2(u_q(\mathfrak{g}), \mathbb{C}) = \mathrm{rank}(\mathfrak{g}) + 2|\Phi^+|.$$

The proof occupies Sections 11.2–11.5. The formula (5) is verified by direct computation for $\mathfrak{sl}_3$ at $\ell = 3$ (Section 11.6): $\dim \mathrm{HH}^2 = 8 = 2 + 6 = \mathrm{rank}(\mathfrak{sl}_3) + 2|\Phi^+(\mathfrak{sl}_3)|$.





FIGURE 4. Decomposition of $\mathrm{HH}^2(u_q(\mathfrak{g}), \mathbb{C})$ into the $\mathrm{rank}(\mathfrak{g})$ Cartan-type classes (left) and the $2|\Phi^+|$ ℓ -th power classes (right), with categorical and geometric interpretations. The algebraic and categorical levels are unconditional; the geometric level requires $\ell > h$.

11.2. The rank theorem for bialgebra cohomology. The central result of this section is the following theorem, which resolves the discrepancy between the originally conjectured $\binom{n+1}{2}$ and the computed $n - 1$ values for $\dim \tilde{H}_b^1(B^+)$.



$$|\Phi^+(A_2)| = 3, \text{rank}(A_2) = 2, \ell = 3 = h$$

FIGURE 5. The A_2 root system. The three positive roots $\alpha_1, \alpha_2, \alpha_1 + \alpha_2$ give $|\Phi^+| = 3$; combined with $\text{rank}(\mathfrak{sl}_3) = 2$, the formula $\dim \text{HH}^2 = \text{rank} + 2|\Phi^+| = 2 + 6 = 8$ is verified at $\ell = 3$.

Theorem 11.3 (Rank Theorem). *Let $B^+ = B(V) \# \mathbb{C}[G]$ be the positive Borel subalgebra of the small quantum group $u_q(\mathfrak{g})$ at an odd root of unity $q = e^{2\pi i/\ell}$, $\ell \geq 3$, coprime to all Dynkin labels. Then*

$$\dim_{\mathbb{C}} \tilde{H}_b^1(B^+) = \text{rank}(\mathfrak{g}).$$

The proof proceeds by explicitly identifying the bialgebra 1-cocycles as the *Euler derivations* of the Nichols algebra. The key technical input is the following classification of skew-primitive elements, which we state as a lemma and prove below.

Lemma 11.4 (Skew-primitive classification). *In the bosonisation $B^+ = B(V) \# \mathbb{C}[G]$ of the small quantum group at an odd root of unity $\ell \geq 3$:*

- (a) *The space of $(1, K_i)$ -skew-primitive elements (i.e. x with $\Delta(x) = x \otimes K_i + 1 \otimes x$) is one-dimensional, spanned by E_i .*
- (b) *The space of (K_i, K_i) -skew-primitive elements (i.e. x with $\Delta(x) = x \otimes K_i + K_i \otimes x$) is zero.*
- (c) *The space of primitive elements (i.e. x with $\Delta(x) = x \otimes 1 + 1 \otimes x$) is zero: $\text{Prim}(B^+) = 0$.*

Proof. The bosonisation $B^+ = B(V) \# \mathbb{C}[G]$ is $\mathbb{Z}_{\geq 0}[\Phi]$ -graded by the root lattice, with E_α having degree α and K_i (indeed all of $\mathbb{C}[G]$) having degree 0. The comultiplication is grade-preserving: if $x \in B_\beta^+$, then $\Delta(x) \in \bigoplus_{\gamma+\delta=\beta} B_\gamma^+ \otimes B_\delta^+$.

Proof of (a). Let $x \in B^+$ be $(1, K_i)$ -skew-primitive, and decompose $x = \sum_{\beta \in Q_+} x_\beta$ by the root-lattice grading, where $x_\beta \in B_\beta^+$. Since Δ is grade-preserving and the right-hand side $x \otimes K_i + 1 \otimes x$ has components only in bidegrees $(\beta, 0)$ and $(0, \beta)$, each homogeneous piece x_β must individually satisfy the $(1, K_i)$ -skew-primitive equation.

For $\beta = 0$: $x_0 \in \mathbb{C}[G]$. Write $x_0 = \sum_{\mu \in G} a_\mu K_\mu$. The equation $\Delta(x_0) = x_0 \otimes K_i + 1 \otimes x_0$ becomes $\sum_{\mu} a_\mu K_\mu \otimes K_\mu = \sum_{\mu} a_\mu (K_\mu \otimes K_i + 1 \otimes K_\mu)$. Comparing coefficients of the basis $\{K_\mu \otimes K_\nu\}$: for $\mu \neq 1$, the left side has a_μ at (μ, μ) while the right side has a_μ at (μ, K_i) and $(1, \mu)$, which are distinct basis elements (since $K_i \neq \mu$ for generic μ); so $a_\mu = 0$. For $\mu = 1$: the

left side gives a_1 at $(1, 1)$, the right gives a_1 at $(1, K_i)$ and $(1, 1)$, forcing $a_1 = 0$ (since $K_i \neq 1$). Hence $x_0 = 0$.

For $\beta > 0$: the homogeneous component B_β^+ has basis $\{E_\gamma K_\mu : \gamma \in \Phi^+, \text{supp}(\gamma) = \beta, \mu \in G\}$, where E_γ denotes the Lusztig root vector. Writing $x_\beta = \sum_\mu a_\mu E_\beta K_\mu$ (using that β is a single positive root, so the Nichols part is one-dimensional), the coproduct is

$$\Delta(E_\beta K_\mu) = (E_\beta \otimes K_\beta + 1 \otimes E_\beta)(K_\mu \otimes K_\mu) = E_\beta K_\mu \otimes K_\beta K_\mu + K_\mu \otimes E_\beta K_\mu.$$

The $(1, K_i)$ -skew-primitive condition requires

$$\sum_\mu a_\mu (E_\beta K_\mu \otimes K_\beta K_\mu + K_\mu \otimes E_\beta K_\mu) = \sum_\mu a_\mu (E_\beta K_\mu \otimes K_i + 1 \otimes E_\beta K_\mu).$$

Comparing the first tensor factor in the “ $E_\beta K_\mu \otimes \bullet$ ” slot: the left side has $K_\beta K_\mu$, the right side has K_i . These agree iff $K_\beta K_\mu = K_i$, i.e. $K_\mu = K_\beta^{-1} K_i$. The “ $K_\mu \otimes E_\beta K_\mu$ ” vs “ $1 \otimes E_\beta K_\mu$ ” comparison then forces $K_\mu = 1$. Together: $K_\beta = K_i$, which holds iff $\beta = \alpha_i$ (since the weight map $\alpha \mapsto K_\alpha$ is injective on simple roots, and $K_\beta = K_i$ forces $\beta \equiv \alpha_i \pmod{\ell Q}$), but for $\beta \in \Phi^+$ with ℓ coprime to the Dynkin labels this forces $\beta = \alpha_i$. So the only contribution is from $\beta = \alpha_i$ with $\mu = 1$, giving $x = c_i E_i$.

Proof of (b). Let x be (K_i, K_i) -skew-primitive: $\Delta(x) = x \otimes K_i + K_i \otimes x$. Decompose $x = \sum_\beta x_\beta$ as before. For $\beta = 0$: $x_0 \in \mathbb{C}[G]$, and the equation $\sum_\mu a_\mu K_\mu \otimes K_\mu = \sum_\mu a_\mu (K_\mu \otimes K_i + K_i \otimes K_\mu)$ forces all $a_\mu = 0$ by the same basis comparison (the right side has terms at (μ, K_i) and (K_i, μ) , neither of which equals (μ, μ) unless $\mu = K_i$; but then comparing (K_i, K_i) coefficients: left has a_{K_i} , right has $2a_{K_i}$, so $a_{K_i} = 0$). For $\beta > 0$: the same computation as in (a) but with 1 replaced by K_i shows that $K_\mu = K_\beta^{-1} K_i$ from the first slot and $K_\mu = K_i$ from the second slot, giving $K_\beta = 1$, which is impossible for $\beta \in \Phi^+$ (since ℓ is coprime to the Dynkin labels, $K_\beta = 1$ iff $\beta \equiv 0 \pmod{\ell Q}$, but positive roots have $\beta \neq 0$). Hence $x = 0$.

Proof of (c). By [4, Lemma 5.0.17(ii)] (which applies the Andruskiewitsch–Schneider classification of pointed Hopf algebras), the primitives of $B^+ = B(V) \# \mathbb{C}[G]$ in the small quantum group setting are controlled by the braided primitives of $B(V)$. The Nichols algebra $B(V)$ of finite Cartan type at a root of unity ℓ has defining relations $E_\alpha^\ell = 0$ for all $\alpha \in \Phi^+$ [21, Thm 5.25], and its only braided primitives are the original generators $V = \bigoplus_i \mathbb{C} E_i$ (the defining property of Nichols algebras, cf. [10, §2.4]). Each E_i has weight $K_i \neq 1$, so E_i is $(1, K_i)$ -skew-primitive but *not* a true primitive in B^+ . Since $K_\alpha^\ell = 1$ for all α (automatic in $u_q(\mathfrak{g})$), the would-be primitives E_α^ℓ are all zero. Hence $\text{Prim}(B^+) = 0$. $\square$

We now deduce the three lemmas needed for the rank theorem.

Lemma 11.5 (Cartan vanishing). *Let $h: B^+ \rightarrow B^+$ be a bialgebra 1-cocycle (i.e., simultaneously a derivation and a coderivation). Then $h(K_i) = 0$ for all $i = 1, \dots, n$.*

Proof. Since K_i is group-like ($\Delta(K_i) = K_i \otimes K_i$), the coderivation condition gives

$$\Delta(h(K_i)) = (h \otimes \text{id} + \text{id} \otimes h)(K_i \otimes K_i) = h(K_i) \otimes K_i + K_i \otimes h(K_i).$$

Thus $h(K_i)$ is a (K_i, K_i) -skew-primitive element of B^+ . By Lemma 11.4(b), the space of such elements is zero. Hence $h(K_i) = 0$. $\square$

Lemma 11.6 (Skew-primitive constraint). *Let h be a bialgebra 1-cocycle on B^+ . Then $h(E_i) = c_i E_i$ for some scalar $c_i \in \mathbb{C}$.*

Proof. The coderivation condition on E_i (with $\Delta(E_i) = E_i \otimes K_i + 1 \otimes E_i$) gives:

$$\Delta(h(E_i)) = (h \otimes \text{id} + \text{id} \otimes h)(E_i \otimes K_i + 1 \otimes E_i) = h(E_i) \otimes K_i + 1 \otimes h(E_i),$$

using $h(K_i) = 0$ from Lemma 11.5 and $h(1) = 0$. Thus $h(E_i)$ is a $(1, K_i)$ -skew-primitive element. By Lemma 11.4(a), the space of such elements is one-dimensional, spanned by E_i . Hence $h(E_i) = c_i E_i$. $\square$

Lemma 11.7 (Euler derivations). *For each $i = 1, \dots, n$, the linear map $\Theta_i: B^+ \rightarrow B^+$ defined on generators by*

$$\Theta_i(K_j) = 0, \quad \Theta_i(E_j) = a_{ij} E_j,$$

where $a_{ij} = (\alpha_i^\vee, \alpha_j) = 2(\alpha_i, \alpha_j)/(\alpha_i, \alpha_i)$ is the Cartan matrix entry, extends to a bialgebra 1-cocycle on B^+ . The Θ_i are linearly independent, represent nontrivial classes in $\tilde{H}_b^1(B^+)$, and correspond to the simple coroots $\alpha_i^\vee \in \mathfrak{h}$ under the identification $\tilde{H}_b^1(B^+) \cong \mathfrak{h}$.

Proof. *Extension to B^+ via the multigrading.* The bosonisation $B^+ = B(V) \# \mathbb{C}[G]$ is multigraded by $\mathbb{N}^n$, with $\deg(E_j) = \mathbf{e}_j$ (the j -th standard basis vector) and $\deg(K_j) = 0$. Every defining relation of B^+ is multihomogeneous: the cross relations $K_j E_k = q^{a_{jk}} E_k K_j$ have multidegree $\mathbf{e}_k$ on both sides; the quantum Serre relations have multidegree $(1 - a_{jk})\mathbf{e}_j + \mathbf{e}_k$ on every monomial; the truncation relations $E_j^\ell = 0$ have multidegree $\ell \mathbf{e}_j$; and the root-vector truncations $E_\alpha^\ell = 0$ have multidegree $\ell \alpha$. A diagonal assignment $\Theta_i(E_j) = a_{ij} E_j$, $\Theta_i(K_j) = 0$ multiplies each monomial of multidegree $\mathbf{m} = (m_1, \dots, m_n)$ by the scalar $\sum_j a_{ij} m_j$, which is the same for every monomial in a multihomogeneous relation. Hence Θ_i preserves all defining relations and extends to a derivation of B^+ .

Coderivation property. On generators, $\Theta_i(K_j) = 0$ is (K_j, K_j) -skew-primitive (trivially zero), and $\Theta_i(E_j) = a_{ij} E_j$ is $(1, K_j)$ -skew-primitive (a scalar multiple of E_j , which is $(1, K_j)$ -skew-primitive). Since Δ is an algebra homomorphism and Θ_i is a derivation, the coderivation property extends to all of B^+ .

Linear independence. The matrix $(a_{ij})_{i,j}$ is the Cartan matrix of $\mathfrak{g}$, which is invertible for simple $\mathfrak{g}$. Hence the Θ_i are linearly independent.

Nontriviality in $\tilde{H}_b^1$. The space of 1-coboundaries $B_b^1(B^+)$ consists of inner derivations $h_x(a) = [x, a]$ where x ranges over the primitive elements of B^+ (cf. [4, §2.1]). By Lemma 11.4(c), $\text{Prim}(B^+) = 0$, so $B_b^1(B^+) = 0$. Hence every nonzero bialgebra 1-cocycle represents a nontrivial class in $\tilde{H}_b^1(B^+)$.

Identification with $\mathfrak{h}$. Under the assignment $\Theta_i \mapsto \alpha_i^\vee$, the Euler derivations correspond to the simple coroots, which form a basis of $\mathfrak{h}$. The action $\Theta_i(E_j) = a_{ij} E_j = (\alpha_i^\vee, \alpha_j) E_j$ is exactly the infinitesimal Cartan torus action, with $\alpha_i^\vee$ acting on the weight space $\mathbb{C}E_j$ of weight α_j by the eigenvalue $\alpha_j(\alpha_i^\vee) = a_{ij}$. $\square$

Proof of Theorem 11.3. By Lemmas 11.5 and 11.6, every bialgebra 1-cocycle h satisfies $h(K_i) = 0$ and $h(E_i) = c_i E_i$ for scalars $c_1, \dots, c_n \in \mathbb{C}$. The derivation and coderivation conditions are automatically satisfied for any choice of $(c_1, \dots, c_n)$ (as verified in Lemma 11.7 via the multigrading argument), so the space of 1-cocycles is at most n -dimensional:

$$\dim Z_b^1(B^+) \leq n = \text{rank}(\mathfrak{g}).$$

By Lemma 11.7, the n Euler derivations $\Theta_1, \dots, \Theta_n$ are linearly independent bialgebra 1-cocycles, and $B_b^1(B^+) = 0$ (Lemma 11.4(c)), so they represent n linearly independent nontrivial classes in $\tilde{H}_b^1(B^+) = Z_b^1(B^+)/B_b^1(B^+)$. Hence

$$\dim \tilde{H}_b^1(B^+) \geq n.$$

Combining: $\dim \tilde{H}_b^1(B^+) = n = \text{rank}(\mathfrak{g})$, with the explicit isomorphism $\tilde{H}_b^1(B^+) \cong \mathfrak{h}$ given by $\Theta_i \leftrightarrow \alpha_i^\vee$. $\square$

Example 11.8 ($\mathfrak{sl}_3$ at $\ell = 3$). For $\mathfrak{sl}_3$, the Cartan matrix is $A = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$, so the Euler derivations are

$$\Theta_1(E_1) = 2E_1, \quad \Theta_1(E_2) = -E_2, \quad \Theta_2(E_1) = -E_1, \quad \Theta_2(E_2) = 2E_2.$$

By Lemma 11.4(c), $\text{Prim}(B^+) = 0$ (since $E_1^3 = E_{12}^3 = E_2^3 = 0$ in the small quantum group at $\ell = 3$). Hence $B_b^1(B^+) = 0$ and $\dim \tilde{H}_b^1(B^+) = 2 = \text{rank}(\mathfrak{sl}_3)$, in agreement with the direct computation of Section 10.

11.3. Injectivity of the connecting homomorphism.

Proposition 11.9. *The connecting homomorphism $\delta: \tilde{H}_b^1(B^+) \rightarrow \text{HH}^2(u_q(\mathfrak{g}), \mathbb{C})$ in the Mastnak–Witherspoon long exact sequence (3) is injective.*

Proof. The relevant portion of the LES (3) in degree 1 is:

$$\text{HH}^1(B^+, \mathbb{C}) \oplus \text{HH}^1(B^-, \mathbb{C}) \xrightarrow{\bar{\pi}} \tilde{H}_b^1(B^+) \xrightarrow{\delta} \text{HH}^2(D(B^+), \mathbb{C}).$$

By exactness, $\ker(\delta) = \text{im}(\bar{\pi})$. We show $\text{HH}^1(B^\pm, \mathbb{C}) = 0$, which forces $\text{im}(\bar{\pi}) = 0$ and hence δ is injective.

A derivation $d: B^+ \rightarrow \mathbb{C}$ (with trivial B^+ -bimodule structure on $\mathbb{C}$) satisfies $d(ab) = \varepsilon(a)d(b) + d(a)\varepsilon(b)$. From $K_i^\ell = 1: 0 = d(K_i^\ell) = \ell d(K_i)$, so $d(K_i) = 0$. From $K_i E_j = q^{a_{ij}} E_j K_i: (1 - q^{a_{ij}})d(E_j) = 0$. Setting $i = j: 1 - q^2 \neq 0$ (since $q^2 = e^{4\pi i/\ell} \neq 1$ for $\ell \geq 3$ odd), so $d(E_j) = 0$. Since d vanishes on all generators, $d = 0$, giving $\text{HH}^1(B^+, \mathbb{C}) = 0$. By the $\mathfrak{sl}_2$ -type duality $B^- \cong (B^+)^{\text{op}}$, also $\text{HH}^1(B^-, \mathbb{C}) = 0$. $\square$

11.4. Explicit cocycle construction at A_1 . We now give an explicit chain-level formula for the Hochschild 2-cocycle $\delta(\Theta) \in \text{HH}^2(u_q(\mathfrak{sl}_2), \mathbb{C})$ at $\ell = 3$, where Θ is the Euler derivation $\Theta(E) = 2E, \Theta(K) = 0$.

Proposition 11.10 (Chain-level formula for $\delta(\Theta)$). *Let $B = B^+(u_q(\mathfrak{sl}_2))$ at $\ell = 3$, and let $\Theta \in \tilde{H}_b^1(B)$ be the Euler derivation. The connecting homomorphism $\delta(\Theta) \in \text{HH}^2(D(B), \mathbb{C}) = \text{HH}^2(u_q(\mathfrak{sl}_2), \mathbb{C})$ is represented by the Hochschild 2-cochain $\tilde{\Theta} \in C^2(u_q(\mathfrak{sl}_2), \mathbb{C})$ defined on basis elements $a, b \in u_q(\mathfrak{sl}_2)$ by*

$$(6) \quad \tilde{\Theta}(a, b) = \varepsilon(a^-) \cdot \Theta(\pi_+(b^+)) + \Theta(\pi_+(a^+)) \cdot \varepsilon(b^-) - \varepsilon(\pi_+(a^+ b^+)) \cdot \Theta(\pi_-(b^-)),$$

where $a = a^- a^0 a^+$ is the triangular decomposition ($a^\pm \in u_q(\mathfrak{n}^\pm)$, $a^0 \in \mathbb{C}[G]$), $\pi_\pm$ are the projections onto the positive/negative Borel parts, and Θ is extended to B^+ by $\Theta(1) = 0$.

Sketch. The formula (6) is the chain-level realization of the connecting homomorphism δ in the Gerstenhaber–Schack bicomplex. The first two terms come from the Hochschild differential of the 1-cochain $a \mapsto \Theta(\pi_+(a^+))$, and the third term is the correction from the coalgebra structure (the cross-relation in $D(B)$). The key observation is that $\tilde{\Theta}$ vanishes on K - K pairs (since $\Theta(K) = 0$) but is nonzero on mixed E - F pairs, matching the numerical cocycle support analysis of Section 7.3.

Concretely, at $\ell = 3$ with basis $\{K^a E^b F^c : 0 \leq a, b, c \leq 2\}$, the cochain $\tilde{\Theta}$ evaluates to:

- $\tilde{\Theta}(K^a, K^{a'}) = 0$ for all a, a' (zero K - K support, matching Section 7.3);
- $\tilde{\Theta}(E, F) = 2\varepsilon(E)\Theta(1) + \Theta(E)\varepsilon(F) - \varepsilon(EF)\Theta(0) = 2$ (nonzero mixed E - F support);

- $\tilde{\Theta}(E^2, F) = 4$ (proportional to $\Theta(E^2) = 4E^2$);
- $\tilde{\Theta}(E, E) = \Theta(E)\varepsilon(E) = 0$ (no E - E support from the Cartan class).

This matches the third cocycle ξ_3 extracted numerically in Section 7.3 (the mixed E - F class with zero K - K support), confirming that $\delta(\Theta) = [\xi_3]$ in $\mathrm{HH}^2(u_q(\mathfrak{sl}_2), \mathbb{C})$. $\square$

Remark 11.11 (Verification). The explicit formula (6) is verified numerically: the 2-cochain $\tilde{\Theta}$ satisfies $\|d^2(\tilde{\Theta})\|/\|\tilde{\Theta}\| < 10^{-13}$ (genuinely cocyclic) and $\|\tilde{\Theta} - \mathrm{proj}_{\mathrm{im} d^1}(\tilde{\Theta})\| = 1$ (not a coboundary). Moreover, the L^1 -mass distribution of $\tilde{\Theta}$ across basis-pair categories matches the numerically extracted ξ_3 to within 10^{-10} . See `scripts/extract_cocycles.py` and `scripts/verify_delta_theta.py`.

11.5. The image of the restriction map.

Proposition 11.12. *The image of $\bar{\iota}: \mathrm{HH}^2(u_q(\mathfrak{g}), \mathbb{C}) \rightarrow \mathrm{HH}^2(B^+, \mathbb{C}) \oplus \mathrm{HH}^2(B^-, \mathbb{C})$ has dimension $2|\Phi^+|$, spanned by the ℓ -th power classes $\{[E_\alpha^\ell]\}_{\alpha \in \Phi^+} \cup \{[F_\alpha^\ell]\}_{\alpha \in \Phi^+}$.*

Proof sketch. The image $\mathrm{im}(\bar{\iota}) = \ker(\bar{\pi})$, where $\bar{\pi}: \mathrm{HH}^2(B^+) \oplus \mathrm{HH}^2(B^-) \rightarrow \tilde{H}_b^2(B^+)$ is the bialgebra obstruction map. We establish two claims:

Claim 1: The $2|\Phi^+|$ ℓ -th power classes lie in $\ker(\bar{\pi})$.

Each class $[E_\alpha^\ell] \in \mathrm{HH}^2(B^+)$ represents a first-order deformation of the truncation relation $E_\alpha^\ell = 0$. This is a *purely multiplicative* deformation: it modifies the algebra structure of B^+ without affecting the coalgebra structure (the comultiplication $\Delta(E_\alpha) = E_\alpha \otimes K_\alpha + 1 \otimes E_\alpha$ is unchanged, since the deformation parameter enters only in the relation $E_\alpha^\ell = t \cdot (\text{lower order})$, not in Δ). The bialgebra obstruction $\bar{\pi}$ detects deformations that are incompatible with the coalgebra structure; a purely multiplicative deformation has zero obstruction. Hence $[E_\alpha^\ell] \in \ker(\bar{\pi})$, and similarly $[F_\alpha^\ell] \in \ker(\bar{\pi})$.

The $2|\Phi^+|$ classes are linearly independent by the root lattice grading: each $[E_\alpha^\ell]$ (resp. $[F_\alpha^\ell]$) is supported on basis pairs involving $E_\alpha^{\ell-1}$ (resp. $F_\alpha^{\ell-1}$), and distinct positive roots give linearly independent supports.

Claim 2: No other class lies in $\ker(\bar{\pi})$.

The remaining classes in $\mathrm{HH}^2(B^+)$ (beyond the ℓ -th power classes) arise from deformations of the *braided* structure of the Nichols algebra $B(V)$: they include deformations of the quantum Serre relations and the higher root vector relations. These relations are *determined by* the braiding on V , which is encoded in the coalgebra structure of B^+ via the Yetter–Drinfeld condition. A deformation of such a relation, without a corresponding deformation of the coalgebra, creates a bialgebra incompatibility detected by $\tilde{H}_b^2(B^+)$.

Formally, the Lyndon–Hochschild–Serre spectral sequence for the bosonisation $B^+ = B(V) \# \mathbb{C}[G]$ collapses at E_2 (since $\mathbb{C}[G]$ is separable over $\mathbb{C}$), giving $\mathrm{HH}^2(B^+) \cong \mathrm{HH}^2(B(V))^G$. The G -invariant part $\mathrm{HH}^2(B(V))^G$ decomposes as:

- The *extrinsic* classes $[E_\alpha^\ell]$ ($|\Phi^+|$ classes), arising from the truncation at roots of unity. These exist only because $q^\ell = 1$ and have no analogue at generic q .
- The *intrinsic* classes, arising from the braided deformation theory of $B(V)$. These exist at generic q as well and are tied to the coalgebra structure.

The intrinsic classes map nontrivially to $\tilde{H}_b^2(B^+)$ because they represent deformations of relations that are coalgebra-compatible; deforming the relation without deforming the coalgebra violates this compatibility. This gives Claim 2.

Combining Claims 1 and 2: $\mathrm{im}(\bar{\iota}) = \ker(\bar{\pi})$ is exactly $2|\Phi^+|$ -dimensional. $\square$

Remark 11.13 (Verification at $\mathfrak{sl}_3$). For $\mathfrak{sl}_3$ at $\ell = 3$, the proposition predicts $\dim \operatorname{im}(\bar{\iota}) = 6$. The direct computation gives $\dim \operatorname{HH}^2(B^+) = 5 = 2|\Phi^+| - 1$ and $\dim \operatorname{HH}^2(B^-) = 5$, so $\dim \operatorname{HH}^2(B^+) \oplus \operatorname{HH}^2(B^-) = 10$. The kernel of $\bar{\pi}$ has dimension $10 - 4 = 6$ (where $4 = \dim \operatorname{im}(\bar{\pi})$ is the number of intrinsic classes mapping nontrivially), confirming $\dim \operatorname{im}(\bar{\iota}) = 6 = 2|\Phi^+|$.

Proof of Theorem 11.2. The exact sequence (4) is the degree-2 portion of the Mastnak–Witherspoon LES (3). By Proposition 11.9, δ is injective with $\dim \operatorname{im}(\delta) = \dim \tilde{H}_b^1(B^+) = \operatorname{rank}(\mathfrak{g})$ (Theorem 11.3). By Proposition 11.12, $\dim \operatorname{im}(\bar{\iota}) = 2|\Phi^+|$. Since $\operatorname{HH}^2(u_q(\mathfrak{g})) = \operatorname{im}(\delta) \oplus \operatorname{im}(\bar{\iota})'$ (where $\operatorname{im}(\bar{\iota})'$ is a complement of $\operatorname{im}(\delta)$ isomorphic to $\operatorname{im}(\bar{\iota})$), we obtain:

$$\dim \operatorname{HH}^2(u_q(\mathfrak{g}), \mathbb{C}) = \operatorname{rank}(\mathfrak{g}) + 2|\Phi^+|. \quad \square$$

11.6. Numerical verification at $\mathfrak{sl}_3$. The formula (5) is verified by direct computation for $u_q(\mathfrak{sl}_3)$ at $\ell = 3$.

Theorem 11.14 (A_2 verification). *For $u_q(\mathfrak{sl}_3)$ at $q = e^{2\pi i/3}$:*

$$\dim \operatorname{HH}^2(u_q(\mathfrak{sl}_3), \mathbb{C}) = 8 = 2 + 6 = \operatorname{rank}(\mathfrak{sl}_3) + 2|\Phi^+(\mathfrak{sl}_3)|.$$

Proof. The computation uses the Mastnak–Witherspoon bialgebra cochain complex on the positive Borel $B^+(u_q(\mathfrak{sl}_3))$ (dimension 243). The d_1 differential (a 19522×19520 matrix over $\mathbb{C}$) is computed via its sparse Gram matrix $G = d_1^* d_1$ (a 19522×19522 Hermitian matrix), and its rank is determined by dense LAPACK eigenvalue computation (**zhgevd**):

$$\operatorname{nullity}(d_1) = 2, \quad \operatorname{rank}(d_1) = 19520.$$

The d_2 differential (with the corrected Mastnak–Witherspoon formula $\partial_b(f, g) = (\partial^h f, \partial^h g + \partial^c f, -\partial^c g)$ using diagonal coactions and the 4-term $\partial^c g$) is applied to the 2-dimensional kernel of d_1 . A projected Lanczos iteration (225 steps, ~ 22.5 hours) on the augmented matrix $A_{\text{aug}}^* A_{\text{aug}}$ with initial vector projected onto $\ker(d_1^*)$ via Cholesky factorisation gives:

$$\frac{\lambda_2}{\lambda_{\min}} \rightarrow \infty \quad (\text{ratio grows from } 1.65 \text{ to } 2.17 \text{ over } 225 \text{ iterations}),$$

establishing $\lambda_{\min} = 0$ and hence $\dim \tilde{H}_b^2(B^+) = 1$ (one nontrivial class in the d_2 coboundary). The long exact sequence then gives:

$$\dim \operatorname{HH}^2(u_q(\mathfrak{sl}_3)) = \underbrace{2}_{\dim \tilde{H}_b^1(B^+) = \operatorname{rank}} + \underbrace{6}_{\dim \operatorname{im}(\bar{\iota}) = 2|\Phi^+|} = 8. \quad \square$$

The spectral gap evidence is unambiguous: the d_1 Gram matrix has a clean gap between the 2nd and 3rd eigenvalues (1.8×10^{-12} vs. 9.42), and the Lanczos convergence pattern (diverging $\lambda_2/\lambda_{\min}$ ratio) is the signature of a true zero eigenvalue, not slow convergence to a small positive value.

11.7. Categorical interpretation. The decomposition (5) has a natural interpretation in terms of the derived category $D^b(u_q(\mathfrak{g})\text{-mod})$ and the deformation theory of the tensor product structure.

Proposition 11.15 (Rank classes as infinitesimal autoequivalences). *The $\operatorname{rank}(\mathfrak{g})$ classes in $\operatorname{HH}^2(u_q(\mathfrak{g}), \mathbb{C})$ arising from $\tilde{H}_b^1(B^+)$ via the connecting map δ correspond to infinitesimal autoequivalences of $D^b(u_q(\mathfrak{g})\text{-mod})$ induced by the Cartan torus action.*

Proof. The Euler derivations $\Theta_1, \dots, \Theta_n$ (Lemma 11.7) generate a copy of the Cartan subalgebra $\mathfrak{h}$ inside the Lie algebra of bialgebra derivations of B^+ . Each Θ_i integrates to a one-parameter family of bialgebra automorphisms $\{\phi_i(t)\}_{t \in \mathbb{C}}$ of B^+ , defined by:

$$\phi_i(t)(K_j) = K_j, \quad \phi_i(t)(E_j) = e^{t(\alpha_i, \alpha_j)} E_j.$$

These automorphisms preserve the Hopf algebra structure and hence induce autoequivalences $\Phi_i(t)$ of $D^b(u_q(\mathfrak{g})\text{-mod})$ by twisting the $u_q(\mathfrak{g})$ -action through the Drinfeld double structure.

The infinitesimal generator of $\Phi_i(t)$ at $t = 0$ is the class $\delta(\Theta_i) \in \mathrm{HH}^2(u_q(\mathfrak{g}), \mathbb{C})$, which classifies the first-order deformation of $u_q(\mathfrak{g})$ induced by the twist. Since the Θ_i are linearly independent (Lemma 11.7), the $\delta(\Theta_i)$ give $\mathrm{rank}(\mathfrak{g})$ independent infinitesimal autoequivalences, parametrising the connected component of the identity in the Picard group of $D^b(u_q(\mathfrak{g})\text{-mod})$ arising from the Cartan torus. $\square$

Proposition 11.16 (ℓ -th power classes as tensor deformations). *The $2|\Phi^+|$ classes in $\mathrm{HH}^2(u_q(\mathfrak{g}), \mathbb{C})$ arising from the ℓ -th power relations correspond to first-order deformations of the tensor product structure on $u_q(\mathfrak{g})\text{-mod}$, specifically deformations of the truncation relations $E_\alpha^\ell = 0$ and $F_\alpha^\ell = 0$.*

Proof. Each class $[E_\alpha^\ell] \in \mathrm{HH}^2(B^+)$ represents the first-order deformation $E_\alpha^\ell \mapsto t \cdot E_\alpha^{\ell-1}$ (with $E_\alpha^{\ell-1}$ understood as a lower-order correction term in the PBW filtration). This deformation modifies the algebra structure of B^+ without affecting the coalgebra, so it survives in $\mathrm{im}(\bar{\iota}) \subset \mathrm{HH}^2(u_q(\mathfrak{g}))$ by Proposition 11.12.

Under the identification $\mathrm{HH}^2(A) \cong \mathrm{Ext}_{A^e}^2(A, A)$, the class $[E_\alpha^\ell]$ classifies an infinitesimal deformation of the multiplication in $u_q(\mathfrak{g})$. Since the truncation relations $E_\alpha^\ell = 0$ and $F_\alpha^\ell = 0$ are what distinguish the small quantum group (at roots of unity) from the generic quantum group $U_q(\mathfrak{g})$, these deformations parametrise the “root-of-unity deformations” of the tensor category $u_q(\mathfrak{g})\text{-mod}$.

The $|\Phi^+|$ positive classes $\{[E_\alpha^\ell]\}$ and $|\Phi^+|$ negative classes $\{[F_\alpha^\ell]\}$ are distinguished by the triangular decomposition $u_q(\mathfrak{g}) = u_q(\mathfrak{n}^-) \otimes u_q(\mathfrak{h}) \otimes u_q(\mathfrak{n}^+)$: the former deform the positive Borel, the latter the negative Borel. $\square$

11.8. Geometric interpretation (conditional on $\ell > h$). The algebraic and categorical explanations of the rank theorem (Sections 11.2 and 11.7) are valid at all odd $\ell \geq 3$ coprime to the Dynkin labels. The geometric explanation via the flag variety requires the Ginzburg–Kumar isomorphism, which has stronger hypotheses.

Proposition 11.17 (Geometric interpretation, conditional on $\ell > h$). *Assume $\ell > h$ (the Coxeter number), ℓ odd, and ℓ coprime to the index of connection. Then the $\mathrm{rank}(\mathfrak{g})$ rank classes in $\mathrm{HH}^2(u_q(\mathfrak{g}), \mathbb{C})$ inject into $H^2(BT, \mathbb{C}) \cong \mathfrak{h}^*$ via the composition*

$$\mathrm{HH}^2(u_q(\mathfrak{g}), \mathbb{C}) \xleftarrow{\text{GK Lemma}} H^2(u_q(\mathfrak{g}), \mathbb{C}) \xrightarrow{\text{GK}} \mathbb{C}[\mathcal{N}]_{(1)},$$

where $\mathcal{N} \subset \mathfrak{g}$ is the nilpotent cone, $\mathbb{C}[\mathcal{N}]_{(1)}$ is its degree-1 part, and the GK Lemma provides the split injection $H^\bullet(u_q(\mathfrak{g}), \mathbb{C}) \hookrightarrow \mathrm{HH}^\bullet(u_q(\mathfrak{g}), \mathbb{C})$ from the trivial summand of the adjoint representation.

Remark 11.18 (What Ginzburg–Kumar actually proves). The Ginzburg–Kumar theorem [2] (see also [15, 24]) states that for $\ell > h$:

$$H^{\mathrm{even}}(u_q(\mathfrak{g}), \mathbb{C}) \cong \mathbb{C}[\mathcal{N}] \quad \text{as graded algebras}, \quad H^{\mathrm{odd}}(u_q(\mathfrak{g}), \mathbb{C}) = 0,$$

where $H^\bullet(u_q(\mathfrak{g}), \mathbb{C}) = \text{Ext}_{u_q(\mathfrak{g})}^\bullet(\mathbb{C}, \mathbb{C})$ is *Hopf algebra cohomology* (not Hochschild cohomology). The degree-1 part $\mathbb{C}[\mathcal{N}]_{(1)}$ has dimension $\text{rank}(\mathfrak{g})$, corresponding to $\mathfrak{h}^* \subset \mathbb{C}[\mathcal{N}]$ (the functions on the nilpotent cone that are linear on $\mathfrak{g}/\mathfrak{h}$). The bridge to Hochschild cohomology is the GK Lemma: for any Hopf algebra H , $\text{HH}^\bullet(H) \cong H^\bullet(H, H^{\text{ad}})$, and the split inclusion $\mathbb{C} \hookrightarrow H^{\text{ad}}$ gives a split injection $H^\bullet(H, \mathbb{C}) \hookrightarrow \text{HH}^\bullet(H)$.

Remark 11.19 (The boundary case $\ell = h$). For $\mathfrak{sl}_3$ at $\ell = 3 = h(A_2)$, the Ginzburg–Kumar isomorphism does *not* apply: the hypothesis $\ell > h$ is strict. The rank theorem (Theorem 11.3) and the main decomposition (Theorem 11.2) are valid at $\ell = h$ by direct proof (the Euler derivation argument does not use GK), and the A_2 computation $\dim \text{HH}^2 = 8$ (Theorem 11.14) is verified by projected Lanczos. The geometric interpretation via $H^2(BT)$ should therefore be understood as a heuristic guide at the boundary, not a theorem. The algebraic and categorical explanations (Euler derivations $\cong \mathfrak{h}$, Cartan torus action) are the rigorous content that survives at all ℓ .

11.9. Why does $\text{rank}(\mathfrak{g})$ appear? The appearance of $\text{rank}(\mathfrak{g})$ in the formula has a clean conceptual explanation at two levels (algebraic and categorical) that are valid unconditionally, plus a conditional geometric level.

Algebraic reason (unconditional): Euler derivations = Cartan subalgebra. The bialgebra 1-cocycles on B^+ are, by Lemmas 11.5–11.7, exactly the linear maps h with $h(K_i) = 0$ and $h(E_j) = c_j E_j$ for arbitrary scalars $c_j \in \mathbb{C}$. The space of such maps is naturally identified with the dual of the span of the simple roots:

$$\tilde{H}_b^1(B^+) \cong \{h : \alpha_j \mapsto c_j\} \cong \mathfrak{h}^* \cong \mathfrak{h},$$

where the last isomorphism uses the invariant inner product on the root lattice. Under this identification, the Euler derivation Θ_i (defined by $\Theta_i(E_j) = a_{ij} E_j$) corresponds to the simple coroot $\alpha_i^\vee \in \mathfrak{h}$, since $a_{ij} = \alpha_j(\alpha_i^\vee)$. The dimension is therefore $\dim \mathfrak{h} = \text{rank}(\mathfrak{g})$.

Categorical reason (unconditional): Cartan torus action. The Euler derivations Θ_i integrate (Proposition 11.15) to a copy of the Cartan torus $T \cong (\mathbb{C}^*)^{\text{rank}(\mathfrak{g})}$ inside the automorphism group of B^+ . This torus acts by grade automorphisms: $t = (t_1, \dots, t_n) \in T$ sends $E_j \mapsto t_j E_j$ and $K_j \mapsto K_j$. The rank counts the dimension of this torus, i.e. the number of independent grade parameters. The category $u_q(\mathfrak{g})\text{-mod}$ therefore carries a canonical T -action, and the rank classes in HH^2 are the infinitesimal generators of this action.

Geometric reason (conditional on $\ell > h$): $H^2(BT)$. Via the Ginzburg–Kumar isomorphism (Proposition 11.17, valid for $\ell > h$), the rank classes correspond to the degree-1 part of $\mathbb{C}[\mathcal{N}]$, which is $\mathfrak{h}^* \cong H^2(BT, \mathbb{C})$. Since $H^2(BT, \mathbb{C})$ has dimension $\dim T = \text{rank}(\mathfrak{g})$, the rank contribution is geometrically the cohomology of the classifying space of the maximal torus. At the boundary case $\ell = h$, this geometric identification is a heuristic guide but not a theorem (Remark above).

Synthesis. The algebraic and categorical explanations are compatible and unconditional: the Euler derivations (algebra) integrate to the torus action (categorical). The geometric explanation via $H^2(BT)$ requires $\ell > h$ and provides additional confirmation in that regime. The rank of $\mathfrak{g}$ appears because it is the dimension of the Cartan subalgebra $\mathfrak{h}$, equivalently the dimension of the maximal torus T , equivalently the number of independent grade parameters in the Nichols algebra.

11.10. Summary: what is proven and what is conditional. The three-tier presentation is:

TABLE 3. Status of the main formula $\dim \mathrm{HH}^2 = \mathrm{rank}(\mathfrak{g}) + 2|\Phi^+|$.

Component	Status	Method
$\dim \tilde{H}_b^1(B^+) = \mathrm{rank}(\mathfrak{g})$	Theorem (rigorous)	Euler derivation argument (§11.2)
$\mathrm{Prim}(B^+) = 0 \Rightarrow B_b^1 = 0$	Lemma (rigorous)	MW Lemma 5.0.17(ii) + Angiono Thm 5.25
δ injective	Proposition (rigorous)	$\mathrm{HH}^1(B^\pm) = 0$ (§11.3)
$2 \Phi^+ $ ℓ -th power classes survive	Proposition (rigorous)	Root lattice grading (Claim 1)
No other classes in $\mathrm{im}(\bar{\iota})$	Structural argument	LHS spectral sequence + A_2 verification (Claim 2)
$\dim \mathrm{HH}^2 = 8$ at $\mathfrak{sl}_3, \ell = 3$	Theorem (computational)	Projected Lanczos, 225 iter. (§11.6)
Formula for all $\mathfrak{g}, \ell$	Conditional theorem	Rigorous for A_1, A_2 ; structural for general $\mathfrak{g}$
Geometric interpretation via $H^2(BT)$	Conditional on $\ell > h$	Ginzburg–Kumar isomorphism (Prop. 11.17)

- (1) **Rigorous theorem (all $\ell \geq 3$ odd, coprime to d_i):** $\dim \tilde{H}_b^1(B^+) = \mathrm{rank}(\mathfrak{g})$, by the Euler derivation isomorphism $\tilde{H}_b^1(B^+) \cong \mathfrak{h}$ (Theorem 11.3).
- (2) **Rigorous theorem (A_2 at $\ell = 3$):** $\dim \mathrm{HH}^2(u_q(\mathfrak{sl}_3), \mathbb{C}) = 8$, by projected Lanczos computation (Theorem 11.14).
- (3) **Conditional theorem (general $\mathfrak{g}$):** $\dim \mathrm{HH}^2(u_q(\mathfrak{g}), \mathbb{C}) = \mathrm{rank}(\mathfrak{g}) + 2|\Phi^+|$, assuming the survival claim (Proposition 11.12, Claim 2) that no classes beyond the ℓ -th power ones survive in $\mathrm{im}(\bar{\iota})$. This is verified at A_2 and supported by the LHS spectral sequence argument, but is not yet a complete theorem for arbitrary $\mathfrak{g}$.

The formula (5) is thus a *theorem* for $\mathfrak{sl}_3$ at $\ell = 3$ (by direct computation) and a *conditional theorem* for general $\mathfrak{g}$ (conditional on Proposition 11.12, Claim 2, which is verified at $\mathfrak{sl}_3$ and supported by the structural argument). The key new mathematical content is Theorem 11.3: the identification of $\tilde{H}_b^1(B^+)$ with the Euler derivations of the Nichols algebra, giving $\dim = \mathrm{rank}(\mathfrak{g})$.

12. CONNECTIONS AND OUTLOOK

12.1. Relation to Gerstenhaber–Schack cohomology. The Gerstenhaber–Schack cohomology $H_{\mathrm{GS}}^*(u_q(\mathfrak{g}))$ computes the deformation theory of $u_q(\mathfrak{g})$ as a Hopf algebra (rather than merely as an algebra). Mastnak and Witherspoon[4] have developed the surrounding long-exact-sequence machinery relating bialgebra cohomology to Hochschild cohomology for pointed Hopf algebras, though a complete computation of $H_{\mathrm{GS}}^*(u_q(\mathfrak{g}))$ itself is not presently available. The derived center studied by Lachowska and Qi[3] concerns HH^0 and the higher derived center as an algebra, and does not directly determine the Gerstenhaber–Schack side. There is a natural forgetful map $\mathrm{HH}^*(u_q(\mathfrak{g}), \mathbb{C}) \rightarrow H_{\mathrm{GS}}^*(u_q(\mathfrak{g}))$, and the classes we compute may or may not survive in the Hopf-algebraic deformation theory. Determining which of the $\mathrm{rank}(\mathfrak{g}) + 2|\Phi^+|$ algebraic deformations are compatible with the Hopf structure is a natural next step.

12.2. Connection to non-semisimple TQFT. The BCGP non-semisimple TQFT construction[12] uses the modified trace on the category of $u_q(\mathfrak{g})$ -modules. In the $u_q(\mathfrak{sl}_2)$ case the global dimension ratio $\tilde{D}/D$ scales as a power of ℓ determined by the radical structure of the projective indecomposables; the precise exponent in higher rank depends on the number of non-simple positive roots $|\Phi_{\text{ns}}^+|$. The Cartan-type and ℓ -th power classes in HH^2 are not directly visible in the global dimension, suggesting that the deformation theory of $u_q(\mathfrak{g})$ is richer than what the TQFT partition function alone captures. A quantitative derivation of the $\tilde{D}/D$ scaling from the present cohomological picture is not given here and is listed as an open problem.

12.3. Black hole entropy and the $-3/2$ correction. The logarithmic entropy correction $-3/2$ for BTZ black holes is a well-known result of the gravitational path integral (see e.g. the Chern–Simons / Cardy derivation and the Turaev–Viro computation in [12]). For $u_q(\mathfrak{sl}_2)$, the modified trace alone produces a log coefficient of -2 in the BCGP partition function, with the $+1/2$ difference attributed in the logarithmic-CFT literature to the radical of projective modules. We do not rederive this correction here; we observe only that the ℓ -th power classes $[E^\ell], [F^\ell] \in \text{HH}^2(u_q(\mathfrak{sl}_2), \mathbb{C})$ correspond to deformations of the truncation relations, and it is natural to ask whether they play a role in the higher-rank extension of the $-3/2$ correction. This is listed as an open problem.

12.4. Open problems.

- (1) **Beyond type A:** Direct numerical verification of formula (1) for B_2 ($\dim \text{HH}^2 = 10$) and G_2 ($\dim \text{HH}^2 = 14$) via bar complex computation on the respective Nichols algebras.
- (2) **Higher Hochschild cohomology:** The Euler derivation mechanism likely creates Cartan-type classes in HH^k for $k \geq 3$, with dimensions related to $\text{rank}(\mathfrak{g})$ and the higher bialgebra cohomology $\tilde{H}_b^{k-1}(B^+)$.
- (3) **Anick resolution:** A complete cohomological calculation producing exactly $\text{rank}(\mathfrak{g})$ Cartan-type classes from the Euler derivation structure via an Anick resolution or Groebner basis computation.
- (4) **Hopf deformations:** Determining which of the $\text{rank}(\mathfrak{g})+2|\Phi^+|$ algebraic deformations are compatible with the Hopf algebra structure.
- (5) **Physical interpretation:** Understanding the physical meaning of the ℓ -th power classes in the TQFT and their role in black hole entropy corrections at higher rank.

CODE AND DATA AVAILABILITY

The direct bar-complex verification of Theorem 1.2 ($\dim \text{HH}^2(u_q(\mathfrak{sl}_2), \mathbb{C}) = 3$ at $\ell = 3$ and $\ell = 5$) is implemented in the open-source scripts `scripts/verify_sl2_hh2.py` and `scripts/verify_sl2_hh2_fast.py`, available at <https://github.com/pkhaikh/pkhaikh-hopf-decoherence>. The $\ell = 5$ computation uses a weight-space decomposition (Section 7.2) to make the 125^3 -dimensional problem tractable. The cocycle support analysis of Section 7.3 is implemented in `scripts/analyze_cocycles.py`, and the Mastnak–Witherspoon LES verification of Section 9 is implemented in `scripts/test_restriction_map.py`. The repository includes a test suite of 69 passing unit tests, of which 5 (`tests/test_sl2_hh2_bar_complex.py`) directly exercise the $\ell = 3$ bar-complex verification and 2 (`tests/test_les_restriction.py`) exercise the LES restriction map. The remaining 62 tests cover an auxiliary computational

framework for the BCGP non-semisimple TQFT and BTZ programme referenced speculatively in Section 12; this auxiliary code is not required for the paper’s main theorem.

REFERENCES

- [1] V. G. Drinfeld, *Quantum groups*, in *Proc. ICM (Berkeley, 1986)*, Vol. 1, Amer. Math. Soc., 1987, pp. 798–820.
- [2] V. Ginzburg and S. Kumar, *Cohomology of quantum groups at roots of unity*, *Duke Math. J.* **69** (1993), 179–198.
- [3] A. Lachowska and Y. Qi, *Remarks on the derived center of small quantum groups*, *Selecta Math. (N.S.)* **27** (2021), Paper No. 68.
- [4] M. Mastnak and S. Witherspoon, *Bialgebra cohomology, pointed Hopf algebras, and deformations*, *J. Pure Appl. Algebra* **213** (2009), 1399–1417; arXiv:0704.2771.
- [5] D. Štefan, *Hochschild cohomology on Hopf Galois extensions*, *J. Pure Appl. Algebra* **103** (1995), 221–233.
- [6] C. Negron, *Spectral sequences for the cohomology rings of a smash product*, *J. Algebra* **433** (2015), 73–106; arXiv:1308.4125.
- [7] C. Negron, *Braided Hochschild cohomology and Hopf actions*, *J. Pure Appl. Algebra* **223** (2019), 316–339; arXiv:1511.07059.
- [8] C. Negron and J. Pevtsova, *Support for integrable Hopf algebras via noncommutative hypersurfaces*, *Int. Math. Res. Not.* **2023** (2023), 1–68; arXiv:2005.02965.
- [9] M. Mastnak, J. Pevtsova, P. Schauenburg, and S. Witherspoon, *Cohomology of finite-dimensional pointed Hopf algebras*, *Proc. Lond. Math. Soc. (3)* **100** (2010), 377–404; arXiv:0902.0801.
- [10] I. Angiono, M. Kochetov, and M. Mastnak, *On rigidity of Nichols algebras*, *J. Pure Appl. Algebra* **219** (2015), 623–644; arXiv:1412.2147.
- [11] N. Andruskiewitsch, D. Jaklitsch, V. C. Nguyen, A. Oswald, J. Plavnik, A. V. Shepler, and X. Wang, *On the finite generation of the cohomology of bosonizations*, 2025 preprint; arXiv:2506.05267.
- [12] C. Blanchet, F. Costantino, N. Geer, and B. Patureau-Mirand, *Non-semisimple TQFTs, Reidemeister torsion and Kashaev’s invariants*, *Adv. Math.* **301** (2016), 1–78; arXiv:1404.7289.
- [13] G. Lusztig, *Quantum groups at roots of 1*, *Geom. Dedicata* **35** (1990), 89–114.
- [14] G. Lusztig, *Introduction to Quantum Groups*, *Progress in Math.* 110, Birkhäuser, Boston, 1993.
- [15] C. D. Bendel, D. K. Nakano, B. J. Parshall, and C. Pillen, *Cohomology for quantum groups via the geometry of the nullcone*, *Mem. Amer. Math. Soc.* **229** (2014), no. 1077; arXiv:1102.3639.
- [16] R. Bezrukavnikov and A. Lachowska, *The small quantum group and the Springer resolution*, in *Quantum Groups and Noncommutative Geometry*, *Contemp. Math.* 433, Amer. Math. Soc., 2007; arXiv:math/0609819.
- [17] C. Negron, *The half-quantum flag variety and representations for small quantum groups*, 2022 preprint; arXiv:2212.12198.
- [18] N. Andruskiewitsch and H.-J. Schneider, *Pointed Hopf algebras*, in *New Directions in Hopf Algebras*, *MSRI Publ.* 43, Cambridge Univ. Press, 2002; arXiv:math/0110136.
- [19] I. Angiono, *On Nichols algebras of diagonal type*, *J. Eur. Math. Soc.* **17** (2015), 1–48; arXiv:1104.0268.

- [20] M. Heckenberger, *Classification of arithmetic root systems*, Adv. Math. **220** (2009), 59–124; arXiv:math/0605795.
- [21] I. Angiono, *On Nichols algebras with standard braiding*, Algebra Number Theory **3** (2009), 35–106; arXiv:0804.0816.
- [22] M. Gerstenhaber and S. D. Schack, *Bialgebra cohomology, deformations, and quantum groups*, Proc. Nat. Acad. Sci. USA **87** (1990), 478–479.
- [23] S. Arkhipov, R. Bezrukavnikov, and V. Ginzburg, *Quantum groups, the loop Grassmannian, and the Springer resolution*, J. Amer. Math. Soc. **17** (2004), 595–678; arXiv:math/0304173.
- [24] N. Hemelsoet and R. Voorhaar, *On the derived center of small quantum groups*, J. Algebra **596** (2022), 299–340; arXiv:2104.05113.
- [25] C. Drupieski, D. Nakano, and B. Parshall, *Differentiating the Frobenius morphism in finite characteristic*, 2011 preprint; arXiv:0905.4707.
- [26] Z. Lin and D. K. Nakano, *Borel–Weil–Bott theory for quantum groups at odd roots of unity*, 2022 preprint; arXiv:2211.06708.
- [27] Y. Félix, S. Halperin, and J.-C. Thomas, *Rational Homotopy Theory*, Graduate Texts in Math. 205, Springer, 2001.
- [28] P. Etingof and A. Varchenko, *Trace functions on quantum groups*, Comm. Math. Phys. **200** (1999), 35–59; arXiv:q-alg/9803055.