FCI Questions

Patryk Kozlowski

April 21, 2023

1 two differences

$$\langle \Psi | V | \Psi(k \rightarrow k', l \rightarrow l') \rangle = v^{\alpha\beta\gamma\delta} \, (-1)^{\varepsilon\left(\kappa_1,...,\kappa_i',...,\kappa_j,...,\kappa_n\right)} \tag{1}$$

$$\langle 0 | \left(\prod_{\kappa=(\kappa_n...\kappa_3)} a_\kappa \right) a_2 a_1 a_\alpha^\dagger a_\beta^\dagger a_\gamma a_\delta a_{1'}^\dagger a_{2'}^\dagger \left(\prod_{\kappa'=(\kappa_3...\kappa_n)} a_{\kappa'}^\dagger \right) | 0 \rangle \tag{2}$$

$$= \langle 0 | \left(\prod_{\kappa=(\kappa_n...\kappa_3)} a_\kappa \right) a_2 a_1 a_\alpha^\dagger a_\beta^\dagger a_\gamma \delta_{\delta\kappa_{1'}} a_{2'}^\dagger \left(\prod_{\kappa'=(\kappa_3...\kappa_n)} a_{\kappa'}^\dagger \right) | 0 \rangle \tag{3}$$

$$- \langle 0 | \left(\prod_{\kappa=(\kappa_n...\kappa_3)} a_\kappa \right) a_2 a_1 a_\alpha^\dagger a_\beta^\dagger a_\gamma a_{1'}^\dagger a_\delta a_{2'}^\dagger \left(\prod_{\kappa'=(\kappa_3...\kappa_n)} a_{\kappa'}^\dagger \right) | 0 \rangle \tag{4}$$

$$= \langle 0 | \left(\prod_{\kappa=(\kappa_n...\kappa_3)} a_\kappa \right) a_2 a_1 a_\alpha^\dagger a_\beta^\dagger \delta_{\gamma\kappa_{2'}} \delta_{\delta\kappa_{1'}} \left(\prod_{\kappa'=(\kappa_3...\kappa_n)} a_{\kappa'}^\dagger \right) | 0 \rangle \tag{5}$$

$$-0 \tag{6}$$

$$- \langle 0 | \left(\prod_{\kappa=(\kappa_n...\kappa_3)} a_\kappa \right) a_2 a_1 a_\alpha^\dagger a_\beta^\dagger \delta_{\gamma\kappa_{1'}} a_\delta a_{2'}^\dagger \left(\prod_{\kappa'=(\kappa_3...\kappa_n)} a_{\kappa'}^\dagger \right) | 0 \rangle \tag{7}$$

$$+0 \tag{8}$$

$$= \delta_{\alpha\kappa_1} \delta_{\beta\kappa_2} \delta_{\gamma\kappa_{2'}} \delta_{\delta\kappa_{1'}} \tag{9}$$

$$- \delta_{\alpha\kappa_1} \delta_{\beta\kappa_2} \delta_{\gamma\kappa_{1'}} \delta_{\delta\kappa_{2'}} \tag{10}$$

$$= v^{122'1'} - v^{121'2'} \tag{11}$$