

# The Bethe-Salpeter Equation (BSE)

Patryk Kozlowski

Caltech

June 3, 2024

# Linear Response Theory

- ▶ **Random Phase Approximation** tells us that<sup>1</sup>

$$\hat{O}^\dagger = \sum_{i,a} a^\dagger_i X_{ia} + \sum_{i,a} i^\dagger_a Y_{ia}, \quad (1)$$

which counts all of the excitations  $X_{ia}$  and de-excitations  $Y_{ia}$  for holes  $i$  and electrons  $a$ .

# Linear Response Theory

- ▶ **Random Phase Approximation** tells us that<sup>1</sup>

$$\hat{O}^\dagger = \sum_{i,a} a^\dagger i X_{ia} + \sum_{i,a} i^\dagger a Y_{ia}, \quad (1)$$

which counts all of the excitations  $X_{ia}$  and de-excitations  $Y_{ia}$  for holes  $i$  and electrons  $a$ .

- ▶ **Problem:** only describes the linear response of single-particle transitions. It does not work well for excitons, where a hole is interacting with an electron, *after* the transition has taken place

# Linear Response Theory

- ▶ **Random Phase Approximation** tells us that<sup>1</sup>

$$\hat{O}^\dagger = \sum_{i,a} a^\dagger i X_{ia} + \sum_{i,a} i^\dagger a Y_{ia}, \quad (1)$$

which counts all of the excitations  $X_{ia}$  and de-excitations  $Y_{ia}$  for holes  $i$  and electrons  $a$ .

- ▶ **Problem:** only describes the linear response of single-particle transitions. It does not work well for excitons, where a hole is interacting with an electron, *after* the transition has taken place
- ▶ **Solution:** BSE within the Green's function *formalism*

# Green's Functions

The single-particle Green's function  $\mathcal{D}$  is given as

$$\mathcal{D}(\mathbf{r}_1, t_1; \mathbf{r}_2, t_2) = -i \left\langle \Psi_0 \left| T \left[ \hat{\psi}(\mathbf{r}_1, t_1) \hat{\psi}^\dagger(\mathbf{r}_2, t_2) \right] \right| \Psi_0 \right\rangle, \quad (2)$$

where  $\hat{\psi}$  is the second quantization field operator, the spacetime coordinates are  $\mathbf{r}_1, t_1$  and  $\mathbf{r}_2, t_2$ ,  $T$  is the time-ordering operator, and  $\Psi_0$  is the ground state wave function.

# Green's Functions

The single-particle Green's function  $\mathcal{D}$  is given as

$$\mathcal{D}(\mathbf{r}_1, t_1; \mathbf{r}_2, t_2) = -i \langle \Psi_0 | T [\hat{\psi}(\mathbf{r}_1, t_1) \hat{\psi}^\dagger(\mathbf{r}_2, t_2)] | \Psi_0 \rangle, \quad (2)$$


where  $\hat{\psi}$  is the second quantization field operator, the spacetime coordinates are  $\mathbf{r}_1, t_1$  and  $\mathbf{r}_2, t_2$ ,  $T$  is the time-ordering operator, and  $\Psi_0$  is the ground state wave function.

**Distinction:** Noninteracting  $G_0$  (think mean field i.e. DFT) versus interacting  $G$  Green's functions


# Self-Energy

Dyson equation<sup>2</sup>:  $\Sigma = G^{-1} - G_0^{-1}$ , where  $\Sigma$  is the self-energy.

Figure: Electron gas propagation<sup>3</sup>



(a) The electron is shot into the gas



(b) The electron creates holes as it moves along



Analogous to

$$\epsilon_{\text{self}} = \epsilon_{\text{quasi}} - \epsilon_{\text{bare}}, \quad (3)$$

So  $\Sigma$  is the difference between the quasi-particle and bare energies; an electron's "clothing."

# GW Approximation



Hedin's equations<sup>4</sup> define a self-consistent procedure, schematically<sup>5</sup> as





# GW Approximation


Hedin's equations<sup>4</sup> define a self-consistent procedure, schematically<sup>5</sup> as



Where is the vertex function  $\Gamma$  neglected?

# GW Approximation

Hedin's equations<sup>4</sup> define a self-consistent procedure, schematically<sup>5</sup> as



Where is the vertex function  $\Gamma$  neglected?

Right diagram constitutes the GW approximation, whereas for the BSE we need to consider the left diagram.

# Bethe-Salpeter Equation

- ▶ Why was linear response theory unable to describe excitons?

# Bethe-Salpeter Equation

- ▶ Why was linear response theory unable to describe excitons?
- ▶ Can define an electron-hole correlation function given by

$$L(1, 2; 1', 2') = -G_2(1, 2; 1', 2') + G(1, 1') G(2, 2') \quad (4)$$

where we have defined a two-particle Green's function  $G_2$ .

# Bethe-Salpeter Equation

- ▶ Why was linear response theory unable to describe excitons?
- ▶ Can define an electron-hole correlation function given by

$$L(1, 2; 1', 2') = -G_2(1, 2; 1', 2') + G(1, 1') G(2, 2') \quad (4)$$

where we have defined a two-particle Green's function  $G_2$ .

- ▶ Now we are able to describe the interaction between an electron and a hole because this is inherently a two-particle process.

## Bibliography

1. Casida, M. E. & Huix-Rotllant, M. Progress in Time-Dependent Density-Functional Theory. en. *Annual Review of Physical Chemistry* **63**. Publisher: Annual Reviews, 287–323. ISSN: 0066-426X, 1545-1593.  
<https://www.annualreviews.org/content/journals/10.1146/annurev-physchem-032511-143803> (2024) (May 2012).
2. Dyson, F. J. The  $SSS$  Matrix in Quantum Electrodynamics. *Physical Review* **75**. Publisher: American Physical Society, 1736–1755.  
<https://link.aps.org/doi/10.1103/PhysRev.75.1736> (June 1949).
3. Mattuck, R. *A Guide to Feynman Diagrams in the Many-body Problem*. ISBN: 978-0-486-67047-8.  
<https://books.google.com/books?id=pe-v8zfxE68C> (Dover Publications, 1992).
4. Hedin, L. New Method for Calculating the One-Particle