

Slides for Patryk's Notes

Patryk Kozlowski

January 30, 2026

1. How to do larger UEG supercell
2. PT2 variants

1. How to do larger UEG supercell

2. PT2 variants

How to do larger UEG supercell

- python vs. cpp
- using Choleskies with $G=0$ included to implement my methods

1. How to do larger UEG supercell
2. PT2 variants

Review of what I do with the PT2+C spectral functions

I presented two varieties of cumulants. The Landau form is

$$C_{pp}(t) \equiv i \int \frac{d\omega}{2\pi} \frac{\Sigma_{pp}(\omega + E_p)}{(\omega + i\eta)^2} e^{-i\omega t} \quad (1)$$

and the time domain Green's function in the cumulant ansatz is

$$G_{pp}(t) = \underbrace{-i\Theta(t)e^{-t(iE_p + \eta)}}_{\mathbf{G}_0} e^{C_{pp}(t)} \quad (2)$$

In 'G₀ via HF', $E_p \equiv \epsilon_p^{HF}$, $\{\epsilon_p^{HF}\} \rightarrow \Sigma_{pp}(\omega)$, while in 'G₀ via SRG', $E_p \equiv \epsilon_p^{SRG-MP2}$, $\{\epsilon_p^{SRG-MP2}\} \rightarrow \Sigma_{pp}(\omega)$. I understand this through

$$\mathbf{G}(\omega) = \mathbf{G}_0(\omega) + \mathbf{G}_0(\omega)\mathbf{\Sigma}(\omega)\mathbf{G}(\omega) \quad (3)$$

so in 1 and 2, the former approach sets $\mathbf{G}_0 \equiv \mathbf{G}_{HF}$ and the latter sets $\mathbf{G}_0 \equiv \mathbf{G}_{SRG-MP2}$. *Loos found that for small molecules, SRG-GW+C can be worse than G₀W₀+C.*

How SRG-MP2 comes about

Derivation can be done, but the final expression is most relevant:

$$\begin{aligned} F_{pq}^{(2)}(s) = & \frac{1}{2} \sum_{ija} \frac{\Delta_{ij}^{pa} + \Delta_{ij}^{qa}}{\left(\Delta_{ij}^{pa}\right)^2 + \left(\Delta_{ij}^{qa}\right)^2} \langle pa|ij\rangle \langle ij|qa\rangle \\ & \times \left[1 - e^{-\left[\left(\Delta_{ij}^{pa}\right)^2 + \left(\Delta_{ij}^{qa}\right)^2\right]s} \right] \\ & + \frac{1}{2} \sum_{abi} \frac{\Delta_{ab}^{pi} + \Delta_{ab}^{qi}}{\left(\Delta_{ab}^{pi}\right)^2 + \left(\Delta_{ab}^{qi}\right)^2} \langle pi|ab\rangle \langle ab|qi\rangle \\ & \times \left[1 - e^{-\left[\left(\Delta_{ab}^{pi}\right)^2 + \left(\Delta_{ab}^{qi}\right)^2\right]s} \right]. \end{aligned} \tag{4}$$

The idea is to add this to your HF Fock matrix and iterate to self-consistency. Tolle found that it is optimal to choose $s = 1e3$ in his IP/EA tests, so that's what I use.

Nooijen's approach

MP2 tells us that we have to diagonalize:

$$A_{pq}^{\text{MP2}}(\omega) = \epsilon_p \delta_{pq} + \frac{1}{2} \sum_{l,c,d} \frac{V_{pl}[cd] V_{cd}[ql]}{\omega + \epsilon_l - \epsilon_c - \epsilon_d} - \frac{1}{2} \sum_{k,l,d} \frac{V_{pd}[kl] V_{kl}[qd]}{\epsilon_k + \epsilon_l - \epsilon_d - \omega} \quad (5)$$

Nooijen starts with CCSD-GF, but then approximates CC amplitudes with MBPT(2) analogues as

$$t_i^a = 0$$
$$t_{ij}^{ab} = \frac{V_{abij}}{\epsilon_i + \epsilon_j - \epsilon_a - \epsilon_b}$$

They call this MBPT(2)-GF and it leads to $A^{\text{MBPT}(2)}(\omega)$. The paper is not about this theory though, but rather they use it as a starting point for further approximations that reduce cost.

Starting point for DSO-GF

They first replace the most demanding 2h1p block of $A^{\text{MBPT}(2)}(\omega)$ by a diagonal form, but this is not size-consistent. But it is when the diagonal is approximated with a MP partitioning as

$$^{\text{DSO}}D_{ij}^b = -\epsilon_i - \epsilon_j + \epsilon_b \quad (10)$$

leading to

$$A_{ij}^{\text{DSO-GF}}(\omega) = U_{ij} - \frac{1}{2} \sum_{k,l,d} \frac{W_{id}[kl] W_{kl}[jd]}{\epsilon_k + \epsilon_l - \epsilon_d - \omega} \quad (6)$$

$$= \epsilon_i \delta_{ij} + \sum_{c,d,l} V_{ilcd} \left(2t_{jl}^{cd} - t_{jl}^{dc} \right) - \frac{1}{2} \sum_{k,l,d} \frac{W_{id}[kl] V_{kl}[jd]}{\epsilon_k + \epsilon_l - \epsilon_d - \omega} \quad (7)$$

$$= \epsilon_i \delta_{ij} + \frac{1}{2} \sum_{l,c,d} \frac{V_{il}[cd] V_{cd}[jl]}{\epsilon_j + \epsilon_l - \epsilon_c - \epsilon_d} - \frac{1}{2} \sum_{k,l,d} \frac{W_{id}[kl] V_{kl}[jd]}{\epsilon_k + \epsilon_l - \epsilon_d - \omega} \quad (8)$$

Approximations they propose

The intermediates which are used to get 8 are defined as

$$U_{ki} = \epsilon_i \delta_{ki} + \sum_{c,d,l} V_{klcd} \left(2t_{il}^{cd} - t_{il}^{dc} \right) \quad (9)$$

$$W_{klid} = V_{klid} \quad (10)$$

$$\begin{aligned} W_{kbij} = & V_{kbij} + \sum_{l,d} V_{klid} \left(2t_{lj}^{db} - t_{jl}^{db} \right) - \sum_{l,d} V_{lkid} t_{lj}^{db} \\ & + \sum_{c,d} V_{bkdc} t_{ji}^{dc} - \sum_{l,c} V_{lkjc} t_{li}^{bc} \end{aligned} \quad (11)$$

As can be seen, 8 has a $W_{id[kl]}$ remaining, which is obtained via 11. They try to approximate this by just the leading term $V_{id[kl]}$ to reduce cost; this is m-DSO-GF in the paper. But their data suggests keeping the full $W_{id[kl]}$ in 8 leads to good results.

- Would it make sense to do self consistency in Nooijen's schemes?
- Would a paper on the full hierarchy of PT2 methods for one-shot cumulant be useful; we have
 - MP2
 - SRG-MP2
 - DSO-GF
 - m-DSO-GF
 - MBPT(2)-GF
 - CCSD-GF
- Is it useful to study UEG for intermediate supercell sizes?