

# A Brief Review of Probability

Patrick Lam

# Outline

Expectation, Variance, and Densities

Important Distributions

Discrete Distributions

Continuous Distributions

# Outline

Expectation, Variance, and Densities

Important Distributions

Discrete Distributions

Continuous Distributions

# Expectation

# Expectation

The expected value of a random variable  $X$  is simply the weighted average of all possible values of  $X$ .

# Expectation

The expected value of a random variable  $X$  is simply the weighted average of all possible values of  $X$ .

Discrete Case:

# Expectation

The expected value of a random variable  $X$  is simply the weighted average of all possible values of  $X$ .

Discrete Case:

$$E(X) = \sum_i x_i P(X = x_i)$$

where  $P(X = x)$  is the probability mass function (PMF).

# Expectation

The expected value of a random variable  $X$  is simply the weighted average of all possible values of  $X$ .

Discrete Case:

$$E(X) = \sum_i x_i P(X = x_i)$$

where  $P(X = x)$  is the probability mass function (PMF).

Continuous Case:



# Expectation

The expected value of a random variable  $X$  is simply the weighted average of all possible values of  $X$ .

Discrete Case:

$$E(X) = \sum_i x_i P(X = x_i)$$

where  $P(X = x)$  is the probability mass function (PMF).

Continuous Case:

$$E(X) = \int_{-\infty}^{\infty} xp(x)dx$$

where  $p(x)$  is the probability density function (PDF).

# Expectation of a Function of a Random Variable

# Expectation of a Function of a Random Variable

Suppose we want to find  $E[g(X)]$ , where  $g(X)$  is any function of  $X$ .

# Expectation of a Function of a Random Variable

Suppose we want to find  $E[g(X)]$ , where  $g(X)$  is any function of  $X$ . We can simply weight the values of  $g(x)$  by the PDF or PMF of  $X$ :

# Expectation of a Function of a Random Variable

Suppose we want to find  $E[g(X)]$ , where  $g(X)$  is any function of  $X$ . We can simply weight the values of  $g(x)$  by the PDF or PMF of  $X$ :

$$E[g(X)] = \sum_i g(x_i)P(X = x_i)$$

for discrete random variables

# Expectation of a Function of a Random Variable

Suppose we want to find  $E[g(X)]$ , where  $g(X)$  is any function of  $X$ . We can simply weight the values of  $g(x)$  by the PDF or PMF of  $X$ :

$$E[g(X)] = \sum_i g(x_i)P(X = x_i)$$

for discrete random variables and

$$E[g(X)] = \int_{-\infty}^{\infty} g(x)p(x)dx$$

for continuous random variables.

# Expectation of a Function of a Random Variable

Suppose we want to find  $E[g(X)]$ , where  $g(X)$  is any function of  $X$ . We can simply weight the values of  $g(x)$  by the PDF or PMF of  $X$ :

$$E[g(X)] = \sum_i g(x_i)P(X = x_i)$$

for discrete random variables and

$$E[g(X)] = \int_{-\infty}^{\infty} g(x)p(x)dx$$

for continuous random variables.

This is sometimes known as the *Law of the Unconscious Statistician* (LOTUS).

# Variance



# Variance

The formula for the variance of a random variable is

$$\text{Var}(X) = E[(X - E(X))^2]$$

# Variance

The formula for the variance of a random variable is

$$\text{Var}(X) = E[(X - E(X))^2]$$

We can find the variance using LOTUS,

# Variance

The formula for the variance of a random variable is

$$\text{Var}(X) = E[(X - E(X))^2]$$

We can find the variance using LOTUS, or we can simplify the formula first.

# Variance

The formula for the variance of a random variable is

$$\text{Var}(X) = E[(X - E(X))^2]$$

We can find the variance using LOTUS, or we can simplify the formula first.

$$\text{Var}(X) = E[(X - E(X))^2]$$

# Variance

The formula for the variance of a random variable is

$$\text{Var}(X) = E[(X - E(X))^2]$$

We can find the variance using LOTUS, or we can simplify the formula first.

$$\begin{aligned}\text{Var}(X) &= E[(X - E(X))^2] \\ &= E[X^2 - 2XE(X) + (E(X))^2]\end{aligned}$$

# Variance

The formula for the variance of a random variable is

$$\text{Var}(X) = E[(X - E(X))^2]$$

We can find the variance using LOTUS, or we can simplify the formula first.

$$\begin{aligned}\text{Var}(X) &= E[(X - E(X))^2] \\ &= E[X^2 - 2XE(X) + (E(X))^2] \\ &= E(X^2) - 2E(X)E[E(X)] + E([E(X)]^2)\end{aligned}$$

# Variance

The formula for the variance of a random variable is

$$\text{Var}(X) = E[(X - E(X))^2]$$

We can find the variance using LOTUS, or we can simplify the formula first.

$$\begin{aligned}\text{Var}(X) &= E[(X - E(X))^2] \\ &= E[X^2 - 2XE(X) + (E(X))^2] \\ &= E(X^2) - 2E(X)E[E(X)] + E([E(X)]^2) \\ &= E(X^2) - 2[E(X)]^2 + [E(X)]^2\end{aligned}$$

# Variance

The formula for the variance of a random variable is

$$\text{Var}(X) = E[(X - E(X))^2]$$

We can find the variance using LOTUS, or we can simplify the formula first.

$$\begin{aligned}\text{Var}(X) &= E[(X - E(X))^2] \\&= E[X^2 - 2XE(X) + (E(X))^2] \\&= E(X^2) - 2E(X)E[E(X)] + E([E(X)]^2) \\&= E(X^2) - 2[E(X)]^2 + [E(X)]^2 \\&= \mathbf{E(X^2) - [E(X)]^2}\end{aligned}$$



# Variance

The formula for the variance of a random variable is

$$\text{Var}(X) = E[(X - E(X))^2]$$

We can find the variance using LOTUS, or we can simplify the formula first.

$$\begin{aligned}\text{Var}(X) &= E[(X - E(X))^2] \\&= E[X^2 - 2XE(X) + (E(X))^2] \\&= E(X^2) - 2E(X)E[E(X)] + E([E(X)]^2) \\&= E(X^2) - 2[E(X)]^2 + [E(X)]^2 \\&= \mathbf{E(X^2) - [E(X)]^2}\end{aligned}$$

We can then find the first part with LOTUS.

# Marginal, Conditional, and Joint Densities

# Marginal, Conditional, and Joint Densities

$$p(x) = \int p(x, y) dy$$

# Marginal, Conditional, and Joint Densities

$$p(x) = \int p(x, y) dy$$

$$p(x, y) = \int p(x, y, z) dz$$

# Marginal, Conditional, and Joint Densities

$$p(x) = \int p(x, y) dy$$

$$p(x, y) = \int p(x, y, z) dz$$

$$p(x|y) = \frac{p(x, y)}{p(y)}$$

# Marginal, Conditional, and Joint Densities

$$p(x) = \int p(x, y) dy$$

$$p(x, y) = \int p(x, y, z) dz$$

$$p(x|y) = \frac{p(x, y)}{p(y)}$$

$$p(x|y, z) = \frac{p(x, y, z)}{p(y, z)}$$

# Marginal, Conditional, and Joint Densities

$$p(x) = \int p(x, y) dy$$

$$p(x, y) = \int p(x, y, z) dz$$

$$p(x|y) = \frac{p(x, y)}{p(y)}$$

$$p(x|y, z) = \frac{p(x, y, z)}{p(y, z)}$$

$$p(x, y) = p(x|y)p(y)$$

# Marginal, Conditional, and Joint Densities

$$p(x) = \int p(x, y) dy$$

$$p(x, y) = \int p(x, y, z) dz$$

$$p(x|y) = \frac{p(x, y)}{p(y)}$$

$$p(x|y, z) = \frac{p(x, y, z)}{p(y, z)}$$

$$\begin{aligned} p(x, y) &= p(x|y)p(y) \\ &= p(y|x)p(x) \end{aligned}$$



# Marginal, Conditional, and Joint Densities

$$p(x) = \int p(x, y) dy$$

$$p(x, y) = \int p(x, y, z) dz$$

$$p(x|y) = \frac{p(x, y)}{p(y)}$$

$$p(x|y, z) = \frac{p(x, y, z)}{p(y, z)}$$

$$\begin{aligned} p(x, y) &= p(x|y)p(y) \\ &= p(y|x)p(x) \end{aligned}$$

$$p(x, y, z) = p(x|y, z)p(y|z)p(z)$$

# Outline

Expectation, Variance, and Densities

Important Distributions

Discrete Distributions

Continuous Distributions

# Outline

Expectation, Variance, and Densities

Important Distributions

Discrete Distributions

Continuous Distributions

# The Bernoulli Distribution

# The Bernoulli Distribution

$$Y \sim \text{Bernoulli}(\pi)$$

# The Bernoulli Distribution

$$Y \sim \text{Bernoulli}(\pi)$$

$$y = 0, 1$$

# The Bernoulli Distribution

$$Y \sim \text{Bernoulli}(\pi)$$

$$y = 0, 1$$

probability of success:  $\pi \in [0, 1]$

# The Bernoulli Distribution

$$Y \sim \text{Bernoulli}(\pi)$$

$$y = 0, 1$$

probability of success:  $\pi \in [0, 1]$

$$p(y|\pi) = \pi^y(1 - \pi)^{(1-y)}$$



# The Bernoulli Distribution

$$Y \sim \text{Bernoulli}(\pi)$$

$$y = 0, 1$$

probability of success:  $\pi \in [0, 1]$

$$p(y|\pi) = \pi^y(1 - \pi)^{(1-y)}$$

$$E(Y) = \pi$$

# The Bernoulli Distribution

$$Y \sim \text{Bernoulli}(\pi)$$

$$y = 0, 1$$

probability of success:  $\pi \in [0, 1]$

$$p(y|\pi) = \pi^y(1 - \pi)^{(1-y)}$$

$$E(Y) = \pi$$

$$\text{Var}(Y) = \pi(1 - \pi)$$

# The Bernoulli Distribution

$$Y \sim \text{Bernoulli}(\pi)$$


$$y = 0, 1$$

probability of success:  $\pi \in [0, 1]$

$$p(y|\pi) = \pi^y(1 - \pi)^{(1-y)}$$

$$E(Y) = \pi$$

$$\text{Var}(Y) = \pi(1 - \pi)$$



# The Binomial Distribution

# The Binomial Distribution

$$Y \sim \text{Binomial}(n, \pi)$$

# The Binomial Distribution

$$Y \sim \text{Binomial}(n, \pi)$$

$$y = 0, 1, \dots, n$$

# The Binomial Distribution

$$Y \sim \text{Binomial}(n, \pi)$$

$$y = 0, 1, \dots, n$$

number of trials:  $n \in \{1, 2, \dots\}$

# The Binomial Distribution

$$Y \sim \text{Binomial}(n, \pi)$$

$$y = 0, 1, \dots, n$$

number of trials:  $n \in \{1, 2, \dots\}$

probability of success:  $\pi \in [0, 1]$



# The Binomial Distribution

$$Y \sim \text{Binomial}(n, \pi)$$

$$y = 0, 1, \dots, n$$

number of trials:  $n \in \{1, 2, \dots\}$

probability of success:  $\pi \in [0, 1]$

$$p(y|\pi) = \binom{n}{y} \pi^y (1 - \pi)^{(n-y)}$$

# The Binomial Distribution

$$Y \sim \text{Binomial}(n, \pi)$$

$$y = 0, 1, \dots, n$$

number of trials:  $n \in \{1, 2, \dots\}$

probability of success:  $\pi \in [0, 1]$

$$p(y|\pi) = \binom{n}{y} \pi^y (1 - \pi)^{(n-y)}$$

$$E(Y) = n\pi$$

# The Binomial Distribution

$$Y \sim \text{Binomial}(n, \pi)$$

$$y = 0, 1, \dots, n$$

number of trials:  $n \in \{1, 2, \dots\}$

probability of success:  $\pi \in [0, 1]$

$$p(y|\pi) = \binom{n}{y} \pi^y (1 - \pi)^{(n-y)}$$

$$E(Y) = n\pi$$

$$\text{Var}(Y) = n\pi(1 - \pi)$$

# The Binomial Distribution

$$Y \sim \text{Binomial}(n, \pi)$$

$$y = 0, 1, \dots, n$$


number of trials:  $n \in \{1, 2, \dots\}$

probability of success:  $\pi \in [0, 1]$

$$p(y|\pi) = \binom{n}{y} \pi^y (1 - \pi)^{(n-y)}$$

$$E(Y) = n\pi$$

$$\text{Var}(Y) = n\pi(1 - \pi)$$



# The Multinomial Distribution

# The Multinomial Distribution

$$Y \sim \text{Multinomial}(n, \pi_1, \dots, \pi_k)$$

# The Multinomial Distribution

$$Y \sim \text{Multinomial}(n, \pi_1, \dots, \pi_k)$$

$$y_j = 0, 1, \dots, n; \quad \sum_{j=1}^k y_j = n$$

# The Multinomial Distribution

$$Y \sim \text{Multinomial}(n, \pi_1, \dots, \pi_k)$$

$$y_j = 0, 1, \dots, n; \quad \sum_{j=1}^k y_j = n$$

number of trials:  $n \in \{1, 2, \dots\}$



# The Multinomial Distribution

$$Y \sim \text{Multinomial}(n, \pi_1, \dots, \pi_k)$$

$$y_j = 0, 1, \dots, n; \quad \sum_{j=1}^k y_j = n$$

number of trials:  $n \in \{1, 2, \dots\}$

probability of success for  $j$ :  $\pi_j \in [0, 1]; \quad \sum_{j=1}^k \pi_j = 1$

# The Multinomial Distribution

$$Y \sim \text{Multinomial}(n, \pi_1, \dots, \pi_k)$$

$$y_j = 0, 1, \dots, n; \quad \sum_{j=1}^k y_j = n$$

number of trials:  $n \in \{1, 2, \dots\}$

probability of success for  $j$ :  $\pi_j \in [0, 1]$ ;  $\sum_{j=1}^k \pi_j = 1$

$$p(\mathbf{y}|n, \boldsymbol{\pi}) = \frac{n!}{y_1! y_2! \dots y_k!} \pi_1^{y_1} \pi_2^{y_2} \dots \pi_k^{y_k}$$

# The Multinomial Distribution

$$Y \sim \text{Multinomial}(n, \pi_1, \dots, \pi_k)$$

$$y_j = 0, 1, \dots, n; \quad \sum_{j=1}^k y_j = n$$

number of trials:  $n \in \{1, 2, \dots\}$

probability of success for  $j$ :  $\pi_j \in [0, 1]$ ;  $\sum_{j=1}^k \pi_j = 1$

$$p(\mathbf{y}|n, \boldsymbol{\pi}) = \frac{n!}{y_1! y_2! \dots y_k!} \pi_1^{y_1} \pi_2^{y_2} \dots \pi_k^{y_k}$$

$$E(Y_j) = n\pi_j$$

# The Multinomial Distribution

$$Y \sim \text{Multinomial}(n, \pi_1, \dots, \pi_k)$$

$$y_j = 0, 1, \dots, n; \quad \sum_{j=1}^k y_j = n$$

number of trials:  $n \in \{1, 2, \dots\}$

probability of success for  $j$ :  $\pi_j \in [0, 1]$ ;  $\sum_{j=1}^k \pi_j = 1$

$$p(\mathbf{y}|n, \boldsymbol{\pi}) = \frac{n!}{y_1! y_2! \dots y_k!} \pi_1^{y_1} \pi_2^{y_2} \dots \pi_k^{y_k}$$

$$E(Y_j) = n\pi_j$$

$$\text{Var}(Y_j) = n\pi_j(1 - \pi_j)$$

# The Multinomial Distribution

$$Y \sim \text{Multinomial}(n, \pi_1, \dots, \pi_k)$$

$$y_j = 0, 1, \dots, n; \quad \sum_{j=1}^k y_j = n$$

number of trials:  $n \in \{1, 2, \dots\}$

probability of success for  $j$ :  $\pi_j \in [0, 1]$ ;  $\sum_{j=1}^k \pi_j = 1$

$$p(\mathbf{y}|n, \boldsymbol{\pi}) = \frac{n!}{y_1! y_2! \dots y_k!} \pi_1^{y_1} \pi_2^{y_2} \dots \pi_k^{y_k}$$

$$E(Y_j) = n\pi_j$$

$$\text{Var}(Y_j) = n\pi_j(1 - \pi_j)$$

$$\text{Cov}(Y_i, Y_j) = -n\pi_i\pi_j$$

# The Poisson Distribution

# The Poisson Distribution

$$Y \sim \text{Poisson}(\lambda)$$

# The Poisson Distribution

$$Y \sim \text{Poisson}(\lambda)$$

$$y = 0, 1, \dots$$



# The Poisson Distribution

$$Y \sim \text{Poisson}(\lambda)$$

$$y = 0, 1, \dots$$

expected number of  
occurrences:  $\lambda > 0$

# The Poisson Distribution

$$Y \sim \text{Poisson}(\lambda)$$

$$y = 0, 1, \dots$$

expected number of  
occurrences:  $\lambda > 0$

$$p(y|\lambda) = \frac{e^{-\lambda} \lambda^y}{y!}$$

# The Poisson Distribution

$$Y \sim \text{Poisson}(\lambda)$$

$$y = 0, 1, \dots$$

expected number of  
occurrences:  $\lambda > 0$

$$p(y|\lambda) = \frac{e^{-\lambda} \lambda^y}{y!}$$

$$E(Y) = \lambda$$

# The Poisson Distribution

$$Y \sim \text{Poisson}(\lambda)$$

$$y = 0, 1, \dots$$

expected number of  
occurrences:  $\lambda > 0$

$$p(y|\lambda) = \frac{e^{-\lambda} \lambda^y}{y!}$$

$$E(Y) = \lambda$$

$$\text{Var}(Y) = \lambda$$

# The Poisson Distribution

$$Y \sim \text{Poisson}(\lambda)$$


$$y = 0, 1, \dots$$

expected number of  
occurrences:  $\lambda > 0$

$$p(y|\lambda) = \frac{e^{-\lambda} \lambda^y}{y!}$$

$$E(Y) = \lambda$$

$$\text{Var}(Y) = \lambda$$



# The Geometric Distribution

# The Geometric Distribution

How many Bernoulli trials until success?

# The Geometric Distribution

How many Bernoulli trials until success?

$$Y \sim \text{Geometric}(\pi)$$



# The Geometric Distribution

How many Bernoulli trials until success?

$$Y \sim \text{Geometric}(\pi)$$

$$y = 1, 2, 3, \dots$$

# The Geometric Distribution

How many Bernoulli trials until success?

$$Y \sim \text{Geometric}(\pi)$$

$$y = 1, 2, 3, \dots$$

probability of success:  $\pi \in [0, 1]$

# The Geometric Distribution

How many Bernoulli trials until success?

$$Y \sim \text{Geometric}(\pi)$$

$$y = 1, 2, 3, \dots$$

probability of success:  $\pi \in [0, 1]$

$$p(y|\pi) = (1 - \pi)^{(y-1)}\pi$$

# The Geometric Distribution

How many Bernoulli trials until success?

$$Y \sim \text{Geometric}(\pi)$$

$$y = 1, 2, 3, \dots$$

probability of success:  $\pi \in [0, 1]$

$$p(y|\pi) = (1 - \pi)^{(y-1)}\pi$$

$$E(Y) = \frac{1}{\pi}$$

# The Geometric Distribution

How many Bernoulli trials until success?

$$Y \sim \text{Geometric}(\pi)$$

$$y = 1, 2, 3, \dots$$

probability of success:  $\pi \in [0, 1]$

$$p(y|\pi) = (1 - \pi)^{(y-1)}\pi$$

$$E(Y) = \frac{1}{\pi}$$

$$\text{Var}(Y) = \frac{1-\pi}{\pi^2}$$

# The Geometric Distribution

How many Bernoulli trials until success?

$$Y \sim \text{Geometric}(\pi)$$


$$y = 1, 2, 3, \dots$$

probability of success:  $\pi \in [0, 1]$

$$p(y|\pi) = (1 - \pi)^{(y-1)}\pi$$

$$E(Y) = \frac{1}{\pi}$$

$$\text{Var}(Y) = \frac{1-\pi}{\pi^2}$$



# Outline

Expectation, Variance, and Densities

Important Distributions

Discrete Distributions

Continuous Distributions

# The Univariate Normal Distribution



# The Univariate Normal Distribution

$$Y \sim \text{Normal}(\mu, \sigma^2)$$

# The Univariate Normal Distribution

$$Y \sim \text{Normal}(\mu, \sigma^2)$$

$$y \in \mathbb{R}$$

# The Univariate Normal Distribution

$$Y \sim \text{Normal}(\mu, \sigma^2)$$

$$y \in \mathbb{R}$$

$$\text{mean: } \mu \in \mathbb{R}$$

# The Univariate Normal Distribution

$$Y \sim \text{Normal}(\mu, \sigma^2)$$

$$y \in \mathbb{R}$$

mean:  $\mu \in \mathbb{R}$

variance:  $\sigma^2 > 0$

# The Univariate Normal Distribution

$$Y \sim \text{Normal}(\mu, \sigma^2)$$

$$y \in \mathbb{R}$$

mean:  $\mu \in \mathbb{R}$

variance:  $\sigma^2 > 0$

$$p(y|\mu, \sigma^2) = \frac{\exp\left(-\frac{(y-\mu)^2}{2\sigma^2}\right)}{\sigma\sqrt{2\pi}}$$

# The Univariate Normal Distribution

$$Y \sim \text{Normal}(\mu, \sigma^2)$$

$$y \in \mathbb{R}$$

mean:  $\mu \in \mathbb{R}$

variance:  $\sigma^2 > 0$

$$p(y|\mu, \sigma^2) = \frac{\exp\left(-\frac{(y-\mu)^2}{2\sigma^2}\right)}{\sigma\sqrt{2\pi}}$$

$$E(Y) = \mu$$

# The Univariate Normal Distribution

$$Y \sim \text{Normal}(\mu, \sigma^2)$$

$$y \in \mathbb{R}$$

mean:  $\mu \in \mathbb{R}$

variance:  $\sigma^2 > 0$

$$p(y|\mu, \sigma^2) = \frac{\exp\left(-\frac{(y-\mu)^2}{2\sigma^2}\right)}{\sigma\sqrt{2\pi}}$$

$$E(Y) = \mu$$

$$\text{Var}(Y) = \sigma^2$$

# The Univariate Normal Distribution

$$Y \sim \text{Normal}(\mu, \sigma^2)$$

$$y \in \mathbb{R}$$


mean:  $\mu \in \mathbb{R}$

variance:  $\sigma^2 > 0$

$$p(y|\mu, \sigma^2) = \frac{\exp\left(-\frac{(y-\mu)^2}{2\sigma^2}\right)}{\sigma\sqrt{2\pi}}$$

$$E(Y) = \mu$$

$$\text{Var}(Y) = \sigma^2$$





# The Multivariate Normal Distribution

# The Multivariate Normal Distribution

$$Y \sim \mathcal{N}(\mu, \Sigma)$$

# The Multivariate Normal Distribution

$$Y \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

$$\mathbf{y} \in \mathbb{R}^k$$

# The Multivariate Normal Distribution

$$Y \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

$$\mathbf{y} \in \mathbb{R}^k$$

mean vector:  $\boldsymbol{\mu} \in \mathbb{R}^k$

# The Multivariate Normal Distribution

$$Y \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

$$\mathbf{y} \in \mathbb{R}^k$$

mean vector:  $\boldsymbol{\mu} \in \mathbb{R}^k$

variance-covariance matrix:  $\boldsymbol{\Sigma}$  positive definite  $k \times k$  matrix

# The Multivariate Normal Distribution

$$Y \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

$$\mathbf{y} \in \mathbb{R}^k$$

mean vector:  $\boldsymbol{\mu} \in \mathbb{R}^k$

variance-covariance matrix:  $\boldsymbol{\Sigma}$  positive definite  $k \times k$  matrix

$$p(\mathbf{y}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = (2\pi)^{-k/2} |\boldsymbol{\Sigma}|^{-1/2} \exp\left(-\frac{1}{2}(\mathbf{y} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1}(\mathbf{y} - \boldsymbol{\mu})\right)$$

# The Multivariate Normal Distribution

$$Y \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

$$\mathbf{y} \in \mathbb{R}^k$$

mean vector:  $\boldsymbol{\mu} \in \mathbb{R}^k$

variance-covariance matrix:  $\boldsymbol{\Sigma}$  positive definite  $k \times k$  matrix

$$p(\mathbf{y}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = (2\pi)^{-k/2} |\boldsymbol{\Sigma}|^{-1/2} \exp\left(-\frac{1}{2}(\mathbf{y} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1}(\mathbf{y} - \boldsymbol{\mu})\right)$$

$$E(Y) = \boldsymbol{\mu}$$

# The Multivariate Normal Distribution

$$Y \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

$$\mathbf{y} \in \mathbb{R}^k$$

mean vector:  $\boldsymbol{\mu} \in \mathbb{R}^k$

variance-covariance matrix:  $\boldsymbol{\Sigma}$  positive definite  $k \times k$  matrix

$$p(\mathbf{y}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = (2\pi)^{-k/2} |\boldsymbol{\Sigma}|^{-1/2} \exp\left(-\frac{1}{2}(\mathbf{y} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1}(\mathbf{y} - \boldsymbol{\mu})\right)$$

$$E(Y) = \boldsymbol{\mu}$$

$$\text{Var}(Y) = \boldsymbol{\Sigma}$$



# The Uniform Distribution

# The Uniform Distribution

$$Y \sim \text{Uniform}(\alpha, \beta)$$

# The Uniform Distribution

$$Y \sim \text{Uniform}(\alpha, \beta)$$

$$y \in [\alpha, \beta]$$

# The Uniform Distribution

$$Y \sim \text{Uniform}(\alpha, \beta)$$

$$y \in [\alpha, \beta]$$

Interval:  $[\alpha, \beta]$ ;  $\beta > \alpha$

# The Uniform Distribution

$$Y \sim \text{Uniform}(\alpha, \beta)$$

$$y \in [\alpha, \beta]$$

Interval:  $[\alpha, \beta]$ ;  $\beta > \alpha$

$$p(y|\alpha, \beta) = \frac{1}{\beta - \alpha}$$

# The Uniform Distribution

$$Y \sim \text{Uniform}(\alpha, \beta)$$

$$y \in [\alpha, \beta]$$

Interval:  $[\alpha, \beta]$ ;  $\beta > \alpha$

$$p(y|\alpha, \beta) = \frac{1}{\beta - \alpha}$$

$$E(Y) = \frac{\alpha + \beta}{2}$$

# The Uniform Distribution

$$Y \sim \text{Uniform}(\alpha, \beta)$$

$$y \in [\alpha, \beta]$$

$$\text{Interval: } [\alpha, \beta]; \quad \beta > \alpha$$

$$p(y|\alpha, \beta) = \frac{1}{\beta - \alpha}$$

$$E(Y) = \frac{\alpha + \beta}{2}$$

$$\text{Var}(Y) = \frac{(\beta - \alpha)^2}{12}$$

# The Beta Distribution



# The Beta Distribution

$$Y \sim \text{Beta}(\alpha, \beta)$$

# The Beta Distribution

$$Y \sim \text{Beta}(\alpha, \beta)$$

$$y \in [0, 1]$$

# The Beta Distribution

$$Y \sim \text{Beta}(\alpha, \beta)$$

$$y \in [0, 1]$$

shape parameters:

$$\alpha > 0; \beta > 0$$

# The Beta Distribution

$$Y \sim \text{Beta}(\alpha, \beta)$$

$$y \in [0, 1]$$

shape parameters:

$$\alpha > 0; \quad \beta > 0$$

$$p(y|\alpha, \beta) = \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} y^{(\alpha-1)} (1-y)^{(\beta-1)}$$

# The Beta Distribution

$$Y \sim \text{Beta}(\alpha, \beta)$$

$$y \in [0, 1]$$

shape parameters:

$$\alpha > 0; \quad \beta > 0$$

$$p(y|\alpha, \beta) = \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} y^{(\alpha-1)} (1-y)^{(\beta-1)}$$

$$E(Y) = \frac{\alpha}{\alpha+\beta}$$

# The Beta Distribution

$$Y \sim \text{Beta}(\alpha, \beta)$$

$$y \in [0, 1]$$


shape parameters:

$$\alpha > 0; \beta > 0$$

$$p(y|\alpha, \beta) = \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} y^{(\alpha-1)}(1-y)^{(\beta-1)}$$

$$E(Y) = \frac{\alpha}{\alpha+\beta}$$

$$\text{Var}(Y) = \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$$



# The Gamma Distribution

# The Gamma Distribution

$$Y \sim \text{Gamma}(\alpha, \beta)$$



# The Gamma Distribution

$$Y \sim \text{Gamma}(\alpha, \beta)$$

$$y > 0$$

# The Gamma Distribution

$$Y \sim \text{Gamma}(\alpha, \beta)$$

$$y > 0$$

shape parameter:  $\alpha > 0$

# The Gamma Distribution

$$Y \sim \text{Gamma}(\alpha, \beta)$$

$$y > 0$$

shape parameter:  $\alpha > 0$

inverse scale parameter:  $\beta > 0$

# The Gamma Distribution

$$Y \sim \text{Gamma}(\alpha, \beta)$$

$$y > 0$$

shape parameter:  $\alpha > 0$

inverse scale parameter:  $\beta > 0$

$$p(y|\alpha, \beta) = \frac{\beta^\alpha}{\Gamma(\alpha)} y^{(\alpha-1)} \exp(-\beta y)$$

# The Gamma Distribution

$$Y \sim \text{Gamma}(\alpha, \beta)$$

$$y > 0$$

shape parameter:  $\alpha > 0$

inverse scale parameter:  $\beta > 0$

$$p(y|\alpha, \beta) = \frac{\beta^\alpha}{\Gamma(\alpha)} y^{(\alpha-1)} \exp(-\beta y)$$

$$E(Y) = \frac{\alpha}{\beta}$$

# The Gamma Distribution

$$Y \sim \text{Gamma}(\alpha, \beta)$$

$$y > 0$$


shape parameter:  $\alpha > 0$

inverse scale parameter:  $\beta > 0$

$$p(y|\alpha, \beta) = \frac{\beta^\alpha}{\Gamma(\alpha)} y^{(\alpha-1)} \exp(-\beta y)$$

$$E(Y) = \frac{\alpha}{\beta}$$

$$\text{Var}(Y) = \frac{\alpha}{\beta^2}$$



# The Inverse Gamma Distribution

# The Inverse Gamma Distribution

Distribution of the Inverse of a Gamma Distribution:



# The Inverse Gamma Distribution

Distribution of the Inverse of a Gamma Distribution: If  $X \sim \text{Gamma}(\alpha, \beta)$ , then  $\frac{1}{X} \sim \text{Invgamma}(\alpha, \beta)$ .

# The Inverse Gamma Distribution

Distribution of the Inverse of a Gamma Distribution: If  $X \sim \text{Gamma}(\alpha, \beta)$ , then  $\frac{1}{X} \sim \text{Invgamma}(\alpha, \beta)$ .

$$Y \sim \text{Invgamma}(\alpha, \beta)$$

# The Inverse Gamma Distribution

Distribution of the Inverse of a Gamma Distribution: If  $X \sim \text{Gamma}(\alpha, \beta)$ , then  $\frac{1}{X} \sim \text{Invgamma}(\alpha, \beta)$ .

$$Y \sim \text{Invgamma}(\alpha, \beta)$$

$$y > 0$$

# The Inverse Gamma Distribution

Distribution of the Inverse of a Gamma Distribution: If  $X \sim \text{Gamma}(\alpha, \beta)$ , then  $\frac{1}{X} \sim \text{Invgamma}(\alpha, \beta)$ .

$$Y \sim \text{Invgamma}(\alpha, \beta)$$

$$y > 0$$

shape parameter:  $\alpha > 0$

# The Inverse Gamma Distribution

Distribution of the Inverse of a Gamma Distribution: If  $X \sim \text{Gamma}(\alpha, \beta)$ , then  $\frac{1}{X} \sim \text{Invgamma}(\alpha, \beta)$ .

$$Y \sim \text{Invgamma}(\alpha, \beta)$$

$$y > 0$$

shape parameter:  $\alpha > 0$

scale parameter:  $\beta > 0$

# The Inverse Gamma Distribution

Distribution of the Inverse of a Gamma Distribution: If  $X \sim \text{Gamma}(\alpha, \beta)$ , then  $\frac{1}{X} \sim \text{Invgamma}(\alpha, \beta)$ .

$$Y \sim \text{Invgamma}(\alpha, \beta)$$

$$y > 0$$

shape parameter:  $\alpha > 0$

scale parameter:  $\beta > 0$

$$p(y|\alpha, \beta) = \frac{\beta^\alpha}{\Gamma(\alpha)} y^{-(\alpha+1)} e^{-\frac{\beta}{y}}$$

# The Inverse Gamma Distribution

Distribution of the Inverse of a Gamma Distribution: If  $X \sim \text{Gamma}(\alpha, \beta)$ , then  $\frac{1}{X} \sim \text{Invgamma}(\alpha, \beta)$ .

$$Y \sim \text{Invgamma}(\alpha, \beta)$$

$$y > 0$$

shape parameter:  $\alpha > 0$

scale parameter:  $\beta > 0$

$$p(y|\alpha, \beta) = \frac{\beta^\alpha}{\Gamma(\alpha)} y^{-(\alpha+1)} e^{-\frac{\beta}{y}}$$

$$E(Y) = \frac{\beta}{\alpha-1} \text{ for } \alpha > 1$$

# The Inverse Gamma Distribution

Distribution of the Inverse of a Gamma Distribution: If  $X \sim \text{Gamma}(\alpha, \beta)$ , then  $\frac{1}{X} \sim \text{Invgamma}(\alpha, \beta)$ .

$$Y \sim \text{Invgamma}(\alpha, \beta)$$

$$y > 0$$

shape parameter:  $\alpha > 0$

scale parameter:  $\beta > 0$

$$p(y|\alpha, \beta) = \frac{\beta^\alpha}{\Gamma(\alpha)} y^{-(\alpha+1)} e^{-\frac{\beta}{y}}$$

$$E(Y) = \frac{\beta}{\alpha-1} \text{ for } \alpha > 1$$

$$\text{Var}(Y) = \frac{\beta^2}{(\alpha-1)^2(\alpha-2)} \text{ for } \alpha > 2$$



# The Inverse Gamma Distribution

Distribution of the Inverse of a Gamma Distribution: If  $X \sim \text{Gamma}(\alpha, \beta)$ , then  $\frac{1}{X} \sim \text{Invgamma}(\alpha, \beta)$ .

$$Y \sim \text{Invgamma}(\alpha, \beta)$$

$$y > 0$$


shape parameter:  $\alpha > 0$

scale parameter:  $\beta > 0$

$$p(y|\alpha, \beta) = \frac{\beta^\alpha}{\Gamma(\alpha)} y^{-(\alpha+1)} e^{-\frac{\beta}{y}}$$

$$E(Y) = \frac{\beta}{\alpha-1} \text{ for } \alpha > 1$$

$$\text{Var}(Y) = \frac{\beta^2}{(\alpha-1)^2(\alpha-2)} \text{ for } \alpha > 2$$



# The Dirichlet Distribution

# The Dirichlet Distribution

$$Y \sim \text{Dirichlet}(\alpha_1, \dots, \alpha_k)$$

# The Dirichlet Distribution

$$Y \sim \text{Dirichlet}(\alpha_1, \dots, \alpha_k)$$

$$y_j \in [0, 1]; \quad \sum_{j=1}^k y_j = 1$$

# The Dirichlet Distribution

$$Y \sim \text{Dirichlet}(\alpha_1, \dots, \alpha_k)$$

$$y_j \in [0, 1]; \quad \sum_{j=1}^k y_j = 1$$

$$\alpha \text{ parameters: } \alpha_j > 0; \quad \sum_{j=1}^k \alpha_j \equiv \alpha_0$$

# The Dirichlet Distribution

$$Y \sim \text{Dirichlet}(\alpha_1, \dots, \alpha_k)$$

$$y_j \in [0, 1]; \quad \sum_{j=1}^k y_j = 1$$

$$\alpha \text{ parameters: } \alpha_j > 0; \quad \sum_{j=1}^k \alpha_j \equiv \alpha_0$$

$$p(\mathbf{y}|\boldsymbol{\alpha}) = \frac{\Gamma(\alpha_1 + \dots + \alpha_k)}{\Gamma(\alpha_1) \dots \Gamma(\alpha_k)} y_1^{\alpha_1-1} \dots y_k^{\alpha_k-1}$$

# The Dirichlet Distribution

$$Y \sim \text{Dirichlet}(\alpha_1, \dots, \alpha_k)$$

$$y_j \in [0, 1]; \quad \sum_{j=1}^k y_j = 1$$

$$\alpha \text{ parameters: } \alpha_j > 0; \quad \sum_{j=1}^k \alpha_j \equiv \alpha_0$$

$$p(\mathbf{y}|\boldsymbol{\alpha}) = \frac{\Gamma(\alpha_1 + \dots + \alpha_k)}{\Gamma(\alpha_1) \dots \Gamma(\alpha_k)} y_1^{\alpha_1-1} \dots y_k^{\alpha_k-1}$$

$$E(Y_j) = \frac{\alpha_j}{\alpha_0}$$

# The Dirichlet Distribution

$$Y \sim \text{Dirichlet}(\alpha_1, \dots, \alpha_k)$$

$$y_j \in [0, 1]; \quad \sum_{j=1}^k y_j = 1$$

$$\alpha \text{ parameters: } \alpha_j > 0; \quad \sum_{j=1}^k \alpha_j \equiv \alpha_0$$

$$p(\mathbf{y}|\boldsymbol{\alpha}) = \frac{\Gamma(\alpha_1 + \dots + \alpha_k)}{\Gamma(\alpha_1) \dots \Gamma(\alpha_k)} y_1^{\alpha_1-1} \dots y_k^{\alpha_k-1}$$

$$E(Y_j) = \frac{\alpha_j}{\alpha_0}$$

$$\text{Var}(Y_j) = \frac{\alpha_j(\alpha_0 - \alpha_j)}{\alpha_0^2(\alpha_0 + 1)}$$



# The Dirichlet Distribution

$$Y \sim \text{Dirichlet}(\alpha_1, \dots, \alpha_k)$$

$$y_j \in [0, 1]; \quad \sum_{j=1}^k y_j = 1$$

$$\alpha \text{ parameters: } \alpha_j > 0; \quad \sum_{j=1}^k \alpha_j \equiv \alpha_0$$

$$p(\mathbf{y}|\boldsymbol{\alpha}) = \frac{\Gamma(\alpha_1 + \dots + \alpha_k)}{\Gamma(\alpha_1) \dots \Gamma(\alpha_k)} y_1^{\alpha_1-1} \dots y_k^{\alpha_k-1}$$

$$E(Y_j) = \frac{\alpha_j}{\alpha_0}$$

$$\text{Var}(Y_j) = \frac{\alpha_j(\alpha_0 - \alpha_j)}{\alpha_0^2(\alpha_0 + 1)}$$

$$\text{Cov}(Y_i, Y_j) = -\frac{\alpha_i \alpha_j}{\alpha_0^2(\alpha_0 + 1)}$$