

# Bayesian Statistics in One Hour

Patrick Lam

# Outline

Introduction

Bayesian Models

Applications

- Missing Data

- Hierarchical Models

# Outline

Introduction

Bayesian Models


Applications

Missing Data



Hierarchical Models

# References




# References






# References



# References



# References



Western, Bruce and Simon Jackman. 1994. "Bayesian Inference for Comparative Research." *American Political Science Review* 88(2): 412-423.

Jackman, Simon. 2000. "Estimation and Inference via Bayesian Simulation: An Introduction to Markov Chain Monte Carlo." *American Journal of Political Science* 44(2): 375-404.

Jackman, Simon. 2000. "Estimation and Inference Are Missing Data Problems: Unifying Social Science Statistics via Bayesian Simulation." *Political Analysis* 8(4): 307-332.



# Introduction

# Introduction

Three general approaches to statistics:

# Introduction

Three general approaches to statistics:

- ▶ frequentist (Neyman-Pearson, hypothesis testing)

# Introduction

Three general approaches to statistics:

- ▶ frequentist (Neyman-Pearson, hypothesis testing)
- ▶ likelihood (what we've been learning all semester)

# Introduction

Three general approaches to statistics:

- ▶ frequentist (Neyman-Pearson, hypothesis testing)
- ▶ likelihood (what we've been learning all semester)
- ▶ Bayesian

# Introduction

Three general approaches to statistics:

- ▶ frequentist (Neyman-Pearson, hypothesis testing)
- ▶ likelihood (what we've been learning all semester)
- ▶ Bayesian

Today's goal: Contrast {frequentist, likelihood} with Bayesian,

# Introduction

Three general approaches to statistics:

- ▶ frequentist (Neyman-Pearson, hypothesis testing)
- ▶ likelihood (what we've been learning all semester)
- ▶ Bayesian

Today's goal: Contrast {frequentist, likelihood} with Bayesian, with emphasis on Bayesian versus likelihood.

# Introduction

Three general approaches to statistics:

- ▶ frequentist (Neyman-Pearson, hypothesis testing)
- ▶ likelihood (what we've been learning all semester)
- ▶ Bayesian

Today's goal: Contrast {frequentist, likelihood} with Bayesian, with emphasis on Bayesian versus likelihood.

We'll go over some of the Bayesian critiques of non-Bayesian analysis



# Introduction

Three general approaches to statistics:

- ▶ frequentist (Neyman-Pearson, hypothesis testing)
- ▶ likelihood (what we've been learning all semester)
- ▶ Bayesian

Today's goal: Contrast {frequentist, likelihood} with Bayesian, with emphasis on Bayesian versus likelihood.

We'll go over some of the Bayesian critiques of non-Bayesian analysis and *non-Bayesian critiques of Bayesian analysis*.

# Probability

# Probability

Objective view of probability (non-Bayesian):

# Probability

Objective view of probability (non-Bayesian):

- ▶ The relative frequency of an outcome of an experiment over repeated runs of the experiment.

# Probability

Objective view of probability (non-Bayesian):

- ▶ The relative frequency of an outcome of an experiment over repeated runs of the experiment.
- ▶ The observed proportion in a population.

# Probability

Objective view of probability (non-Bayesian):

- ▶ The relative frequency of an outcome of an experiment over repeated runs of the experiment.
- ▶ The observed proportion in a population.

Subjective view of probability (Bayesian):

# Probability

Objective view of probability (non-Bayesian):

- ▶ The relative frequency of an outcome of an experiment over repeated runs of the experiment.
- ▶ The observed proportion in a population.

Subjective view of probability (Bayesian):

- ▶ Individual's degree of belief in a statement

# Probability

Objective view of probability (non-Bayesian):

- ▶ The relative frequency of an outcome of an experiment over repeated runs of the experiment.
- ▶ The observed proportion in a population.

Subjective view of probability (Bayesian):

- ▶ Individual's degree of belief in a statement
- ▶ Defined personally



# Probability

Objective view of probability (non-Bayesian):

- ▶ The relative frequency of an outcome of an experiment over repeated runs of the experiment.
- ▶ The observed proportion in a population.

Subjective view of probability (Bayesian):

- ▶ Individual's degree of belief in a statement
- ▶ Defined personally (how much money would you wager on an outcome?)

# Probability

## Objective view of probability (non-Bayesian):

- ▶ The relative frequency of an outcome of an experiment over repeated runs of the experiment.
- ▶ The observed proportion in a population.

## Subjective view of probability (Bayesian):

- ▶ Individual's degree of belief in a statement
- ▶ Defined personally (how much money would you wager on an outcome?)
- ▶ Can be influenced in many ways (personal beliefs, prior evidence)

# Probability

Objective view of probability (non-Bayesian):

- ▶ The relative frequency of an outcome of an experiment over repeated runs of the experiment.
- ▶ The observed proportion in a population.

Subjective view of probability (Bayesian):

- ▶ Individual's degree of belief in a statement
- ▶ Defined personally (how much money would you wager on an outcome?)
- ▶ Can be influenced in many ways (personal beliefs, prior evidence)

Bayesian statistics is convenient because it does not require repeated sampling or large  $n$  assumptions.

# Maximum Likelihood

# Maximum Likelihood

$$\begin{aligned} p(\theta|y) &= \frac{p(y|\theta)p(\theta)}{p(y)} \\ &= p(y|\theta)k(y) \end{aligned}$$

# Maximum Likelihood

$$\begin{aligned} p(\theta|y) &= \frac{p(y|\theta)p(\theta)}{p(y)} \\ &= p(y|\theta)k(y) \\ &\propto p(y|\theta) \end{aligned}$$

# Maximum Likelihood

$$\begin{aligned} p(\theta|y) &= \frac{p(y|\theta)p(\theta)}{p(y)} \\ &= p(y|\theta)k(y) \\ &\propto p(y|\theta) \end{aligned}$$

$$L(\theta|y) = p(y|\theta)$$

# Maximum Likelihood

$$\begin{aligned} p(\theta|y) &= \frac{p(y|\theta)p(\theta)}{p(y)} \\ &= p(y|\theta)k(y) \\ &\propto p(y|\theta) \end{aligned}$$

$$L(\theta|y) = p(y|\theta)$$

There is a fixed, true value of  $\theta$ ,



# Maximum Likelihood

$$\begin{aligned} p(\theta|y) &= \frac{p(y|\theta)p(\theta)}{p(y)} \\ &= p(y|\theta)k(y) \\ &\propto p(y|\theta) \end{aligned}$$

$$L(\theta|y) = p(y|\theta)$$

There is a fixed, true value of  $\theta$ , and we maximize the likelihood to estimate  $\theta$  and make assumptions to generate uncertainty about our estimate of  $\theta$ .

# Bayesian

# Bayesian

$$p(\theta|y) = \frac{p(y|\theta)p(\theta)}{p(y)}$$

# Bayesian

$$\begin{aligned} p(\theta|y) &= \frac{p(y|\theta)p(\theta)}{p(y)} \\ &\propto p(y|\theta)p(\theta) \end{aligned}$$

# Bayesian

$$\begin{aligned} p(\theta|y) &= \frac{p(y|\theta)p(\theta)}{p(y)} \\ &\propto p(y|\theta)p(\theta) \end{aligned}$$

- $\theta$  is a random variable.

# Bayesian

$$\begin{aligned} p(\theta|y) &= \frac{p(y|\theta)p(\theta)}{p(y)} \\ &\propto p(y|\theta)p(\theta) \end{aligned}$$

- ▶  $\theta$  is a random variable.
  - ▶  $\theta$  is stochastic and changes from time to time.

# Bayesian

$$\begin{aligned} p(\theta|y) &= \frac{p(y|\theta)p(\theta)}{p(y)} \\ &\propto p(y|\theta)p(\theta) \end{aligned}$$

- ▶  $\theta$  is a random variable.
  - ▶  $\theta$  is stochastic and changes from time to time.
  - ▶  $\theta$  is truly fixed, but we want to reflect our uncertainty about it.

# Bayesian

$$\begin{aligned} p(\theta|y) &= \frac{p(y|\theta)p(\theta)}{p(y)} \\ &\propto p(y|\theta)p(\theta) \end{aligned}$$

- ▶  $\theta$  is a random variable.
  - ▶  $\theta$  is stochastic and changes from time to time.
  - ▶  $\theta$  is truly fixed, but we want to reflect our uncertainty about it.
- ▶ We have a **prior** subjective belief about  $\theta$ , which we update with the data to form **posterior** beliefs about  $\theta$ .



# Bayesian

$$\begin{aligned} p(\theta|y) &= \frac{p(y|\theta)p(\theta)}{p(y)} \\ &\propto p(y|\theta)p(\theta) \end{aligned}$$

- ▶  $\theta$  is a random variable.
  - ▶  $\theta$  is stochastic and changes from time to time.
  - ▶  $\theta$  is truly fixed, but we want to reflect our uncertainty about it.
- ▶ We have a **prior** subjective belief about  $\theta$ , which we update with the data to form **posterior** beliefs about  $\theta$ .
- ▶ The **posterior** is a probability distribution that must integrate to 1.

# Bayesian

$$\begin{aligned} p(\theta|y) &= \frac{p(y|\theta)p(\theta)}{p(y)} \\ &\propto p(y|\theta)p(\theta) \end{aligned}$$

- ▶  $\theta$  is a random variable.
  - ▶  $\theta$  is stochastic and changes from time to time.
  - ▶  $\theta$  is truly fixed, but we want to reflect our uncertainty about it.
- ▶ We have a **prior** subjective belief about  $\theta$ , which we update with the data to form **posterior** beliefs about  $\theta$ .
- ▶ The **posterior** is a probability distribution that must integrate to 1.
- ▶ The **prior** is usually a probability distribution that integrates to 1 (proper prior).

# $\theta$ as Fixed versus as a Random Variable

# $\theta$ as Fixed versus as a Random Variable

Non-Bayesian approach ( $\theta$  fixed):

## $\theta$ as Fixed versus as a Random Variable

Non-Bayesian approach ( $\theta$  fixed):

- ▶ Estimate  $\theta$  with measures of uncertainty (SE, CIs)

## $\theta$ as Fixed versus as a Random Variable

Non-Bayesian approach ( $\theta$  fixed):

- ▶ Estimate  $\theta$  with measures of uncertainty (SE, CIs)
- ▶ 95% Confidence Interval: 95% of the time,  $\theta$  is in the 95% interval that is estimated each time.

## $\theta$ as Fixed versus as a Random Variable

Non-Bayesian approach ( $\theta$  fixed):

- ▶ Estimate  $\theta$  with measures of uncertainty (SE, CIs)
- ▶ 95% Confidence Interval: 95% of the time,  $\theta$  is in the 95% interval that is estimated each time.
  - ▶  $P(\theta \in 95\% \text{ CI}) =$

## $\theta$ as Fixed versus as a Random Variable

Non-Bayesian approach ( $\theta$  fixed):

- ▶ Estimate  $\theta$  with measures of uncertainty (SE, CIs)
- ▶ 95% Confidence Interval: 95% of the time,  $\theta$  is in the 95% interval that is estimated each time.
  - ▶  $P(\theta \in 95\% \text{ CI}) = 0 \text{ or } 1$



# $\theta$ as Fixed versus as a Random Variable

Non-Bayesian approach ( $\theta$  fixed):

- ▶ Estimate  $\theta$  with measures of uncertainty (SE, CIs)
- ▶ 95% Confidence Interval: 95% of the time,  $\theta$  is in the 95% interval that is estimated each time.
  - ▶  $P(\theta \in 95\% \text{ CI}) = 0 \text{ or } 1$
- ▶  $P(\theta > 2) =$

## $\theta$ as Fixed versus as a Random Variable

Non-Bayesian approach ( $\theta$  fixed):

- ▶ Estimate  $\theta$  with measures of uncertainty (SE, CIs)
- ▶ 95% Confidence Interval: 95% of the time,  $\theta$  is in the 95% interval that is estimated each time.
  - ▶  $P(\theta \in 95\% \text{ CI}) = 0 \text{ or } 1$
- ▶  $P(\theta > 2) = 0 \text{ or } 1$

## $\theta$ as Fixed versus as a Random Variable

Non-Bayesian approach ( $\theta$  fixed):

- ▶ Estimate  $\theta$  with measures of uncertainty (SE, CIs)
- ▶ 95% Confidence Interval: 95% of the time,  $\theta$  is in the 95% interval that is estimated each time.
  - ▶  $P(\theta \in 95\% \text{ CI}) = 0 \text{ or } 1$
- ▶  $P(\theta > 2) = 0 \text{ or } 1$

Bayesian approach ( $\theta$  random):

# $\theta$ as Fixed versus as a Random Variable

Non-Bayesian approach ( $\theta$  fixed):

- ▶ Estimate  $\theta$  with measures of uncertainty (SE, CIs)
- ▶ 95% Confidence Interval: 95% of the time,  $\theta$  is in the 95% interval that is estimated each time.
  - ▶  $P(\theta \in 95\% \text{ CI}) = 0 \text{ or } 1$
- ▶  $P(\theta > 2) = 0 \text{ or } 1$

Bayesian approach ( $\theta$  random):

- ▶ Find the **posterior** distribution of  $\theta$ .

# $\theta$ as Fixed versus as a Random Variable

Non-Bayesian approach ( $\theta$  fixed):

- ▶ Estimate  $\theta$  with measures of uncertainty (SE, CIs)
- ▶ 95% Confidence Interval: 95% of the time,  $\theta$  is in the 95% interval that is estimated each time.
  - ▶  $P(\theta \in 95\% \text{ CI}) = 0 \text{ or } 1$
- ▶  $P(\theta > 2) = 0 \text{ or } 1$

Bayesian approach ( $\theta$  random):

- ▶ Find the **posterior** distribution of  $\theta$ .
- ▶ Take quantities of interest from the distribution (posterior mean, posterior SD, posterior credible intervals)

# $\theta$ as Fixed versus as a Random Variable

Non-Bayesian approach ( $\theta$  fixed):

- ▶ Estimate  $\theta$  with measures of uncertainty (SE, CIs)
- ▶ 95% Confidence Interval: 95% of the time,  $\theta$  is in the 95% interval that is estimated each time.
  - ▶  $P(\theta \in 95\% \text{ CI}) = 0 \text{ or } 1$
- ▶  $P(\theta > 2) = 0 \text{ or } 1$

Bayesian approach ( $\theta$  random):

- ▶ Find the **posterior** distribution of  $\theta$ .
- ▶ Take quantities of interest from the distribution (posterior mean, posterior SD, posterior credible intervals)
- ▶ We can make probability statements regarding  $\theta$ .

## $\theta$ as Fixed versus as a Random Variable

Non-Bayesian approach ( $\theta$  fixed):

- ▶ Estimate  $\theta$  with measures of uncertainty (SE, CIs)
- ▶ 95% Confidence Interval: 95% of the time,  $\theta$  is in the 95% interval that is estimated each time.
  - ▶  $P(\theta \in 95\% \text{ CI}) = 0 \text{ or } 1$
- ▶  $P(\theta > 2) = 0 \text{ or } 1$

Bayesian approach ( $\theta$  random):

- ▶ Find the **posterior** distribution of  $\theta$ .
- ▶ Take quantities of interest from the distribution (posterior mean, posterior SD, posterior credible intervals)
- ▶ We can make probability statements regarding  $\theta$ .
  - ▶ 95% Credible Interval:

# $\theta$ as Fixed versus as a Random Variable

Non-Bayesian approach ( $\theta$  fixed):

- ▶ Estimate  $\theta$  with measures of uncertainty (SE, CIs)
- ▶ 95% Confidence Interval: 95% of the time,  $\theta$  is in the 95% interval that is estimated each time.
  - ▶  $P(\theta \in 95\% \text{ CI}) = 0 \text{ or } 1$
- ▶  $P(\theta > 2) = 0 \text{ or } 1$

Bayesian approach ( $\theta$  random):

- ▶ Find the **posterior** distribution of  $\theta$ .
- ▶ Take quantities of interest from the distribution (posterior mean, posterior SD, posterior credible intervals)
- ▶ We can make probability statements regarding  $\theta$ .
  - ▶ 95% Credible Interval:  $P(\theta \in 95\% \text{ CI}) = 0.95$



# $\theta$ as Fixed versus as a Random Variable

Non-Bayesian approach ( $\theta$  fixed):

- ▶ Estimate  $\theta$  with measures of uncertainty (SE, CIs)
- ▶ 95% Confidence Interval: 95% of the time,  $\theta$  is in the 95% interval that is estimated each time.
  - ▶  $P(\theta \in 95\% \text{ CI}) = 0 \text{ or } 1$
- ▶  $P(\theta > 2) = 0 \text{ or } 1$

Bayesian approach ( $\theta$  random):

- ▶ Find the **posterior** distribution of  $\theta$ .
- ▶ Take quantities of interest from the distribution (posterior mean, posterior SD, posterior credible intervals)
- ▶ We can make probability statements regarding  $\theta$ .
  - ▶ 95% Credible Interval:  $P(\theta \in 95\% \text{ CI}) = 0.95$
  - ▶  $P(\theta > 2) = (0, 1)$

# Critiques

# Critiques

$$\text{Posterior} = \text{Evidence} \times \text{Prior}$$

# Critiques

$$\text{Posterior} = \text{Evidence} \times \text{Prior}$$

NB: *Bayesians introduce **priors** that are not justifiable.*

# Critiques

$$\text{Posterior} = \text{Evidence} \times \text{Prior}$$

NB: *Bayesians introduce **priors** that are not justifiable.*

B: Non-Bayesians are just doing Bayesian statistics with uninformative priors, which may be equally unjustifiable.

# Critiques

$$\text{Posterior} = \text{Evidence} \times \text{Prior}$$

NB: *Bayesians introduce **priors** that are not justifiable.*

B: Non-Bayesians are just doing Bayesian statistics with uninformative priors, which may be equally unjustifiable.

NB: *Unjustified Bayesian **priors** are driving the results.*

# Critiques

$$\text{Posterior} = \text{Evidence} \times \text{Prior}$$

NB: *Bayesians introduce **priors** that are not justifiable.*

B: Non-Bayesians are just doing Bayesian statistics with uninformative priors, which may be equally unjustifiable.

NB: *Unjustified Bayesian **priors** are driving the results.*

B: Bayesian results  $\approx$  non-Bayesian results as  $n$  gets larger (the data overwhelm the prior).

# Critiques

$$\text{Posterior} = \text{Evidence} \times \text{Prior}$$

NB: *Bayesians introduce **priors** that are not justifiable.*

B: Non-Bayesians are just doing Bayesian statistics with uninformative priors, which may be equally unjustifiable.

NB: *Unjustified Bayesian **priors** are driving the results.*

B: Bayesian results  $\approx$  non-Bayesian results as  $n$  gets larger (the data overwhelm the prior).

NB: *Bayesian is too hard. Why use it?*



# Critiques

$$\text{Posterior} = \text{Evidence} \times \text{Prior}$$

NB: *Bayesians introduce **priors** that are not justifiable.*

B: Non-Bayesians are just doing Bayesian statistics with uninformative priors, which may be equally unjustifiable.

NB: *Unjustified Bayesian **priors** are driving the results.*

B: Bayesian results  $\approx$  non-Bayesian results as  $n$  gets larger (the data overwhelm the prior).

NB: *Bayesian is too hard. Why use it?*

B: Bayesian methods allow us to easily estimate models that are too hard to estimate (cannot computationally find the MLE) or unidentified (no unique MLE exists) with non-Bayesian methods.

# Critiques

$$\text{Posterior} = \text{Evidence} \times \text{Prior}$$

NB: *Bayesians introduce **priors** that are not justifiable.*

B: Non-Bayesians are just doing Bayesian statistics with uninformative priors, which may be equally unjustifiable.

NB: *Unjustified Bayesian **priors** are driving the results.*

B: Bayesian results  $\approx$  non-Bayesian results as  $n$  gets larger (the data overwhelm the prior).

NB: *Bayesian is too hard. Why use it?*

B: Bayesian methods allow us to easily estimate models that are too hard to estimate (cannot computationally find the MLE) or unidentified (no unique MLE exists) with non-Bayesian methods. Bayesian methods also allow us to incorporate prior/qualitative information into the model.

# Outline

Introduction

Bayesian Models

Applications

Missing Data

Hierarchical Models

# Running a Model

# Running a Model

Non-Bayesian:

# Running a Model

Non-Bayesian:

1. Specify a probability model  
(distribution for  $Y$ ).

# Running a Model

Non-Bayesian:

1. Specify a probability model (distribution for  $Y$ ).
2. Find MLE  $\hat{\theta}$  and measures of uncertainty (SE, CI).  
Assume  $\hat{\theta}$  follows a (multivariate) normal distribution.

# Running a Model

Non-Bayesian:

1. Specify a probability model (distribution for  $Y$ ).
2. Find MLE  $\hat{\theta}$  and measures of uncertainty (SE, CI).  
Assume  $\hat{\theta}$  follows a (multivariate) normal distribution.
3. Estimate quantities of interest analytically or via simulation.



# Running a Model

## Non-Bayesian:

1. Specify a probability model (distribution for  $Y$ ).
2. Find MLE  $\hat{\theta}$  and measures of uncertainty (SE, CI).  
Assume  $\hat{\theta}$  follows a (multivariate) normal distribution.
3. Estimate quantities of interest analytically or via simulation.

## Bayesian:

1. Specify a probability model (distribution for  $Y$  and **priors** on  $\theta$ ).

# Running a Model

## Non-Bayesian:

1. Specify a probability model (distribution for  $Y$ ).
2. Find MLE  $\hat{\theta}$  and measures of uncertainty (SE, CI). Assume  $\hat{\theta}$  follows a (multivariate) normal distribution.
3. Estimate quantities of interest analytically or via simulation.

## Bayesian:

1. Specify a probability model (distribution for  $Y$  and **priors** on  $\theta$ ).
2. Solve for **posterior** and summarize it (mean, SD, credible interval, etc.). We can do both analytically or via simulation.

# Running a Model

## Non-Bayesian:

1. Specify a probability model (distribution for  $Y$ ).
2. Find MLE  $\hat{\theta}$  and measures of uncertainty (SE, CI). Assume  $\hat{\theta}$  follows a (multivariate) normal distribution.
3. Estimate quantities of interest analytically or via simulation.

## Bayesian:

1. Specify a probability model (distribution for  $Y$  and **priors** on  $\theta$ ).
2. Solve for **posterior** and summarize it (mean, SD, credible interval, etc.). We can do both analytically or via simulation.
3. Estimate quantities of interest analytically or via simulation.

There is a Bayesian way to do any non-Bayesian parametric model.

# A Simple (Beta-Binomial) Model

# A Simple (Beta-Binomial) Model

The Los Angeles Lakers play 82 games during a regular NBA season.

## A Simple (Beta-Binomial) Model

The Los Angeles Lakers play 82 games during a regular NBA season. In the 2008-2009 season, they won 65 games.

## A Simple (Beta-Binomial) Model

The Los Angeles Lakers play 82 games during a regular NBA season. In the 2008-2009 season, they won 65 games. Suppose the Lakers win each game with probability  $\pi$ .

## A Simple (Beta-Binomial) Model

The Los Angeles Lakers play 82 games during a regular NBA season. In the 2008-2009 season, they won 65 games. Suppose the Lakers win each game with probability  $\pi$ . Estimate  $\pi$ .



## A Simple (Beta-Binomial) Model

The Los Angeles Lakers play 82 games during a regular NBA season. In the 2008-2009 season, they won 65 games. Suppose the Lakers win each game with probability  $\pi$ . Estimate  $\pi$ .

We have 82 Bernoulli observations or one observation  $Y$ , where

$$Y \sim \text{Binomial}(n, \pi)$$

with  $n = 82$ .

## A Simple (Beta-Binomial) Model

The Los Angeles Lakers play 82 games during a regular NBA season. In the 2008-2009 season, they won 65 games. Suppose the Lakers win each game with probability  $\pi$ . Estimate  $\pi$ .

We have 82 Bernoulli observations or one observation  $Y$ , where

$$Y \sim \text{Binomial}(n, \pi)$$

with  $n = 82$ .

Assumptions:

# A Simple (Beta-Binomial) Model

The Los Angeles Lakers play 82 games during a regular NBA season. In the 2008-2009 season, they won 65 games. Suppose the Lakers win each game with probability  $\pi$ . Estimate  $\pi$ .

We have 82 Bernoulli observations or one observation  $Y$ , where

$$Y \sim \text{Binomial}(n, \pi)$$

with  $n = 82$ .

Assumptions:

- Each game is a Bernoulli trial.

# A Simple (Beta-Binomial) Model

The Los Angeles Lakers play 82 games during a regular NBA season. In the 2008-2009 season, they won 65 games. Suppose the Lakers win each game with probability  $\pi$ . Estimate  $\pi$ .

We have 82 Bernoulli observations or one observation  $Y$ , where

$$Y \sim \text{Binomial}(n, \pi)$$

with  $n = 82$ .

Assumptions:

- ▶ Each game is a Bernoulli trial.
- ▶ The Lakers have the same probability of winning each game.

# A Simple (Beta-Binomial) Model

The Los Angeles Lakers play 82 games during a regular NBA season. In the 2008-2009 season, they won 65 games. Suppose the Lakers win each game with probability  $\pi$ . Estimate  $\pi$ .

We have 82 Bernoulli observations or one observation  $Y$ , where

$$Y \sim \text{Binomial}(n, \pi)$$

with  $n = 82$ .

Assumptions:

- ▶ Each game is a Bernoulli trial.
- ▶ The Lakers have the same probability of winning each game.
- ▶ The outcomes of the games are independent.

We can use the beta distribution as a **prior** for  $\pi$  since it has support over  $[0,1]$ .

We can use the beta distribution as a **prior** for  $\pi$  since it has support over  $[0,1]$ .

$$p(\pi|y) \propto p(y|\pi)p(\pi)$$

We can use the beta distribution as a **prior** for  $\pi$  since it has support over  $[0,1]$ .

$$\begin{aligned} p(\pi|y) &\propto p(y|\pi)p(\pi) \\ &= \text{Binomial}(n, \pi) \times \text{Beta}(\alpha, \beta) \end{aligned}$$



We can use the beta distribution as a **prior** for  $\pi$  since it has support over  $[0,1]$ .

$$\begin{aligned} p(\pi|y) &\propto p(y|\pi)p(\pi) \\ &= \text{Binomial}(n, \pi) \times \text{Beta}(\alpha, \beta) \\ &= \binom{n}{y} \pi^y (1 - \pi)^{(n-y)} \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \pi^{(\alpha-1)} (1 - \pi)^{(\beta-1)} \end{aligned}$$

We can use the beta distribution as a **prior** for  $\pi$  since it has support over  $[0,1]$ .

$$\begin{aligned} p(\pi|y) &\propto p(y|\pi)p(\pi) \\ &= \text{Binomial}(n, \pi) \times \text{Beta}(\alpha, \beta) \\ &= \binom{n}{y} \pi^y (1 - \pi)^{(n-y)} \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \pi^{(\alpha-1)} (1 - \pi)^{(\beta-1)} \\ &\propto \pi^y (1 - \pi)^{(n-y)} \pi^{(\alpha-1)} (1 - \pi)^{(\beta-1)} \end{aligned}$$

We can use the beta distribution as a **prior** for  $\pi$  since it has support over  $[0,1]$ .

$$\begin{aligned} p(\pi|y) &\propto p(y|\pi)p(\pi) \\ &= \text{Binomial}(n, \pi) \times \text{Beta}(\alpha, \beta) \\ &= \binom{n}{y} \pi^y (1 - \pi)^{(n-y)} \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \pi^{(\alpha-1)} (1 - \pi)^{(\beta-1)} \\ &\propto \pi^y (1 - \pi)^{(n-y)} \pi^{(\alpha-1)} (1 - \pi)^{(\beta-1)} \end{aligned}$$

$$p(\pi|y) \propto \pi^{y+\alpha-1} (1 - \pi)^{n-y+\beta-1}$$

We can use the beta distribution as a **prior** for  $\pi$  since it has support over  $[0,1]$ .

$$\begin{aligned} p(\pi|y) &\propto p(y|\pi)p(\pi) \\ &= \text{Binomial}(n, \pi) \times \text{Beta}(\alpha, \beta) \\ &= \binom{n}{y} \pi^y (1 - \pi)^{(n-y)} \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \pi^{(\alpha-1)} (1 - \pi)^{(\beta-1)} \\ &\propto \pi^y (1 - \pi)^{(n-y)} \pi^{(\alpha-1)} (1 - \pi)^{(\beta-1)} \end{aligned}$$

$$p(\pi|y) \propto \pi^{y+\alpha-1} (1 - \pi)^{n-y+\beta-1}$$

The **posterior** distribution is simply a **Beta**( $y + \alpha, n - y + \beta$ ) distribution.

We can use the beta distribution as a **prior** for  $\pi$  since it has support over  $[0,1]$ .

$$\begin{aligned} p(\pi|y) &\propto p(y|\pi)p(\pi) \\ &= \text{Binomial}(n, \pi) \times \text{Beta}(\alpha, \beta) \\ &= \binom{n}{y} \pi^y (1 - \pi)^{(n-y)} \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \pi^{(\alpha-1)} (1 - \pi)^{(\beta-1)} \\ &\propto \pi^y (1 - \pi)^{(n-y)} \pi^{(\alpha-1)} (1 - \pi)^{(\beta-1)} \end{aligned}$$

$$p(\pi|y) \propto \pi^{y+\alpha-1} (1 - \pi)^{n-y+\beta-1}$$

The **posterior** distribution is simply a  $\text{Beta}(y + \alpha, n - y + \beta)$  distribution. Effectively, our **prior** is just adding  $\alpha - 1$  successes and  $\beta - 1$  failures to the dataset.

We can use the beta distribution as a **prior** for  $\pi$  since it has support over  $[0,1]$ .

$$\begin{aligned} p(\pi|y) &\propto p(y|\pi)p(\pi) \\ &= \text{Binomial}(n, \pi) \times \text{Beta}(\alpha, \beta) \\ &= \binom{n}{y} \pi^y (1 - \pi)^{(n-y)} \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \pi^{(\alpha-1)} (1 - \pi)^{(\beta-1)} \\ &\propto \pi^y (1 - \pi)^{(n-y)} \pi^{(\alpha-1)} (1 - \pi)^{(\beta-1)} \end{aligned}$$

$$p(\pi|y) \propto \pi^{y+\alpha-1} (1 - \pi)^{n-y+\beta-1}$$

The **posterior** distribution is simply a  $\text{Beta}(y + \alpha, n - y + \beta)$  distribution. Effectively, our **prior** is just adding  $\alpha - 1$  successes and  $\beta - 1$  failures to the dataset.

*Bayesian **priors** are just adding pseudo observations to the data.*

Since we know the **posterior** is a **Beta( $y + \alpha, n - y + \beta$ )** distribution, we can summarize it analytically or via simulation with the following quantities:

Since we know the **posterior** is a **Beta( $y + \alpha, n - y + \beta$ )** distribution, we can summarize it analytically or via simulation with the following quantities:

- ▶ posterior mean



Since we know the **posterior** is a **Beta( $y + \alpha, n - y + \beta$ )** distribution, we can summarize it analytically or via simulation with the following quantities:

- ▶ posterior mean
- ▶ posterior standard deviation


Since we know the **posterior** is a **Beta( $y + \alpha, n - y + \beta$ )** distribution, we can summarize it analytically or via simulation with the following quantities:

- ▶ posterior mean
- ▶ posterior standard deviation
- ▶ posterior credible intervals (credible sets)


Since we know the **posterior** is a **Beta( $y + \alpha, n - y + \beta$ )** distribution, we can summarize it analytically or via simulation with the following quantities:

- ▶ posterior mean
- ▶ posterior standard deviation
- ▶ posterior credible intervals (credible sets)
- ▶ highest posterior density region


Uninformative Beta(1,1) Prior




Beta(2,12) Prior



Uninformative Beta(1,1) Prior (n=1000)



Beta(2,12) Prior (n=1000)



In the previous model, we had

$$\text{Beta}(y + \alpha, n - y + \beta) = \frac{\text{Binomial}(n, \pi) \times \text{Beta}(\alpha, \beta)}{p(y)}$$

In the previous model, we had

$$\text{Beta}(y + \alpha, n - y + \beta) = \frac{\text{Binomial}(n, \pi) \times \text{Beta}(\alpha, \beta)}{p(y)}$$

We knew that the likelihood  $\times$  prior produced something that looked like a Beta distribution up to a constant of proportionality.

In the previous model, we had

$$\text{Beta}(y + \alpha, n - y + \beta) = \frac{\text{Binomial}(n, \pi) \times \text{Beta}(\alpha, \beta)}{p(y)}$$

We knew that the likelihood  $\times$  prior produced something that looked like a Beta distribution up to a constant of proportionality.

Since the posterior must be a probability distribution, we know that it is a Beta distribution and we can easily solve for the normalizing constant

In the previous model, we had

$$\text{Beta}(y + \alpha, n - y + \beta) = \frac{\text{Binomial}(n, \pi) \times \text{Beta}(\alpha, \beta)}{p(y)}$$

We knew that the likelihood  $\times$  prior produced something that looked like a Beta distribution up to a constant of proportionality.

Since the posterior must be a probability distribution, we know that it is a Beta distribution and we can easily solve for the normalizing constant (although we don't need to since we already have the posterior).



In the previous model, we had

$$\text{Beta}(y + \alpha, n - y + \beta) = \frac{\text{Binomial}(n, \pi) \times \text{Beta}(\alpha, \beta)}{p(y)}$$

We knew that the likelihood  $\times$  **prior** produced something that looked like a Beta distribution up to a **constant** of proportionality.

Since the **posterior** must be a probability distribution, we know that it is a Beta distribution and we can easily solve for the **normalizing constant** (although we don't need to since we already have the **posterior**).

When the **posterior** is the same distribution family as the **prior**, we have **conjugacy**.

In the previous model, we had

$$\text{Beta}(y + \alpha, n - y + \beta) = \frac{\text{Binomial}(n, \pi) \times \text{Beta}(\alpha, \beta)}{p(y)}$$

We knew that the likelihood  $\times$  **prior** produced something that looked like a Beta distribution up to a **constant** of proportionality.

Since the **posterior** must be a probability distribution, we know that it is a Beta distribution and we can easily solve for the **normalizing constant** (although we don't need to since we already have the **posterior**).

When the **posterior** is the same distribution family as the **prior**, we have **conjugacy**.

Conjugate models are great because we can find the exact **posterior**,

In the previous model, we had

$$\text{Beta}(y + \alpha, n - y + \beta) = \frac{\text{Binomial}(n, \pi) \times \text{Beta}(\alpha, \beta)}{p(y)}$$

We knew that the likelihood  $\times$  **prior** produced something that looked like a Beta distribution up to a **constant** of proportionality.

Since the **posterior** must be a probability distribution, we know that it is a Beta distribution and we can easily solve for the **normalizing constant** (although we don't need to since we already have the **posterior**).

When the **posterior** is the same distribution family as the **prior**, we have **conjugacy**.

Conjugate models are great because we can find the exact **posterior**, but we almost never have conjugacy except in very simple models.

# What Happens When We Don't Have Conjugacy?

# What Happens When We Don't Have Conjugacy?

Consider a Poisson regression model with Normal priors on  $\beta$ .

# What Happens When We Don't Have Conjugacy?

Consider a Poisson regression model with Normal priors on  $\beta$ .

$$p(\beta|\mathbf{y}) \propto \prod_{i=1}^n \text{Poisson}(\lambda_i) \times \text{Normal}(\mu, \Sigma)$$

# What Happens When We Don't Have Conjugacy?

Consider a Poisson regression model with Normal priors on  $\beta$ .

$$p(\beta|\mathbf{y}) \propto \prod_{i=1}^n \text{Poisson}(\lambda_i) \times \text{Normal}(\mu, \Sigma)$$
$$\lambda_i = \exp(\mathbf{x}_i\beta)$$

# What Happens When We Don't Have Conjugacy?

Consider a Poisson regression model with Normal priors on  $\beta$ .

$$p(\beta|\mathbf{y}) \propto \prod_{i=1}^n \text{Poisson}(\lambda_i) \times \text{Normal}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$
$$\lambda_i = \exp(\mathbf{x}_i\boldsymbol{\beta})$$

$$p(\beta|\mathbf{y}) \propto \prod_{i=1}^n \frac{\exp(-e^{\mathbf{x}_i\boldsymbol{\beta}}) \exp(\mathbf{x}_i\boldsymbol{\beta})^{y_i}}{y_i!} \times$$
$$\frac{1}{(2\pi)^{d/2}|\boldsymbol{\Sigma}|^{1/2}} \exp\left(\frac{1}{2}(\boldsymbol{\beta} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\boldsymbol{\beta} - \boldsymbol{\mu})\right)$$



# What Happens When We Don't Have Conjugacy?

Consider a Poisson regression model with Normal priors on  $\beta$ .

$$p(\beta|\mathbf{y}) \propto \prod_{i=1}^n \text{Poisson}(\lambda_i) \times \text{Normal}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$
$$\lambda_i = \exp(\mathbf{x}_i\beta)$$

$$p(\beta|\mathbf{y}) \propto \prod_{i=1}^n \frac{\exp(-e^{\mathbf{x}_i\beta}) \exp(\mathbf{x}_i\beta)^{y_i}}{y_i!} \times$$
$$\frac{1}{(2\pi)^{d/2} |\boldsymbol{\Sigma}|^{1/2}} \exp\left(\frac{1}{2}(\beta - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\beta - \boldsymbol{\mu})\right)$$

Likelihood  $\times$  prior doesn't look like any distribution we know (non-conjugacy)

# What Happens When We Don't Have Conjugacy?

Consider a Poisson regression model with Normal priors on  $\beta$ .

$$p(\beta|\mathbf{y}) \propto \prod_{i=1}^n \text{Poisson}(\lambda_i) \times \text{Normal}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$
$$\lambda_i = \exp(\mathbf{x}_i\boldsymbol{\beta})$$

$$p(\beta|\mathbf{y}) \propto \prod_{i=1}^n \frac{\exp(-e^{\mathbf{x}_i\boldsymbol{\beta}}) \exp(\mathbf{x}_i\boldsymbol{\beta})^{y_i}}{y_i!} \times$$
$$\frac{1}{(2\pi)^{d/2}|\boldsymbol{\Sigma}|^{1/2}} \exp\left(\frac{1}{2}(\boldsymbol{\beta} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\boldsymbol{\beta} - \boldsymbol{\mu})\right)$$

Likelihood  $\times$  prior doesn't look like any distribution we know (non-conjugacy) and normalizing constant is too hard to find,

# What Happens When We Don't Have Conjugacy?

Consider a Poisson regression model with Normal priors on  $\beta$ .

$$p(\beta|\mathbf{y}) \propto \prod_{i=1}^n \text{Poisson}(\lambda_i) \times \text{Normal}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$
$$\lambda_i = \exp(\mathbf{x}_i\beta)$$

$$p(\beta|\mathbf{y}) \propto \prod_{i=1}^n \frac{\exp(-e^{\mathbf{x}_i\beta}) \exp(\mathbf{x}_i\beta)^{y_i}}{y_i!} \times$$
$$\frac{1}{(2\pi)^{d/2} |\boldsymbol{\Sigma}|^{1/2}} \exp\left(-\frac{1}{2}(\beta - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\beta - \boldsymbol{\mu})\right)$$

Likelihood  $\times$  prior doesn't look like any distribution we know (non-conjugacy) and normalizing constant is too hard to find, so how do we find our posterior?

# MCMC to the Rescue

# MCMC to the Rescue

Ideal Goal:

# MCMC to the Rescue

Ideal Goal: Produce **independent** draws from our **posterior** distribution via simulation and summarize the **posterior** by using those draws.

# MCMC to the Rescue

Ideal Goal: Produce **independent** draws from our **posterior** distribution via simulation and summarize the **posterior** by using those draws.

**Markov Chain Monte Carlo (MCMC)**: a class of algorithms that produce a **chain** of **simulated draws** from a distribution where **each** draw is **dependent** on the previous draw.

# MCMC to the Rescue

Ideal Goal: Produce **independent** draws from our **posterior** distribution via simulation and summarize the **posterior** by using those draws.

**Markov Chain Monte Carlo (MCMC)**: a class of algorithms that produce a **chain** of **simulated draws** from a distribution where **each draw is dependent** on the previous draw.

Theory:



# MCMC to the Rescue

Ideal Goal: Produce **independent** draws from our **posterior** distribution via simulation and summarize the **posterior** by using those draws.

**Markov Chain Monte Carlo (MCMC)**: a class of algorithms that produce a **chain** of **simulated draws** from a distribution where **each draw is dependent on the previous draw**.

Theory: If our chain satisfies some basic conditions, then the chain will **eventually converge** to a stationary distribution (in our case, the **posterior**)

# MCMC to the Rescue

Ideal Goal: Produce **independent** draws from our **posterior** distribution via simulation and summarize the **posterior** by using those draws.

**Markov Chain Monte Carlo (MCMC)**: a class of algorithms that produce a **chain** of **simulated draws** from a distribution where **each draw is dependent on the previous draw**.

Theory: If our chain satisfies some basic conditions, then the chain will **eventually converge** to a stationary distribution (in our case, the **posterior**) and we have approximate draws from the **posterior**.

# MCMC to the Rescue

Ideal Goal: Produce **independent** draws from our **posterior** distribution via simulation and summarize the **posterior** by using those draws.

**Markov Chain Monte Carlo (MCMC)**: a class of algorithms that produce a **chain** of **simulated draws** from a distribution where **each draw is dependent on the previous draw**.

Theory: If our chain satisfies some basic conditions, then the chain will **eventually converge** to a stationary distribution (in our case, the **posterior**) and we have approximate draws from the **posterior**.

*But there is no way to know for sure whether our chain has converged.*

# Algorithm 1: Gibbs Sampler

# Algorithm 1: Gibbs Sampler

Let  $\boldsymbol{\theta}^t = (\theta_1^t, \dots, \theta_k^t)$  be the  $t$ th draw of our parameter vector  $\boldsymbol{\theta}$ .

## Algorithm 1: Gibbs Sampler

Let  $\boldsymbol{\theta}^t = (\theta_1^t, \dots, \theta_k^t)$  be the  $t$ th draw of our parameter vector  $\boldsymbol{\theta}$ .

Draw a new vector  $\boldsymbol{\theta}^{t+1}$  from the following distributions:

# Algorithm 1: Gibbs Sampler

Let  $\boldsymbol{\theta}^t = (\theta_1^t, \dots, \theta_k^t)$  be the  $t$ th draw of our parameter vector  $\boldsymbol{\theta}$ .

Draw a new vector  $\boldsymbol{\theta}^{t+1}$  from the following distributions:

$$\theta_1^{t+1} \sim p(\theta_1 | \theta_2^t, \dots, \theta_k^t, \mathbf{y})$$

# Algorithm 1: Gibbs Sampler

Let  $\boldsymbol{\theta}^t = (\theta_1^t, \dots, \theta_k^t)$  be the  $t$ th draw of our parameter vector  $\boldsymbol{\theta}$ .

Draw a new vector  $\boldsymbol{\theta}^{t+1}$  from the following distributions:

$$\theta_1^{t+1} \sim p(\theta_1 | \theta_2^t, \dots, \theta_k^t, \mathbf{y})$$

$$\theta_2^{t+1} \sim p(\theta_2 | \theta_1^{t+1}, \dots, \theta_k^t, \mathbf{y})$$



# Algorithm 1: Gibbs Sampler

Let  $\boldsymbol{\theta}^t = (\theta_1^t, \dots, \theta_k^t)$  be the  $t$ th draw of our parameter vector  $\boldsymbol{\theta}$ .

Draw a new vector  $\boldsymbol{\theta}^{t+1}$  from the following distributions:

$$\begin{aligned}\theta_1^{t+1} &\sim p(\theta_1 | \theta_2^t, \dots, \theta_k^t, \mathbf{y}) \\ \theta_2^{t+1} &\sim p(\theta_2 | \theta_1^{t+1}, \dots, \theta_k^t, \mathbf{y}) \\ &\vdots\end{aligned}$$

# Algorithm 1: Gibbs Sampler

Let  $\boldsymbol{\theta}^t = (\theta_1^t, \dots, \theta_k^t)$  be the  $t$ th draw of our parameter vector  $\boldsymbol{\theta}$ .

Draw a new vector  $\boldsymbol{\theta}^{t+1}$  from the following distributions:

$$\theta_1^{t+1} \sim p(\theta_1 | \theta_2^t, \dots, \theta_k^t, \mathbf{y})$$

$$\theta_2^{t+1} \sim p(\theta_2 | \theta_1^{t+1}, \dots, \theta_k^t, \mathbf{y})$$

$$\vdots$$

$$\theta_k^{t+1} \sim p(\theta_k | \theta_1^{t+1}, \dots, \theta_{k-1}^{t+1}, \mathbf{y})$$

# Algorithm 1: Gibbs Sampler

Let  $\boldsymbol{\theta}^t = (\theta_1^t, \dots, \theta_k^t)$  be the  $t$ th draw of our parameter vector  $\boldsymbol{\theta}$ .

Draw a new vector  $\boldsymbol{\theta}^{t+1}$  from the following distributions:

$$\begin{aligned}\theta_1^{t+1} &\sim p(\theta_1 | \theta_2^t, \dots, \theta_k^t, \mathbf{y}) \\ \theta_2^{t+1} &\sim p(\theta_2 | \theta_1^{t+1}, \dots, \theta_k^t, \mathbf{y}) \\ &\vdots \\ \theta_k^{t+1} &\sim p(\theta_k | \theta_1^{t+1}, \dots, \theta_{k-1}^{t+1}, \mathbf{y})\end{aligned}$$

Repeat  $m$  times to get  $m$  draws of our parameters from the approximate **posterior** (assuming convergence).

# Algorithm 1: Gibbs Sampler

Let  $\theta^t = (\theta_1^t, \dots, \theta_k^t)$  be the  $t$ th draw of our parameter vector  $\theta$ .

Draw a new vector  $\theta^{t+1}$  from the following distributions:

$$\begin{aligned}\theta_1^{t+1} &\sim p(\theta_1 | \theta_2^t, \dots, \theta_k^t, \mathbf{y}) \\ \theta_2^{t+1} &\sim p(\theta_2 | \theta_1^{t+1}, \dots, \theta_k^t, \mathbf{y}) \\ &\vdots \\ \theta_k^{t+1} &\sim p(\theta_k | \theta_1^{t+1}, \dots, \theta_{k-1}^{t+1}, \mathbf{y})\end{aligned}$$

Repeat  $m$  times to get  $m$  draws of our parameters from the approximate **posterior** (assuming convergence).

Requires that we know the conditional distributions for each  $\theta$ .

# Algorithm 1: Gibbs Sampler

Let  $\theta^t = (\theta_1^t, \dots, \theta_k^t)$  be the  $t$ th draw of our parameter vector  $\theta$ .

Draw a new vector  $\theta^{t+1}$  from the following distributions:

$$\begin{aligned}\theta_1^{t+1} &\sim p(\theta_1 | \theta_2^t, \dots, \theta_k^t, \mathbf{y}) \\ \theta_2^{t+1} &\sim p(\theta_2 | \theta_1^{t+1}, \dots, \theta_k^t, \mathbf{y}) \\ &\vdots \\ \theta_k^{t+1} &\sim p(\theta_k | \theta_1^{t+1}, \dots, \theta_{k-1}^{t+1}, \mathbf{y})\end{aligned}$$

Repeat  $m$  times to get  $m$  draws of our parameters from the approximate **posterior** (assuming convergence).

Requires that we know the conditional distributions for each  $\theta$ .  
What if we don't?

## Algorithm 2: Metropolis-Hastings

## Algorithm 2: Metropolis-Hastings

Draw a new vector  $\theta^{t+1}$  in the following way:

## Algorithm 2: Metropolis-Hastings

Draw a new vector  $\boldsymbol{\theta}^{t+1}$  in the following way:

1. Specify a jumping distribution  $J_{t+1}(\boldsymbol{\theta}^* | \boldsymbol{\theta}^t)$



## Algorithm 2: Metropolis-Hastings

Draw a new vector  $\theta^{t+1}$  in the following way:

1. Specify a jumping distribution  $J_{t+1}(\theta^*|\theta^t)$  (usually a symmetric distribution such as the multivariate normal).

## Algorithm 2: Metropolis-Hastings

Draw a new vector  $\theta^{t+1}$  in the following way:

1. Specify a jumping distribution  $J_{t+1}(\theta^*|\theta^t)$  (usually a symmetric distribution such as the multivariate normal).
2. Draw a proposed parameter vector  $\theta^*$  from the jumping distribution.

## Algorithm 2: Metropolis-Hastings

Draw a new vector  $\theta^{t+1}$  in the following way:

1. Specify a jumping distribution  $J_{t+1}(\theta^*|\theta^t)$  (usually a symmetric distribution such as the multivariate normal).
2. Draw a proposed parameter vector  $\theta^*$  from the jumping distribution.
3. Accept  $\theta^*$  as  $\theta^{t+1}$  with probability  $\min(r, 1)$ , where

## Algorithm 2: Metropolis-Hastings

Draw a new vector  $\theta^{t+1}$  in the following way:

1. Specify a jumping distribution  $J_{t+1}(\theta^*|\theta^t)$  (usually a symmetric distribution such as the multivariate normal).
2. Draw a proposed parameter vector  $\theta^*$  from the jumping distribution.
3. Accept  $\theta^*$  as  $\theta^{t+1}$  with probability  $\min(r, 1)$ , where

$$r = \frac{p(\mathbf{y}|\theta^*)p(\theta^*)}{p(\mathbf{y}|\theta^t)p(\theta^t)}$$

## Algorithm 2: Metropolis-Hastings

Draw a new vector  $\theta^{t+1}$  in the following way:

1. Specify a jumping distribution  $J_{t+1}(\theta^*|\theta^t)$  (usually a symmetric distribution such as the multivariate normal).
2. Draw a proposed parameter vector  $\theta^*$  from the jumping distribution.
3. Accept  $\theta^*$  as  $\theta^{t+1}$  with probability  $\min(r, 1)$ , where

$$r = \frac{p(\mathbf{y}|\theta^*)p(\theta^*)}{p(\mathbf{y}|\theta^t)p(\theta^t)}$$

If  $\theta^*$  is rejected, then  $\theta^{t+1} = \theta^t$ .

## Algorithm 2: Metropolis-Hastings

Draw a new vector  $\theta^{t+1}$  in the following way:

1. Specify a jumping distribution  $J_{t+1}(\theta^*|\theta^t)$  (usually a symmetric distribution such as the multivariate normal).
2. Draw a proposed parameter vector  $\theta^*$  from the jumping distribution.
3. Accept  $\theta^*$  as  $\theta^{t+1}$  with probability  $\min(r, 1)$ , where

$$r = \frac{p(\mathbf{y}|\theta^*)p(\theta^*)}{p(\mathbf{y}|\theta^t)p(\theta^t)}$$

If  $\theta^*$  is rejected, then  $\theta^{t+1} = \theta^t$ .

Repeat  $m$  times to get  $m$  draws of our parameters from the approximate **posterior** (assuming convergence).

## Algorithm 2: Metropolis-Hastings

Draw a new vector  $\theta^{t+1}$  in the following way:

1. Specify a jumping distribution  $J_{t+1}(\theta^*|\theta^t)$  (usually a symmetric distribution such as the multivariate normal).
2. Draw a proposed parameter vector  $\theta^*$  from the jumping distribution.
3. Accept  $\theta^*$  as  $\theta^{t+1}$  with probability  $\min(r, 1)$ , where

$$r = \frac{p(\mathbf{y}|\theta^*)p(\theta^*)}{p(\mathbf{y}|\theta^t)p(\theta^t)}$$

If  $\theta^*$  is rejected, then  $\theta^{t+1} = \theta^t$ .

Repeat  $m$  times to get  $m$  draws of our parameters from the approximate **posterior** (assuming convergence).

M-H always works, but can be very slow.

# Outline

Introduction

Bayesian Models

Applications

Missing Data

Hierarchical Models



# Missing Data

# Missing Data

Suppose we have missing data.

# Missing Data

Suppose we have missing data. Define  $D$  as our data matrix and  $M$  as our missingness matrix.

# Missing Data

Suppose we have missing data. Define  $D$  as our data matrix and  $M$  as our missingness matrix.

$$D = \begin{pmatrix} y & x_1 & x_2 & x_3 \\ 1 & 2.5 & 432 & 0 \\ 5 & 3.2 & 543 & 1 \\ 2 & ? & 219 & ? \\ ? & 1.9 & ? & 1 \\ ? & 1.2 & 108 & 0 \\ ? & 7.7 & 95 & 1 \end{pmatrix}$$

$$M = \begin{pmatrix} y & x_1 & x_2 & x_3 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 \end{pmatrix}$$

# Missing Data

Suppose we have missing data. Define  $D$  as our data matrix and  $M$  as our missingness matrix.

$$D = \begin{pmatrix} y & x_1 & x_2 & x_3 \\ 1 & 2.5 & 432 & 0 \\ 5 & 3.2 & 543 & 1 \\ 2 & ? & 219 & ? \\ ? & 1.9 & ? & 1 \\ ? & 1.2 & 108 & 0 \\ ? & 7.7 & 95 & 1 \end{pmatrix}$$

$$M = \begin{pmatrix} y & x_1 & x_2 & x_3 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 \end{pmatrix}$$

One non-Bayesian approach to dealing with missing data is **multiple imputation**.

# Multiple Imputation

# Multiple Imputation

We need a method for filling in the missing cells of  $D$  as a first step **before** we go to the analysis stage.

# Multiple Imputation

We need a method for filling in the missing cells of  $D$  as a first step **before** we go to the analysis stage.

1. Assume a joint distribution for  $D_{obs}$  and  $M$ :



# Multiple Imputation

We need a method for filling in the missing cells of  $D$  as a first step **before** we go to the analysis stage.

1. Assume a joint distribution for  $D_{obs}$  and  $M$ :

$$p(D_{obs}, M | \phi, \gamma) = \int p(D | \phi) p(M | D_{obs}, D_{mis}, \gamma) dD_{mis}$$

# Multiple Imputation

We need a method for filling in the missing cells of  $D$  as a first step **before** we go to the analysis stage.

1. Assume a joint distribution for  $D_{obs}$  and  $M$ :

$$p(D_{obs}, M|\phi, \gamma) = \int p(D|\phi)p(M|D_{obs}, D_{mis}, \gamma)dD_{mis}$$

If we assume MAR, then

$$p(D_{obs}, M|\phi, \gamma) \propto \int p(D|\phi)dD_{mis}$$

2. Find  $\phi$ , which characterizes the full  $D$ .

2. Find  $\phi$ , which characterizes the full  $D$ .

$$L(\phi|D_{obs}) = \prod_{i=1}^n \int p(D_{i,obs}|\phi) dD_{mis}$$

2. Find  $\phi$ , which characterizes the full  $D$ .

$$L(\phi|D_{obs}) = \prod_{i=1}^n \int p(D_{i,obs}|\phi) dD_{mis}$$

How do we do this integral?

2. Find  $\phi$ , which characterizes the full  $D$ .

$$L(\phi|D_{obs}) = \prod_{i=1}^n \int p(D_{i,obs}|\phi) dD_{mis}$$

How do we do this integral?

Assume Normality since the marginals of a multivariate Normal are Normal.

2. Find  $\phi$ , which characterizes the full  $D$ .

$$L(\phi|D_{obs}) = \prod_{i=1}^n \int p(D_{i,obs}|\phi) dD_{mis}$$

How do we do this integral?

Assume Normality since the marginals of a multivariate Normal are Normal.

$$L(\mu, \Sigma|D_{obs}) = \prod_{i=1}^n N(D_{i,obs}|\mu_{obs}, \Sigma_{obs})$$

2. Find  $\phi$ , which characterizes the full  $D$ .

$$L(\phi|D_{obs}) = \prod_{i=1}^n \int p(D_{i,obs}|\phi) dD_{mis}$$

How do we do this integral?

Assume Normality since the marginals of a multivariate Normal are Normal.

$$L(\mu, \Sigma|D_{obs}) = \prod_{i=1}^n N(D_{i,obs}|\mu_{obs}, \Sigma_{obs})$$

3. Find  $\hat{\mu}, \hat{\Sigma}$  and its distribution via EM algorithm and bootstrapping.



2. Find  $\phi$ , which characterizes the full  $D$ .

$$L(\phi|D_{obs}) = \prod_{i=1}^n \int p(D_{i,obs}|\phi) dD_{mis}$$

How do we do this integral?

Assume Normality since the marginals of a multivariate Normal are Normal.

$$L(\mu, \Sigma|D_{obs}) = \prod_{i=1}^n N(D_{i,obs}|\mu_{obs}, \Sigma_{obs})$$

3. Find  $\hat{\mu}, \hat{\Sigma}$  and its distribution via EM algorithm and bootstrapping.
4. Draw  $m$   $\mu, \Sigma$  values, then use them to predict values for  $D_{mis}$ .

What we end up with is  $m$  datasets with missing values imputed.

What we end up with is  $m$  datasets with missing values imputed.

We then run our regular analyses on the  $m$  datasets and combine the results using Rubin's rule.

# Bayesian Approach to Missing Data

# Bayesian Approach to Missing Data

Bayesian paradigm: Everything unobserved is a random variable.

# Bayesian Approach to Missing Data

Bayesian paradigm: Everything unobserved is a random variable.

So we can set up the missing data as a “parameter” that we need to find.

# Bayesian Approach to Missing Data

Bayesian paradigm: Everything unobserved is a random variable.

So we can set up the missing data as a “parameter” that we need to find.

$$p(D_{mis}, \phi, \gamma | D_{obs}, M) \propto p(D_{obs} | D_{mis}, \phi) p(D_{mis} | \phi) p(M | D_{obs}, D_{mis}, \gamma) \\ p(\gamma) p(\phi)$$

# Bayesian Approach to Missing Data

Bayesian paradigm: Everything unobserved is a random variable.

So we can set up the missing data as a “parameter” that we need to find.

$$p(D_{mis}, \phi, \gamma | D_{obs}, M) \propto p(D_{obs} | D_{mis}, \phi) p(D_{mis} | \phi) p(M | D_{obs}, D_{mis}, \gamma) \\ p(\gamma) p(\phi)$$

If we assume MAR:



# Bayesian Approach to Missing Data

Bayesian paradigm: Everything unobserved is a random variable.

So we can set up the missing data as a “parameter” that we need to find.

$$p(D_{mis}, \phi, \gamma | D_{obs}, M) \propto p(D_{obs} | D_{mis}, \phi) p(D_{mis} | \phi) p(M | D_{obs}, D_{mis}, \gamma) \\ p(\gamma) p(\phi)$$

If we assume MAR:

$$p(D_{mis}, \phi, \gamma | D_{obs}, M) \propto p(D_{obs} | D_{mis}, \phi) p(D_{mis} | \phi) p(\phi)$$

# Bayesian Approach to Missing Data

Bayesian paradigm: Everything unobserved is a random variable.

So we can set up the missing data as a “parameter” that we need to find.

$$p(D_{mis}, \phi, \gamma | D_{obs}, M) \propto p(D_{obs} | D_{mis}, \phi) p(D_{mis} | \phi) p(M | D_{obs}, D_{mis}, \gamma) \\ p(\gamma) p(\phi)$$

If we assume MAR:

$$p(D_{mis}, \phi, \gamma | D_{obs}, M) \propto p(D_{obs} | D_{mis}, \phi) p(D_{mis} | \phi) p(\phi)$$

Use Gibbs Sampling or M-H to sample both  $D_{mis}$  and  $\phi$ .

# Bayesian Approach to Missing Data

Bayesian paradigm: Everything unobserved is a random variable.

So we can set up the missing data as a “parameter” that we need to find.

$$p(D_{mis}, \phi, \gamma | D_{obs}, M) \propto p(D_{obs} | D_{mis}, \phi) p(D_{mis} | \phi) p(M | D_{obs}, D_{mis}, \gamma) \\ p(\gamma) p(\phi)$$

If we assume MAR:

$$p(D_{mis}, \phi, \gamma | D_{obs}, M) \propto p(D_{obs} | D_{mis}, \phi) p(D_{mis} | \phi) p(\phi)$$

Use Gibbs Sampling or M-H to sample both  $D_{mis}$  and  $\phi$ .

We don't have to assume normality of  $D$  to integrate over  $D_{mis}$ .

# Bayesian Approach to Missing Data

Bayesian paradigm: Everything unobserved is a random variable.

So we can set up the missing data as a “parameter” that we need to find.

$$p(D_{mis}, \phi, \gamma | D_{obs}, M) \propto p(D_{obs} | D_{mis}, \phi) p(D_{mis} | \phi) p(M | D_{obs}, D_{mis}, \gamma) \\ p(\gamma) p(\phi)$$

If we assume MAR:

$$p(D_{mis}, \phi, \gamma | D_{obs}, M) \propto p(D_{obs} | D_{mis}, \phi) p(D_{mis} | \phi) p(\phi)$$

Use Gibbs Sampling or M-H to sample both  $D_{mis}$  and  $\phi$ .

We don't have to assume normality of  $D$  to integrate over  $D_{mis}$ .  
We can just drop the draws of  $D_{mis}$ .

We can also incorporate both imputation and analyses in the same model.

$$p(\theta, D_{mis}, \phi, \gamma | D_{obs}, M) \propto p(\mathbf{y}_{obs}, \mathbf{y}_{mis} | \theta) p(D_{obs} | D_{mis}, \phi) p(D_{mis} | \phi) \\ p(\phi) p(\theta)$$

We can also incorporate both imputation and analyses in the same model.

$$p(\theta, D_{mis}, \phi, \gamma | D_{obs}, M) \propto p(\mathbf{y}_{obs}, \mathbf{y}_{mis} | \theta) p(D_{obs} | D_{mis}, \phi) p(D_{mis} | \phi) \\ p(\phi) p(\theta)$$

Again, find the **posterior** via Gibbs Sampling or M-H.

We can also incorporate both imputation and analyses in the same model.

$$p(\theta, D_{mis}, \phi, \gamma | D_{obs}, M) \propto p(\mathbf{y}_{obs}, \mathbf{y}_{mis} | \theta) p(D_{obs} | D_{mis}, \phi) p(D_{mis} | \phi) \\ p(\phi) p(\theta)$$

Again, find the **posterior** via Gibbs Sampling or M-H.

Moral: We can easily set up an application specific Bayesian model to incorporate missing data.

# Multilevel Data



# Multilevel Data

Suppose we have data in which we have  $i = 1, \dots, n$  observations that belong to one of  $j = 1, \dots, J$  groups.

# Multilevel Data

Suppose we have data in which we have  $i = 1, \dots, n$  observations that belong to one of  $j = 1, \dots, J$  groups.

Examples:

# Multilevel Data

Suppose we have data in which we have  $i = 1, \dots, n$  observations that belong to one of  $j = 1, \dots, J$  groups.

Examples:

- ▶ students within schools

# Multilevel Data

Suppose we have data in which we have  $i = 1, \dots, n$  observations that belong to one of  $j = 1, \dots, J$  groups.

Examples:

- ▶ students within schools
- ▶ multiple observations per country

# Multilevel Data

Suppose we have data in which we have  $i = 1, \dots, n$  observations that belong to one of  $j = 1, \dots, J$  groups.

Examples:

- ▶ students within schools
- ▶ multiple observations per country
- ▶ districts within states

# Multilevel Data

Suppose we have data in which we have  $i = 1, \dots, n$  observations that belong to one of  $j = 1, \dots, J$  groups.

Examples:

- ▶ students within schools
- ▶ multiple observations per country
- ▶ districts within states

We can have covariates on multiple levels.

# Multilevel Data

Suppose we have data in which we have  $i = 1, \dots, n$  observations that belong to one of  $j = 1, \dots, J$  groups.

Examples:

- ▶ students within schools
- ▶ multiple observations per country
- ▶ districts within states

We can have covariates on multiple levels. How do we deal with this type of data?

# The Fixed Effects Model



# The Fixed Effects Model

We can let the intercept  $\alpha$  vary by group:

# The Fixed Effects Model

We can let the intercept  $\alpha$  vary by group:

$$y_i = \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i$$

# The Fixed Effects Model

We can let the intercept  $\alpha$  vary by group:

$$y_i = \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i$$

This is known as the **fixed effects** model.

# The Fixed Effects Model

We can let the intercept  $\alpha$  vary by group:

$$y_i = \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i$$

This is known as the **fixed effects** model.

This is equivalent to estimating dummy variables for  $J - 1$  groups.

# The Fixed Effects Model

We can let the intercept  $\alpha$  vary by group:

$$y_i = \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i$$

This is known as the **fixed effects** model.

This is equivalent to estimating dummy variables for  $J - 1$  groups.

We can use this model if we think that there is something inherent about the groups that affects our dependent variable.

# The Fixed Effects Model

We can let the intercept  $\alpha$  vary by group:

$$y_i = \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i$$

This is known as the **fixed effects** model.

This is equivalent to estimating dummy variables for  $J - 1$  groups.

We can use this model if we think that there is something inherent about the groups that affects our dependent variable.

However, the fixed effects model involves estimating many parameters, and also cannot take into account group-level covariates.

# Hierarchical Model

# Hierarchical Model

A more flexible alternative is to use a **hierarchical model**, also known as a multilevel model, mixed effects model, or random effects model.



# Hierarchical Model

A more flexible alternative is to use a **hierarchical model**, also known as a multilevel model, mixed effects model, or random effects model.

$$y_i = \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i$$

# Hierarchical Model

A more flexible alternative is to use a **hierarchical model**, also known as a multilevel model, mixed effects model, or random effects model.

$$\begin{aligned}y_i &= \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i \\ \alpha_j &\sim N(\alpha, \sigma_\alpha^2)\end{aligned}$$

# Hierarchical Model

A more flexible alternative is to use a **hierarchical model**, also known as a multilevel model, mixed effects model, or random effects model.

$$\begin{aligned}y_i &= \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i \\ \alpha_j &\sim N(\alpha, \sigma_\alpha^2)\end{aligned}$$

Instead of assuming a completely different intercept for each group, we can assume that the intercepts are drawn from a common (Normal) distribution.

We can incorporate group-level covariates in the following way:

We can incorporate group-level covariates in the following way:

$$y_i = \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i$$

We can incorporate group-level covariates in the following way:

$$\begin{aligned}y_i &= \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i \\ \alpha_j &\sim N(\gamma_0 + u_{j1}\gamma_1, \sigma_\alpha^2)\end{aligned}$$

We can incorporate group-level covariates in the following way:

$$\begin{aligned}y_i &= \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i \\ \alpha_j &\sim N(\gamma_0 + u_{j1}\gamma_1, \sigma_\alpha^2)\end{aligned}$$

or equivalently

We can incorporate group-level covariates in the following way:

$$\begin{aligned}y_i &= \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i \\ \alpha_j &\sim N(\gamma_0 + u_{j1}\gamma_1, \sigma_\alpha^2)\end{aligned}$$

or equivalently

$$y_i = \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i$$



We can incorporate group-level covariates in the following way:

$$\begin{aligned}y_i &= \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i \\ \alpha_j &\sim N(\gamma_0 + u_{j1}\gamma_1, \sigma_\alpha^2)\end{aligned}$$

or equivalently

$$\begin{aligned}y_i &= \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i \\ \alpha_j &= \gamma_0 + u_{j1}\gamma_1 + \eta_j\end{aligned}$$

We can incorporate group-level covariates in the following way:

$$\begin{aligned}y_i &= \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i \\ \alpha_j &\sim N(\gamma_0 + u_{j1}\gamma_1, \sigma_\alpha^2)\end{aligned}$$

or equivalently

$$\begin{aligned}y_i &= \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i \\ \alpha_j &= \gamma_0 + u_{j1}\gamma_1 + \eta_j \\ \eta_j &\sim N(0, \sigma_\alpha^2)\end{aligned}$$

We can incorporate group-level covariates in the following way:

$$\begin{aligned}y_i &= \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i \\ \alpha_j &\sim N(\gamma_0 + u_{j1}\gamma_1, \sigma_\alpha^2)\end{aligned}$$

or equivalently

$$\begin{aligned}y_i &= \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i \\ \alpha_j &= \gamma_0 + u_{j1}\gamma_1 + \eta_j \\ \eta_j &\sim N(0, \sigma_\alpha^2)\end{aligned}$$

This is a relatively difficult model to estimate using non-Bayesian methods. The `lme4()` package in R can do it.

# Bayesian Hierarchical Model

# Bayesian Hierarchical Model

$$y_i = \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i$$

# Bayesian Hierarchical Model

$$\begin{aligned}y_i &= \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i \\ \alpha_j &\sim N(\gamma_0 + u_{j1}\gamma_1, \sigma_\alpha^2)\end{aligned}$$

# Bayesian Hierarchical Model

$$\begin{aligned}y_i &= \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i \\ \alpha_j &\sim N(\gamma_0 + u_{j1}\gamma_1, \sigma_\alpha^2)\end{aligned}$$

We can do hierarchical models easily using Bayesian methods.

# Bayesian Hierarchical Model

$$\begin{aligned}y_i &= \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i \\ \alpha_j &\sim N(\gamma_0 + u_{j1}\gamma_1, \sigma_\alpha^2)\end{aligned}$$

We can do hierarchical models easily using Bayesian methods.

$$p(\alpha, \beta, \gamma | \mathbf{y}) \propto p(\mathbf{y} | \alpha, \beta, \gamma) p(\alpha | \gamma) p(\gamma) p(\beta)$$



# Bayesian Hierarchical Model

$$\begin{aligned}y_i &= \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i \\ \alpha_j &\sim N(\gamma_0 + u_{j1}\gamma_1, \sigma_\alpha^2)\end{aligned}$$

We can do hierarchical models easily using Bayesian methods.

$$p(\alpha, \beta, \gamma | \mathbf{y}) \propto p(\mathbf{y} | \alpha, \beta, \gamma) p(\alpha | \gamma) p(\gamma) p(\beta)$$

Solve for the joint **posterior** using Gibbs Sampling or M-H.

# Bayesian Hierarchical Model

$$\begin{aligned}y_i &= \alpha_{j[i]} + x_{i1}\beta_1 + x_{i2}\beta_2 + \epsilon_i \\ \alpha_j &\sim N(\gamma_0 + u_{j1}\gamma_1, \sigma_\alpha^2)\end{aligned}$$

We can do hierarchical models easily using Bayesian methods.

$$p(\alpha, \beta, \gamma | \mathbf{y}) \propto p(\mathbf{y} | \alpha, \beta, \gamma) p(\alpha | \gamma) p(\gamma) p(\beta)$$

Solve for the joint **posterior** using Gibbs Sampling or M-H.

We incorporate data with more than two levels easily as well.

# Conclusion

# Conclusion

There are pros and cons to using Bayesian statistics.

# Conclusion

There are pros and cons to using Bayesian statistics.

Pros:

# Conclusion

There are pros and cons to using Bayesian statistics.

Pros:

- ▶ Incorporate outside/prior knowledge

# Conclusion

There are pros and cons to using Bayesian statistics.

Pros:

- ▶ Incorporate outside/prior knowledge
- ▶ Estimate much more difficult models

# Conclusion

There are pros and cons to using Bayesian statistics.

Pros:

- ▶ Incorporate outside/prior knowledge
- ▶ Estimate much more difficult models
- ▶ CI have more intuitive meaning



# Conclusion

There are pros and cons to using Bayesian statistics.

Pros:

- ▶ Incorporate outside/prior knowledge
- ▶ Estimate much more difficult models
- ▶ CI have more intuitive meaning
- ▶ Helps with unidentified models

# Conclusion

There are pros and cons to using Bayesian statistics.

Pros:

- ▶ Incorporate outside/prior knowledge
- ▶ Estimate much more difficult models
- ▶ CI have more intuitive meaning
- ▶ Helps with unidentified models

Cons:

# Conclusion

There are pros and cons to using Bayesian statistics.

## Pros:

- ▶ Incorporate outside/prior knowledge
- ▶ Estimate much more difficult models
- ▶ CI have more intuitive meaning
- ▶ Helps with unidentified models

## Cons:

- ▶ It's hard(er)

# Conclusion

There are pros and cons to using Bayesian statistics.

## Pros:

- ▶ Incorporate outside/prior knowledge
- ▶ Estimate much more difficult models
- ▶ CI have more intuitive meaning
- ▶ Helps with unidentified models

## Cons:

- ▶ It's hard(er)
- ▶ Computationally intensive

# Conclusion

There are pros and cons to using Bayesian statistics.

## Pros:

- ▶ Incorporate outside/prior knowledge
- ▶ Estimate much more difficult models
- ▶ CI have more intuitive meaning
- ▶ Helps with unidentified models

## Cons:

- ▶ It's hard(er)
- ▶ Computationally intensive
- ▶ Need defense of priors

# Conclusion

There are pros and cons to using Bayesian statistics.

## Pros:

- ▶ Incorporate outside/prior knowledge
- ▶ Estimate much more difficult models
- ▶ CI have more intuitive meaning
- ▶ Helps with unidentified models

## Cons:

- ▶ It's hard(er)
- ▶ Computationally intensive
- ▶ Need defense of priors
- ▶ No guarantee of MCMC convergence

Statistical packages for Bayesian are also less developed (MCMCpack() in R, WinBUGS, JAGS).