

Probability Theory

Patrick Lam

Outline

Probability

Random Variables

Simulation

Important Distributions

- Discrete Distributions

- Continuous Distributions

Most Basic Definition of Probability:

Most Basic Definition of Probability:

$$\frac{\text{number of successes}}{\text{number of possible occurrences}}$$

Three Axioms of Probability

Three Axioms of Probability

Let S be the sample space and A be an event in S .

Three Axioms of Probability

Let S be the sample space and A be an event in S .

1. For any event A , $P(A) \geq 0$.

Three Axioms of Probability

Let S be the sample space and A be an event in S .

1. For any event A , $P(A) \geq 0$.
2. $P(S) = 1$.

Three Axioms of Probability

Let S be the sample space and A be an event in S .

1. For any event A , $P(A) \geq 0$.
2. $P(S) = 1$.
3. If A_1, A_2, \dots, A_n are mutually disjoint, then

$$P(A_1 \cup A_2 \cup \dots \cup A_n) = P(A_1) + P(A_2) + \dots + P(A_n)$$

Three Axioms of Probability

Let S be the sample space and A be an event in S .

1. For any event A , $P(A) \geq 0$.
2. $P(S) = 1$.
3. If A_1, A_2, \dots, A_n are mutually disjoint, then

$$P(A_1 \cup A_2 \cup \dots \cup A_n) = P(A_1) + P(A_2) + \dots + P(A_n)$$

The three axioms imply:

Three Axioms of Probability

Let S be the sample space and A be an event in S .

1. For any event A , $P(A) \geq 0$.
2. $P(S) = 1$.
3. If A_1, A_2, \dots, A_n are mutually disjoint, then

$$P(A_1 \cup A_2 \cup \dots \cup A_n) = P(A_1) + P(A_2) + \dots + P(A_n)$$

The three axioms imply:

► $P(\emptyset) = 0$

Three Axioms of Probability

Let S be the sample space and A be an event in S .

1. For any event A , $P(A) \geq 0$.
2. $P(S) = 1$.
3. If A_1, A_2, \dots, A_n are mutually disjoint, then

$$P(A_1 \cup A_2 \cup \dots \cup A_n) = P(A_1) + P(A_2) + \dots + P(A_n)$$

The three axioms imply:

- ▶ $P(\emptyset) = 0$
- ▶ $P(A^c) = 1 - P(A)$

Three Axioms of Probability

Let S be the sample space and A be an event in S .

1. For any event A , $P(A) \geq 0$.
2. $P(S) = 1$.
3. If A_1, A_2, \dots, A_n are mutually disjoint, then

$$P(A_1 \cup A_2 \cup \dots \cup A_n) = P(A_1) + P(A_2) + \dots + P(A_n)$$

The three axioms imply:

- ▶ $P(\emptyset) = 0$
- ▶ $P(A^c) = 1 - P(A)$
- ▶ For any event A , $0 \leq P(A) \leq 1$.

Three Axioms of Probability

Let S be the sample space and A be an event in S .

1. For any event A , $P(A) \geq 0$.
2. $P(S) = 1$.
3. If A_1, A_2, \dots, A_n are mutually disjoint, then

$$P(A_1 \cup A_2 \cup \dots \cup A_n) = P(A_1) + P(A_2) + \dots + P(A_n)$$

The three axioms imply:

- ▶ $P(\emptyset) = 0$
- ▶ $P(A^c) = 1 - P(A)$
- ▶ For any event A , $0 \leq P(A) \leq 1$.
- ▶ If $A \subset B$, then $P(A) \leq P(B)$.

Three Axioms of Probability

Let S be the sample space and A be an event in S .

1. For any event A , $P(A) \geq 0$.
2. $P(S) = 1$.
3. If A_1, A_2, \dots, A_n are mutually disjoint, then

$$P(A_1 \cup A_2 \cup \dots \cup A_n) = P(A_1) + P(A_2) + \dots + P(A_n)$$

The three axioms imply:

- ▶ $P(\emptyset) = 0$
- ▶ $P(A^c) = 1 - P(A)$
- ▶ For any event A , $0 \leq P(A) \leq 1$.
- ▶ If $A \subset B$, then $P(A) \leq P(B)$.
- ▶ For any two events A and B ,
 $P(A \cup B) = P(A) + P(B) - P(A \cap B)$.

Conditional Probability

Conditional Probability

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Conditional Probability

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Multiplicative Law of Probability:

Conditional Probability

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Multiplicative Law of Probability:

$$P(A \cap B) = P(B|A)P(A) = P(A|B)P(B)$$

Conditional Probability

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Multiplicative Law of Probability:

$$P(A \cap B) = P(B|A)P(A) = P(A|B)P(B)$$

Bayes' Rule:

Conditional Probability

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Multiplicative Law of Probability:

$$P(A \cap B) = P(B|A)P(A) = P(A|B)P(B)$$

Bayes' Rule:

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

Conditional Probability

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Multiplicative Law of Probability:

$$P(A \cap B) = P(B|A)P(A) = P(A|B)P(B)$$

Bayes' Rule:

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

Law of Total Probability:

Conditional Probability

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Multiplicative Law of Probability:

$$P(A \cap B) = P(B|A)P(A) = P(A|B)P(B)$$

Bayes' Rule:

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

Law of Total Probability:

$$P(B) = \sum_{i=1}^n P(B|A_i)P(A_i)$$

Independence

A and B are independent if

$$P(AB) = P(A)P(B)$$

Independence

A and B are independent if

$$P(AB) = P(A)P(B)$$

If A and B are independent, then

$$P(A|B) = \frac{P(AB)}{P(B)}$$

Independence

A and B are independent if

$$P(AB) = P(A)P(B)$$

If A and B are independent, then

$$\begin{aligned} P(A|B) &= \frac{P(AB)}{P(B)} \\ &= \frac{P(A)P(B)}{P(B)} \end{aligned}$$

Independence

A and B are independent if

$$P(AB) = P(A)P(B)$$

If A and B are independent, then

$$\begin{aligned}P(A|B) &= \frac{P(AB)}{P(B)} \\&= \frac{P(A)P(B)}{P(B)} \\&= P(A)\end{aligned}$$

Independence

A and B are independent if

$$P(AB) = P(A)P(B)$$

If A and B are independent, then

$$\begin{aligned}P(A|B) &= \frac{P(AB)}{P(B)} \\&= \frac{P(A)P(B)}{P(B)} \\&= P(A)\end{aligned}$$

Conditional Independence:

$$P(AB|C) = P(A|C)P(B|C)$$

Outline

Probability

Random Variables

Simulation

Important Distributions

Discrete Distributions

Continuous Distributions

Random Variables

Random Variables

A **random variable** is a function that takes a random experiment and assigns a number to the outcome of the experiment.

Random Variables

A **random variable** is a function that takes a random experiment and assigns a number to the outcome of the experiment.

Outcome values are assigned probabilities by a probability mass function (for discrete RV) or probability density function (for continuous RV).


Probability Mass Function

Probability Mass Function

$$P(Y = y)$$

Probability Mass Function

$$P(Y = y)$$




Probability Density Function

Probability Density Function

$$P(Y \in A) = \int_A f(y) dy$$

Probability Density Function

$$P(Y \in A) = \int_A f(y) dy$$



Characteristics of all PDFs and PMFs:

Characteristics of all PDFs and PMFs:

- ▶ Area under the curve must integrate to 1

Characteristics of all PDFs and PMFs:

- ▶ Area under the curve must integrate to 1
- ▶ $P(Y = y) \geq 0$ for all Y

Characteristics of all PDFs and PMFs:

- ▶ Area under the curve must integrate to 1
- ▶ $P(Y = y) \geq 0$ for all Y
- ▶ The support is all y 's where $P(Y = y) > 0$.

Marginal, Conditional, and Joint Densities

Marginal, Conditional, and Joint Densities

$$f(x) = \int f(x, y) dy$$

Marginal, Conditional, and Joint Densities

$$f(x) = \int f(x, y) dy$$

$$f(x, y) = \int f(x, y, z) dz$$

Marginal, Conditional, and Joint Densities

$$f(x) = \int f(x, y) dy$$

$$f(x, y) = \int f(x, y, z) dz$$

$$f(x|y) = \frac{f(x, y)}{f(y)}$$

Marginal, Conditional, and Joint Densities

$$f(x) = \int f(x, y) dy$$

$$f(x, y) = \int f(x, y, z) dz$$

$$f(x|y) = \frac{f(x, y)}{f(y)}$$

$$f(x|y, z) = \frac{f(x, y, z)}{f(y, z)}$$

Marginal, Conditional, and Joint Densities

$$f(x) = \int f(x, y) dy$$

$$f(x, y) = \int f(x, y, z) dz$$

$$f(x|y) = \frac{f(x, y)}{f(y)}$$

$$f(x|y, z) = \frac{f(x, y, z)}{f(y, z)}$$

$$f(x, y) = f(x|y)f(y)$$

Marginal, Conditional, and Joint Densities

$$f(x) = \int f(x, y) dy$$

$$f(x, y) = \int f(x, y, z) dz$$

$$f(x|y) = \frac{f(x, y)}{f(y)}$$

$$f(x|y, z) = \frac{f(x, y, z)}{f(y, z)}$$

$$\begin{aligned} f(x, y) &= f(x|y)f(y) \\ &= f(y|x)f(x) \end{aligned}$$

Marginal, Conditional, and Joint Densities

$$f(x) = \int f(x, y) dy$$

$$f(x, y) = \int f(x, y, z) dz$$

$$f(x|y) = \frac{f(x, y)}{f(y)}$$

$$f(x|y, z) = \frac{f(x, y, z)}{f(y, z)}$$

$$\begin{aligned} f(x, y) &= f(x|y)f(y) \\ &= f(y|x)f(x) \end{aligned}$$

$$f(x, y, z) = f(x|y, z)f(y|z)f(z)$$

Expectation

Expectation

Discrete Case:

Expectation

Discrete Case:

$$E(X) = \sum_i x_i P(X = x_i)$$

where $P(X = x)$ is the probability mass function (PMF).

Expectation

Discrete Case:

$$E(X) = \sum_i x_i P(X = x_i)$$

where $P(X = x)$ is the probability mass function (PMF).

Continuous Case:

Expectation

Discrete Case:

$$E(X) = \sum_i x_i P(X = x_i)$$

where $P(X = x)$ is the probability mass function (PMF).

Continuous Case:

$$E(X) = \int_{-\infty}^{\infty} xf(x)dx$$

where $f(x)$ is the probability density function (PDF).

Expectation of a Function of a Random Variable

Expectation of a Function of a Random Variable

$$E[g(X)] = \sum_i g(x_i)P(X = x_i)$$

for discrete random variables

Expectation of a Function of a Random Variable

$$E[g(X)] = \sum_i g(x_i)P(X = x_i)$$

for discrete random variables and

$$E[g(X)] = \int_{-\infty}^{\infty} g(x)f(x)dx$$

for continuous random variables.

Expectation of a Function of a Random Variable

$$E[g(X)] = \sum_i g(x_i)P(X = x_i)$$

for discrete random variables and

$$E[g(X)] = \int_{-\infty}^{\infty} g(x)f(x)dx$$

for continuous random variables.

Variance

Variance

$$\text{Var}(X) = E[(X - E(X))^2]$$

Variance

$$\text{Var}(X) = E[(X - E(X))^2]$$

$$\text{Var}(X) = E[(X - E(X))^2]$$

Variance

$$\text{Var}(X) = E[(X - E(X))^2]$$

$$\begin{aligned}\text{Var}(X) &= E[(X - E(X))^2] \\ &= E[X^2 - 2XE(X) + (E(X))^2]\end{aligned}$$

Variance

$$\text{Var}(X) = E[(X - E(X))^2]$$

$$\begin{aligned}\text{Var}(X) &= E[(X - E(X))^2] \\ &= E[X^2 - 2XE(X) + (E(X))^2] \\ &= E(X^2) - 2E(X)E[E(X)] + E([E(X)]^2)\end{aligned}$$

Variance

$$\text{Var}(X) = E[(X - E(X))^2]$$

$$\begin{aligned}\text{Var}(X) &= E[(X - E(X))^2] \\ &= E[X^2 - 2XE(X) + (E(X))^2] \\ &= E(X^2) - 2E(X)E[E(X)] + E([E(X)]^2) \\ &= E(X^2) - 2[E(X)]^2 + [E(X)]^2\end{aligned}$$

Variance

$$\text{Var}(X) = E[(X - E(X))^2]$$

$$\begin{aligned}\text{Var}(X) &= E[(X - E(X))^2] \\&= E[X^2 - 2XE(X) + (E(X))^2] \\&= E(X^2) - 2E(X)E[E(X)] + E([E(X)]^2) \\&= E(X^2) - 2[E(X)]^2 + [E(X)]^2 \\&= \mathbf{E(X^2) - [E(X)]^2}\end{aligned}$$

Variance

$$\text{Var}(X) = E[(X - E(X))^2]$$

$$\begin{aligned}\text{Var}(X) &= E[(X - E(X))^2] \\&= E[X^2 - 2XE(X) + (E(X))^2] \\&= E(X^2) - 2E(X)E[E(X)] + E([E(X)]^2) \\&= E(X^2) - 2[E(X)]^2 + [E(X)]^2 \\&= \mathbf{E(X^2) - [E(X)]^2}\end{aligned}$$

Outline

Probability

Random Variables

Simulation

Important Distributions

Discrete Distributions

Continuous Distributions

Monte Carlo Simulation

Monte Carlo Simulation

All the simulations we will be doing in class is what we call **Monte Carlo simulation**.


Monte Carlo Simulation

All the simulations we will be doing in class is what we call **Monte Carlo simulation**.



Monte Carlo Simulation

All the simulations we will be doing in class is what we call **Monte Carlo simulation**.



Fancy way of saying we will simulate random draws to calculate quantities of interest.


Simulating from a Random Variable

Simulating from a Random Variable

Suppose we have a random variable X that follows a $\text{Beta}(\alpha, \beta)$ distribution.


Simulating from a Random Variable

Suppose we have a random variable X that follows a $\text{Beta}(\alpha, \beta)$ distribution.



Simulating from a Random Variable

Suppose we have a random variable X that follows a $\text{Beta}(\alpha, \beta)$ distribution.



$$f(x|\alpha, \beta) = \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{(\alpha-1)} (1-x)^{(\beta-1)}$$

Let $\alpha = 2$ and $\beta = 3$. Find $E(X)$.

Let $\alpha = 2$ and $\beta = 3$. Find $E(X)$.

$$E(X) = \int_0^1 xf(x)dx$$

Let $\alpha = 2$ and $\beta = 3$. Find $E(X)$.

$$\begin{aligned} E(X) &= \int_0^1 xf(x)dx \\ &= \int_0^1 x \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{(\alpha-1)}(1-x)^{(\beta-1)} \end{aligned}$$

Let $\alpha = 2$ and $\beta = 3$. Find $E(X)$.

$$\begin{aligned} E(X) &= \int_0^1 xf(x)dx \\ &= \int_0^1 x \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{(\alpha-1)}(1-x)^{(\beta-1)} \\ &= \int_0^1 x \frac{\Gamma(2+3)}{\Gamma(2)\Gamma(3)} x^{(2-1)}(1-x)^{(3-1)} dx \end{aligned}$$

Let $\alpha = 2$ and $\beta = 3$. Find $E(X)$.

$$\begin{aligned} E(X) &= \int_0^1 xf(x)dx \\ &= \int_0^1 x \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{(\alpha-1)}(1-x)^{(\beta-1)} \\ &= \int_0^1 x \frac{\Gamma(2 + 3)}{\Gamma(2)\Gamma(3)} x^{(2-1)}(1-x)^{(3-1)} dx \end{aligned}$$

We can ask R to help us integrate.

Let $\alpha = 2$ and $\beta = 3$. Find $E(X)$.

$$\begin{aligned} E(X) &= \int_0^1 xf(x)dx \\ &= \int_0^1 x \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{(\alpha-1)}(1-x)^{(\beta-1)} \\ &= \int_0^1 x \frac{\Gamma(2+3)}{\Gamma(2)\Gamma(3)} x^{(2-1)}(1-x)^{(3-1)}dx \end{aligned}$$

We can ask R to help us integrate.

```
> ex.beta.func <- function(x, alpha, beta) {  
+   x * gamma(alpha + beta)/(gamma(alpha) * gamma(beta)) * x^(alpha -  
+     1) * (1 - x)^(beta - 1)  
+ }  
> e.x <- integrate(Vectorize(ex.beta.func), lower = 0, upper = 1,  
+   alpha = 2, beta = 3)$value  
> e.x  
[1] 0.4
```

Or we can use simulation, which is much easier.

Or we can use simulation, which is much easier.

```
> x.draws <- rbeta(10000, shape1 = 2, shape2 = 3)
> sim.e.x <- mean(x.draws)
> sim.e.x

[1] 0.3987049
```

Or we can use simulation, which is much easier.

```
> x.draws <- rbeta(10000, shape1 = 2, shape2 = 3)
> sim.e.x <- mean(x.draws)
> sim.e.x

[1] 0.3987049
```

We can find all kinds of quantities of interest (variance, quantiles, etc.) by just doing it on the simulated draws rather than doing complicated integrals.

Or we can use simulation, which is much easier.

```
> x.draws <- rbeta(10000, shape1 = 2, shape2 = 3)
> sim.e.x <- mean(x.draws)
> sim.e.x

[1] 0.3987049
```

We can find all kinds of quantities of interest (variance, quantiles, etc.) by just doing it on the simulated draws rather than doing complicated integrals.

Why does this work?

Monte Carlo Integration

Monte Carlo Integration

What we just did was called **Monte Carlo Integration**, which means exactly what it sounds like (doing integrals via Monte Carlo simulation).

Monte Carlo Integration

What we just did was called **Monte Carlo Integration**, which means exactly what it sounds like (doing integrals via Monte Carlo simulation).

If we need to take an integral of the following form:

$$I = \int g(x)p(x)dx$$

Monte Carlo Integration

What we just did was called **Monte Carlo Integration**, which means exactly what it sounds like (doing integrals via Monte Carlo simulation).

If we need to take an integral of the following form:

$$I = \int g(x)p(x)dx$$

Monte Carlo Integration allows us to approximate it by simulating M values from $f(x)$ and calculating:

$$\hat{I}_M = \frac{1}{M} \sum_{i=1}^M g(x^{(i)})$$

Monte Carlo Integration

What we just did was called **Monte Carlo Integration**, which means exactly what it sounds like (doing integrals via Monte Carlo simulation).

If we need to take an integral of the following form:

$$I = \int g(x)p(x)dx$$

Monte Carlo Integration allows us to approximate it by simulating M values from $f(x)$ and calculating:

$$\hat{I}_M = \frac{1}{M} \sum_{i=1}^M g(x^{(i)})$$

By the Strong Law of Large Numbers, our estimate \hat{I}_M is a simulation consistent estimator of I as $M \rightarrow \infty$

Monte Carlo Integration

What we just did was called **Monte Carlo Integration**, which means exactly what it sounds like (doing integrals via Monte Carlo simulation).

If we need to take an integral of the following form:

$$I = \int g(x)p(x)dx$$

Monte Carlo Integration allows us to approximate it by simulating M values from $f(x)$ and calculating:

$$\hat{I}_M = \frac{1}{M} \sum_{i=1}^M g(x^{(i)})$$

By the Strong Law of Large Numbers, our estimate \hat{I}_M is a simulation consistent estimator of I as $M \rightarrow \infty$ (our estimate gets better as we increase the number of simulations).

Let Y be another random variable where $Y = e^X$. Find $E(Y)$.

Let Y be another random variable where $Y = e^X$. Find $E(Y)$.

$$E(Y) = \int_0^1 e^x \frac{\Gamma(2+3)}{\Gamma(2)\Gamma(3)} x^{(2-1)} (1-x)^{(3-1)} dx$$

Let Y be another random variable where $Y = e^X$. Find $E(Y)$.

$$E(Y) = \int_0^1 e^x \frac{\Gamma(2+3)}{\Gamma(2)\Gamma(3)} x^{(2-1)} (1-x)^{(3-1)} dx$$

Via simulation:

Let Y be another random variable where $Y = e^X$. Find $E(Y)$.

$$E(Y) = \int_0^1 e^x \frac{\Gamma(2+3)}{\Gamma(2)\Gamma(3)} x^{(2-1)} (1-x)^{(3-1)} dx$$

Via simulation:

```
> e.y <- mean(exp(x.draws))  
> e.y  
[1] 1.520348
```


Let Y be another random variable where $Y = e^X$. Find $E(Y)$.

$$E(Y) = \int_0^1 e^x \frac{\Gamma(2+3)}{\Gamma(2)\Gamma(3)} x^{(2-1)} (1-x)^{(3-1)} dx$$

Via simulation:

```
> e.y <- mean(exp(x.draws))  
> e.y  
[1] 1.520348
```

Note that $E(g(X)) \neq g(E(X))$.

Let Y be another random variable where $Y = e^X$. Find $E(Y)$.

$$E(Y) = \int_0^1 e^x \frac{\Gamma(2+3)}{\Gamma(2)\Gamma(3)} x^{(2-1)} (1-x)^{(3-1)} dx$$

Via simulation:

```
> e.y <- mean(exp(x.draws))  
> e.y  
[1] 1.520348
```

Note that $E(g(X)) \neq g(E(X))$. In fact, $E(g(X)) \geq g(E(X))$ by Jensen's Inequality.

Let Y be another random variable where $Y = e^X$. Find $E(Y)$.

$$E(Y) = \int_0^1 e^x \frac{\Gamma(2+3)}{\Gamma(2)\Gamma(3)} x^{(2-1)} (1-x)^{(3-1)} dx$$

Via simulation:

```
> e.y <- mean(exp(x.draws))  
> e.y  
[1] 1.520348
```

Note that $E(g(X)) \neq g(E(X))$. In fact, $E(g(X)) \geq g(E(X))$ by Jensen's Inequality.

Monte Carlo Integration tells us we need $E(g(X))$.

Take home point:

Take home point:

If we can somehow generate random draws from the distribution of a random variable, we can calculate complicated integrals (mean, variance, functions of RV) easily by simulation.

Take home point:

If we can somehow generate random draws from the distribution of a random variable, we can calculate complicated integrals (mean, variance, functions of RV) easily by simulation.

This seems trivial but is one of the foundations of statistics, especially Bayesian statistics.

Simulating Probability Problems

Simulating Probability Problems

A related application of simulation is to help us solve probability problems.

Simulating Probability Problems

A related application of simulation is to help us solve probability problems.

General idea:

Simulating Probability Problems

A related application of simulation is to help us solve probability problems.

General idea:

1. Suppose we have an experiment where we want to know the probability of success. Simulate from the population many times.

Simulating Probability Problems

A related application of simulation is to help us solve probability problems.

General idea:

1. Suppose we have an experiment where we want to know the probability of success. Simulate from the population many times.
2. For each simulation, conduct the experiment and see whether there is success.

Simulating Probability Problems

A related application of simulation is to help us solve probability problems.

General idea:

1. Suppose we have an experiment where we want to know the probability of success. Simulate from the population many times.
2. For each simulation, conduct the experiment and see whether there is success.
3. The proportion of simulations that achieve success is the probability of success.

An Example

An Example

Suppose we have two urns containing marbles. The first urn contains 6 red marbles and 4 green marbles and the second urn contains 9 red marbles and 1 green marble. Take one marble from the first urn (without looking at it) and put it in the second urn. Then take one marble from the second urn (again without looking at it) and put it in the first urn. What is the probability of now drawing a red marble from the first urn?

An Example

Suppose we have two urns containing marbles. The first urn contains 6 red marbles and 4 green marbles and the second urn contains 9 red marbles and 1 green marble. Take one marble from the first urn (without looking at it) and put it in the second urn. Then take one marble from the second urn (again without looking at it) and put it in the first urn. What is the probability of now drawing a red marble from the first urn?

```
> urn.func <- function(n.sims, urn1, urn2) {  
+   final.draws <- c()  
+   for (i in 1:n.sims) {  
+     draw1 <- sample(urn1, 1)  
+     draw2 <- sample(c(urn2, draw1), 1)  
+     final.draws[i] <- sample(c(urn1, draw2), 1)  
+   }  
+   prob <- mean(final.draws)  
+   return(prob)  
+ }  
> urn.func(n.sims = 10000, urn1 = c(rep(1, 6), rep(0, 4)), urn2 = c(rep(1,  
+   9), 0))
```

```
[1] 0.6293
```

An Example

Suppose we have two urns containing marbles. The first urn contains 6 red marbles and 4 green marbles and the second urn contains 9 red marbles and 1 green marble. Take one marble from the first urn (without looking at it) and put it in the second urn. Then take one marble from the second urn (again without looking at it) and put it in the first urn. What is the probability of now drawing a red marble from the first urn?

```
> urn.func <- function(n.sims, urn1, urn2) {  
+   final.draws <- c()  
+   for (i in 1:n.sims) {  
+     draw1 <- sample(urn1, 1)  
+     draw2 <- sample(c(urn2, draw1), 1)  
+     final.draws[i] <- sample(c(urn1, draw2), 1)  
+   }  
+   prob <- mean(final.draws)  
+   return(prob)  
+ }  
> urn.func(n.sims = 10000, urn1 = c(rep(1, 6), rep(0, 4)), urn2 = c(rep(1,  
+   9), 0))
```

```
[1] 0.6293
```

$$\left(\frac{6}{10}\right)\left(\frac{10}{11}\right)\left(\frac{6}{10}\right) + \left(\frac{6}{10}\right)\left(\frac{1}{11}\right)\left(\frac{5}{10}\right) + \left(\frac{4}{10}\right)\left(\frac{9}{11}\right)\left(\frac{7}{10}\right) + \left(\frac{4}{10}\right)\left(\frac{2}{11}\right)\left(\frac{6}{10}\right) \approx 0.63$$

Another Example

Another Example

Suppose we have two urns containing marbles. The first urn contains $10 - g$ red marbles and g green marbles and the second urn contains 9 red marbles and 1 green marble. Take one marble from the first urn (without looking at it) and put it in the second urn. Then take one marble from the second urn (again without looking at it) and put it in the first urn. What is the minimum g such that the probability of now drawing a red marble is less than 0.5?

Another Example

Suppose we have two urns containing marbles. The first urn contains $10 - g$ red marbles and g green marbles and the second urn contains 9 red marbles and 1 green marble. Take one marble from the first urn (without looking at it) and put it in the second urn. Then take one marble from the second urn (again without looking at it) and put it in the first urn. What is the minimum g such that the probability of now drawing a red marble is less than 0.5?

```
> urn.func2 <- function(n.sims, urn2, p) {  
+   final.draws <- c()  
+   g <- 0  
+   urn1 <- c(rep(1, 10 - g), rep(0, g))  
+   prob <- 1  
+   while (prob >= p) {  
+     for (i in 1:n.sims) {  
+       draw1 <- sample(urn1, 1)  
+       draw2 <- sample(c(urn2, draw1), 1)  
+       final.draws[i] <- sample(c(urn1, draw2), 1)  
+     }  
+     prob <- mean(final.draws)  
+     g <- g + 1  
+     urn1 <- c(rep(1, 10 - g), rep(0, g))  
+   }  
+   g <- g - 1  
+   return(g)  
+ }  
> urn.func2(1000, urn2 = c(rep(1, 9), 0), p = 0.5)
```

```
[1] 6
```

Outline

Probability

Random Variables

Simulation

Important Distributions

Discrete Distributions

Continuous Distributions

Outline

Probability

Random Variables

Simulation

Important Distributions

Discrete Distributions

Continuous Distributions

The Bernoulli Distribution

The Bernoulli Distribution

$$Y \sim \text{Bernoulli}(\pi)$$

The Bernoulli Distribution

$$Y \sim \text{Bernoulli}(\pi)$$

$$y = 0, 1$$

The Bernoulli Distribution

$$Y \sim \text{Bernoulli}(\pi)$$

$$y = 0, 1$$

probability of success: $\pi \in [0, 1]$

The Bernoulli Distribution

$$Y \sim \text{Bernoulli}(\pi)$$

$$y = 0, 1$$

probability of success: $\pi \in [0, 1]$

$$p(y|\pi) = \pi^y(1 - \pi)^{(1-y)}$$

The Bernoulli Distribution

$$Y \sim \text{Bernoulli}(\pi)$$

$$y = 0, 1$$

probability of success: $\pi \in [0, 1]$

$$p(y|\pi) = \pi^y(1 - \pi)^{(1-y)}$$

$$E(Y) = \pi$$

The Bernoulli Distribution

$$Y \sim \text{Bernoulli}(\pi)$$

$$y = 0, 1$$

probability of success: $\pi \in [0, 1]$

$$p(y|\pi) = \pi^y(1 - \pi)^{(1-y)}$$

$$E(Y) = \pi$$

$$\text{Var}(Y) = \pi(1 - \pi)$$

The Bernoulli Distribution

$$Y \sim \text{Bernoulli}(\pi)$$


$$y = 0, 1$$

probability of success: $\pi \in [0, 1]$

$$p(y|\pi) = \pi^y(1 - \pi)^{(1-y)}$$

$$E(Y) = \pi$$

$$\text{Var}(Y) = \pi(1 - \pi)$$



The Binomial Distribution

The Binomial Distribution

$$Y \sim \text{Binomial}(n, \pi)$$

The Binomial Distribution

$$Y \sim \text{Binomial}(n, \pi)$$

$$y = 0, 1, \dots, n$$

The Binomial Distribution

$$Y \sim \text{Binomial}(n, \pi)$$

$$y = 0, 1, \dots, n$$

number of trials: $n \in \{1, 2, \dots\}$

The Binomial Distribution

$$Y \sim \text{Binomial}(n, \pi)$$

$$y = 0, 1, \dots, n$$

number of trials: $n \in \{1, 2, \dots\}$

probability of success: $\pi \in [0, 1]$

The Binomial Distribution

$$Y \sim \text{Binomial}(n, \pi)$$

$$y = 0, 1, \dots, n$$

number of trials: $n \in \{1, 2, \dots\}$

probability of success: $\pi \in [0, 1]$

$$p(y|\pi) = \binom{n}{y} \pi^y (1 - \pi)^{(n-y)}$$

The Binomial Distribution

$$Y \sim \text{Binomial}(n, \pi)$$

$$y = 0, 1, \dots, n$$

number of trials: $n \in \{1, 2, \dots\}$

probability of success: $\pi \in [0, 1]$

$$p(y|\pi) = \binom{n}{y} \pi^y (1 - \pi)^{(n-y)}$$

$$E(Y) = n\pi$$

The Binomial Distribution

$$Y \sim \text{Binomial}(n, \pi)$$

$$y = 0, 1, \dots, n$$

number of trials: $n \in \{1, 2, \dots\}$

probability of success: $\pi \in [0, 1]$

$$p(y|\pi) = \binom{n}{y} \pi^y (1 - \pi)^{(n-y)}$$

$$E(Y) = n\pi$$

$$\text{Var}(Y) = n\pi(1 - \pi)$$

The Binomial Distribution

$$Y \sim \text{Binomial}(n, \pi)$$

$$y = 0, 1, \dots, n$$


number of trials: $n \in \{1, 2, \dots\}$

probability of success: $\pi \in [0, 1]$

$$p(y|\pi) = \binom{n}{y} \pi^y (1 - \pi)^{(n-y)}$$

$$E(Y) = n\pi$$

$$\text{Var}(Y) = n\pi(1 - \pi)$$



The Multinomial Distribution

The Multinomial Distribution

$$Y \sim \text{Multinomial}(n, \pi_1, \dots, \pi_k)$$

The Multinomial Distribution

$$Y \sim \text{Multinomial}(n, \pi_1, \dots, \pi_k)$$

$$y_j = 0, 1, \dots, n; \quad \sum_{j=1}^k y_j = n$$

The Multinomial Distribution

$$Y \sim \text{Multinomial}(n, \pi_1, \dots, \pi_k)$$

$$y_j = 0, 1, \dots, n; \quad \sum_{j=1}^k y_j = n$$

number of trials: $n \in \{1, 2, \dots\}$

The Multinomial Distribution

$$Y \sim \text{Multinomial}(n, \pi_1, \dots, \pi_k)$$

$$y_j = 0, 1, \dots, n; \quad \sum_{j=1}^k y_j = n$$

number of trials: $n \in \{1, 2, \dots\}$

probability of success for j : $\pi_j \in [0, 1]$; $\sum_{j=1}^k \pi_j = 1$

The Multinomial Distribution

$$Y \sim \text{Multinomial}(n, \pi_1, \dots, \pi_k)$$

$$y_j = 0, 1, \dots, n; \quad \sum_{j=1}^k y_j = n$$

number of trials: $n \in \{1, 2, \dots\}$

probability of success for j : $\pi_j \in [0, 1]$; $\sum_{j=1}^k \pi_j = 1$

$$p(\mathbf{y}|n, \boldsymbol{\pi}) = \frac{n!}{y_1! y_2! \dots y_k!} \pi_1^{y_1} \pi_2^{y_2} \dots \pi_k^{y_k}$$

The Multinomial Distribution

$$Y \sim \text{Multinomial}(n, \pi_1, \dots, \pi_k)$$

$$y_j = 0, 1, \dots, n; \quad \sum_{j=1}^k y_j = n$$

number of trials: $n \in \{1, 2, \dots\}$

probability of success for j : $\pi_j \in [0, 1]$; $\sum_{j=1}^k \pi_j = 1$

$$p(\mathbf{y}|n, \boldsymbol{\pi}) = \frac{n!}{y_1! y_2! \dots y_k!} \pi_1^{y_1} \pi_2^{y_2} \dots \pi_k^{y_k}$$

$$E(Y_j) = n\pi_j$$

The Multinomial Distribution

$$Y \sim \text{Multinomial}(n, \pi_1, \dots, \pi_k)$$

$$y_j = 0, 1, \dots, n; \quad \sum_{j=1}^k y_j = n$$

number of trials: $n \in \{1, 2, \dots\}$

probability of success for j : $\pi_j \in [0, 1]$; $\sum_{j=1}^k \pi_j = 1$

$$p(\mathbf{y}|n, \boldsymbol{\pi}) = \frac{n!}{y_1! y_2! \dots y_k!} \pi_1^{y_1} \pi_2^{y_2} \dots \pi_k^{y_k}$$

$$E(Y_j) = n\pi_j$$

$$\text{Var}(Y_j) = n\pi_j(1 - \pi_j)$$

The Multinomial Distribution

$$Y \sim \text{Multinomial}(n, \pi_1, \dots, \pi_k)$$

$$y_j = 0, 1, \dots, n; \quad \sum_{j=1}^k y_j = n$$

number of trials: $n \in \{1, 2, \dots\}$

probability of success for j : $\pi_j \in [0, 1]$; $\sum_{j=1}^k \pi_j = 1$

$$p(\mathbf{y}|n, \boldsymbol{\pi}) = \frac{n!}{y_1! y_2! \dots y_k!} \pi_1^{y_1} \pi_2^{y_2} \dots \pi_k^{y_k}$$

$$E(Y_j) = n\pi_j$$

$$\text{Var}(Y_j) = n\pi_j(1 - \pi_j)$$

$$\text{Cov}(Y_i, Y_j) = -n\pi_i\pi_j$$

The Poisson Distribution

The Poisson Distribution

$$Y \sim \text{Poisson}(\lambda)$$

The Poisson Distribution

$$Y \sim \text{Poisson}(\lambda)$$

$$y = 0, 1, \dots$$

The Poisson Distribution

$$Y \sim \text{Poisson}(\lambda)$$

$$y = 0, 1, \dots$$

expected number of
occurrences: $\lambda > 0$

The Poisson Distribution

$$Y \sim \text{Poisson}(\lambda)$$

$$y = 0, 1, \dots$$

expected number of
occurrences: $\lambda > 0$

$$p(y|\lambda) = \frac{e^{-\lambda} \lambda^y}{y!}$$

The Poisson Distribution

$$Y \sim \text{Poisson}(\lambda)$$

$$y = 0, 1, \dots$$

expected number of
occurrences: $\lambda > 0$

$$p(y|\lambda) = \frac{e^{-\lambda} \lambda^y}{y!}$$

$$E(Y) = \lambda$$

The Poisson Distribution

$$Y \sim \text{Poisson}(\lambda)$$

$$y = 0, 1, \dots$$

expected number of
occurrences: $\lambda > 0$

$$p(y|\lambda) = \frac{e^{-\lambda} \lambda^y}{y!}$$

$$E(Y) = \lambda$$

$$\text{Var}(Y) = \lambda$$

The Poisson Distribution

$$Y \sim \text{Poisson}(\lambda)$$


$$y = 0, 1, \dots$$

expected number of
occurrences: $\lambda > 0$

$$p(y|\lambda) = \frac{e^{-\lambda} \lambda^y}{y!}$$

$$E(Y) = \lambda$$

$$\text{Var}(Y) = \lambda$$



The Geometric Distribution

The Geometric Distribution

How many Bernoulli trials until success?

The Geometric Distribution

How many Bernoulli trials until success?

$$Y \sim \text{Geometric}(\pi)$$

The Geometric Distribution

How many Bernoulli trials until success?

$$Y \sim \text{Geometric}(\pi)$$

$$y = 1, 2, 3, \dots$$

The Geometric Distribution

How many Bernoulli trials until success?

$$Y \sim \text{Geometric}(\pi)$$

$$y = 1, 2, 3, \dots$$

probability of success: $\pi \in [0, 1]$

The Geometric Distribution

How many Bernoulli trials until success?

$$Y \sim \text{Geometric}(\pi)$$

$$y = 1, 2, 3, \dots$$

probability of success: $\pi \in [0, 1]$

$$p(y|\pi) = (1 - \pi)^{(y-1)}\pi$$

The Geometric Distribution

How many Bernoulli trials until success?

$$Y \sim \text{Geometric}(\pi)$$

$$y = 1, 2, 3, \dots$$

probability of success: $\pi \in [0, 1]$

$$p(y|\pi) = (1 - \pi)^{(y-1)}\pi$$

$$E(Y) = \frac{1}{\pi}$$

The Geometric Distribution

How many Bernoulli trials until success?

$$Y \sim \text{Geometric}(\pi)$$

$$y = 1, 2, 3, \dots$$

probability of success: $\pi \in [0, 1]$

$$p(y|\pi) = (1 - \pi)^{(y-1)}\pi$$

$$E(Y) = \frac{1}{\pi}$$

$$\text{Var}(Y) = \frac{1-\pi}{\pi^2}$$

The Geometric Distribution

How many Bernoulli trials until success?

$$Y \sim \text{Geometric}(\pi)$$


$$y = 1, 2, 3, \dots$$

probability of success: $\pi \in [0, 1]$

$$p(y|\pi) = (1 - \pi)^{(y-1)}\pi$$

$$E(Y) = \frac{1}{\pi}$$

$$\text{Var}(Y) = \frac{1-\pi}{\pi^2}$$



Outline

Probability

Random Variables

Simulation

Important Distributions

Discrete Distributions

Continuous Distributions

The Univariate Normal Distribution

The Univariate Normal Distribution

$$Y \sim \text{Normal}(\mu, \sigma^2)$$

The Univariate Normal Distribution

$$Y \sim \text{Normal}(\mu, \sigma^2)$$

$$y \in \mathbb{R}$$

The Univariate Normal Distribution

$$Y \sim \text{Normal}(\mu, \sigma^2)$$

$$y \in \mathbb{R}$$

$$\text{mean: } \mu \in \mathbb{R}$$

The Univariate Normal Distribution

$$Y \sim \text{Normal}(\mu, \sigma^2)$$

$$y \in \mathbb{R}$$

mean: $\mu \in \mathbb{R}$

variance: $\sigma^2 > 0$

The Univariate Normal Distribution

$$Y \sim \text{Normal}(\mu, \sigma^2)$$

$$y \in \mathbb{R}$$

mean: $\mu \in \mathbb{R}$

variance: $\sigma^2 > 0$

$$p(y|\mu, \sigma^2) = \frac{\exp\left(-\frac{(y-\mu)^2}{2\sigma^2}\right)}{\sigma\sqrt{2\pi}}$$

The Univariate Normal Distribution

$$Y \sim \text{Normal}(\mu, \sigma^2)$$

$$y \in \mathbb{R}$$

mean: $\mu \in \mathbb{R}$

variance: $\sigma^2 > 0$

$$p(y|\mu, \sigma^2) = \frac{\exp\left(-\frac{(y-\mu)^2}{2\sigma^2}\right)}{\sigma\sqrt{2\pi}}$$

$$E(Y) = \mu$$

The Univariate Normal Distribution

$$Y \sim \text{Normal}(\mu, \sigma^2)$$

$$y \in \mathbb{R}$$

$$\text{mean: } \mu \in \mathbb{R}$$

$$\text{variance: } \sigma^2 > 0$$

$$p(y|\mu, \sigma^2) = \frac{\exp\left(-\frac{(y-\mu)^2}{2\sigma^2}\right)}{\sigma\sqrt{2\pi}}$$

$$E(Y) = \mu$$

$$\text{Var}(Y) = \sigma^2$$

The Univariate Normal Distribution

$$Y \sim \text{Normal}(\mu, \sigma^2)$$

$$y \in \mathbb{R}$$


mean: $\mu \in \mathbb{R}$

variance: $\sigma^2 > 0$

$$p(y|\mu, \sigma^2) = \frac{\exp\left(-\frac{(y-\mu)^2}{2\sigma^2}\right)}{\sigma\sqrt{2\pi}}$$

$$E(Y) = \mu$$

$$\text{Var}(Y) = \sigma^2$$



The Multivariate Normal Distribution

The Multivariate Normal Distribution

$$Y \sim \mathcal{N}(\mu, \Sigma)$$

The Multivariate Normal Distribution

$$Y \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

$$\mathbf{y} \in \mathbb{R}^k$$

The Multivariate Normal Distribution

$$Y \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

$$\mathbf{y} \in \mathbb{R}^k$$

mean vector: $\boldsymbol{\mu} \in \mathbb{R}^k$

The Multivariate Normal Distribution

$$Y \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

$$\mathbf{y} \in \mathbb{R}^k$$

mean vector: $\boldsymbol{\mu} \in \mathbb{R}^k$

variance-covariance matrix: $\boldsymbol{\Sigma}$ positive definite $k \times k$ matrix

The Multivariate Normal Distribution

$$Y \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

$$\mathbf{y} \in \mathbb{R}^k$$

mean vector: $\boldsymbol{\mu} \in \mathbb{R}^k$

variance-covariance matrix: $\boldsymbol{\Sigma}$ positive definite $k \times k$ matrix

$$p(\mathbf{y}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = (2\pi)^{-k/2} |\boldsymbol{\Sigma}|^{-1/2} \exp\left(-\frac{1}{2}(\mathbf{y} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1}(\mathbf{y} - \boldsymbol{\mu})\right)$$

The Multivariate Normal Distribution

$$Y \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

$$\mathbf{y} \in \mathbb{R}^k$$

mean vector: $\boldsymbol{\mu} \in \mathbb{R}^k$

variance-covariance matrix: $\boldsymbol{\Sigma}$ positive definite $k \times k$ matrix

$$p(\mathbf{y}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = (2\pi)^{-k/2} |\boldsymbol{\Sigma}|^{-1/2} \exp\left(-\frac{1}{2}(\mathbf{y} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\mathbf{y} - \boldsymbol{\mu})\right)$$

$$E(Y) = \boldsymbol{\mu}$$

The Multivariate Normal Distribution

$$Y \sim \mathcal{N}(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

$$\mathbf{y} \in \mathbb{R}^k$$

mean vector: $\boldsymbol{\mu} \in \mathbb{R}^k$

variance-covariance matrix: $\boldsymbol{\Sigma}$ positive definite $k \times k$ matrix

$$p(\mathbf{y}|\boldsymbol{\mu}, \boldsymbol{\Sigma}) = (2\pi)^{-k/2} |\boldsymbol{\Sigma}|^{-1/2} \exp\left(-\frac{1}{2}(\mathbf{y} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1}(\mathbf{y} - \boldsymbol{\mu})\right)$$

$$E(Y) = \boldsymbol{\mu}$$

$$\text{Var}(Y) = \boldsymbol{\Sigma}$$

The Uniform Distribution

The Uniform Distribution

$$Y \sim \text{Uniform}(\alpha, \beta)$$

The Uniform Distribution

$$Y \sim \text{Uniform}(\alpha, \beta)$$

$$y \in [\alpha, \beta]$$

The Uniform Distribution

$$Y \sim \text{Uniform}(\alpha, \beta)$$

$$y \in [\alpha, \beta]$$

Interval: $[\alpha, \beta]$; $\beta > \alpha$

The Uniform Distribution

$$Y \sim \text{Uniform}(\alpha, \beta)$$

$$y \in [\alpha, \beta]$$

Interval: $[\alpha, \beta]$; $\beta > \alpha$

$$p(y|\alpha, \beta) = \frac{1}{\beta - \alpha}$$

The Uniform Distribution

$$Y \sim \text{Uniform}(\alpha, \beta)$$

$$y \in [\alpha, \beta]$$

$$\text{Interval: } [\alpha, \beta]; \quad \beta > \alpha$$

$$p(y|\alpha, \beta) = \frac{1}{\beta - \alpha}$$

$$E(Y) = \frac{\alpha + \beta}{2}$$

The Uniform Distribution

$$Y \sim \text{Uniform}(\alpha, \beta)$$

$$y \in [\alpha, \beta]$$

Interval: $[\alpha, \beta]$; $\beta > \alpha$

$$p(y|\alpha, \beta) = \frac{1}{\beta - \alpha}$$

$$E(Y) = \frac{\alpha + \beta}{2}$$

$$\text{Var}(Y) = \frac{(\beta - \alpha)^2}{12}$$

The Beta Distribution

The Beta Distribution

$$Y \sim \text{Beta}(\alpha, \beta)$$

The Beta Distribution

$$Y \sim \text{Beta}(\alpha, \beta)$$

$$y \in [0, 1]$$

The Beta Distribution

$$Y \sim \text{Beta}(\alpha, \beta)$$

$$y \in [0, 1]$$

shape parameters:

$$\alpha > 0; \beta > 0$$

The Beta Distribution

$$Y \sim \text{Beta}(\alpha, \beta)$$

$$y \in [0, 1]$$

shape parameters:

$$\alpha > 0; \quad \beta > 0$$

$$p(y|\alpha, \beta) = \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} y^{(\alpha-1)} (1-y)^{(\beta-1)}$$

The Beta Distribution

$$Y \sim \text{Beta}(\alpha, \beta)$$

$$y \in [0, 1]$$

shape parameters:

$$\alpha > 0; \quad \beta > 0$$

$$p(y|\alpha, \beta) = \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} y^{(\alpha-1)} (1-y)^{(\beta-1)}$$

$$E(Y) = \frac{\alpha}{\alpha+\beta}$$

The Beta Distribution

$$Y \sim \text{Beta}(\alpha, \beta)$$

$$y \in [0, 1]$$


shape parameters:

$$\alpha > 0; \beta > 0$$

$$p(y|\alpha, \beta) = \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} y^{(\alpha-1)}(1-y)^{(\beta-1)}$$

$$E(Y) = \frac{\alpha}{\alpha+\beta}$$

$$\text{Var}(Y) = \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$$



The Gamma Distribution

The Gamma Distribution

$$Y \sim \text{Gamma}(\alpha, \beta)$$

The Gamma Distribution

$$Y \sim \text{Gamma}(\alpha, \beta)$$

$$y > 0$$

The Gamma Distribution

$$Y \sim \text{Gamma}(\alpha, \beta)$$

$$y > 0$$

shape parameter: $\alpha > 0$

The Gamma Distribution

$$Y \sim \text{Gamma}(\alpha, \beta)$$

$$y > 0$$

shape parameter: $\alpha > 0$

inverse scale parameter: $\beta > 0$

The Gamma Distribution

$$Y \sim \text{Gamma}(\alpha, \beta)$$

$$y > 0$$

shape parameter: $\alpha > 0$

inverse scale parameter: $\beta > 0$

$$p(y|\alpha, \beta) = \frac{\beta^\alpha}{\Gamma(\alpha)} y^{(\alpha-1)} \exp(-\beta y)$$

The Gamma Distribution

$$Y \sim \text{Gamma}(\alpha, \beta)$$

$$y > 0$$

shape parameter: $\alpha > 0$

inverse scale parameter: $\beta > 0$

$$p(y|\alpha, \beta) = \frac{\beta^\alpha}{\Gamma(\alpha)} y^{(\alpha-1)} \exp(-\beta y)$$

$$E(Y) = \frac{\alpha}{\beta}$$

The Gamma Distribution

$$Y \sim \text{Gamma}(\alpha, \beta)$$

$$y > 0$$


shape parameter: $\alpha > 0$

inverse scale parameter: $\beta > 0$

$$p(y|\alpha, \beta) = \frac{\beta^\alpha}{\Gamma(\alpha)} y^{(\alpha-1)} \exp(-\beta y)$$

$$E(Y) = \frac{\alpha}{\beta}$$

$$\text{Var}(Y) = \frac{\alpha}{\beta^2}$$



The Inverse Gamma Distribution

The Inverse Gamma Distribution

Distribution of the Inverse of a Gamma Distribution:

The Inverse Gamma Distribution

Distribution of the Inverse of a Gamma Distribution: If $X \sim \text{Gamma}(\alpha, \beta)$, then $\frac{1}{X} \sim \text{Invgamma}(\alpha, \beta)$.

The Inverse Gamma Distribution

Distribution of the Inverse of a Gamma Distribution: If $X \sim \text{Gamma}(\alpha, \beta)$, then $\frac{1}{X} \sim \text{Invgamma}(\alpha, \beta)$.

$$Y \sim \text{Invgamma}(\alpha, \beta)$$

The Inverse Gamma Distribution

Distribution of the Inverse of a Gamma Distribution: If $X \sim \text{Gamma}(\alpha, \beta)$, then $\frac{1}{X} \sim \text{Invgamma}(\alpha, \beta)$.

$$Y \sim \text{Invgamma}(\alpha, \beta)$$

$$y > 0$$

The Inverse Gamma Distribution

Distribution of the Inverse of a Gamma Distribution: If $X \sim \text{Gamma}(\alpha, \beta)$, then $\frac{1}{X} \sim \text{Invgamma}(\alpha, \beta)$.

$$Y \sim \text{Invgamma}(\alpha, \beta)$$

$$y > 0$$

shape parameter: $\alpha > 0$

The Inverse Gamma Distribution

Distribution of the Inverse of a Gamma Distribution: If $X \sim \text{Gamma}(\alpha, \beta)$, then $\frac{1}{X} \sim \text{Invgamma}(\alpha, \beta)$.

$$Y \sim \text{Invgamma}(\alpha, \beta)$$

$$y > 0$$

shape parameter: $\alpha > 0$

scale parameter: $\beta > 0$

The Inverse Gamma Distribution

Distribution of the Inverse of a Gamma Distribution: If $X \sim \text{Gamma}(\alpha, \beta)$, then $\frac{1}{X} \sim \text{Invgamma}(\alpha, \beta)$.

$$Y \sim \text{Invgamma}(\alpha, \beta)$$

$$y > 0$$

shape parameter: $\alpha > 0$

scale parameter: $\beta > 0$

$$p(y|\alpha, \beta) = \frac{\beta^\alpha}{\Gamma(\alpha)} y^{-(\alpha+1)} e^{-\frac{\beta}{y}}$$

The Inverse Gamma Distribution

Distribution of the Inverse of a Gamma Distribution: If $X \sim \text{Gamma}(\alpha, \beta)$, then $\frac{1}{X} \sim \text{Invgamma}(\alpha, \beta)$.

$$Y \sim \text{Invgamma}(\alpha, \beta)$$

$$y > 0$$

shape parameter: $\alpha > 0$

scale parameter: $\beta > 0$

$$p(y|\alpha, \beta) = \frac{\beta^\alpha}{\Gamma(\alpha)} y^{-(\alpha+1)} e^{-\frac{\beta}{y}}$$

$$E(Y) = \frac{\beta}{\alpha-1} \text{ for } \alpha > 1$$

The Inverse Gamma Distribution

Distribution of the Inverse of a Gamma Distribution: If $X \sim \text{Gamma}(\alpha, \beta)$, then $\frac{1}{X} \sim \text{Invgamma}(\alpha, \beta)$.

$$Y \sim \text{Invgamma}(\alpha, \beta)$$

$$y > 0$$

shape parameter: $\alpha > 0$

scale parameter: $\beta > 0$

$$p(y|\alpha, \beta) = \frac{\beta^\alpha}{\Gamma(\alpha)} y^{-(\alpha+1)} e^{-\frac{\beta}{y}}$$

$$E(Y) = \frac{\beta}{\alpha-1} \text{ for } \alpha > 1$$

$$\text{Var}(Y) = \frac{\beta^2}{(\alpha-1)^2(\alpha-2)} \text{ for } \alpha > 2$$

The Inverse Gamma Distribution

Distribution of the Inverse of a Gamma Distribution: If $X \sim \text{Gamma}(\alpha, \beta)$, then $\frac{1}{X} \sim \text{Invgamma}(\alpha, \beta)$.

$$Y \sim \text{Invgamma}(\alpha, \beta)$$

$$y > 0$$


shape parameter: $\alpha > 0$

scale parameter: $\beta > 0$

$$p(y|\alpha, \beta) = \frac{\beta^\alpha}{\Gamma(\alpha)} y^{-(\alpha+1)} e^{-\frac{\beta}{y}}$$

$$E(Y) = \frac{\beta}{\alpha-1} \text{ for } \alpha > 1$$

$$\text{Var}(Y) = \frac{\beta^2}{(\alpha-1)^2(\alpha-2)} \text{ for } \alpha > 2$$



The Dirichlet Distribution

The Dirichlet Distribution

$$Y \sim \text{Dirichlet}(\alpha_1, \dots, \alpha_k)$$

The Dirichlet Distribution

$$Y \sim \text{Dirichlet}(\alpha_1, \dots, \alpha_k)$$

$$y_j \in [0, 1]; \quad \sum_{j=1}^k y_j = 1$$

The Dirichlet Distribution

$$Y \sim \text{Dirichlet}(\alpha_1, \dots, \alpha_k)$$

$$y_j \in [0, 1]; \quad \sum_{j=1}^k y_j = 1$$

$$\alpha \text{ parameters: } \alpha_j > 0; \quad \sum_{j=1}^k \alpha_j \equiv \alpha_0$$

The Dirichlet Distribution

$$Y \sim \text{Dirichlet}(\alpha_1, \dots, \alpha_k)$$

$$y_j \in [0, 1]; \quad \sum_{j=1}^k y_j = 1$$

$$\alpha \text{ parameters: } \alpha_j > 0; \quad \sum_{j=1}^k \alpha_j \equiv \alpha_0$$

$$p(\mathbf{y}|\boldsymbol{\alpha}) = \frac{\Gamma(\alpha_1 + \dots + \alpha_k)}{\Gamma(\alpha_1) \dots \Gamma(\alpha_k)} y_1^{\alpha_1-1} \dots y_k^{\alpha_k-1}$$

The Dirichlet Distribution

$$Y \sim \text{Dirichlet}(\alpha_1, \dots, \alpha_k)$$

$$y_j \in [0, 1]; \quad \sum_{j=1}^k y_j = 1$$

$$\alpha \text{ parameters: } \alpha_j > 0; \quad \sum_{j=1}^k \alpha_j \equiv \alpha_0$$

$$p(\mathbf{y}|\boldsymbol{\alpha}) = \frac{\Gamma(\alpha_1 + \dots + \alpha_k)}{\Gamma(\alpha_1) \dots \Gamma(\alpha_k)} y_1^{\alpha_1-1} \dots y_k^{\alpha_k-1}$$

$$E(Y_j) = \frac{\alpha_j}{\alpha_0}$$

The Dirichlet Distribution

$$Y \sim \text{Dirichlet}(\alpha_1, \dots, \alpha_k)$$

$$y_j \in [0, 1]; \quad \sum_{j=1}^k y_j = 1$$

$$\alpha \text{ parameters: } \alpha_j > 0; \quad \sum_{j=1}^k \alpha_j \equiv \alpha_0$$

$$p(\mathbf{y}|\boldsymbol{\alpha}) = \frac{\Gamma(\alpha_1 + \dots + \alpha_k)}{\Gamma(\alpha_1) \dots \Gamma(\alpha_k)} y_1^{\alpha_1-1} \dots y_k^{\alpha_k-1}$$

$$E(Y_j) = \frac{\alpha_j}{\alpha_0}$$

$$\text{Var}(Y_j) = \frac{\alpha_j(\alpha_0 - \alpha_j)}{\alpha_0^2(\alpha_0 + 1)}$$

The Dirichlet Distribution

$$Y \sim \text{Dirichlet}(\alpha_1, \dots, \alpha_k)$$

$$y_j \in [0, 1]; \quad \sum_{j=1}^k y_j = 1$$

$$\alpha \text{ parameters: } \alpha_j > 0; \quad \sum_{j=1}^k \alpha_j \equiv \alpha_0$$

$$p(\mathbf{y}|\boldsymbol{\alpha}) = \frac{\Gamma(\alpha_1 + \dots + \alpha_k)}{\Gamma(\alpha_1) \dots \Gamma(\alpha_k)} y_1^{\alpha_1-1} \dots y_k^{\alpha_k-1}$$

$$E(Y_j) = \frac{\alpha_j}{\alpha_0}$$

$$\text{Var}(Y_j) = \frac{\alpha_j(\alpha_0 - \alpha_j)}{\alpha_0^2(\alpha_0 + 1)}$$

$$\text{Cov}(Y_i, Y_j) = -\frac{\alpha_i \alpha_j}{\alpha_0^2(\alpha_0 + 1)}$$