

Delta Method and Generalized Linear Models

Patrick Lam

Suppose we have a random variable X , and we know $E(X)$ and $\text{Var}(X)$.

Suppose we have a random variable X , and we know $E(X)$ and $\text{Var}(X)$.

Let Y be another random variable such that $Y = g(X)$.

Suppose we have a random variable X , and we know $E(X)$ and $\text{Var}(X)$.

Let Y be another random variable such that $Y = g(X)$.

How do we find $E(Y)$ and $\text{Var}(Y)$?

Suppose we have a random variable X , and we know $E(X)$ and $\text{Var}(X)$.

Let Y be another random variable such that $Y = g(X)$.

How do we find $E(Y)$ and $\text{Var}(Y)$? (useful for reparameterizations and quantities of interest)

Suppose we have a random variable X , and we know $E(X)$ and $\text{Var}(X)$.

Let Y be another random variable such that $Y = g(X)$.

How do we find $E(Y)$ and $\text{Var}(Y)$? (useful for reparameterizations and quantities of interest)

- Find $E(Y)$ via $\int_{-\infty}^{\infty} g(x)p(x)dx$

Suppose we have a random variable X , and we know $E(X)$ and $\text{Var}(X)$.

Let Y be another random variable such that $Y = g(X)$.

How do we find $E(Y)$ and $\text{Var}(Y)$? (useful for reparameterizations and quantities of interest)

- ▶ Find $E(Y)$ via $\int_{-\infty}^{\infty} g(x)p(x)dx$
- ▶ Simulation

Suppose we have a random variable X , and we know $E(X)$ and $\text{Var}(X)$.

Let Y be another random variable such that $Y = g(X)$.

How do we find $E(Y)$ and $\text{Var}(Y)$? (useful for reparameterizations and quantities of interest)

- ▶ Find $E(Y)$ via $\int_{-\infty}^{\infty} g(x)p(x)dx$
- ▶ Simulation
- ▶ Delta Method

If $g(X)$ is linear...

If $g(X)$ is linear...

this is pretty straightforward:

If $g(X)$ is linear...

this is pretty straightforward:

$$Y = a + bX$$

If $g(X)$ is linear...

this is pretty straightforward:

$$\begin{aligned} Y &= a + bX \\ E(Y) &= a + bE(X) \end{aligned}$$

If $g(X)$ is linear...

this is pretty straightforward:

$$Y = a + bX$$

$$E(Y) = a + bE(X)$$

$$\text{Var}(Y) = b^2 \text{Var}(X)$$

If $g(X)$ is linear...

this is pretty straightforward:

$$\begin{aligned} Y &= a + bX \\ E(Y) &= a + bE(X) \\ \text{Var}(Y) &= b^2\text{Var}(X) \end{aligned}$$

What if $g(X)$ isn't linear?

If $g(X)$ is linear...

this is pretty straightforward:

$$\begin{aligned} Y &= a + bX \\ E(Y) &= a + bE(X) \\ \text{Var}(Y) &= b^2\text{Var}(X) \end{aligned}$$

What if $g(X)$ isn't linear?

We will use the **delta method**, which relies on a linear approximation to $g(X)$ near the mean of X .

Delta Method

Delta Method

Denote μ_X as the mean of X .

Delta Method

Denote μ_X as the mean of X . We use a first-order Taylor series approximation around μ_X :

Delta Method

Denote μ_X as the mean of X . We use a first-order Taylor series approximation around μ_X :

$$Y = g(X)$$

Delta Method

Denote μ_X as the mean of X . We use a first-order Taylor series approximation around μ_X :

$$\begin{aligned} Y &= g(X) \\ &\approx g(\mu_X) + (X - \mu_X)g'(\mu_X) \end{aligned}$$

Delta Method

Denote μ_X as the mean of X . We use a first-order Taylor series approximation around μ_X :

$$\begin{aligned} Y &= g(X) \\ &\approx g(\mu_X) + (X - \mu_X)g'(\mu_X) \end{aligned}$$

$$E(Y) \approx E[g(\mu_X)] + E[(X - \mu_X)g'(\mu_X)]$$

Delta Method

Denote μ_X as the mean of X . We use a first-order Taylor series approximation around μ_X :

$$\begin{aligned} Y &= g(X) \\ &\approx g(\mu_X) + (X - \mu_X)g'(\mu_X) \end{aligned}$$

$$\begin{aligned} E(Y) &\approx E[g(\mu_X)] + E[(X - \mu_X)g'(\mu_X)] \\ &= g(\mu_X) \text{ since } E(X - \mu_X) = 0 \end{aligned}$$

Delta Method

Denote μ_X as the mean of X . We use a first-order Taylor series approximation around μ_X :

$$\begin{aligned} Y &= g(X) \\ &\approx g(\mu_X) + (X - \mu_X)g'(\mu_X) \end{aligned}$$

$$\begin{aligned} E(Y) &\approx E[g(\mu_X)] + E[(X - \mu_X)g'(\mu_X)] \\ &= g(\mu_X) \text{ since } E(X - \mu_X) = 0 \end{aligned}$$

$$\text{Var}(Y) \approx \text{Var}[g(\mu_X)] + \text{Var}[(X - \mu_X)g'(\mu_X)]$$

Delta Method

Denote μ_X as the mean of X . We use a first-order Taylor series approximation around μ_X :

$$\begin{aligned} Y &= g(X) \\ &\approx g(\mu_X) + (X - \mu_X)g'(\mu_X) \end{aligned}$$

$$\begin{aligned} E(Y) &\approx E[g(\mu_X)] + E[(X - \mu_X)g'(\mu_X)] \\ &= g(\mu_X) \text{ since } E(X - \mu_X) = 0 \end{aligned}$$

$$\begin{aligned} \text{Var}(Y) &\approx \text{Var}[g(\mu_X)] + \text{Var}[(X - \mu_X)g'(\mu_X)] \\ &= 0 + \text{Var}[Xg'(\mu_X)] - \text{Var}[\mu_X g'(\mu_X)] \end{aligned}$$

Delta Method

Denote μ_X as the mean of X . We use a first-order Taylor series approximation around μ_X :

$$\begin{aligned} Y &= g(X) \\ &\approx g(\mu_X) + (X - \mu_X)g'(\mu_X) \end{aligned}$$

$$\begin{aligned} E(Y) &\approx E[g(\mu_X)] + E[(X - \mu_X)g'(\mu_X)] \\ &= g(\mu_X) \text{ since } E(X - \mu_X) = 0 \end{aligned}$$

$$\begin{aligned} \text{Var}(Y) &\approx \text{Var}[g(\mu_X)] + \text{Var}[(X - \mu_X)g'(\mu_X)] \\ &= 0 + \text{Var}[Xg'(\mu_X)] - \text{Var}[\mu_X g'(\mu_X)] \\ &= \text{Var}(X)[g'(\mu_X)]^2 \end{aligned}$$

One More Step...

One More Step...

We have

$$E(Y) \approx g(\mu_X)$$

One More Step...

We have

$$E(Y) \approx g(\mu_X)$$

but we know generally (from Jensen's inequality)

$$E(g(X)) \neq g(E(X))$$

One More Step...

We have

$$E(Y) \approx g(\mu_X)$$

but we know generally (from Jensen's inequality)

$$E(g(X)) \neq g(E(X))$$

so we use the second order Taylor expansion for $E(Y)$

One More Step...

We have

$$E(Y) \approx g(\mu_X)$$

but we know generally (from Jensen's inequality)

$$E(g(X)) \neq g(E(X))$$

so we use the second order Taylor expansion for $E(Y)$

$$Y \approx g(\mu_X) + (X - \mu_X)g'(\mu_X) + \frac{1}{2}(X - \mu_X)^2 g''(\mu_X)$$

One More Step...

We have

$$E(Y) \approx g(\mu_X)$$

but we know generally (from Jensen's inequality)

$$E(g(X)) \neq g(E(X))$$

so we use the second order Taylor expansion for $E(Y)$

$$Y \approx g(\mu_X) + (X - \mu_X)g'(\mu_X) + \frac{1}{2}(X - \mu_X)^2 g''(\mu_X)$$

$$E(Y) \approx E[g(\mu_X)] + E[(X - \mu_X)g'(\mu_X)] + \frac{1}{2}E[(X - \mu_X)^2 g''(\mu_X)]$$

One More Step...

We have

$$E(Y) \approx g(\mu_X)$$

but we know generally (from Jensen's inequality)

$$E(g(X)) \neq g(E(X))$$

so we use the second order Taylor expansion for $E(Y)$

$$Y \approx g(\mu_X) + (X - \mu_X)g'(\mu_X) + \frac{1}{2}(X - \mu_X)^2 g''(\mu_X)$$

$$\begin{aligned} E(Y) &\approx E[g(\mu_X)] + E[(X - \mu_X)g'(\mu_X)] + \frac{1}{2}E[(X - \mu_X)^2 g''(\mu_X)] \\ &= g(\mu_X) + \frac{1}{2}\text{Var}(X)g''(\mu_X) \end{aligned}$$

So we have

$$\begin{aligned} E(Y) &\approx g(\mu_X) + \frac{1}{2}\text{Var}(X)g''(\mu_X) \\ \text{Var}(Y) &\approx \text{Var}(X)[g'(\mu_X)]^2 \end{aligned}$$

So we have

$$\begin{aligned} E(Y) &\approx g(\mu_X) + \frac{1}{2}\text{Var}(X)g''(\mu_X) \\ \text{Var}(Y) &\approx \text{Var}(X)[g'(\mu_X)]^2 \end{aligned}$$

How good the approximations are depend on how nonlinear $g(X)$ is in the neighborhood of μ_X and on the size of $\text{Var}(X)$.

Generalized Linear Models

Generalized Linear Models

All of the models we've talked about so far (and for the rest of the class) belong to a class called **generalized linear models (GLM)**.

Generalized Linear Models

All of the models we've talked about so far (and for the rest of the class) belong to a class called **generalized linear models (GLM)**.

Three elements of a GLM:

Generalized Linear Models

All of the models we've talked about so far (and for the rest of the class) belong to a class called **generalized linear models (GLM)**.

Three elements of a GLM:

- ▶ A distribution for Y

Generalized Linear Models

All of the models we've talked about so far (and for the rest of the class) belong to a class called **generalized linear models (GLM)**.

Three elements of a GLM:

- ▶ A distribution for Y
- ▶ A linear predictor $X\beta$

Generalized Linear Models

All of the models we've talked about so far (and for the rest of the class) belong to a class called **generalized linear models (GLM)**.

Three elements of a GLM:

- ▶ A distribution for Y
- ▶ A linear predictor $X\beta$
- ▶ A link function that relates the linear predictor to the mean of the distribution.

Generalized Linear Models

All of the models we've talked about so far (and for the rest of the class) belong to a class called **generalized linear models (GLM)**.

Three elements of a GLM:

- ▶ A distribution for Y
- ▶ A linear predictor $X\beta$
- ▶ A link function that relates the linear predictor to the mean of the distribution.

Steps to running a GLM:

1. Specify a distribution for Y

1. Specify a distribution for Y

Assume our data was generated from some distribution.

1. Specify a distribution for Y

Assume our data was generated from some distribution.

Examples:

1. Specify a distribution for Y

Assume our data was generated from some distribution.

Examples:

- ▶ Continuous and Unbounded:

1. Specify a distribution for Y

Assume our data was generated from some distribution.

Examples:

- ▶ Continuous and Unbounded: Normal

1. Specify a distribution for Y

Assume our data was generated from some distribution.

Examples:

- ▶ Continuous and Unbounded: Normal
- ▶ Binary:

1. Specify a distribution for Y

Assume our data was generated from some distribution.

Examples:

- ▶ Continuous and Unbounded: Normal
- ▶ Binary: Bernoulli

1. Specify a distribution for Y

Assume our data was generated from some distribution.

Examples:

- ▶ Continuous and Unbounded: Normal
- ▶ Binary: Bernoulli
- ▶ Event Count:

1. Specify a distribution for Y

Assume our data was generated from some distribution.

Examples:

- ▶ Continuous and Unbounded: Normal
- ▶ Binary: Bernoulli
- ▶ Event Count: Poisson

1. Specify a distribution for Y

Assume our data was generated from some distribution.

Examples:

- ▶ Continuous and Unbounded: Normal
- ▶ Binary: Bernoulli
- ▶ Event Count: Poisson
- ▶ Duration:

1. Specify a distribution for Y

Assume our data was generated from some distribution.

Examples:

- ▶ Continuous and Unbounded: Normal
- ▶ Binary: Bernoulli
- ▶ Event Count: Poisson
- ▶ Duration: Exponential

1. Specify a distribution for Y

Assume our data was generated from some distribution.

Examples:

- ▶ Continuous and Unbounded: Normal
- ▶ Binary: Bernoulli
- ▶ Event Count: Poisson
- ▶ Duration: Exponential
- ▶ Ordered Categories:

1. Specify a distribution for Y

Assume our data was generated from some distribution.

Examples:

- ▶ Continuous and Unbounded: Normal
- ▶ Binary: Bernoulli
- ▶ Event Count: Poisson
- ▶ Duration: Exponential
- ▶ Ordered Categories: Normal with observation mechanism

1. Specify a distribution for Y

Assume our data was generated from some distribution.

Examples:

- ▶ Continuous and Unbounded: Normal
- ▶ Binary: Bernoulli
- ▶ Event Count: Poisson
- ▶ Duration: Exponential
- ▶ Ordered Categories: Normal with observation mechanism
- ▶ Unordered Categories:

1. Specify a distribution for Y

Assume our data was generated from some distribution.

Examples:

- ▶ Continuous and Unbounded: Normal
- ▶ Binary: Bernoulli
- ▶ Event Count: Poisson
- ▶ Duration: Exponential
- ▶ Ordered Categories: Normal with observation mechanism
- ▶ Unordered Categories: Multinomial

2. Specify a linear predictor

2. Specify a linear predictor

$$X\beta = \beta_0 + x_1\beta_1 + x_2\beta_2 + \cdots + x_k\beta_k$$

3. Specify a link function

3. Specify a link function

The link function relates the linear predictor to the mean of the distribution for Y .

3. Specify a link function

The link function relates the linear predictor to the mean of the distribution for Y .

Let $g(\cdot)$ be the link function and let $E(Y) = \theta$ be the mean of distribution for Y .

3. Specify a link function

The link function relates the linear predictor to the mean of the distribution for Y .

Let $g(\cdot)$ be the link function and let $E(Y) = \theta$ be the mean of distribution for Y .

$$g(\theta) = X\beta$$

3. Specify a link function

The link function relates the linear predictor to the mean of the distribution for Y .

Let $g(\cdot)$ be the link function and let $E(Y) = \theta$ be the mean of distribution for Y .

$$\begin{aligned}g(\theta) &= X\beta \\ \theta &= g^{-1}(X\beta)\end{aligned}$$

3. Specify a link function

The link function relates the linear predictor to the mean of the distribution for Y .

Let $g(\cdot)$ be the link function and let $E(Y) = \theta$ be the mean of distribution for Y .

$$\begin{aligned}g(\theta) &= X\beta \\ \theta &= g^{-1}(X\beta)\end{aligned}$$

Note that we usually use the **inverse link function** $g^{-1}(X\beta)$ rather than the link function.

3. Specify a link function

The link function relates the linear predictor to the mean of the distribution for Y .

Let $g(\cdot)$ be the link function and let $E(Y) = \theta$ be the mean of distribution for Y .

$$\begin{aligned}g(\theta) &= X\beta \\ \theta &= g^{-1}(X\beta)\end{aligned}$$

Note that we usually use the **inverse link function** $g^{-1}(X\beta)$ rather than the link function.

This is the systematic component that we've been talking about all along.

Example Link Functions

Example Link Functions

Identity:

Example Link Functions

Identity:

- ▶ Link: $\mu = X\beta$

Example Link Functions

Identity:

- ▶ Link: $\mu = X\beta$
- ▶ Inverse Link: $\mu = X\beta$

Example Link Functions

Identity:

- ▶ Link: $\mu = X\beta$
- ▶ Inverse Link: $\mu = X\beta$

Inverse:

Example Link Functions

Identity:

- ▶ Link: $\mu = X\beta$
- ▶ Inverse Link: $\mu = X\beta$

Inverse:

- ▶ Link: $\lambda^{-1} = X\beta$

Example Link Functions

Identity:

- ▶ Link: $\mu = X\beta$
- ▶ Inverse Link: $\mu = X\beta$

Inverse:

- ▶ Link: $\lambda^{-1} = X\beta$
- ▶ Inverse Link: $\lambda = (X\beta)^{-1}$

Example Link Functions

Identity:

- ▶ Link: $\mu = X\beta$
- ▶ Inverse Link: $\mu = X\beta$

Inverse:

- ▶ Link: $\lambda^{-1} = X\beta$
- ▶ Inverse Link: $\lambda = (X\beta)^{-1}$

Logit:

Example Link Functions

Identity:

- ▶ Link: $\mu = X\beta$
- ▶ Inverse Link: $\mu = X\beta$

Inverse:

- ▶ Link: $\lambda^{-1} = X\beta$
- ▶ Inverse Link: $\lambda = (X\beta)^{-1}$

Logit:

- ▶ Link: $\ln\left(\frac{\pi}{1-\pi}\right) = X\beta$

Example Link Functions

Identity:

- ▶ Link: $\mu = X\beta$
- ▶ Inverse Link: $\mu = X\beta$

Inverse:

- ▶ Link: $\lambda^{-1} = X\beta$
- ▶ Inverse Link: $\lambda = (X\beta)^{-1}$

Logit:

- ▶ Link: $\ln\left(\frac{\pi}{1-\pi}\right) = X\beta$
- ▶ Inverse Link: $\pi = \frac{1}{1+e^{-X\beta}}$

Example Link Functions

Identity:

- ▶ Link: $\mu = X\beta$
- ▶ Inverse Link: $\mu = X\beta$

Inverse:

- ▶ Link: $\lambda^{-1} = X\beta$
- ▶ Inverse Link: $\lambda = (X\beta)^{-1}$

Logit:

- ▶ Link: $\ln\left(\frac{\pi}{1-\pi}\right) = X\beta$
- ▶ Inverse Link: $\pi = \frac{1}{1+e^{-X\beta}}$

Probit:

Example Link Functions

Identity:

- ▶ Link: $\mu = X\beta$
- ▶ Inverse Link: $\mu = X\beta$

Inverse:

- ▶ Link: $\lambda^{-1} = X\beta$
- ▶ Inverse Link: $\lambda = (X\beta)^{-1}$

Logit:

- ▶ Link: $\ln\left(\frac{\pi}{1-\pi}\right) = X\beta$
- ▶ Inverse Link: $\pi = \frac{1}{1+e^{-X\beta}}$

Probit:

- ▶ Link: $\Phi^{-1}(\pi) = X\beta$

Example Link Functions

Identity:

- ▶ Link: $\mu = X\beta$
- ▶ Inverse Link: $\mu = X\beta$

Inverse:

- ▶ Link: $\lambda^{-1} = X\beta$
- ▶ Inverse Link: $\lambda = (X\beta)^{-1}$

Logit:

- ▶ Link: $\ln\left(\frac{\pi}{1-\pi}\right) = X\beta$
- ▶ Inverse Link: $\pi = \frac{1}{1+e^{-X\beta}}$

Probit:

- ▶ Link: $\Phi^{-1}(\pi) = X\beta$
- ▶ Inverse Link: $\pi = \Phi(X\beta)$

Example Link Functions

Identity:

- ▶ Link: $\mu = X\beta$
- ▶ Inverse Link: $\mu = X\beta$

Inverse:

- ▶ Link: $\lambda^{-1} = X\beta$
- ▶ Inverse Link: $\lambda = (X\beta)^{-1}$

Logit:

- ▶ Link: $\ln\left(\frac{\pi}{1-\pi}\right) = X\beta$
- ▶ Inverse Link: $\pi = \frac{1}{1+e^{-X\beta}}$

Probit:

- ▶ Link: $\Phi^{-1}(\pi) = X\beta$
- ▶ Inverse Link: $\pi = \Phi(X\beta)$

Log:

Example Link Functions

Identity:

- ▶ Link: $\mu = X\beta$
- ▶ Inverse Link: $\mu = X\beta$

Inverse:

- ▶ Link: $\lambda^{-1} = X\beta$
- ▶ Inverse Link: $\lambda = (X\beta)^{-1}$

Logit:

- ▶ Link: $\ln\left(\frac{\pi}{1-\pi}\right) = X\beta$
- ▶ Inverse Link: $\pi = \frac{1}{1+e^{-X\beta}}$

Probit:

- ▶ Link: $\Phi^{-1}(\pi) = X\beta$
- ▶ Inverse Link: $\pi = \Phi(X\beta)$

Log:

- ▶ Link: $\ln(\lambda) = X\beta$

Example Link Functions

Identity:

- ▶ Link: $\mu = X\beta$
- ▶ Inverse Link: $\mu = X\beta$

Inverse:

- ▶ Link: $\lambda^{-1} = X\beta$
- ▶ Inverse Link: $\lambda = (X\beta)^{-1}$

Logit:

- ▶ Link: $\ln\left(\frac{\pi}{1-\pi}\right) = X\beta$
- ▶ Inverse Link: $\pi = \frac{1}{1+e^{-X\beta}}$

Probit:

- ▶ Link: $\Phi^{-1}(\pi) = X\beta$
- ▶ Inverse Link: $\pi = \Phi(X\beta)$

Log:

- ▶ Link: $\ln(\lambda) = X\beta$
- ▶ Inverse Link: $\lambda = \exp(X\beta)$

4. Estimate Parameters via ML

4. Estimate Parameters via ML

Do it.

5. Quantities of Interest

5. Quantities of Interest

1. Simulate parameters from multivariate normal.

5. Quantities of Interest

1. Simulate parameters from multivariate normal.
2. Run $X\beta$ through inverse link function to get expected values.

5. Quantities of Interest

1. Simulate parameters from multivariate normal.
2. Run $X\beta$ through inverse link function to get expected values.
3. Draw from distribution of Y for predicted values.

Binary Dependent Variable

Let our dependent variable be a binary random variable that can take on values of either 0 or 1.

1. Specify a distribution for Y

1. Specify a distribution for Y

$$Y_i \sim \text{Bernoulli}(\pi_i)$$

1. Specify a distribution for Y

$$Y_i \sim \text{Bernoulli}(\pi_i)$$

$$p(y|\boldsymbol{\pi}) = \prod_{i=1}^n \pi_i^{y_i} (1 - \pi_i)^{1-y_i}$$

1. Specify a distribution for Y

$$Y_i \sim \text{Bernoulli}(\pi_i)$$
$$p(y|\boldsymbol{\pi}) = \prod_{i=1}^n \pi_i^{y_i} (1 - \pi_i)^{1-y_i}$$

2. Specify a linear predictor:

1. Specify a distribution for Y

$$Y_i \sim \text{Bernoulli}(\pi_i)$$
$$p(y|\boldsymbol{\pi}) = \prod_{i=1}^n \pi_i^{y_i} (1 - \pi_i)^{1-y_i}$$

2. Specify a linear predictor: $X\beta$

3. Specify a link (or inverse link) function.

3. Specify a link (or inverse link) function.

► Logit: $\pi_i = \frac{1}{1+e^{-x_i\beta}}$

3. Specify a link (or inverse link) function.

- ▶ Logit: $\pi_i = \frac{1}{1+e^{-x_i\beta}}$
- ▶ Probit: $\pi_i = \Phi(x_i\beta)$

3. Specify a link (or inverse link) function.

- ▶ Logit: $\pi_i = \frac{1}{1 + e^{-x_i\beta}}$
- ▶ Probit: $\pi_i = \Phi(x_i\beta)$
- ▶ Complementary Log-log (cloglog): $\pi_i = 1 - \exp(-\exp(X\beta))$

3. Specify a link (or inverse link) function.

- ▶ Logit: $\pi_i = \frac{1}{1 + e^{-x_i\beta}}$
- ▶ Probit: $\pi_i = \Phi(x_i\beta)$
- ▶ Complementary Log-log (cloglog): $\pi_i = 1 - \exp(-\exp(X\beta))$
- ▶ Scobit: $\pi_i = (1 + e^{-x_i\beta})^{-\alpha}$

3. Specify a link (or inverse link) function.

- ▶ Logit: $\pi_i = \frac{1}{1+e^{-x_i\beta}}$
- ▶ Probit: $\pi_i = \Phi(x_i\beta)$
- ▶ Complementary Log-log (cloglog): $\pi_i = 1 - \exp(-\exp(X\beta))$
- ▶ Scobit: $\pi_i = (1 + e^{-x_i\beta})^{-\alpha}$

4. Estimate parameters via ML.

3. Specify a link (or inverse link) function.

- ▶ Logit: $\pi_i = \frac{1}{1+e^{-x_i\beta}}$
- ▶ Probit: $\pi_i = \Phi(x_i\beta)$
- ▶ Complementary Log-log (cloglog): $\pi_i = 1 - \exp(-\exp(X\beta))$
- ▶ Scobit: $\pi_i = (1 + e^{-x_i\beta})^{-\alpha}$

4. Estimate parameters via ML.

$$l(\beta|\mathbf{y}) = \sum_{i=1}^n y_i \ln \left(\frac{1}{1 + e^{-x_i\beta}} \right) + (1 - y_i) \ln \left(1 - \frac{1}{1 + e^{-x_i\beta}} \right)$$

3. Specify a link (or inverse link) function.

- ▶ Logit: $\pi_i = \frac{1}{1 + e^{-x_i\beta}}$
- ▶ Probit: $\pi_i = \Phi(x_i\beta)$
- ▶ Complementary Log-log (cloglog): $\pi_i = 1 - \exp(-\exp(X\beta))$
- ▶ Scobit: $\pi_i = (1 + e^{-x_i\beta})^{-\alpha}$

4. Estimate parameters via ML.


$$l(\beta|\mathbf{y}) = \sum_{i=1}^n y_i \ln \left(\frac{1}{1 + e^{-x_i\beta}} \right) + (1 - y_i) \ln \left(1 - \frac{1}{1 + e^{-x_i\beta}} \right)$$

5. Simulate Quantities of Interest

Identification


Identification

Suppose we have the following log-likelihood function:



Identification

Suppose we have the following log-likelihood function:



What's wrong with this?

The Identification Problem

The Identification Problem

There are more than one set of parameters that give the same maximum likelihood value, so our model is **unidentified**.

The Identification Problem

There are more than one set of parameters that give the same maximum likelihood value, so our model is **unidentified**.

Ordered Probit/Logit:

The Identification Problem

There are more than one set of parameters that give the same maximum likelihood value, so our model is **unidentified**.

Ordered Probit/Logit:

- ▶ If we estimate all the β s and τ s, we can get many sets of parameters that have the same likelihood.

The Identification Problem

There are more than one set of parameters that give the same maximum likelihood value, so our model is **unidentified**.

Ordered Probit/Logit:

- ▶ If we estimate all the β s and τ s, we can get many sets of parameters that have the same likelihood.
- ▶ We can make our model identified in two ways:

The Identification Problem

There are more than one set of parameters that give the same maximum likelihood value, so our model is **unidentified**.

Ordered Probit/Logit:

- ▶ If we estimate all the β s and τ s, we can get many sets of parameters that have the same likelihood.
- ▶ We can make our model identified in two ways:
 - ▶ Fix $\beta_0 = 0$ and estimate all the τ s (basically don't estimate an intercept)

The Identification Problem

There are more than one set of parameters that give the same maximum likelihood value, so our model is **unidentified**.

Ordered Probit/Logit:

- ▶ If we estimate all the β s and τ s, we can get many sets of parameters that have the same likelihood.
- ▶ We can make our model identified in two ways:
 - ▶ Fix $\beta_0 = 0$ and estimate all the τ s (basically don't estimate an intercept)
 - ▶ Fix $\tau_1 = 0$ and estimate an intercept.