

Task1: Oscar student problem ¶

Hypothesis:

Null(Ho)= The class average score be 71 and above


Alternate(H1)= The class average might be lower

We use T statistics since number of samples less than 30 and also we don't know population standard deviation

```
In [1]: 1 from scipy.stats import t,f,f_oneway
2 import numpy as np
3 import matplotlib.pyplot as plt

In [2]: 1 scores=[61,93,75,67,84,95]
2 mean_s=np.mean(scores)
3 alpha=0.1
4 dof=6-1
5 s_samp=np.std(scores)
6
7 t_crit=t.ppf(1-alpha,dof)
8
9 t_stats=(mean_s-71)/(s_samp/(dof)**0.5)
10
11 p_val=t.sf(t_stats,dof)
12
13 print("The test critical value is = ",t_crit)
14 print("The test statistics for given sample = ",t_stats)
15 print("The p value for this sample = ",p_val)
16
17
18 nr=np.arange(-4,4,.05)
19 plt.plot(nr,t.pdf(nr,5),color="purple")
20 plt.title("Normal distribution plot")
21 plt.show()
```

The test critical value is = 1.4758840487820273
The test statistics for given sample = 1.4430501569317138
The p value for this sample = 0.10429845495147679



```
In [3]: 1 if t_stats > t_crit and p_val<alpha :
2     print("Reject Null Hypothesis")
3 else :
4     print("Accept Null Hypothesis")
```

Accept Null Hypothesis

Thus from our findings and hypothesis test we can see that we are getting acceptance of null hypothesis so there is quite a chance that oscar will get an mark of 71 with 90% confidence level and Ho is True


Task2: Oscar wishes to identify whether two batches can get same average of marks

Ho(Null)= The null hypothesis is considered as both means of classes are equal

H1(Alternate)= The alternate hypothesis is considered as either its higher or lower and cannot be same

```
In [4]: 1 b1=[9,9,9,36,45,48,51,57,69,72,96]
2 b2=[60,39,39,60,79,96,69,60,75,45,90]
3
4 x1=np.mean(b1)
5 x2=np.mean(b2)
6
7 n1=11
8 n2=11
9
10 v1=(np.std(b1))**2
11 v2=(np.std(b2))**2
12
13 alpha=0.1
14 dof=22-2
15
16 t_crit=t.ppf(1-alpha/2,dof)
17 print("The test critical value is = ",t_crit)
18
19 numerator = abs(x1-x2)
20 denominator = (v1*(n1-1) + v2*(n2-1))**0.5*(1/n1 +1/n2)**0.5
21
22 t_stats=numerator/denominator
23 print("The test statistics values is = ",t_stats)
24
25 p_val= t.sf(abs(t_stats), dof)*2
26 print("The p value for this samples ", p_val)
27
28 nr=np.arange(-4,4,.05)
29 plt.plot(nr,t.pdf(nr,21),color="orange")
30 plt.title("Normal distribution plot")
31 plt.show()
```

The test critical value is = 1.7247182429207857
The test statistics values is = 0.4346146276529854
The p value for this samples 0.6684953273749834



```
In [5]: 1 if (t_stats > t_crit) and (p_val<alpha) :
2     print("Reject Null Hypothesis")
3 else :
4     print("Accept Null Hypothesis")
5 print()
6
7 lhs=abs(x1 - x2) - 1.645*((v1/n1 + v2/n2))
8 rhs=(x1 - x2) + 1.645*((v1/n1 + v2/n2))
9 print("The difference between two means = ",rhs+lhs)
```

Accept Null Hypothesis

The difference between two means = 0.0

Decision made:

Since we are accepting Null hypothesis in this case we can conclude that the means for both the batches are same and do not differ

Task3: Oscar wishes to identify the same as task 2 for task 3 with 4 batches


Hypothesis :

Null(Ho): The means of all the samples remain the same

H1(Alternate): The alternate hypothesis is to claim the means are not equal to each other

```
In [6]: 1 b3=[38,61,62,89,86,50,85,55,80,69,95]
2 b4=[45,55,57,88,78,60,35,97,58,88,78]
3 dfc = 4-1
4 dfe = 44-4
5
6 f_critical =f.ppf(1-0.01,dfc,dfe) # 99% confidence Level
7 print('F-critical for the 4 samples =', f_critical)
8
9 f_statistics, pvalue= f_oneway(b1,b2,b3,b4)
10 print('F-statistics for the 4 samples =', f_statistics)
11 print("p value for 4 samples = ",pvalue)
12
13 fig = plt.figure(figsize =(10,7))
14 data=[b1,b2,b3,b4]
15 plt.boxplot(data, showmeans= True)
16 plt.xticks([1,2,3,4], [ 'batch-1','batch-2','batch-3','batch-4'])
17 plt.show()
18
19
```

F-critical for the 4 samples = 4.312569212492142
F-statistics for the 4 samples = 2.8403108468764224
p value for 4 samples = 0.04991227386422173



```
In [7]: 1 if f_statistics > f_critical and pvalue< 0.01 :
2     print("Reject Null Hypothesis")
3 else :
4     print("Accept Null Hypothesis")
```

Accept Null Hypothesis

Decision :

From above hypothesis null we can see that its acceptable so we can state that the averages of all the batches are same