

Spectra of graphs

Andries E. Brouwer

Willem H. Haemers

Contents

1	Graph spectrum	11
1.1	Matrices associated to a graph	11
1.2	The spectrum of a graph	11
1.2.1	Characteristic polynomial	13
1.3	The spectrum of an undirected graph	13
1.3.1	Regular graphs	13
1.3.2	Complements	14
1.3.3	Walks	14
1.3.4	Diameter	14
1.3.5	Spanning trees	14
1.3.6	Bipartite graphs	15
1.3.7	Connectedness	16
1.4	Spectrum of some graphs	17
1.4.1	The complete graph	17
1.4.2	The complete bipartite graph	17
1.4.3	The cycle	17
1.4.4	The path	18
1.4.5	Line graphs	18
1.4.6	Cartesian products	18
1.4.7	Kronecker products and bipartite double	19
1.4.8	Cayley graphs	19
1.5	Decompositions	19
1.5.1	Decomposing K_{10} into Petersen graphs	20
1.5.2	Decomposing K_n into complete bipartite graphs	20
1.6	Automorphisms	20
1.7	Algebraic connectivity	21
1.8	Cospectral graphs	22
1.8.1	The 4-cube	22
1.8.2	Seidel switching	23
1.8.3	Godsil-McKay switching	23
1.8.4	Reconstruction	24
1.9	Very small graphs	24
1.10	Exercises	25
2	Linear algebra	27
2.1	Simultaneous diagonalization	27
2.2	Perron-Frobenius Theory	27
2.3	Equitable partitions	30

2.3.1	Equitable and almost equitable partitions of graphs . . .	30
2.4	The Rayleigh quotient	31
2.5	Interlacing	31
2.6	Schur's inequality	33
2.7	Schur complements	33
2.8	The Courant-Weyl inequalities	34
2.9	Gram matrices	35
2.10	Diagonally dominant matrices	36
2.10.1	Geršgorin circles	36
2.11	Projections	37
2.12	Exercises	37
3	Graph spectrum (II)	39
3.1	The largest eigenvalue	39
3.1.1	Graphs with largest eigenvalue at most 2	40
3.1.2	Subdividing an edge	41
3.1.3	The Kelmans operation	42
3.2	Interlacing	42
3.3	Regular graphs	43
3.4	Bipartite graphs	43
3.5	Classification of integral cubic graphs	44
3.6	The largest Laplace eigenvalue	47
3.7	Laplace eigenvalues and degrees	48
3.8	The Grone-Morris Conjecture	50
3.8.1	Threshold graphs	50
3.8.2	Proof of the Grone-Morris Conjecture	50
3.9	The Laplacian for hypergraphs	53
3.10	Applications of eigenvectors	55
3.10.1	Ranking	55
3.10.2	Google Page rank	55
3.10.3	Cutting	56
3.10.4	Graph drawing	56
3.10.5	Clustering	57
3.10.6	Searching an eigenspace	58
3.11	Exercises	58
4	Substructures	61
4.1	Cliques and cocliques	61
4.1.1	Using weighted adjacency matrices	62
4.2	Chromatic number	62
4.2.1	Using weighted adjacency matrices	64
4.2.2	Rank and chromatic number	64
4.3	Shannon capacity	64
4.4	Stars and star complements	65
4.5	The second largest eigenvalue	66
4.5.1	Bounds for the second largest eigenvalue	66
4.5.2	Large regular subgraphs are connected	67
4.5.3	Randomness	67
4.5.4	Expansion	68
4.5.5	Toughness and Hamiltonicity	69

4.5.6	The Petersen graph is not Hamiltonian	70
4.5.7	Diameter bound	70
4.6	Separation	70
4.6.1	Bandwidth	72
4.6.2	Perfect matchings	72
4.7	Block Designs	75
4.8	Polarities	77
4.9	Exercises	77
5	Trees	79
5.1	Characteristic polynomials of trees	79
5.2	Eigenvectors and multiplicities	81
5.3	Sign patterns of eigenvectors of graphs	82
5.4	Sign patterns of eigenvectors of trees	83
5.5	The spectral center of a tree	84
5.6	Integral trees	85
5.7	Exercises	85
6	Groups and graphs	87
6.1	$\Gamma(G, H, S)$	87
6.2	Spectrum	87
6.3	Covers	88
6.4	Cayley sum graphs	90
6.4.1	(3,6)-fullerenes	90
6.5	Exercises	91
7	Euclidean representations	93
7.1	Examples	93
7.2	Euclidean representation	93
7.3	Root lattices	94
7.4	Cameron-Goethals-Seidel-Shult	99
7.5	Exercises	100
8	Strongly regular graphs	101
8.1	Strongly regular graphs	101
8.1.1	Simple examples	101
8.1.2	The Paley graphs	102
8.1.3	Adjacency matrix	103
8.1.4	Imprimitive graphs	103
8.1.5	Parameters	103
8.1.6	The half case and cyclic strongly regular graphs	104
8.1.7	Strongly regular graphs without triangles	104
8.1.8	Further parameter restrictions	105
8.1.9	Strongly regular graphs from permutation groups	106
8.1.10	Strongly regular graphs from quasisymmetric designs	107
8.1.11	Symmetric 2-designs from strongly regular graphs	107
8.1.12	Latin square graphs	107
8.1.13	Partial Geometries	108
8.2	Strongly regular graphs with eigenvalue -2	109
8.3	Connectivity	110

8.4	Cocliques and colorings	112
8.5	Automorphisms	113
8.6	Generalized quadrangles	114
8.6.1	Parameters	114
8.6.2	Constructions of generalized quadrangles	115
8.6.3	Strongly regular graphs from generalized quadrangles	116
8.6.4	Generalized quadrangles with lines of size 3	116
8.7	The (81,20,1,6) strongly regular graph	117
8.7.1	Descriptions	117
8.7.2	Uniqueness	118
8.7.3	Independence and chromatic numbers	119
8.8	Strongly regular graphs and 2-weight codes	120
8.8.1	Codes, graphs and projective sets	120
8.8.2	The correspondence between linear codes and subsets of a projective space	120
8.8.3	The correspondence between projective two-weight codes, subsets of a projective space with two intersection num- bers, and affine strongly regular graphs	121
8.8.4	Duality for affine strongly regular graphs	123
8.8.5	Cyclotomy	124
8.9	Table	126
8.10	Exercises	131
9	Regular two-graphs	133
9.1	Strong graphs	133
9.2	Two-graphs	134
9.3	Regular two-graphs	135
9.3.1	Related strongly regular graphs	137
9.3.2	The regular two-graph on 276 points	138
9.3.3	Coherent subsets	138
9.3.4	Completely regular two-graphs	138
9.4	Conference matrices	139
9.5	Hadamard matrices	140
9.6	Equiangular lines	142
9.6.1	Equiangular lines in \mathbb{R}^d and two-graphs	142
9.6.2	Bounds on equiangular sets of lines in \mathbb{R}^d or \mathbb{C}^d	143
9.6.3	Bounds on sets of lines with few angles and sets of vectors with few distances	144
10	Association schemes	147
10.1	Definition	147
10.2	The Bose-Mesner algebra	148
10.3	The Linear Programming Bound	150
10.4	The Krein parameters	151
10.5	Automorphisms	152
10.5.1	The Moore graph on 3250 vertices	153
10.6	P - and Q -polynomial association schemes	153
10.7	Exercises	155

11 Distance regular graphs	157
11.1 Parameters	157
11.2 Spectrum	157
11.3 Examples	158
11.3.1 Hamming graphs	158
11.3.2 Johnson graphs	159
11.3.3 Grassmann graphs	159
11.3.4 Van Dam-Koolen graphs	159
11.4 Bannai-Ito conjecture	160
11.5 Connectedness	160
11.6 Degree of eigenvalues	160
11.7 Moore graphs and generalized polygons	161
11.8 Primitivity	161
11.9 Euclidean representations	162
11.10 Extremality	162
11.11 Exercises	163
12 p-ranks	165
12.1 Reduction mod p	165
12.2 The minimal polynomial	166
12.3 Bounds for the p -rank	166
12.4 Interesting primes p	167
12.5 Adding a multiple of J	167
12.6 Paley graphs	168
12.7 Strongly regular graphs	169
12.8 Smith Normal Form	171
12.8.1 Smith Normal Form and spectrum	172
13 Spectral characterizations	175
13.1 Generalized adjacency matrices	175
13.2 Constructing cospectral graphs	176
13.2.1 Trees	177
13.2.2 Partial linear spaces	177
13.2.3 GM switching	178
13.2.4 Sunada's method	179
13.3 Enumeration	179
13.3.1 Lower bounds	179
13.3.2 Computer results	180
13.4 DS graphs	181
13.4.1 Spectrum and structure	181
13.4.2 Some DS graphs	183
13.4.3 Line graphs	185
13.5 Distance-regular graphs	187
13.5.1 Strongly regular DS graphs	187
13.5.2 Distance-regularity from the spectrum	188
13.5.3 Distance-regular DS graphs	189
13.6 The method of Wang & Xu	192
13.7 Exercises	193

14 Graphs with few eigenvalues	195
14.1 Regular graphs with four eigenvalues	195
14.2 Three Laplacian eigenvalues	196
14.3 Other matrices with at most three eigenvalues	198
14.3.1 Few Seidel eigenvalues	198
14.3.2 Three adjacency eigenvalues	199
14.3.3 Three signless Laplacian eigenvalues	200
14.4 Exercises	201
References	203
Author Index	223
Subject Index	226

Algebraic graph theory is the branch of mathematics that studies graphs by using algebraic properties of associated matrices. More in particular, *spectral graph theory* studies the relation between graph properties and the spectrum of the adjacency matrix or Laplace matrix. And the theory of *association schemes* and *coherent configurations* studies the algebra generated by associated matrices.

Chapter 1

Graph spectrum

This chapter presents some simple results on graph spectra. We assume the reader to be familiar with elementary linear algebra and graph theory. Throughout J will denote the all-1 matrix, and $\mathbf{1}$ is the all-1 vector.

1.1 Matrices associated to a graph

Let Γ be a graph without multiple edges. The *adjacency matrix* of Γ is the 0-1 matrix A indexed by the vertex set $V\Gamma$ of Γ , where $A_{xy} = 1$ when there is an edge from x to y in Γ and $A_{xy} = 0$ otherwise. Occasionally we consider multigraphs (possibly with loops) in which case A_{xy} equals the number of edges from x to y .

Let Γ be an undirected graph without loops. The (vertex-edge) *incidence matrix* of Γ is the 0-1 matrix M , with rows indexed by the vertices and columns indexed by the edges, where $M_{xe} = 1$ when vertex x is an endpoint of edge e .

Let Γ be a directed graph without loops. The *directed incidence matrix* of Γ is the 0-1 matrix N , with rows indexed by the vertices and columns by the edges, where $N_{xe} = -1, 1, 0$ when x is the head of e , the tail of e , or not on e , respectively.

Let Γ be an undirected graph without loops. The *Laplace matrix* of Γ is the matrix L indexed by the vertex set of Γ , with zero row sums, where $L_{xy} = -A_{xy}$ for $x \neq y$. If D is the diagonal matrix, indexed by the vertex set of Γ such that D_{xx} is the degree (valency) of x , then $L = D - A$. The matrix $Q = D + A$ is called the *signless Laplace matrix* of Γ .

An important property of the Laplace matrix L and the signless Laplace matrix Q is that they are positive semidefinite. Indeed, one has $Q = MM^\top$ and $L = NN^\top$ if M is the incidence matrix of Γ and N the directed incidence matrix of the directed graph obtained by orienting the edges of Γ in an arbitrary way. It follows that for any vector u one has $u^\top Lu = \sum_{xy} (u_x - u_y)^2$ and $u^\top Qu = \sum_{xy} (u_x + u_y)^2$, where the sum is over the edges of Γ .

1.2 The spectrum of a graph

The (ordinary) *spectrum* of a finite graph Γ is by definition the spectrum of the adjacency matrix A , that is, its set of eigenvalues together with their multiplic-

ities. The *Laplace spectrum* of a finite undirected graph without loops is the spectrum of the Laplace matrix L .

The rows and columns of a matrix of order n are numbered from 1 to n , while A is indexed by the vertices of Γ , so that writing down A requires one to assign some numbering to the vertices. However, the spectrum of the matrix obtained does not depend on the numbering chosen. It is the spectrum of the linear transformation A on the vector space K^X of maps from X into K , where X is the vertex set, and K is some field such as \mathbb{R} or \mathbb{C} .

The *characteristic polynomial* of Γ is that of A , that is, the polynomial p_A defined by $p_A(\theta) = \det(\theta I - A)$.

Example Let Γ be the path P_3 with three vertices and two edges. Assigning some arbitrary order to the three vertices of Γ , we find that the adjacency matrix A becomes one of

$$\begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix} \text{ or } \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \text{ or } \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}.$$

The characteristic polynomial is $p_A(\theta) = \theta^3 - 2\theta$. The spectrum is $\sqrt{2}$, 0 , $-\sqrt{2}$. The eigenvectors are:

$$\begin{array}{ccc} \sqrt{2} & 2 & \sqrt{2} \\ \circ & \text{---} & \circ & \text{---} & \circ \end{array} \quad \begin{array}{ccc} 1 & 0 & -1 \\ \circ & \text{---} & \circ & \text{---} & \circ \end{array} \quad \begin{array}{ccc} \sqrt{2} & -2 & \sqrt{2} \\ \circ & \text{---} & \circ & \text{---} & \circ \end{array}.$$

(Here, for an eigenvector u , we write u_x as a label at the vertex x ; one has $Au = \theta u$ if and only if $\sum_{y \leftarrow x} u_y = \theta u_x$ for all x .) The Laplace matrix L is one of

$$\begin{bmatrix} 2 & -1 & -1 \\ -1 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix} \text{ or } \begin{bmatrix} 1 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 1 \end{bmatrix} \text{ or } \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \\ -1 & -1 & 2 \end{bmatrix}.$$

Its eigenvalues are 0 , 1 and 3 . The Laplace eigenvectors are:

$$\begin{array}{ccc} 1 & 1 & 1 \\ \circ & \text{---} & \circ & \text{---} & \circ \end{array} \quad \begin{array}{ccc} 1 & 0 & -1 \\ \circ & \text{---} & \circ & \text{---} & \circ \end{array} \quad \begin{array}{ccc} 1 & -2 & 1 \\ \circ & \text{---} & \circ & \text{---} & \circ \end{array}.$$

(One has $Lu = \theta u$ if and only if $\sum_{y \sim x} u_y = (d_x - \theta)u_x$ for all x , where d_x is the degree of the vertex x .)

Example Let Γ be the directed triangle with adjacency matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}.$$

Then A has characteristic polynomial $p_A(\theta) = \theta^3 - 1$ and spectrum $1, \omega, \omega^2$, where ω is a primitive cube root of unity.

Example Let Γ be the directed graph with two vertices and a single directed edge. Then $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ with $p_A(\theta) = \theta^2$. So A has the eigenvalue 0 with geometric multiplicity (that is, the dimension of the corresponding eigenspace) equal to 1 and algebraic multiplicity (that is, its multiplicity as a root of the polynomial p_A) equal to 2 .

1.2.1 Characteristic polynomial

Let Γ be a directed graph on n vertices. For any directed subgraph C of Γ that is a union of directed cycles, let $c(C)$ be its number of cycles. Then the characteristic polynomial $p_A(t) = \det(tI - A)$ of Γ can be expanded as $\sum c_i t^{n-i}$, where $c_i = \sum_C (-1)^{c(C)}$, with C running over all regular directed subgraphs with in- and outdegree 1 on i vertices.

(Indeed, this is just a reformulation of the definition of the determinant as $\det M = \sum_{\sigma} \text{sgn}(\sigma) M_{1\sigma(1)} \dots M_{n\sigma(n)}$. Note that when the permutation σ with $n-i$ fixed points is written as a product of non-identity cycles, its sign is $(-1)^e$ where e is the number of even cycles in this product. Since the number of odd non-identity cycles is congruent to $i \pmod{2}$, we have $\text{sgn}(\sigma) = (-1)^{i+c(\sigma)}$.)

For example, the directed triangle has $c_0 = 1$, $c_3 = -1$. Directed edges that do not occur in directed cycles do not influence the (ordinary) spectrum.

The same description of $p_A(t)$ holds for undirected graphs (with each edge viewed as a pair of opposite directed edges).

Since $\frac{d}{dt} \det(tI - A) = \sum_x \det(tI - A_x)$ where A_x is the submatrix of A obtained by deleting row and column x , it follows that $p'_A(t)$ is the sum of the characteristic polynomials of all single-vertex-deleted subgraphs of Γ .

1.3 The spectrum of an undirected graph

Suppose Γ is undirected and simple with n vertices. Since A is real and symmetric, all its eigenvalues are real. Also, for each eigenvalue θ , its algebraic multiplicity coincides with its geometric multiplicity, so that we may omit the adjective and just speak about ‘multiplicity’. Conjugate algebraic integers have the same multiplicity. Since A has zero diagonal, its trace $\text{tr } A$, and hence the sum of the eigenvalues is zero.

Similarly, L is real and symmetric, so that the Laplace spectrum is real. Moreover, L is positive semidefinite and singular, so we may denote the eigenvalues by μ_1, \dots, μ_n , where $0 = \mu_1 \leq \mu_2 \leq \dots \leq \mu_n$. The sum of these eigenvalues is $\text{tr } L$, which is twice the number of edges of Γ .

Finally, also Q has real spectrum and nonnegative eigenvalues (but is not necessarily singular). We have $\text{tr } Q = \text{tr } L$.

1.3.1 Regular graphs

A graph Γ is called *regular* of degree (or valency) k , when every vertex has precisely k neighbors. So, Γ is regular of degree k precisely when its adjacency matrix A has row sums k , i.e., when $A\mathbf{1} = k\mathbf{1}$ (or $AJ = kJ$).

If Γ is regular of degree k , then for every eigenvalue θ we have $|\theta| \leq k$. (One way to see this, is by observing that if $|t| > k$ then the matrix $tI - A$ is strictly diagonally dominant, and hence nonsingular, so that t is not an eigenvalue of A .)

If Γ is regular of degree k , then $L = kI - A$. It follows that if Γ has ordinary eigenvalues $k = \theta_1 \geq \dots \geq \theta_n$ and Laplace eigenvalues $0 = \mu_1 \leq \mu_2 \leq \dots \leq \mu_n$, then $\theta_i = k - \mu_i$ for $i = 1, \dots, n$. The eigenvalues of $Q = kI + A$ are $2k, k + \theta_2, \dots, k + \theta_n$.

1.3.2 Complements

The *complement* $\bar{\Gamma}$ of Γ is the graph with the same vertex set as Γ , where two distinct vertices are adjacent whenever they are nonadjacent in Γ . So, if Γ has adjacency matrix A , then $\bar{\Gamma}$ has adjacency matrix $\bar{A} = J - I - A$ and Laplace matrix $\bar{L} = nI - J - L$.

Because eigenvectors of L are also eigenvectors of J , the eigenvalues of \bar{L} are $0, n - \mu_n, \dots, n - \mu_2$. (In particular, $\mu_n \leq n$.)

If Γ is regular we have a similar result for the ordinary eigenvalues: if Γ is k -regular with eigenvalues $\theta_1 \geq \dots \geq \theta_n$, then the eigenvalues of the complement are $n - k - 1, -1 - \theta_n, \dots, -1 - \theta_2$.

1.3.3 Walks

From the spectrum one can read off the number of closed walks of a given length.

Proposition 1.3.1 *Let h be a nonnegative integer. Then $(A^h)_{xy}$ is the number of walks of length h from x to y . In particular, $(A^2)_{xx}$ is the degree of the vertex x , and $\text{tr } A^2$ equals twice the number of edges of Γ ; similarly, $\text{tr } A^3$ is six times the number of triangles in Γ .*

1.3.4 Diameter

We saw that all eigenvalues of a single directed edge are zero. For undirected graphs this does not happen.

Proposition 1.3.2 *Let Γ be an undirected graph. All its eigenvalues are zero if and only if Γ has no edges. The same holds for the Laplace eigenvalues and the signless Laplace eigenvalues.*

More generally, we find a lower bound for the diameter:

Proposition 1.3.3 *Let Γ be a connected graph with diameter d . Then Γ has at least $d + 1$ distinct eigenvalues, at least $d + 1$ distinct Laplace eigenvalues, and at least $d + 1$ distinct signless Laplace eigenvalues.*

Proof. Let M be any nonnegative symmetric matrix with rows and columns indexed by $V\Gamma$ and such that for distinct vertices x, y we have $M_{xy} > 0$ if and only if $x \sim y$. Let the distinct eigenvalues of M be $\theta_1, \dots, \theta_t$. Then $(M - \theta_1 I) \cdots (M - \theta_t I) = 0$, so that M^t is a linear combination of I, M, \dots, M^{t-1} . But if $d(x, y) = t$ for two vertices x, y of Γ , then $(M^i)_{xy} = 0$ for $0 \leq i \leq t-1$ and $(M^t)_{xy} > 0$, contradiction. Hence $t > d$. This applies to $M = A$, to $M = nI - L$ and to $M = Q$, where A is the adjacency matrix, L is the Laplace matrix and Q is the signless Laplace matrix of Γ . \square

Distance-regular graphs, discussed in Chapter 11, have equality here. For an upper bound on the diameter, see §4.5.7.

1.3.5 Spanning trees

From the Laplace spectrum of a graph one can determine the number of spanning trees (which will be nonzero only if the graph is connected).

Proposition 1.3.4 *Let Γ be an undirected (multi)graph with at least one vertex, and Laplace matrix L with eigenvalues $0 = \mu_1 \leq \mu_2 \leq \dots \leq \mu_n$. Let ℓ_{xy} be the (x, y) -cofactor of L . Then the number N of spanning trees of Γ equals*

$$N = \ell_{xy} = \det(L + \frac{1}{n^2}J) = \frac{1}{n}\mu_2\cdots\mu_n \text{ for any } x, y \in V\Gamma.$$

(The (i, j) -cofactor of a matrix M is by definition $(-1)^{i+j} \det M(i, j)$, where $M(i, j)$ is the matrix obtained from M by deleting row i and column j . Note that ℓ_{xy} does not depend on an ordering of the vertices of Γ .)

Proof. Let L^S , for $S \subseteq V\Gamma$, denote the matrix obtained from L by deleting the rows and columns indexed by S , so that $\ell_{xx} = \det L^{\{x\}}$. The equality $N = \ell_{xx}$ follows by induction on n , and for fixed $n > 1$ on the number of edges incident with x . Indeed, if $n = 1$ then $\ell_{xx} = 1$. Otherwise, if x has degree 0 then $\ell_{xx} = 0$ since $L^{\{x\}}$ has zero row sums. Finally, if xy is an edge, then deleting this edge from Γ diminishes ℓ_{xx} by $\det L^{\{x, y\}}$, which by induction is the number of spanning trees of Γ with edge xy contracted, which is the number of spanning trees containing the edge xy . This shows $N = \ell_{xx}$.

Now $\det(tI - L) = t \prod_{i=2}^n (t - \mu_i)$ and $(-1)^{n-1} \mu_2 \cdots \mu_n$ is the coefficient of t , that is, is $\frac{d}{dt} \det(tI - L)|_{t=0}$. But $\frac{d}{dt} \det(tI - L) = \sum_x \det(tI - L^{\{x\}})$ so that $\mu_2 \cdots \mu_n = \sum_x \ell_{xx} = nN$.

Since the sum of the columns of L is zero, so that one column is minus the sum of the other columns, we have $\ell_{xx} = \ell_{xy}$ for any x, y . Finally, the eigenvalues of $L + \frac{1}{n^2}J$ are $\frac{1}{n}$ and μ_2, \dots, μ_n , so $\det(L + \frac{1}{n^2}J) = \frac{1}{n}\mu_2 \cdots \mu_n$. \square

For example, the multigraph of valency k on 2 vertices has Laplace matrix $L = \begin{bmatrix} k & -k \\ -k & k \end{bmatrix}$ so that $\mu_1 = 0$, $\mu_2 = 2k$, and $N = \frac{1}{2} \cdot 2k = k$.

If we consider the complete graph K_n , then $\mu_2 = \dots = \mu_n = n$, and therefore K_n has $N = n^{n-2}$ spanning trees. This formula is due to Cayley [78]. Proposition 1.3.4 is implicit in Kirchhoff [216] and known as the *Matrix-Tree Theorem*.

There is a ‘1-line proof’ of the above result using the *Cauchy-Binet formula*.

Proposition 1.3.5 (Cauchy-Binet) *Let A and B be $m \times n$ matrices. Then*

$$\det AB^\top = \sum_S \det A_S \det B_S$$

where the sum is over the $\binom{n}{m}$ m -subsets S of the set of columns, and A_S (B_S) is the square submatrix of order m of A (resp. B) with columns indexed by S .

2nd proof of Proposition 1.3.4 (sketch) Let N_x be the directed incidence matrix of Γ , with row x deleted. Then $\ell_{xx} = \det N_x N_x^\top$. Apply Cauchy-Binet to get ℓ_{xx} as a sum of squares of determinants of size $n-1$. These determinants vanish unless the set S of columns is the set of edges of a spanning tree, in which case the determinant is ± 1 . \square

1.3.6 Bipartite graphs

A graph Γ is called *bipartite* when its vertex set can be partitioned into two disjoint parts X_1, X_2 such that all edges of Γ meet both X_1 and X_2 . The

adjacency matrix of a bipartite graph has the form $A = \begin{bmatrix} 0 & B \\ B^\top & 0 \end{bmatrix}$. It follows that the spectrum of a bipartite graph is symmetric w.r.t. 0: if $\begin{bmatrix} u \\ v \end{bmatrix}$ is an eigenvector with eigenvalue θ , then $\begin{bmatrix} u \\ -v \end{bmatrix}$ is an eigenvector with eigenvalue $-\theta$. (The converse also holds, see Proposition 3.4.1.)

One has $\text{rk } A = 2 \text{rk } B$. If $n_i = |X_i|$ ($i = 1, 2$) and $n_1 \geq n_2$, then $\text{rk } A \leq 2n_2$, so that Γ has eigenvalue 0 with multiplicity at least $n_1 - n_2$.

One cannot, in general, recognize bipartiteness from the Laplace or signless Laplace spectrum. For example, $K_{1,3}$ and $K_1 + K_3$ have the same signless Laplace spectrum and only the former is bipartite. And Figure 13.4 gives an example of a bipartite and non-bipartite graph with the same Laplace spectrum. However, by Proposition 1.3.10 below, a graph is bipartite precisely when its Laplace spectrum and signless Laplace spectrum coincide.

1.3.7 Connectedness

The spectrum of a disconnected graph is easily found from the spectra of its connected components:

Proposition 1.3.6 *Let Γ be a graph with connected components Γ_i ($1 \leq i \leq s$). Then the spectrum of Γ is the union of the spectra of Γ_i (and multiplicities are added). The same holds for the Laplace and the signless Laplace spectrum. \square*

Proposition 1.3.7 *The multiplicity of 0 as a Laplace eigenvalue of an undirected graph Γ equals the number of connected components of Γ .*

Proof. We have to show that a connected graph has Laplace eigenvalue 0 with multiplicity 1. As we saw earlier, $L = NN^\top$, where N is the incidence matrix of an orientation of Γ . Now $Lu = 0$ is equivalent to $N^\top u = 0$ (since $0 = u^\top Lu = \|N^\top u\|^2$), that is, for every edge the vector u takes the same value on both endpoints. Since Γ is connected that means that u is constant. \square

Proposition 1.3.8 *Let the undirected graph Γ be regular of valency k . Then k is the largest eigenvalue of Γ , and its multiplicity equals the number of connected components of Γ .*

Proof. We have $L = kI - A$. \square

One cannot see from the spectrum alone whether a (nonregular) graph is connected: both $K_{1,4}$ and $K_1 + C_4$ have spectrum $2^1, 0^3, (-2)^1$ (we write multiplicities as exponents). And both \hat{E}_6 and $K_1 + C_6$ have spectrum $2^1, 1^2, 0, (-1)^2, (-2)^1$.

Proposition 1.3.9 *The multiplicity of 0 as a signless Laplace eigenvalue of an undirected graph Γ equals the number of bipartite connected components of Γ .*

Proof. Let M be the vertex-edge incidence matrix of Γ , so that $Q = MM^\top$. If $MM^\top u = 0$ then $M^\top u = 0$, so that $u_x = -u_y$ for all edges xy , and the support of u is the union of a number of bipartite components of Γ . \square




Figure 1.1: Two pairs of cospectral graphs

Proposition 1.3.10 *A graph Γ is bipartite if and only if the Laplace spectrum and the signless Laplace spectrum of Γ are equal.*

Proof. If Γ is bipartite, the Laplace matrix L and the signless Laplace matrix Q are similar by a diagonal matrix D with diagonal entries ± 1 (that is, $Q = DLD^{-1}$). Therefore Q and L have the same spectrum. Conversely, if both spectra are the same, then by Propositions 1.3.7 and 1.3.9 the number of connected components equals the number of bipartite components. Hence Γ is bipartite. \square

1.4 Spectrum of some graphs

In this section we discuss some special graphs and their spectra. All graphs in this section are finite, undirected and simple. Observe that the all-1 matrix J of order n has rank 1, and that the all-1 vector $\mathbf{1}$ is an eigenvector with eigenvalue n . So the spectrum is $n^1, 0^{n-1}$. (Here and throughout we write multiplicities as exponents where that is convenient and no confusion seems likely.)

1.4.1 The complete graph

Let Γ be the complete graph K_n on n vertices. Its adjacency matrix is $A = J - I$, and the spectrum is $(n-1)^1, (-1)^{n-1}$. The Laplace matrix is $nI - J$, which has spectrum $0^1, n^{n-1}$.

1.4.2 The complete bipartite graph

The spectrum of the complete bipartite graph $K_{m,n}$ is $\pm\sqrt{mn}, 0^{m+n-2}$. The Laplace spectrum is $0^1, m^{n-1}, n^{m-1}, (m+n)^1$.

1.4.3 The cycle

Let Γ be the directed n -cycle D_n . Eigenvectors are $(1, \zeta, \zeta^2, \dots, \zeta^{n-1})^\top$ where $\zeta^n = 1$, and the corresponding eigenvalue is ζ . Thus, the spectrum consists precisely of the complex n -th roots of unity $e^{2\pi i j/n}$ ($j = 0, \dots, n-1$).

Now consider the undirected n -cycle C_n . If B is the adjacency matrix of D_n , then $A = B + B^\top$ is the adjacency matrix of C_n . We find the same eigenvectors as before, with eigenvalues $\zeta + \zeta^{-1}$, so that the spectrum consists of the numbers $2\cos(2\pi j/n)$ ($j = 0, \dots, n-1$).

This graph is regular of valency 2, so the Laplace spectrum consists of the numbers $2 - 2\cos(2\pi j/n)$ ($j = 0, \dots, n-1$).

1.4.4 The path

Let Γ be the undirected path P_n with n vertices. The ordinary spectrum consists of the numbers $2 \cos(\pi j/(n+1))$ ($j = 1, \dots, n$). The Laplace spectrum is $2 - 2 \cos(\pi j/n)$ ($j = 0, \dots, n-1$).

The ordinary spectrum follows by looking at C_{2n+2} . If $u(\zeta) = (1, \zeta, \zeta^2, \dots, \zeta^{2n+1})^\top$ is an eigenvector of C_{2n+2} , where $\zeta^{2n+2} = 1$, then $u(\zeta)$ and $u(\zeta^{-1})$ have the same eigenvalue $2 \cos(\pi j/(n+1))$, and hence so has $u(\zeta) - u(\zeta^{-1})$. This latter vector has two zero coordinates distance $n+1$ apart and (for $\zeta \neq \pm 1$) induces an eigenvector on the two paths obtained by removing the two points where it is zero.

Eigenvectors of L with eigenvalue $2 - \zeta - \zeta^{-1}$ are $(1 + \zeta^{2n-1}, \dots, \zeta^j + \zeta^{2n-1-j}, \dots, \zeta^{n-1} + \zeta^n)$ where $\zeta^{2n} = 1$. One can check this directly, or consider P_n the result of folding C_{2n} where the folding has no fixed vertices. An eigenvector of C_{2n} that is constant on the preimages of the folding yields an eigenvector of P_n with the same eigenvalue.

1.4.5 Line graphs

The *line graph* $L(\Gamma)$ of Γ is the graph with the edge set of Γ as vertex set, where two vertices are adjacent if the corresponding edges of Γ have an endpoint in common. If N is the incidence matrix of Γ , then $N^\top N - 2I$ is the adjacency matrix of $L(\Gamma)$. Since $N^\top N$ is positive semidefinite, the eigenvalues of a line graph are not smaller than -2 . We have an explicit formula for the eigenvalues of $L(\Gamma)$ in terms of the signless Laplace eigenvalues of Γ .

Proposition 1.4.1 *Suppose Γ has m edges, and let $\rho_1 \geq \dots \geq \rho_r$ be the positive signless Laplace eigenvalues of Γ , then the eigenvalues of $L(\Gamma)$ are $\theta_i = \rho_i - 2$ for $i = 1, \dots, r$, and $\theta_i = -2$ if $r < i \leq m$.*

Proof. The signless Laplace matrix Q of Γ , and the adjacency matrix B of $L(\Gamma)$ satisfy $Q = NN^\top$ and $B + 2I = N^\top N$. Because NN^\top and $N^\top N$ have the same nonzero eigenvalues (multiplicities included), the result follows. \square

Example Since the path P_n has line graph P_{n-1} , and is bipartite, the Laplacian and the signless Laplacian eigenvalues of P_n are $2 + 2 \cos \frac{\pi i}{n}$, $i = 1, \dots, n$.

Corollary 1.4.2 *If Γ is a k -regular graph ($k \geq 2$) with n vertices, $e = kn/2$ edges and eigenvalues θ_i ($i = 1, \dots, n$), then $L(\Gamma)$ is $(2k-2)$ -regular with eigenvalues $\theta_i + k - 2$ ($i = 1, \dots, n$) and $e - n$ times -2 .* \square

The line graph of the complete graph K_n ($n \geq 2$), is known as the *triangular graph* $T(n)$. It has spectrum $2(n-2)^1, (n-4)^{n-1}, (-2)^{n(n-3)/2}$. The line graph of the regular complete bipartite graph $K_{m,m}$ ($m \geq 2$), is known as the *lattice graph* $L_2(m)$. It has spectrum $2(m-1)^1, (m-2)^{2m-2}, (-2)^{(m-1)^2}$. These two families of graphs, and their complements, are examples of strongly regular graphs, which will be the subject of Chapter 8. The complement of $T(5)$ is the famous *Petersen graph*. It has spectrum $3^1 1^5 (-2)^4$.

1.4.6 Cartesian products

Consider the Cartesian product $\Gamma \times \Delta$ of two graphs Γ and Δ , where the vertex set is the Cartesian product of the vertex sets of the factors, and two vertices are adjacent where they agree in one coordinate and are adjacent in the other.

If u and v are eigenvectors for Γ and Δ with ordinary or Laplace eigenvalues θ and η , respectively, then the vector w defined by $w_{(x,y)} = u_x v_y$ is an eigenvector of $\Gamma \times \Delta$ with ordinary or Laplace eigenvalue $\theta + \eta$.

For example, $L_2(m) = K_m \times K_m$.

For example, the *hypercube* 2^n , also called Q_n , is the Cartesian product of n factors K_2 . The spectrum of K_2 is $1, -1$, and hence the spectrum of 2^n consists of the numbers $n - 2i$ with multiplicity $\binom{n}{i}$ ($i = 0, 1, \dots, n$).

1.4.7 Kronecker products and bipartite double

Given graphs Γ and Δ with vertex sets V and W , respectively, their *Kronecker product* (sometimes called *conjunction*) $\Gamma \otimes \Delta$ is the graph with vertex set $V \times W$, where $(v, w) \sim (v', w')$ when $v \sim v'$ and $w \sim w'$. The adjacency matrix of $\Gamma \otimes \Delta$ is the Kronecker product of the adjacency matrices of Γ and Δ .

If u and v are eigenvectors for Γ and Δ with eigenvalues θ and η , respectively, then the vector $w = u \otimes v$ (with $w_{(x,y)} = u_x v_y$) is an eigenvector of $\Gamma \otimes \Delta$ with eigenvalue $\theta\eta$. Thus, the spectrum of $\Gamma \otimes \Delta$ consists of the products of the eigenvalues of Γ and Δ .

Given a graph Γ , its *bipartite double* is the graph $\Gamma \otimes K_2$ (with for each vertex x of Γ two vertices x' and x'' , and for each edge xy of Γ two edges $x'y''$ and $x''y'$). If Γ is bipartite, its double is just the union of two disjoint copies. If Γ is connected and not bipartite, then its double is connected and bipartite. If Γ has spectrum Φ , then $\Gamma \otimes K_2$ has spectrum $\Phi \cup -\Phi$.

1.4.8 Cayley graphs

Let G be an abelian group and $S \subseteq G$. The *Cayley graph* on G with difference set S is the (directed) graph Γ with vertex set G and edge set $E = \{(x, y) \mid y - x \in S\}$. Now Γ is regular with in- and outvalency $|S|$. The graph Γ will be undirected when $S = -S$.

It is easy to compute the spectrum of finite Cayley graphs. Let χ be a character of G , that is, a map $\chi : G \rightarrow \mathbb{C}^*$ such that $\chi(x + y) = \chi(x)\chi(y)$. Then $\sum_{y \sim x} \chi(y) = (\sum_{s \in S} \chi(s))\chi(x)$ so that the vector $(\chi(x))_{x \in G}$ is a right eigenvector of the adjacency matrix A of Γ with eigenvalue $\chi(S) := \sum_{s \in S} \chi(s)$. The $n = |G|$ distinct characters give independent eigenvectors, so one obtains the entire spectrum in this way.

For example, the directed pentagon (with in- and outvalency 1) is a Cayley graph for $G = \mathbb{Z}_5$ and $S = \{1\}$. The characters of G are the maps $i \mapsto \zeta^i$ for some fixed 5-th root of unity ζ . Hence the directed pentagon has spectrum $\{\zeta \mid \zeta^5 = 1\}$.

The undirected pentagon (with valency 2) is the Cayley graph for $G = \mathbb{Z}_5$ and $S = \{-1, 1\}$. The spectrum of the pentagon becomes $\{\zeta + \zeta^{-1} \mid \zeta^5 = 1\}$, that is, consists of 2 and $\frac{1}{2}(-1 \pm \sqrt{5})$ (both with multiplicity 2).

1.5 Decompositions

Here we present two non-trivial applications of linear algebra to graph decompositions.

1.5.1 Decomposing K_{10} into Petersen graphs

An amusing application ([30, 269]) is the following. Can the edges of the complete graph K_{10} be colored with three colors such that each color induces a graph isomorphic to the Petersen graph? K_{10} has 45 edges, 9 on each vertex, and the Petersen graph has 15 edges, 3 on each vertex, so at first sight this might seem possible. Let the adjacency matrices of the three color classes be P_1 , P_2 and P_3 , so that $P_1 + P_2 + P_3 = J - I$. If P_1 and P_2 are Petersen graphs, they both have a 5-dimensional eigenspace for eigenvalue 1, contained in the 9-space $\mathbf{1}^\perp$. Therefore, there is a common 1-eigenvector u and $P_3 u = (J - I)u - P_1 u - P_2 u = -3u$ so that u is an eigenvector for P_3 with eigenvalue -3 . But the Petersen graph does not have eigenvalue -3 , so the result of removing two edge-disjoint Petersen graphs from K_{10} is not a Petersen graph. (In fact, it follows that P_3 is connected and bipartite.)

1.5.2 Decomposing K_n into complete bipartite graphs

A famous result is the fact that for any edge-decomposition of K_n into complete bipartite graphs one needs to use at least $n - 1$ summands. Since K_n has eigenvalue -1 with multiplicity $n - 1$ this follows directly from the following:

Proposition 1.5.1 (H. S. Witsenhausen; Graham & Pollak [162]) *Suppose a graph Γ with adjacency matrix A has an edge decomposition into r complete bipartite graphs. Then $r \geq n_+(A)$ and $r \geq n_-(A)$, where $n_+(A)$ and $n_-(A)$ are the numbers of positive (negative) eigenvalues of A .*

Proof. Let the i -th complete bipartite graph have a bipartition where u_i and v_i are the characteristic vectors of both sides of the bipartition, so that its adjacency matrix is $D_i = u_i v_i^\top + v_i u_i^\top$, and $A = \sum D_i$. Let w be a vector orthogonal to all u_i . Then $w^\top A w = 0$ and it follows that w cannot be chosen in the span of eigenvectors of A with positive (negative) eigenvalue. \square

1.6 Automorphisms

An *automorphism* of a graph Γ is a permutation π of its point set X such that $x \sim y$ if and only if $\pi(x) \sim \pi(y)$. Given π , we have a linear transformation P_π on V defined by $(P_\pi(u))_x = u_{\pi(x)}$ for $u \in V$, $x \in X$. That π is an automorphism is expressed by $AP_\pi = P_\pi A$. It follows that P_π preserves the eigenspace V_θ for each eigenvalue θ of A .

More generally, if G is a group of automorphisms of Γ then we find a linear representation of degree $m(\theta) = \dim V_\theta$ of G .

We denote the group of all automorphisms of Γ by $\text{Aut } \Gamma$. One would expect that when $\text{Aut } \Gamma$ is large, then $m(\theta)$ tends to be large, so that Γ has only few distinct eigenvalues. And indeed the arguments below will show that a transitive group of automorphisms does not go very well together with simple eigenvalues.

Suppose $\dim V_\theta = 1$, say $V_\theta = \langle u \rangle$. Since P_π preserves V_θ we must have $P_\pi u = \pm u$. So either u is constant on the orbits of π , or π has even order, $P_\pi(u) = -u$, and u is constant on the orbits of π^2 . For the Perron-Frobenius eigenvector (cf. §2.2) we are always in the former case.

Corollary 1.6.1 *If all eigenvalues are simple, then $\text{Aut } \Gamma$ is an elementary abelian 2-group.*

Proof. If π has order larger than two, then there are two distinct vertices x, y in an orbit of π^2 , and all eigenvectors have identical x - and y -coordinates, a contradiction. \square

Corollary 1.6.2 *Let $\text{Aut } \Gamma$ be transitive on X . (Then Γ is regular of degree k , say.)*

(i) *If $m(\theta) = 1$ for some eigenvalue $\theta \neq k$, then $v = |X|$ is even, and $\theta \equiv k \pmod{2}$. If $\text{Aut } \Gamma$ is moreover edge-transitive then Γ is bipartite and $\theta = -k$.*

(ii) *If $m(\theta) = 1$ for two distinct eigenvalues $\theta \neq k$, then $v \equiv 0 \pmod{4}$.*

(iii) *If $m(\theta) = 1$ for all eigenvalues θ , then Γ has at most two vertices.*

Proof. (i) Suppose $V_\theta = \langle u \rangle$. Then u induces a partition of X into two equal parts: $X = X_+ \cup X_-$, where $u_x = a$ for $x \in X_+$ and $u_x = -a$ for $x \in X_-$. Now $\theta = k - 2|\Gamma(x) \cap X_-|$ for $x \in X_+$.

(ii) If $m(k) = m(\theta) = m(\theta') = 1$, then we find 3 pairwise orthogonal (± 1) -vectors, and a partition of X into four equal parts.

(iii) There are not enough integers $\theta \equiv k \pmod{2}$ between $-k$ and k . \square

For more details, see Cvetković, Doob & Sachs [106], Chapter 5.

1.7 Algebraic connectivity

Let Γ be a graph with at least two vertices. The second smallest Laplace eigenvalue $\mu_2(\Gamma)$ is called the *algebraic connectivity* of the graph Γ . This concept was introduced by Fiedler [141]. Now, by Proposition 1.3.7, $\mu_2(\Gamma) \geq 0$, with equality if and only if Γ is disconnected.

The algebraic connectivity is monotone: it does not decrease when edges are added to the graph:

Proposition 1.7.1 *Let Γ and Δ be two edge-disjoint graphs on the same vertex set, and $\Gamma \cup \Delta$ their union. We have $\mu_2(\Gamma \cup \Delta) \geq \mu_2(\Gamma) + \mu_2(\Delta) \geq \mu_2(\Gamma)$.*

Proof. Use that $\mu_2(\Gamma) = \min_u \{u^\perp L u \mid (u, u) = 1, (u, \mathbf{1}) = 0\}$. \square

The algebraic connectivity is a lower bound for the vertex connectivity:

Proposition 1.7.2 *Let Γ be a graph with vertex set X . Suppose $D \subset X$ is a set of vertices such that the subgraph induced by Γ on $X \setminus D$ is disconnected. Then $|D| \geq \mu_2(\Gamma)$.*

Proof. By monotonicity we may assume that Γ contains all edges between D and $X \setminus D$. Now a nonzero vector u that is 0 on D and constant on each component of $X \setminus D$ and satisfies $(u, \mathbf{1}) = 0$, is a Laplace eigenvector with Laplace eigenvalue $|D|$. \square

1.8 Cospectral graphs

As noted above (in §1.3.7), there exist pairs of nonisomorphic graphs with the same spectrum. Graphs with the same (adjacency) spectrum are called *cospectral* (or *isospectral*). The two graphs of Figure 1.2 below are nonisomorphic and cospectral. Both graphs are regular, which means that they are also cospectral for the Laplace matrix, and any other linear combination of A , I , and J , including the Seidel matrix (see §1.8.2), and the adjacency matrix of the complement.



Figure 1.2: Two cospectral regular graphs
(Spectrum: $4, 1, (-1)^4, \pm\sqrt{5}, \frac{1}{2}(1 \pm \sqrt{17})$)

Let us give some more examples and families of examples. A more extensive discussion is found in Chapter 13.

1.8.1 The 4-cube

The hypercube 2^n is determined by its spectrum for $n < 4$, but not for $n \geq 4$. Indeed, there are precisely two graphs with spectrum $4^1, 2^4, 0^6, (-2)^4, (-4)^1$ (Hoffman [194]). Consider the two binary codes of word length 4 and dimension 3 given by $C_1 = \mathbf{1}^\perp$ and $C_2 = (0111)^\perp$. Construct a bipartite graph, where one class of the bipartition consists of the pairs $(i, x) \in \{1, 2, 3, 4\} \times \{0, 1\}$ of coordinate position and value, and the other class of the bipartition consists of the code words, and code word u is adjacent to the pairs (i, u_i) for $i \in \{1, 2, 3, 4\}$. For the code C_1 this yields the 4-cube (tesseract), and for C_2 we get its unique cospectral mate.




Figure 1.3: Tesseract and cospectral switched version

1.8.2 Seidel switching


The *Seidel adjacency matrix* of a graph Γ with adjacency matrix A is the matrix S defined by

$$S_{uv} = \begin{cases} 0 & \text{if } u = v \\ -1 & \text{if } u \sim v \\ 1 & \text{if } u \not\sim v \end{cases}$$

so that $S = J - I - 2A$. The *Seidel spectrum* of a graph is the spectrum of its Seidel adjacency matrix. For a regular graph on n vertices with valency k and other eigenvalues θ , the Seidel spectrum consists of $n - 1 - 2k$ and the values $-1 - 2\theta$.

Let Γ have vertex set X , and let $Y \subset X$. Let D be the diagonal matrix indexed by X with $D_{xx} = -1$ for $x \in Y$, and $D_{xx} = 1$ otherwise. Then DSD has the same spectrum as S . It is the Seidel adjacency matrix of the graph obtained from Γ by leaving adjacency and nonadjacency inside Y and $X \setminus Y$ as it was, and interchanging adjacency and nonadjacency between Y and $X \setminus Y$. This new graph, Seidel-cospectral with Γ , is said to be obtained by *Seidel switching* with respect to the set of vertices Y .

Being related by Seidel switching is an equivalence relation, and the equivalence classes are called *switching classes*. Here are the three switching classes of graphs with 4 vertices.



The Seidel matrix of the complementary graph $\bar{\Gamma}$ is $-S$, so that a graph and its complement have opposite Seidel eigenvalues.

If two regular graphs of the same valency are Seidel cospectral, then they are also cospectral.

Figure 1.2 shows an example of two cospectral graphs related by Seidel switching (with respect to the four corners). These graphs are nonisomorphic: they have different local structure.

The Seidel adjacency matrix plays a rôle in the description of regular two-graphs (see §§9.1–9.3) and equiangular lines (see §9.6).

1.8.3 Godsil-McKay switching

Let Γ be a graph with vertex set X , and let $\{C_1, \dots, C_t, D\}$ be a partition of X such that $\{C_1, \dots, C_t\}$ is an equitable partition of $X \setminus D$ (that is, any two vertices in C_i have the same number of neighbors in C_j for all i, j), and for every $x \in D$ and every $i \in \{1, \dots, t\}$ the vertex x has either 0 , $\frac{1}{2}|C_i|$ or $|C_i|$ neighbors in C_i . Construct a new graph Γ' by interchanging adjacency and nonadjacency between $x \in D$ and the vertices in C_i whenever x has $\frac{1}{2}|C_i|$ neighbors in C_i . Then Γ and Γ' are cospectral ([156]).

Indeed, let Q_m be the matrix $\frac{2}{m}J - I$ of order m , so that $Q_m^2 = I$. Let $n_i = |C_i|$. Then the adjacency matrix A' of Γ' is found to be QAQ where Q is the block diagonal matrix with blocks Q_{n_i} ($1 \leq i \leq t$) and I (of order $|D|$).

The same argument also applies to the complementary graphs, so that also the complements of Γ and Γ' are cospectral. Thus, for example, the second pair

of graphs in Figure 1.1 is related by GM-switching, and hence has cospectral complements. The first pair does not have cospectral complements and hence does not arise by GM-switching.
















The 4-cube and its cospectral mate (Figure 1.3) can be obtained from each other by GM-switching with respect to the neighborhood of a vertex. Figure 1.2 is also an example of GM-switching. Indeed, when two regular graphs of the same degree are related by Seidel switching, the switch is also a case of GM-switching.

1.8.4 Reconstruction

The famous Kelly-Ulam conjecture (1941) asks whether a graph Γ can be reconstructed when the (isomorphism types of) the n vertex-deleted graphs $\Gamma \setminus x$ are given. The conjecture is still open (see Bondy [29] for a discussion), but Tutte [297] showed that one can reconstruct the characteristic polynomial of Γ , so that any counterexample to the reconstruction conjecture must be a pair of cospectral graphs.

1.9 Very small graphs

Let us give various spectra for the graphs on at most 4 vertices. The columns with heading A, L, Q, S give the spectrum for the adjacency matrix, the Laplace matrix $L = D - A$ (where D is the diagonal matrix of degrees), the signless Laplace matrix $Q = D + A$ and the Seidel matrix $S = J - I - 2A$.

label	picture	A	L	Q	S
0.1					
1.1		0	0	0	0
2.1		1, -1	0, 2	2, 0	-1, 1
2.2		0, 0	0, 0	0, 0	-1, 1
3.1		2, -1, -1	0, 3, 3	4, 1, 1	-2, 1, 1
3.2		$\sqrt{2}, 0, -\sqrt{2}$	0, 1, 3	3, 1, 0	-1, -1, 2
3.3		1, 0, -1	0, 0, 2	2, 0, 0	-2, 1, 1
3.4		0, 0, 0	0, 0, 0	0, 0, 0	-1, -1, 2
4.1		3, -1, -1, -1	0, 4, 4, 4	6, 2, 2, 2	-3, 1, 1, 1
4.2		$\rho, 0, -1, 1-\rho$	0, 2, 4, 4	$2+2\tau, 2, 2, 4-2\tau$	$-\sqrt{5}, -1, 1, \sqrt{5}$
4.3		2, 0, 0, -2	0, 2, 2, 4	4, 2, 2, 0	-1, -1, -1, 3
4.4		$\theta_1, \theta_2, -1, \theta_3$	0, 1, 3, 4	$2+\rho, 2, 1, 3-\rho$	$-\sqrt{5}, -1, 1, \sqrt{5}$
4.5		$\sqrt{3}, 0, 0, -\sqrt{3}$	0, 1, 1, 4	4, 1, 1, 0	-1, -1, -1, 3
4.6		$\tau, \tau-1, 1-\tau, -\tau$	$0, 4-\alpha, 2, \alpha$	$\alpha, 2, 4-\alpha, 0$	$-\sqrt{5}, -1, 1, \sqrt{5}$
4.7		2, 0, -1, -1	0, 0, 3, 3	4, 1, 1, 0	-3, 1, 1, 1

continued...

label	picture	A	L	Q	S
4.8		$\sqrt{2}, 0, 0, -\sqrt{2}$	$0, 0, 1, 3$	$3, 1, 0, 0$	$-\sqrt{5}, -1, 1, \sqrt{5}$
4.9		$1, 1, -1, -1$	$0, 0, 2, 2$	$2, 2, 0, 0$	$-3, 1, 1, 1$
4.10		$1, 0, 0, -1$	$0, 0, 0, 2$	$2, 0, 0, 0$	$-\sqrt{5}, -1, 1, \sqrt{5}$
4.11		$0, 0, 0, 0$	$0, 0, 0, 0$	$0, 0, 0, 0$	$-1, -1, -1, 3$

Table 1.1: Spectra of very small graphs

Here $\alpha = 2 + \sqrt{2}$ and $\tau = (1 + \sqrt{5})/2$ and $\rho = (1 + \sqrt{17})/2$ and $\theta_1 \approx 2.17009$, $\theta_2 \approx 0.31111$, $\theta_3 \approx -1.48119$ are the three roots of $\theta^3 - \theta^2 - 3\theta + 1 = 0$.

1.10 Exercises

Exercise 1 Show that no graph has eigenvalue $-1/2$. Show that no undirected graph has eigenvalue $\sqrt{2} + \sqrt{5}$. (Hint: consider the algebraic conjugates of this number.)

Exercise 2 Let Γ be an undirected graph with eigenvalues $\theta_1, \dots, \theta_n$. Show that for any two vertices a and b of Γ there are constants c_1, \dots, c_n such that the number of walks of length h from a to b equals $\sum c_i \theta_i^h$ for all h .

Exercise 3 Let Γ be a directed graph with constant outdegree $k > 0$ and without directed 2-cycles. Show that Γ has a non-real eigenvalue.

Exercise 4 (i) Let Γ be a directed graph on n vertices, such that there is a h with the property that for any two vertices a and b (distinct or not) there is a unique directed path of length h from a to b . Prove that Γ has constant in-degree and out-degree k , where $n = k^h$, and has spectrum $k^1 \ 0^{n-1}$.

(ii) The *de Bruijn graph* of order m is the directed graph with as vertices the 2^m binary sequences of length m , where there is an arrow from $a_1 \dots a_m$ to $b_1 \dots b_m$ when the tail $a_2 \dots a_m$ of the first equals the head $b_1 \dots b_{m-1}$ of the second. (For $m = 0$ a single vertex with two loops.) Determine the spectrum of the de Bruijn graph.

(iii) A *de Bruijn cycle* of order $m \geq 1$ ([64, 65, 145]) is a circular arrangement of 2^m zeros and ones such that each binary sequence of length m occurs once in this cycle. (In other words, it is a Hamiltonian cycle in the de Bruijn graph of order m , an Eulerian cycle in the de Bruijn graph of order $m - 1$.) Show that there are precisely $2^{2^{m-1}-m}$ de Bruijn cycles of order m .

Exercise 5 ([37, 265]) Let Γ be a *tournament*, that is, a directed graph in which there is precisely one edge between any two distinct vertices, in other words, of which the adjacency matrix A satisfies $A^\top + A = J - I$.

(i) Show that all eigenvalues have real part not less than $-1/2$.

(ii) The tournament Γ is called *transitive* if (x, z) is an edge whenever both (x, y) and (y, z) are edges. Show that all eigenvalues of a transitive tournament are zero.

- (iii) The tournament Γ is called *regular* when each vertex has the same number of out-arrows. Clearly, when there are n vertices, this number of out-arrows is $(n-1)/2$. Show that all eigenvalues θ have real part at most $(n-1)/2$, and that $\operatorname{Re}(\theta) = (n-1)/2$ occurs if and only if Γ is regular (and then $\theta = (n-1)/2$).
- (iv) Show that A either has full rank n or has rank $n-1$, and that A has full rank when Γ is regular and $n > 1$.

(Hint: for a vector u , consider the expression $\bar{u}^\top (A^\top + A)u$.)

Exercise 6 Let Γ be bipartite and consider its line graph $L(\Gamma)$.

- (i) Show that Γ admits a directed incidence matrix N such that $N^\top N - 2I$ is the adjacency matrix of $L(\Gamma)$.
- (ii) Give a relation between the Laplace eigenvalues of Γ and the ordinary eigenvalues of $L(\Gamma)$.
- (iii) Verify this relation in case Γ is the path P_n .

Exercise 7 ([94]) Verify (see §1.2.1) that both graphs pictured here have characteristic polynomial $t^4(t^4 - 7t^2 + 9)$, so that these two trees are cospectral.



Note how the coefficients of the characteristic polynomial of a tree count partial matchings (sets of pairwise disjoint edges) in the tree.

Exercise 8 ([14]) Verify that both graphs pictured here have characteristic polynomial $(t-1)(t+1)^2(t^3 - t^2 - 5t + 1)$ by computing eigenvectors and eigenvalues. Use the observation (§1.6) that the image of an eigenvector under an automorphism is again an eigenvector. In particular, when two vertices x, y are interchanged by an involution (automorphism of order 2), then a basis of the eigenspace exists consisting of vectors where the x - and y -coordinates are either equal or opposite.



Exercise 9 Let the *cone* over a graph Γ be the graph obtained by adding a new vertex and joining that to all vertices of Γ . If Γ is regular of valency k on n vertices, then show that the cone over Γ has characteristic polynomial $p(x) = (x^2 - kx - n)p_\Gamma(x)/(x - k)$.

Exercise 10 Show that the Seidel adjacency matrix S of a graph on n vertices has rank $n-1$ or n . (Hint: $\det S \equiv n-1 \pmod{2}$.)

Exercise 11 Prove that the complete graph K_{55} is not the union of three copies of the triangular graph $T(11)$.

Chapter 2

Linear algebra

In this chapter we present some less elementary, but relevant results from linear algebra.

2.1 Simultaneous diagonalization

Let V be a complex vector space with finite dimension, and fix a basis. Then we can define an inner product on V by putting $(x, y) = \sum \bar{x}_i y_i = \bar{x}^\top y$ for $x, y \in V$, where the bar denotes complex conjugation. If A is *Hermitean*, i.e., if $(Ax, y) = (x, Ay)$ for all $x, y \in V$, then all eigenvalues of A are real, and V admits an orthonormal basis of eigenvectors of A .

Proposition 2.1.1 *Suppose \mathcal{A} is a collection of commuting Hermitean linear transformations on V (i.e., $AB = BA$ for $A, B \in \mathcal{A}$), then V has a basis consisting of common eigenvectors of all $A \in \mathcal{A}$.*

Proof. Induction on $\dim V$. If each $A \in \mathcal{A}$ is a multiple of the identity I , then all is clear. Otherwise, let $A \in \mathcal{A}$ not be a multiple of I . If $Au = \theta u$ and $B \in \mathcal{A}$, then $A(Bu) = BAu = \theta Bu$ so that B acts as a linear transformation on the eigenspace V_θ for the eigenvalue θ of A . By the induction hypothesis we can choose a basis consisting of common eigenvectors for each $B \in \mathcal{A}$ in each eigenspace. The union of these bases is the basis of V we were looking for. \square

Given a square matrix A , we can regard A as a linear transformation on a vector space (with fixed basis). Hence the above concepts apply. The matrix A will be Hermitean precisely when $A = \bar{A}^\top$; in particular, a real symmetric matrix is Hermitean.

2.2 Perron-Frobenius Theory

Let T be a real $n \times n$ matrix with nonnegative entries. T is called *primitive* if for some k we have $T^k > 0$; T is called *irreducible* if for all i, j there is a k such that $(T^k)_{ij} > 0$.

Here, for a matrix (or vector) A , $A > 0$ (≥ 0) means that all its entries are positive (nonnegative).

The matrix $T = (t_{ij})$ is irreducible if and only if the directed graph Γ_T with vertices $\{1, \dots, n\}$ and edges (i, j) whenever $t_{ij} > 0$ is strongly connected.

(A directed graph (X, E) is *strongly connected* if for any two vertices x, y there is a directed path from x to y , i.e., there are vertices $x_0 = x, x_1, \dots, x_m = y$ such that $(x_{i-1}, x_i) \in E$ for $1 \leq i \leq m$.)

Note that if T is irreducible, then $I + T$ is primitive.

The *period* d of an irreducible matrix T is the greatest common divisor of the integers k for which $(T^k)_{ii} > 0$. It is independent of the i chosen.

Theorem 2.2.1 *Let $T \geq 0$ be irreducible. Then there is a (unique) positive real number θ_0 with the following properties:*

- (i) *There is a real vector $x_0 > 0$ with $Tx_0 = \theta_0 x_0$.*
- (ii) *θ_0 has geometric and algebraic multiplicity one.*
- (iii) *For each eigenvalue θ of T we have $|\theta| \leq \theta_0$. If T is primitive, then $|\theta| = \theta_0$ implies $\theta = \theta_0$. In general, if T has period d , then T has precisely d eigenvalues θ with $|\theta| = \theta_0$, namely $\theta = \theta_0 e^{2\pi i j/d}$ for $j = 0, 1, \dots, d-1$. In fact the entire spectrum of T is invariant under rotation of the complex plane over an angle $2\pi/d$ about the origin.*
- (iv) *Any nonnegative left or right eigenvector of T has eigenvalue θ_0 . More generally, if $x \geq 0$, $x \neq 0$ and $Tx \leq \theta x$, then $x > 0$ and $\theta \geq \theta_0$; moreover, $\theta = \theta_0$ if and only if $Tx = \theta x$.*
- (v) *If $0 \leq S \leq T$ or if S is a principal minor of T , and S has eigenvalue σ , then $|\sigma| \leq \theta_0$; if $|\sigma| = \theta_0$, then $S = T$.*
- (vi) *Given a complex matrix S , let $|S|$ denote the matrix with elements $|S|_{ij} = |S_{ij}|$. If $|S| \leq T$ and S has eigenvalue σ , then $|\sigma| \leq \theta_0$. If equality holds, then $|S| = T$, and there are a diagonal matrix E with diagonal entries of absolute value 1 and a constant c of absolute value 1, such that $S = cETE^{-1}$.*

Proof. (i) Let $P = (I + T)^{n-1}$. Then $P > 0$ and $PT = TP$. Let $B = \{x \mid x \geq 0 \text{ and } x \neq 0\}$. Define for $x \in B$:

$$\theta(x) = \max \{\theta \mid \theta \in \mathbb{R}, \theta x \leq Tx\} = \min \left\{ \frac{(Tx)_i}{x_i} \mid 1 \leq i \leq n, x_i \neq 0 \right\}.$$

Now $\theta(\alpha x) = \theta(x)$ for $\alpha \in \mathbb{R}$, $\alpha > 0$, and $(x \leq y, x \neq y \text{ implies } Px < Py)$, so $\theta(Px) \geq \theta(x)$; in fact $\theta(Px) > \theta(x)$ unless x is an eigenvector of T . Put $C = \{x \mid x \geq 0 \text{ and } \|x\| = 1\}$. Then since C is compact and $\theta(\cdot)$ is continuous on $P[C]$ (but not in general on C !), there is an $x_0 \in P[C]$ such that

$$\theta_0 := \sup_{x \in B} \theta(x) = \sup_{x \in C} \theta(x) = \sup_{x \in P[C]} \theta(x) = \theta(x_0).$$

Now $x_0 > 0$ and x_0 is an eigenvector of T , so $Tx_0 = \theta_0 x_0$, and $\theta_0 > 0$.

(ii) For a vector $x = (x_1, \dots, x_n)^\top$, write $x_+ = (|x_1|, \dots, |x_n|)^\top$. If $Tx = \theta x$, then by the triangle inequality we have $Tx_+ \geq |\theta|x_+$. For nonzero x this means $|\theta| \leq \theta(x_+) \leq \theta_0$. If, for some vector $z \in B$, we have $Tz \geq \theta_0 z$, then z is eigenvector of T (otherwise $\theta(Pz) > \theta_0$), and since $0 < Pz = (1 + \theta_0)^{n-1} z$ we

have $z > 0$. Now if x is any eigenvector of T with eigenvalue θ_0 , then if x is real consider $y = x_0 + \varepsilon x$, where ε is chosen such that $y \geq 0$ but not $y > 0$; now $y \notin B$, i.e., $y = 0$, so x is a multiple of x_0 . In the general case, both the real and imaginary parts of x are multiples of x_0 . This shows that the eigenspace of θ_0 has dimension 1, i.e., that the geometric multiplicity of θ_0 is 1. We shall look at the algebraic multiplicity later.

(iii) We have seen $|\theta| \leq \theta_0$. If $|\theta| = \theta_0$ and $Tx = \theta x$, then $Tx_+ = \theta_0 x_+$ and we had equality in the triangle inequality $|\sum_j t_{ij}x_j| \leq \sum_j t_{ij}|x_j|$; this means that all numbers $t_{ij}x_j$ ($1 \leq j \leq n$) have the same angular part (argument). If T is primitive, then we can apply this reasoning with T^k instead of T , where $T^k > 0$, and conclude that all x_j have the same angular part. Consequently, in this case x is a multiple of a real vector and may be taken real, nonnegative. Now $Tx = \theta x$ shows that θ is real, and $|\theta| = \theta_0$ that $\theta = \theta_0$. In the general case, T^d is a direct sum of primitive matrices $T^{(0)}, \dots, T^{(d-1)}$, and if $x = (x^{(0)}, \dots, x^{(d-1)})$ is the corresponding decomposition of an eigenvector of T (with eigenvalue θ), then $(x^{(0)}, \zeta x^{(1)}, \dots, \zeta^{d-1} x^{(d-1)})$ also is an eigenvector of T , with eigenvalue $\zeta\theta$, for any d -th root of unity ζ . (Here we assume that the $T^{(i)}$ are ordered in such a way that in Γ_T the arrows point from the subset corresponding to $T^{(i)}$ to the subset corresponding to $T^{(i+1)}$.) Since T^d has a unique eigenvalue of maximum modulus (let $T_{(i)}^{(i+1)}$ be the (nonsquare) submatrix of T describing the arrows in Γ_T between the subset corresponding to $T^{(i)}$ to the subset corresponding to $T^{(i+1)}$; then $T^{(i)} = \prod_{j=0}^{d-1} T_{(i+j)}^{(i+j+1)}$ and if $T^{(i)}z = \gamma z$, $z > 0$ then $T^{(i-1)}z' = \gamma z'$ where $z' = T_{(i-1)}^{(i)}z \neq 0$, so that all $T^{(i)}$ have the same eigenvalue of maximum modulus), it follows that T has precisely d such eigenvalues.

(iv) Doing the above for left eigenvectors instead of right ones, we find $y_0 > 0$ with $y_0^\top T = \eta_0 y_0^\top$. If $Tx = \theta x$ and $y^\top T = \eta y^\top$, then $\eta y^\top x = y^\top Tx = \theta y^\top x$. It follows that either $\theta = \eta$ or $y^\top x = 0$. Taking $y \in B$, $x = x_0$ or $x \in B$, $y = y_0$ we see that $\theta = \eta (= \theta_0 = \eta_0)$. Similarly, if $Tx \leq \theta x$, $x \in B$ then $\theta_0 y_0^\top x = y_0^\top Tx \leq \theta y_0^\top x$ so that $\theta_0 \leq \theta$; also $0 < Px \leq (1 + \theta)^{n-1}x$, so $x > 0$. If $\theta = \theta_0$, then $y_0^\top (Tx - \theta x) = 0$ so $Tx = \theta x$.

(v) If $s \neq 0$, $Ss = \sigma s$, then $Ts_+ \geq Ss_+ \geq |\sigma|s_+$, so $|\sigma| \leq \theta_0$. But if $|\sigma| = \theta_0$ then s_+ is eigenvector of T and $s_+ > 0$ and $(T - S)s_+ = 0$, so $S = T$.

(vi) If $s \neq 0$, $Ss = \sigma s$, then $Ts_+ \geq |S|s_+ \geq |\sigma|s_+$, so $|\sigma| \leq \theta_0$, and if $|\sigma| = \theta_0$ then s_+ is eigenvector of T and $s_+ > 0$ and $|S| = T$. Equality in $|S|s_+ = |\sigma|s_+$ means that $|\sum S_{ij}s_j| = \sum |S_{ij}| \cdot |s_j|$, so that given i all $S_{ij}s_j$ have the same angular part. Let $E_{ii} = s_i/|s_i|$ and $c = \sigma/|\sigma|$. Then $S_{ij} = cE_{ii}E_{jj}^{-1}|S_{ij}|$.

(vii) Finally, in order to prove that θ_0 is a simple root of χ_T , the characteristic polynomial of T , we have to show that $\frac{d}{d\theta}\chi_T(\theta)$ is nonzero for $\theta = \theta_0$. But $\chi_T(\theta) = \det(\theta I - T)$ and $\frac{d}{d\theta}\chi_T(\theta) = \sum_i \det(\theta I - T_{ii})$, and by (v) we have $\det(\theta I - T_{ii}) > 0$ for $\theta = \theta_0$. \square

Remark In case $T \geq 0$ but T not necessarily irreducible, we can say the following.

(i) The spectral radius θ_0 of T is an eigenvalue, and there are nonnegative left and right eigenvectors corresponding to it.

(ii) If $|S| \leq T$ and S has eigenvalue σ , then $|\sigma| \leq \theta_0$.

(**Proof.** (i) Use continuity arguments; (ii) the old proof still applies.)

For more details, see the exposition of the Perron-Frobenius theory in Gantmacher [149], Chapter XIII; cf. also Varga [299], Marcus & Minc [238], Berman & Plemmons [22], Seneta [279, Ch. 1], or Horn & Johnson [200, Ch. 8].

2.3 Equitable partitions

Suppose A is a symmetric real matrix whose rows and columns are indexed by $X = \{1, \dots, n\}$. Let $\{X_1, \dots, X_m\}$ be a partition of X . The *characteristic matrix* S is the $n \times m$ matrix whose j^{th} column is the characteristic vector of X_j ($j = 1, \dots, m$). Define $n_i = |X_i|$ and $K = \text{diag}(n_1, \dots, n_m)$. Let A be partitioned according to $\{X_1, \dots, X_m\}$, that is

$$A = \begin{bmatrix} A_{1,1} & \dots & A_{1,m} \\ \vdots & & \vdots \\ A_{m,1} & \dots & A_{m,m} \end{bmatrix},$$

wherein $A_{i,j}$ denotes the submatrix (block) of A formed by rows in X_i and the columns in X_j . Let $b_{i,j}$ denote the average row sum of $A_{i,j}$. Then the matrix $B = (b_{i,j})$ is called the *quotient matrix*. We easily have

$$KB = S^\top AS, \quad S^\top S = K.$$

If the row sum of each block $A_{i,j}$ is constant then the partition is called *equitable* (or *regular*) and we have $A_{i,j}\mathbf{1} = b_{i,j}\mathbf{1}$ for $i, j = 1, \dots, m$, so

$$AS = SB.$$

The following result is well-known and useful.

Lemma 2.3.1 *If, for an equitable partition, v is an eigenvector of B for an eigenvalue λ , then Sv is an eigenvector of A for the same eigenvalue λ .*

Proof. $Bv = \lambda v$ implies $ASv = SBv = \lambda Sv$. □

In the situation of this lemma, the spectrum of A consists of the spectrum of the quotient matrix B (with eigenvectors in the column space of S , i.e., constant on the parts of the partition) together with the eigenvalues belonging to eigenvectors orthogonal to the columns of S (i.e., summing to zero on each part of the partition). These latter eigenvalues remain unchanged if the blocks $A_{i,j}$ are replaced by $A_{i,j} + c_{i,j}J$ for certain constants $c_{i,j}$.

2.3.1 Equitable and almost equitable partitions of graphs

If in the above the matrix A is the adjacency matrix (or the Laplace matrix) of a graph, then an equitable partition of the matrix A is a partition of the vertex set into parts X_i such that each vertex in X_i has the same number $b_{i,j}$ of neighbours in part X_j , for any j (or any $j \neq i$). Such partitions are called (almost) equitable partitions of the graph.

For example, the adjacency matrix of the complete bipartite graph $K_{p,q}$ has an equitable partition with $m = 2$. The quotient matrix B equals $\begin{bmatrix} 0 & p \\ q & 0 \end{bmatrix}$ and has eigenvalues $\pm\sqrt{pq}$, which are the nonzero eigenvalues of $K_{p,q}$.

More generally, consider the *join* Γ of two vertex-disjoint graphs Γ_1 and Γ_2 , the graph obtained by inserting all possible edges between Γ_1 and Γ_2 . If Γ_1 and Γ_2 have n_1 resp. n_2 vertices and are both regular, say of valency k_1 resp. k_2 , and have spectra Φ_1 resp. Φ_2 , then Γ has spectrum $\Phi = (\Phi_1 \setminus \{k_1\}) \cup (\Phi_2 \setminus \{k_2\}) \cup \{k', k''\}$ where k', k'' are the two eigenvalues of

$$\begin{bmatrix} k_1 & n_2 \\ n_1 & k_2 \end{bmatrix}.$$

Indeed, we have an equitable partition of the adjacency matrix of Γ with the above quotient matrix. The eigenvalues that do not belong to the quotient coincide with those of the disjoint union of Γ_1 and Γ_2 .

2.4 The Rayleigh quotient

Let A be a real symmetric matrix and let u be a nonzero vector. The *Rayleigh quotient* of u w.r.t. A is defined as

$$\frac{u^\top A u}{u^\top u}.$$

Let u_1, \dots, u_n be an orthonormal set of eigenvectors of A , say with $Au_i = \theta_i u_i$, where $\theta_1 \geq \dots \geq \theta_n$. If $u = \sum \alpha_i u_i$ then $u^\top u = \sum \alpha_i^2$ and $u^\top A u = \sum \alpha_i^2 \theta_i$. It follows that

$$\frac{u^\top A u}{u^\top u} \geq \theta_i \text{ if } u \in \langle u_1, \dots, u_i \rangle$$

and

$$\frac{u^\top A u}{u^\top u} \leq \theta_i \text{ if } u \in \langle u_1, \dots, u_{i-1} \rangle^\perp.$$

In both cases, equality implies that u is a θ_i -eigenvector of A . Conversely, one has

Theorem 2.4.1 (Courant-Fischer) *Let W be an i -subspace of V . Then*

$$\theta_i \geq \min_{u \in W, u \neq 0} \frac{u^\top A u}{u^\top u}$$

and

$$\theta_{i+1} \leq \max_{u \in W^\perp, u \neq 0} \frac{u^\top A u}{u^\top u}.$$

2.5 Interlacing

Consider two sequences of real numbers: $\theta_1 \geq \dots \geq \theta_n$, and $\eta_1 \geq \dots \geq \eta_m$ with $m < n$. The second sequence is said to *interlace* the first one whenever

$$\theta_i \geq \eta_i \geq \theta_{n-m+i}, \text{ for } i = 1, \dots, m.$$

The interlacing is *tight* if there exist an integer $k \in [0, m]$ such that

$$\theta_i = \eta_i \text{ for } 1 \leq i \leq k \text{ and } \theta_{n-m+i} = \eta_i \text{ for } k+1 \leq i \leq m.$$

If $m = n - 1$, the interlacing inequalities become $\theta_1 \geq \eta_1 \geq \theta_2 \geq \eta_2 \geq \dots \geq \eta_m \geq \theta_n$, which clarifies the name. Godsil [154] reserves the name ‘interlacing’ for this particular case and calls it generalized interlacing otherwise.

Theorem 2.5.1 *Let S be a real $n \times m$ matrix such that $S^\top S = I$. Let A be a real symmetric matrix of order n with eigenvalues $\theta_1 \geq \dots \geq \theta_n$. Define $B = S^\top AS$ and let B have eigenvalues $\eta_1 \geq \dots \geq \eta_m$ and respective eigenvectors $v_1 \dots v_m$.*

- (i) *The eigenvalues of B interlace those of A .*
- (ii) *If $\eta_i = \theta_i$ or $\eta_i = \theta_{n-m+i}$ for some $i \in [1, m]$, then B has a η_i -eigenvector v such that Sv is a η_i -eigenvector of A .*
- (iii) *If for some integer l , $\eta_i = \theta_i$, for $i = 1, \dots, l$ (or $\eta_i = \theta_{n-m+i}$ for $i = l, \dots, m$), then Sv_i is a η_i -eigenvector of A for $i = 1, \dots, l$ (respectively $i = l, \dots, m$).*
- (iv) *If the interlacing is tight, then $SB = AS$.*

Proof. Let u_1, \dots, u_n be an orthonormal set of eigenvectors of the matrix A , where $Au_i = \theta_i u_i$. For each $i \in [1, m]$, take a nonzero vector s_i in

$$\langle v_1, \dots, v_i \rangle \cap \langle S^\top u_1, \dots, S^\top u_{i-1} \rangle^\perp. \quad (2.1)$$

Then $Ss_i \in \langle u_1, \dots, u_{i-1} \rangle^\perp$, hence by Rayleigh's principle,

$$\theta_i \geq \frac{(Ss_i)^\top A(Ss_i)}{(Ss_i)^\top (Ss_i)} = \frac{s_i^\top B s_i}{s_i^\top s_i} \geq \eta_i,$$

and similarly (or by applying the above inequality to $-A$ and $-B$) we get $\theta_{n-m+i} \leq \eta_i$, proving (i). If $\theta_i = \eta_i$, then s_i and Ss_i are θ_i -eigenvectors of B and A , respectively, proving (ii). We prove (iii) by induction on l . Assume $Sv_i = u_i$ for $i = 1, \dots, l-1$. Then we may take $s_l = v_l$ in (2.1), but in proving (ii) we saw that Ss_l is a θ_l -eigenvector of A . (The statement between parentheses follows by considering $-A$ and $-B$.) Thus we have (iii). Let the interlacing be tight. Then by (iii), Sv_1, \dots, Sv_m is an orthonormal set of eigenvectors of A for the eigenvalues η_1, \dots, η_m . So we have $SBv_i = \eta_i Sv_i = ASv_i$, for $i = 1, \dots, m$. Since the vectors v_i form a basis, it follows that $SB = AS$. \square

If we take $S = [I \ 0]^\top$, then B is just a principal submatrix of A and we have the following corollary.

Corollary 2.5.2 *If B is a principal submatrix of a symmetric matrix A , then the eigenvalues of B interlace the eigenvalues of A .*

The theorem requires the columns of S to be orthonormal. If one has a situation with orthogonal but not necessarily orthonormal vectors, some scaling is required.

Corollary 2.5.3 *Let A be a real symmetric matrix of order n . Let x_1, \dots, x_m be nonzero orthogonal real vectors of order n . Define a matrix $C = (c_{ij})$ by $c_{ij} = \frac{1}{\|x_i\|^2} x_i^\top A x_j$.*

- (i) *The eigenvalues of C interlace the eigenvalues of A .*
- (ii) *If the interlacing is tight, then $Ax_j = \sum c_{ij} x_i$ for all j .*

(iii) Let $x = \sum x_j$. The number $r := \frac{x^\top Ax}{x^\top x}$ lies between the smallest and largest eigenvalue of C . If x is an eigenvector of A with eigenvalue θ , then also C has an eigenvalue θ (for eigenvector $\mathbf{1}$).

Proof. Let K be the diagonal matrix with $K_{ii} = \|x_i\|$. Let R be the $n \times m$ matrix with columns x_j , and put $S = RK^{-1}$. Then $S^\top S = I$, and the theorem applies with $B = S^\top AS = CKK^{-1}$. If interlacing is tight we have $AR = RC$. With $x = \sum x_j = R\mathbf{1}$ and $y = K\mathbf{1}$, we have $\frac{x^\top Ax}{x^\top x} = \frac{y^\top By}{y^\top y}$. \square

In particular, this applies when the x_i are the characteristic vectors of a partition (or just a collection of pairwise disjoint subsets).

Corollary 2.5.4 *Let C be the quotient matrix of a symmetric matrix A whose rows and columns are partitioned according to a partitioning $\{X_1, \dots, X_m\}$.*

(i) *The eigenvalues of C interlace the eigenvalues of A .*

(ii) *If the interlacing is tight, then the partition is equitable.* \square

Theorem 2.5.1(i) is a classical result; see Courant & Hilbert [98], Vol. 1, Ch. I. For the special case of a principal submatrix (Corollary 2.5.2), the result even goes back to Cauchy and is therefore often referred to as Cauchy interlacing. Interlacing for the quotient matrix (Corollary 2.5.4) is especially applicable to combinatorial structures (as we shall see). Payne (see, for instance, [256]) has applied the extremal inequalities $\theta_1 \geq \eta_i \geq \theta_n$ to finite geometries several times. He attributes the method to Higman and Sims and therefore calls it the *Higman-Sims technique*.

Remark This theorem generalizes directly to complex Hermitean matrices instead of real symmetric matrices (with conjugate transpose instead of transpose) with virtually the same proof.

For more detailed eigenvalue inequalities, see Haemers [175], [177].

2.6 Schur's inequality

Theorem 2.6.1 (Schur [266]) *Let A be a real symmetric matrix with eigenvalues $\theta_1 \geq \theta_2 \geq \dots \geq \theta_n$ and diagonal elements $d_1 \geq d_2 \geq \dots \geq d_n$. Then $\sum_{i=1}^t d_i \leq \sum_{i=1}^t \theta_i$ for $1 \leq t \leq n$.*

Proof. Let B be the principal submatrix of A obtained by deleting the rows and columns containing d_{t+1}, \dots, d_n . If B has eigenvalues η_i ($1 \leq i \leq t$) then by interlacing $\sum_{i=1}^t d_i = \text{tr } B = \sum_{i=1}^t \eta_i \leq \sum_{i=1}^t \theta_i$. \square

Remark Again 'real symmetric' can be replaced by 'Hermitean'.

2.7 Schur complements

In this section, the square matrix

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$$

is a square partitioned matrix (over any field), where A_{11} is nonsingular. The *Schur complement* A/A_{11} of A_{11} in A is the matrix $A_{22} - A_{21}A_{11}^{-1}A_{12}$. The following result is a straightforward but important consequence from the definition.

Theorem 2.7.1 (see [311]) *The Schur complement A/A_{11} satisfies*

$$(i) \begin{bmatrix} I & O \\ -A_{21}A_{11}^{-1} & I \end{bmatrix} \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \begin{bmatrix} I & -A_{11}^{-1}A_{12} \\ O & I \end{bmatrix} = \begin{bmatrix} A_{11} & O \\ O & A/A_{11} \end{bmatrix},$$

$$(ii) \det(A/A_{11}) = \det A / \det A_{11},$$

$$(iii) \operatorname{rk} A = \operatorname{rk} A_{11} + \operatorname{rk}(A/A_{11}).$$

Corollary 2.7.2 *If $\operatorname{rk} A = \operatorname{rk} A_{11}$, then $A_{22} = A_{21}A_{11}^{-1}A_{12}$.*

2.8 The Courant-Weyl inequalities

Denote the eigenvalues of a Hermitean matrix A , arranged in nonincreasing order, by $\lambda_i(A)$.

Theorem 2.8.1 *Let A and B be Hermitean matrices of order n , and let $1 \leq i, j \leq n$.*

$$(i) \text{ If } i + j - 1 \leq n \text{ then } \lambda_{i+j-1}(A + B) \leq \lambda_i(A) + \lambda_j(B).$$

$$(ii) \text{ If } i + j - n \geq 1 \text{ then } \lambda_i(A) + \lambda_j(B) \leq \lambda_{i+j-n}(A + B).$$

$$(iii) \text{ If } B \text{ is positive semidefinite, then } \lambda_i(A + B) \geq \lambda_i(A).$$

Proof. (i) Let u_1, \dots, u_n and v_1, \dots, v_n be orthonormal sets of eigenvectors of A resp. B , with $Au_i = \lambda_i(A)u_i$ and $Bv_j = \lambda_j(B)v_j$. Let $U = \langle u_h \mid 1 \leq h \leq i-1 \rangle$ and $V = \langle v_h \mid 1 \leq h \leq j-1 \rangle$, and $W = U + V$. For $w \in W^\perp$ we have $w^\top(A + B)w \leq (\lambda_i(A) + \lambda_j(B))w^\top w$. It follows that the space spanned by eigenvectors of $A + B$ with eigenvalue larger than $\lambda_i(A) + \lambda_j(B)$ has dimension at most $i + j - 2$.

(ii) Apply (i) to $-A$ and $-B$. (iii) Apply the case $j = n$ of (ii). \square

Ky Fan [138] shows that $\lambda(A) + \lambda(B)$ dominates $\lambda(A + B)$:

Theorem 2.8.2 *Let A and B be Hermitean matrices of order n . Then for all t , $0 \leq t \leq n$, we have $\sum_{i=1}^t \lambda_i(A + B) \leq \sum_{i=1}^t \lambda_i(A) + \sum_{i=1}^t \lambda_i(B)$.*

Proof. $\sum_{i=1}^t \lambda_i(A) = \max \operatorname{tr}(U^*AU)$, where the maximum is over all $n \times t$ matrices U with $U^*U = I$. \square

2.9 Gram matrices

Real symmetric $n \times n$ -matrices G are in bijective correspondence with quadratic forms q on \mathbb{R}^n via the relation

$$q(x) = x^\top Gx \quad (x \in \mathbb{R}^n).$$

Two quadratic forms q and q' on \mathbb{R}^n are *congruent*, i.e., there is a nonsingular $n \times n$ -matrix S such that $q(x) = q'(Sx)$ for all $x \in \mathbb{R}^n$, if and only if their corresponding matrices G and G' satisfy $G = S^\top G' S$. Moreover, this occurs for some S if and only if G and G' have the same rank and the same number of nonnegative eigenvalues—this is Sylvester [290]’s ‘law of inertia for quadratic forms’, cf. Gantmacher [149], Vol. 1, Chapter X, §2. We shall now be concerned with matrices that have nonnegative eigenvalues only.

Lemma 2.9.1 *Let G be a real symmetric $n \times n$ -matrix. Equivalent are:*

- (i) *For all $x \in \mathbb{R}^n$, $x^\top Gx \geq 0$.*
- (ii) *All eigenvalues of G are nonnegative.*
- (iii) *G can be written as $G = H^\top H$, with H an $m \times n$ matrix, where m is the rank of G .*

Proof. There is an orthogonal matrix Q and a diagonal matrix D whose nonzero entries are the eigenvalues of G such that $G = Q^\top DQ$. If (ii) holds, then $x^\top Gx = (Qx)^\top D(Qx) \geq 0$ implies (i). Conversely, (ii) follows from (i) by choosing x to be an eigenvector. If $G = H^\top H$ then $x^\top Gx = \|Hx\|^2 \geq 0$, so (iii) implies (i). Finally, let $E = D^{1/2}$ be the diagonal matrix that squares to D , and let F be the $m \times n$ matrix obtained from E by dropping the zero rows. Then $G = Q^\top E^\top EQ = Q^\top F^\top FQ = H^\top H$, so that (ii) implies (iii). \square

A symmetric $n \times n$ -matrix G satisfying (i) or (ii) is called *positive semidefinite*. It is called *positive definite* when $x^\top Gx = 0$ implies $x = 0$, or, equivalently, when all its eigenvalues are positive. For any collection X of vectors of \mathbb{R}^m , we define its *Gram matrix* as the square matrix G indexed by X whose (x, y) -entry G_{xy} is the inner product $(x, y) = x^\top y$. This matrix is always positive semidefinite, and it is definite if and only if the vectors in X are linearly independent. (Indeed, if $n = |X|$, and we use H to denote the $m \times n$ -matrix whose columns are the vectors of X , then $G = H^\top H$, and $x^\top Gx = \|Hx\|^2 \geq 0$.)

Lemma 2.9.2 *Let N be a real $m \times n$ matrix. Then the matrices NN^\top and $N^\top N$ have the same nonzero eigenvalues (including multiplicities). Moreover, $\text{rk } NN^\top = \text{rk } N^\top N = \text{rk } N$.*

Proof. Let θ be a nonzero eigenvalue of NN^\top . The map $u \mapsto N^\top u$ is an isomorphism from the θ -eigenspace of NN^\top onto the θ -eigenspace of $N^\top N$. Indeed, if $NN^\top u = \theta u$ then $N^\top NN^\top u = \theta N^\top u$ and $N^\top u$ is nonzero for nonzero u since $NN^\top u = \theta u$. The final sentence follows since $\text{rk } N^\top N \leq \text{rk } N$, but if $N^\top Nx = 0$ then $\|Nx\|^2 = x^\top N^\top Nx = 0$, so that $Nx = 0$. \square

2.10 Diagonally dominant matrices

A *diagonally dominant* matrix is a complex matrix B with the property that we have $|b_{ii}| \geq \sum_{j \neq i} |b_{ij}|$ for all i . When all these inequalities are strict, the matrix is called *strictly diagonally dominant*.

Lemma 2.10.1 (i) *A strictly diagonally dominant complex matrix is nonsingular.*

(ii) *A symmetric diagonally dominant real matrix with nonnegative diagonal entries is positive semidefinite.*

(iii) *Let B be a symmetric real matrix with nonnegative row sums and non-positive off-diagonal entries. Define a graph Γ on the index set of the rows of B , where two distinct indices i, j are adjacent when $b_{ij} \neq 0$. The multiplicity of the eigenvalue 0 of B equals the number of connected components C of Γ such that all rows $i \in C$ have zero row sum.*

Proof. Let $B = (b_{ij})$ be diagonally dominant, and let u be an eigenvector, say, with $Bu = bu$. Let $|u_i|$ be maximal among the $|u_j|$. Then $(b_{ii} - b)u_i = -\sum_{j \neq i} b_{ij}u_j$. In all cases the result follows by comparing the absolute values of both sides.

In order to prove (i), assume that B is singular, and that $Bu = 0$. Take absolute values on both sides. We find $|b_{ii}| \cdot |u_i| \leq \sum_{j \neq i} |b_{ij}| \cdot |u_j| \leq \sum_{j \neq i} |b_{ij}| |u_i| < |b_{ii}| \cdot |u_i|$. Contradiction.

For (ii), assume that B has a negative eigenvalue b . Then $(b_{ii} - b) \cdot |u_i| \leq |b_{ii}| \cdot |u_i|$. Contradiction.

For (iii), take $b = 0$ again, and see how equality could hold everywhere in $b_{ii} \cdot |u_i| \leq \sum_{j \neq i} |b_{ij}| \cdot |u_j| \leq \sum_{j \neq i} |b_{ij}| |u_i| \leq b_{ii} \cdot |u_i|$. We see that u must be constant on the connected components of Γ , and zero where row sums are nonzero. \square

2.10.1 Geršgorin circles

The above can be greatly generalized. Let $B(c, r) = \{z \in \mathbb{C} \mid |z - c| \leq r\}$ be the closed ball in \mathbb{C} with center c and radius r .

Proposition 2.10.2 *Let $A = (a_{ij})$ be a complex matrix of order n , and λ an eigenvalue of A . Put $r_i = \sum_{j \neq i} |a_{ij}|$. Then for some i we have $\lambda \in B(a_{ii}, r_i)$. If C is a connected component of $\bigcup_i B(a_{ii}, r_i)$ that contains m of the a_{ii} , then C contains m eigenvalues of A .*

Proof. If $Au = \lambda u$, then $(\lambda - a_{ii})u_i = \sum_{j \neq i} a_{ij}u_j$. Let i be an index for which $|u_i|$ is maximal. Then $|\lambda - a_{ii}| \cdot |u_i| \leq \sum_{j \neq i} |a_{ij}| \cdot |u_j| \leq \sum_{j \neq i} |a_{ij}| |u_i|$ so that $\lambda \in B(a_{ii}, r_i)$. For the second part, use that the eigenvalues are continuous functions of the matrix elements. Let $A(\varepsilon)$ be the matrix with the same diagonal as A and with off-diagonal entries εa_{ij} , so that $A = A(1)$. Then $A(0)$ has eigenvalues a_{ii} , and for $0 \leq \varepsilon \leq 1$ the matrix $A(\varepsilon)$ has eigenvalues inside $\bigcup_i B(a_{ii}, r_i)$. \square

This result is due to Geršgorin [151]. A book-length treatment was given by Varga [300].

2.11 Projections

Lemma 2.11.1 Let $P = \begin{bmatrix} Q & N \\ N^\top & R \end{bmatrix}$ be a real symmetric matrix of order n with two eigenvalues a and b , partitioned with square Q and R . Let Q have h eigenvalues θ_j distinct from a and b . Then R has h eigenvalues $a+b-\theta_j$ distinct from a and b , and $h = m_P(a) - m_Q(a) - m_R(a) = m_P(b) - m_Q(b) - m_R(b)$, where $m_M(\eta)$ denotes the multiplicity of the eigenvalue η of M .

Proof. W.l.o.g. $a = 1$ and $b = 0$ so that P is a projection and $P^2 = P$. Now if $Qu = \theta u$ then $Rv = (1 - \theta)v$ for $v = N^\top u$ and $NN^\top u = \theta(1 - \theta)u$, so that the eigenvalues of Q and R different from 0 and 1 correspond 1-1. The rest follows by taking traces: $0 = \text{tr } P - \text{tr } Q - \text{tr } R = m_P(1) - m_Q(1) - m_R(1) - h$. \square

2.12 Exercises

Exercise 1 Consider a symmetric $n \times n$ matrix A with eigenvalues $\theta_1, \dots, \theta_n$. Suppose A has an equitable partition $\{X_1, \dots, X_m\}$, where all classes have equal size. Let S and B be the characteristic matrix and the quotient matrix of this partition, respectively. Assume that $\theta_1, \dots, \theta_n$ are ordered such that $\theta_1, \dots, \theta_m$ are the eigenvalues of B . Prove that A and SS^\top commute and give an expression for the eigenvalues of $A + \alpha SS^\top$ for $\alpha \in \mathbb{R}$.

Exercise 2 Let B denote the quotient matrix of a symmetric matrix A whose rows and columns are partitioned according to a partitioning $\{X_1, \dots, X_m\}$.

- (i) Give an example, where the eigenvalues of B are a sub(multi)set of the eigenvalues of A , whilst the partition is not equitable.
- (ii) Give an example where the partition is equitable, whilst the interlacing is not tight.

Exercise 3 Let Γ be an undirected graph with smallest eigenvalue -1 . Show that Γ is the disjoint union of complete graphs.

Chapter 3

Graph spectrum (II)

In this chapter we apply the linear algebra from the previous chapter to graph spectra.

3.1 The largest eigenvalue

The largest eigenvalue of a graph is also known as its *spectral radius* or *index*.

The basic information about the largest eigenvalue of a (possibly directed) graph is provided by Perron-Frobenius theory.

Proposition 3.1.1 *Each graph Γ has a real eigenvalue θ_0 with nonnegative real corresponding eigenvector, and such that for each eigenvalue θ we have $|\theta| \leq \theta_0$. The value $\theta_0(\Gamma)$ does not increase when vertices or edges are removed from Γ .*

Assume that Γ is strongly connected. Then

- (i) θ_0 has multiplicity 1.
- (ii) If Γ is primitive (strongly connected, and such that not all cycles have a length that is a multiple of some integer $d > 1$), then $|\theta| < \theta_0$ for all eigenvalues θ different from θ_0 .
- (iii) The value $\theta_0(\Gamma)$ decreases when vertices or edges are removed from Γ . \square

Now let Γ be undirected. By Perron-Frobenius theory and interlacing we find an upper and lower bound for the largest eigenvalue of a connected graph. (Note that A is irreducible if and only if Γ is connected.)

Proposition 3.1.2 *Let Γ be a connected graph with largest eigenvalue θ_1 . If Γ is regular of valency k , then $\theta_1 = k$. Otherwise, we have $k_{\min} < \bar{k} < \theta_1 < k_{\max}$ where k_{\min} , k_{\max} and \bar{k} are the minimum, maximum and average degree.*

Proof. Let $\mathbf{1}$ be the vector with all entries equal to 1. Then $A\mathbf{1} \leq k_{\max}\mathbf{1}$, and by Theorem 2.2.1(iv) we have $\theta_1 \leq k_{\max}$ with equality if and only if $A\mathbf{1} = \theta_1\mathbf{1}$, that is, if and only if Γ is regular of degree θ_1 .

Now consider the partition of the vertex set consisting of a single part. By Corollary 2.5.4 we have $\bar{k} \leq \theta_1$ with equality if and only if Γ is regular. \square

For not necessarily connected graphs, we have $\bar{k} \leq \theta_1 \leq k_{\max}$, and $\bar{k} = \theta_1$ if and only if Γ is regular. If $\theta_1 = k_{\max}$ then we only know that Γ has a regular component with this valency, but Γ need not be regular itself.


As was noted already in Proposition 3.1.1, the largest eigenvalue of a connected graph decreases strictly when an edge is removed.

3.1.1 Graphs with largest eigenvalue at most 2





As an example of the application of Theorem 2.2.1 we can mention:

Theorem 3.1.3 (Smith [285], cf. Lemmens & Seidel [221]). *The only connected graphs having largest eigenvalue 2 are the following graphs (the number of vertices is one more than the index given).*

\hat{A}_n ($n \geq 2$)




\hat{D}_n ($n \geq 4$)



For each graph, the corresponding eigenvector is indicated by the integers at the vertices. Moreover, each connected graph with largest eigenvalue less than 2 is a subgraph of one of the above graphs, i.e., one of the graphs $A_n = P_n$, the path with n vertices ($n \geq 1$), or

D_n ($n \geq 4$)



Finally, each connected graph with largest eigenvalue more than 2 contains one of \hat{A}_n , \hat{D}_n , \hat{E}_6 , \hat{E}_7 , \hat{E}_8 as a subgraph.

Proof. The vectors indicated are eigenvectors for the eigenvalue 2. Therefore, \hat{A}_n , \hat{D}_n and \hat{E}_m ($m = 6, 7, 8$) have largest eigenvalue 2. Any graph containing one of these as an induced proper subgraph has an eigenvalue larger than 2. So, if Γ has largest eigenvalue at most 2 and is not one of \hat{A}_n or \hat{D}_n , then Γ is a

tree without vertices of degree at least 4 and with at most one vertex of degree three, and the result easily follows. \square

These graphs occur as the Dynkin diagrams and extended Dynkin diagrams of finite Coxeter groups, cf. [35, 48, 203]. Let us give their eigenvalues:

The eigenvalues of A_n are $2 \cos i\pi/(n+1)$ ($i = 1, 2, \dots, n$).

The eigenvalues of D_n are 0 and $2 \cos i\pi/(2n-2)$ ($i = 1, 3, 5, \dots, 2n-3$).

The eigenvalues of E_6 are $2 \cos i\pi/12$ ($i = 1, 4, 5, 7, 8, 11$).

The eigenvalues of E_7 are $2 \cos i\pi/18$ ($i = 1, 5, 7, 9, 11, 13, 17$).

The eigenvalues of E_8 are $2 \cos i\pi/30$ ($i = 1, 7, 11, 13, 17, 19, 23, 29$).

(Indeed, these eigenvalues are $2 \cos(d_i - 1)\pi/h$ ($1 \leq i \leq n$) where h is the Coxeter number, and the d_i are the degrees, cf. [48, pp. 84, 308]. Note that in all cases the largest eigenvalue is $2 \cos \pi/h$.)

The eigenvalues of \hat{D}_n are 2, 0, 0, -2 and $2 \cos i\pi/(n-2)$ ($i = 1, \dots, n-3$).

The eigenvalues of \hat{E}_6 are 2, 1, 1, 0, -1 , -1 , -2 .

The eigenvalues of \hat{E}_7 are 2, $\sqrt{2}$, 1, 0, 0, -1 , $-\sqrt{2}$, -2 .

The eigenvalues of \hat{E}_8 are 2, τ , 1, τ^{-1} , 0, $-\tau^{-1}$, -1 , $-\tau$, -2 .

Remark It is possible to go a little bit further, and find all graphs with largest eigenvalue at most $\sqrt{2} + \sqrt{5} \approx 2.05817$, cf. Brouwer & Neumaier [59]. For the graphs with largest eigenvalue at most $\frac{3}{2}\sqrt{2} \approx 2.12132$, see Woo & Neumaier [308] and Cioabă, van Dam, Koolen & Lee [91].

3.1.2 Subdividing an edge

Let Γ be a graph on n vertices, and consider the graph Γ' on $n+1$ vertices obtained from Γ by *subdividing* an edge e (that is, by replacing the edge $e = xy$ by the two edges xz and zy where z is a new vertex). The result below relates the largest eigenvalue of Γ and Γ' .

We say that e lies on an *endpath* if $\Gamma \setminus e$ (the graph on n vertices obtained by removing the edge e from Γ) is disconnected, and one of its connected components is a path.

Proposition 3.1.4 (Hoffman-Smith [197]) *Let Γ be a connected graph, and let the graph Γ' be obtained from Γ by subdividing an edge e . Let Γ and Γ' have largest eigenvalues λ and λ' , respectively. Then if e lies on an endpath, we have $\lambda' > \lambda$, and otherwise $\lambda' \leq \lambda$, with equality only when both equal 2.*

Proof. If e lies on an endpath, then Γ is obtained from Γ' by removing a leaf vertex, and $\lambda < \lambda'$ follows by Proposition 3.1.1. Suppose e is not on an endpath. By Theorem 3.1.3, $\lambda \geq 2$. Let A and A' be the adjacency matrices of Γ and Γ' , so that $Au = \lambda u$ for some vector $u > 0$. We use Theorem 2.2.1 (iv) and conclude $\lambda' \leq \lambda$ from the existence of a nonzero vector v with $v \geq 0$ and $A'v \leq \lambda v$. Such a vector v can be constructed as follows. If z is the new point on the edge $e = xy$, then we can take $v_p = u_p$ for $p \neq z$, and $v_p = \min(u_x, u_y)$, provided that $\lambda v_p \geq u_x + u_y$. Suppose not. W.l.o.g., assume $u_x \leq u_y$, so that $\lambda u_x < u_x + u_y$, and hence $u_x < u_y$. We have $0 \leq \sum_{p \sim x, p \neq y} u_p = \lambda u_x - u_y < u_x$. If x has degree 2 in Γ , say $x \sim p, y$, then replace $e = xy$ by $e = px$ to decrease the values of u on the end points of e —this does not change Γ' . If x has degree $m > 2$, then construct v by $v_x = \lambda u_x - u_y$ and $v_z = u_x$ and $v_p = u_p$ for $p \neq x, z$. We have to check that $\lambda v_x \geq v_x + u_x$, but this follows from $\lambda v_x = \lambda \sum_{p \sim x, p \neq y} u_p \geq (m-1)u_x \geq 2u_x > v_x + u_x$. \square

3.1.3 The Kelmans operation

As we saw, adding edges causes the largest eigenvalue to increase. The operation described below (due to Kelmans [215]) only moves edges, but also increases θ_1 .

Given a graph Γ and two specified vertices u, v construct a new graph Γ' by replacing the edge vx by a new edge ux for all x such that $v \sim x \not\sim u$. The new graph Γ' obtained in this way has the same number of vertices and edges as the old graph, and all vertices different from u, v retain their valency. The vertices u, v are adjacent in Γ' if and only if they are adjacent in Γ . An isomorphic graph is obtained if the rôles of u and v are interchanged: if $N(u)$ and $N(v)$ are the sets of neighbours of u, v distinct from u and v , then in the resulting graph the corresponding sets are $N(u) \cup N(v)$ and $N(u) \cap N(v)$.

If $\bar{\Gamma}$ denotes the complementary graph of Γ then also $\bar{\Gamma}'$ is obtained by a Kelmans operation from $\bar{\Gamma}$.

Proposition 3.1.5 (Csikvári [101]) *Let Γ be a graph, and let Γ' be obtained from Γ by a Kelmans operation. Then $\theta_1(\Gamma) \leq \theta_1(\Gamma')$. (And hence also $\theta_1(\bar{\Gamma}) \leq \theta_1(\bar{\Gamma}')$.)*

Proof. Let A and A' be the adjacency matrices of Γ and Γ' , and let $Ax = \theta_1 x$ where $x \geq 0$, $x^\top x = 1$. W.l.o.g., let $x_u \geq x_v$. Then $\theta_1(\Gamma') \geq x^\top A'x = x^\top Ax + 2(x_u - x_v) \sum_{w \in N(v) \setminus N(u)} x_w \geq \theta_1(\Gamma)$. \square

Csikvári continues and uses this to show that $\theta_1(\Gamma) + \theta_1(\bar{\Gamma}) \leq \frac{1}{2}(1 + \sqrt{3})n$.

Earlier, Brualdi & Hoffman [63] had observed that a graph with maximal spectral radius ρ among the graphs with a given number of vertices and edges has a vertex ordering such that if $x \sim y$ and $z \leq x$, $w \leq y$, $z \neq w$, then $z \sim w$. Rowlinson [263] calls the adjacency matrices of these graphs (ordered this way) *stepwise* and proves that the maximal value of ρ among the graphs on n vertices and e edges is obtained by taking $K_m + (n - m)K_1$, where m is minimal such that $\binom{m}{2} \geq e$, and removing $\binom{m}{2} - e$ edges on a single vertex.

It follows from the above proposition that a graph with maximal $\theta_1(\Gamma) + \theta_1(\bar{\Gamma})$ has stepwise matrix. It is conjectured that in fact $\theta_1(\Gamma) + \theta_1(\bar{\Gamma}) < \frac{4}{3}n - 1$.

3.2 Interlacing

By Perron-Frobenius theory, the largest eigenvalue of a connected graph goes down when one removes an edge or a vertex. Interlacing also gives information about what happens with the other eigenvalues.

The pictures for A and L differ. The eigenvalues for the adjacency matrix A show nice interlacing behavior when one removes a vertex, but not when an edge is removed. (Cf. §1.9.) The Laplace eigenvalues behave well in both cases. For A an eigenvalue can go both up or down when an edge is removed. For L it cannot increase.

Proposition 3.2.1 (i) *Let Γ be a graph and Δ an induced subgraph. Then the eigenvalues of Δ interlace those of Γ ,*

(ii) *Let Γ be a graph and let Δ be a subgraph, not necessarily induced, on m vertices. Then the i -th largest Laplace eigenvalue of Δ is not larger than the i -th largest Laplace eigenvalue of Γ ($1 \leq i \leq m$), and the i -th largest signless Laplace*

eigenvalue of Δ is not larger than the i -th largest signless Laplace eigenvalue of Γ ($1 \leq i \leq m$).

Proof. Part (i) is immediate from Corollary 2.5.2. For part (ii), recall that we have $L = NN^\top$ when N is the directed point-edge incidence matrix obtained by orienting the edges of Γ arbitrarily, and that NN^\top and $N^\top N$ have the same nonzero eigenvalues. Removing an edge from Γ corresponds to removing a column from N , and leads to a principal submatrix of $N^\top N$, and interlacing holds. Removing an isolated vertex from Γ corresponds to removing a Laplace eigenvalue 0. The same proof applies to the signless Laplace matrix. \square

3.3 Regular graphs

It is possible to see from the spectrum whether a graph is regular:

Proposition 3.3.1 *Let Γ be a graph with eigenvalues $k = \theta_1 \geq \theta_2 \geq \dots \geq \theta_v$. Equivalent are:*

(i) Γ is regular (of degree k),

(ii) $AJ = kJ$,

(iii) $\sum \theta_i^2 = kv$.

Proof. We have seen that (i) and (ii) are equivalent. Also, if Γ is regular of degree k , then $\sum \theta_i^2 = \text{tr } A^2 = kv$. Conversely, if (iii) holds, then $\bar{k} = v^{-1} \sum \theta_i^2 = \theta_1$ and, by Proposition 3.1.2, Γ is regular. \square

As we saw above in §1.3.7, it is also possible to see from the spectrum whether a graph is regular and connected. However, for nonregular graphs it is not possible to see from the spectrum whether it is connected.

A very useful characterization of regular connected graphs was given by Hoffman [194]:

Proposition 3.3.2 *The graph Γ is regular and connected if and only if there exists a polynomial p such that $J = p(A)$.*

Proof. If $J = p(A)$, then J commutes with A and hence Γ is regular (and clearly also connected). Conversely, let Γ be connected and regular. Choose a basis such that the commuting matrices A and J become diagonal. Then J becomes $\text{diag}(v, 0, \dots, 0)$ and A becomes $\text{diag}(k, \theta_2, \dots, \theta_v)$. Hence, if we put $f(x) = \prod_{i=2}^v (x - \theta_i)$, then $J = vf(A)/f(k)$, so that $p(x) = vf(x)/f(k)$ satisfies the requirements. \square

3.4 Bipartite graphs

Among the connected graphs Γ , those with imprimitive A are precisely the bipartite graphs (and for these, A has period 2). Consequently we find from Theorem 2.2.1(iii):

Proposition 3.4.1 (i) *A graph Γ is bipartite if and only if for each eigenvalue θ of Γ , $-\theta$ is also an eigenvalue, with the same multiplicity.*

(ii) *If Γ is connected with largest eigenvalue θ_1 , then Γ is bipartite if and only if $-\theta_1$ is an eigenvalue of Γ .*

Proof. For connected graphs all is clear from the the Perron-Frobenius theorem. That gives (ii) and (by taking unions) the ‘only if’ part of (i). For the ‘if’ part of (i), let θ_1 be the spectral radius of Γ . Then some connected component of Γ has eigenvalues θ_1 and $-\theta_1$, and hence is bipartite. Removing its contribution to the spectrum of Γ we see by induction on the number of components that all components are bipartite. \square

3.5 Classification of integral cubic graphs

As an application of Proposition 3.3.2, let us classify the cubic graphs (graphs, that are regular of valency 3) with integral spectrum. The result is due to Bussemaker & Cvetković [68]. See also Schwenk [268]. There are 13 examples.

case	v	spectrum	description
i	6	$\pm 3, 0^4$	$K_{3,3}$
ii	8	$\pm 3, (\pm 1)^3$	2^3
iii	10	$\pm 3, \pm 2, (\pm 1)^2, 0^2$	$K_{2,3}^* \otimes K_2$
iv	12	$\pm 3, (\pm 2)^2, \pm 1, 0^4$	$C_6 \times K_2$
v	20	$\pm 3, (\pm 2)^4, (\pm 1)^5$	$\Pi \otimes K_2$
vi	20	$\pm 3, (\pm 2)^4, (\pm 1)^5$	$T^* \otimes K_2$
vii	24	$\pm 3, (\pm 2)^6, (\pm 1)^3, 0^4$	$\Sigma \otimes K_2$
viii	30	$\pm 3, (\pm 2)^9, 0^{10}$	$GQ(2, 2)$
ix	4	$3, (-1)^3$	K_4
x	6	$3, 1, 0^2, (-2)^2$	$K_3 \times K_2$
xi	10	$3, 1^5, (-2)^4$	Π
xii	10	$3, 2, 1^3, (-1)^2, (-2)^3$	$(\Pi \otimes K_2)/\sigma$
xiii	12	$3, 2^3, 0^2, (-1)^3, (-2)^3$	Σ

A quotient of the hexagonal grid

Let us describe a graph that comes up in the classification. Take a tetrahedron and cut off each corner. Our graph Σ is the 1-skeleton of the resulting polytope, or, equivalently, the result of replacing each vertex of K_4 by a triangle (a $Y - \Delta$ operation). It can also be described as the line graph of the graph obtained from K_4 by subdividing each edge. The bipartite double $\Sigma \otimes K_2$ of Σ is more beautiful (for example, its group is a factor 6 larger than that of Σ), and can be described as the quotient $\Lambda/6\Lambda$ of the hexagonal grid $\Lambda = \langle a + b\omega \mid a, b \in \mathbb{Z}, a + b = 0, 1 \pmod{3} \rangle$ in the complex plane, with $\omega^2 + \omega + 1 = 0$. Now Σ is found e.g. as $\Lambda/\langle 3a + 6b\omega \mid a, b \in \mathbb{Z} \rangle$.

Cubic graphs with loops

For a graph Γ where all vertices have degree 2 or 3, let Γ^* be the cubic graph (with loops) obtained by adding a loop at each vertex of degree 2. Note that the

sum of the eigenvalues of Γ^* , the trace of its adjacency matrix, is the number of loops.

The graph $K_{2,3}^*$ has spectrum $3, 1, 1, 0, -2$.

Let T be the graph on the singletons and pairs in a 4-set, where adjacency is inclusion. Then T^* has spectrum $3^1, 2^3, 1^2, (-1)^3, (-2)^1$.

The classification

Let Π be the Petersen graph and Σ, T the graphs described above.

We split the result into two propositions, one for the bipartite and one for the nonbipartite case.

Proposition 3.5.1 *Let Γ be a connected bipartite cubic graph such that all of its eigenvalues are integral. Then Γ is one of 8 possible graphs, namely (i) $K_{3,3}$, (ii) 2^3 , (iii) $K_{2,3}^* \otimes K_2$, (iv) $C_6 \times K_2$, (v) the Desargues graph (that is, the bipartite double $\Pi \otimes K_2$ of the Petersen graph Π), (vi) T^* (cospectral with the previous), (vii) the bipartite double of Σ , (viii) the point-line incidence graph of the generalized quadrangle of order 2 (that is, the unique 3-regular bipartite graph with diameter 4 and girth 8, also known as Tutte's 8-cage).*

Proof. Let Γ have spectrum $(\pm 3)^1(\pm 2)^a(\pm 1)^b 0^{2c}$ (with multiplicities written as exponents).

The total number of vertices is $v = 2 + 2a + 2b + 2c$. The total number of edges is $\frac{3}{2}v = \frac{1}{2}\text{tr}A^2 = 9 + 4a + b$ (so that $2b + 3c = 6 + a$). The total number of quadrangles is $q = 9 - a - b$, as one finds by computing $\text{tr}A^4 = 15v + 8q = 2(81 + 16a + b)$. The total number of hexagons is $h = 10 + 2b - 2c$, found similarly by computing $\text{tr}A^6 = 87v + 96q + 12h = 2(729 + 64a + b)$.

Somewhat more detailed, let q_u be the number of quadrangles on the vertex u , and q_{uv} the number of quadrangles on the edge uv , and similarly for h . Let uv be an edge. Then $A_{uv} = 1$ and $(A^3)_{uv} = 5 + q_{uv}$ and $(A^5)_{uv} = 29 + 2q_u + 2q_v + 6q_{uv} + h_{uv}$.

The Hoffman polynomial $A(A+3I)(A^2-I)(A^2-4I)$ defines a rank 1 matrix with eigenvalue 720, so that $A(A+3I)(A^2-I)(A^2-4I) = \frac{720}{v}J$ and in particular $v|240$ since for an edge xy the xy entry of $\frac{720}{v}J$ must be divisible by 3. This leaves for (a, b, c, v) the possibilities a) $(0, 0, 2, 6)$, b) $(0, 3, 0, 8)$, c) $(1, 2, 1, 10)$, d) $(2, 1, 2, 12)$, e) $(3, 3, 1, 16)$, f) $(4, 5, 0, 20)$, g) $(5, 1, 3, 20)$, h) $(6, 3, 2, 24)$, i) $(9, 0, 5, 30)$.

In case a) we have $K_{3,3}$, case (i) of the theorem.

In case b) we have the cube 2^3 , case (ii).

In case c) we have a graph of which the bipartite complement has spectrum $2^2 1^2 0^2 (-1)^2 (-2)^2$ hence is the disjoint union of a 4-cycle and a 6-cycle, case (iii).

In case d) we have $q = 6$ and $h = 8$. Let uv be an edge, and evaluate $A(A+3I)(A^2-I)(A^2-4I) = 60J$ at the uv position to find $(A^5 - 5A^3 + 4A)_{uv} = 20$ and $2q_u + 2q_v + q_{uv} + h_{uv} = 12$. It follows that uv cannot lie in 3 or more quadrangles. Suppose u lies in (at least) 3 quadrangles. Then for each neighbor x of u we have $2q_x + h_{ux} = 4$ so that $q_x = 2$ and $h_{ux} = 0$. The mod 2 sum of two quadrangles on u is not a hexagon, and it follows that we have a $K_{2,3}$ on points u, w, x, y, z (with u, w adjacent to x, y, z). The six quadrangles visible in the $K_{2,3}$ on u, w, x, y, z contribute $6 + 4 + 2 + 0$ to $2q_u + 2q_x + q_{ux} + h_{ux} = 12$, and it follows that there are no further quadrangles or hexagons on these points.

So the three further neighbors p, q, r of x, y, z are distinct and have no common neighbors, impossible since $v = 12$. So, no vertex is in 3 or more quadrangles, and hence every vertex u is in precisely 2 quadrangles. These two quadrangles have an edge uu' in common, and we find an involution interchanging each u and u' , and preserving the graph. It follows that we either have $C_6 \times K_2$ (and this has the desired spectrum, it is case (iv)), or a twisted version, but that has only 6 hexagons.

In case e) we have $v = 16$ vertices. For any vertex x , Hoffman's polynomial yields $(A^6)_{xx} - 5(A^4)_{xx} + 4(A^2)_{xx} = 45$. On the other hand, $(A^{2i})_{xx}$ is odd for each i , since each walk of length $2i$ from x to x can be paired with the reverse walk, so that the parity of $(A^{2i})_{xx}$ is that of the number of self-reverse walks $x \dots zwz \dots x$ which is 3^i . Contradiction.

In case f) we have $v = 20, q = 0, h = 20$. Since $c = 0$ we can omit the factor A from Hoffman's polynomial and find $(A + 3I)(A^2 - I)(A^2 - 4I) = 12J$. If u, w have even distance, then $(A^4 - 5A^2 + 4I)_{uw} = 4$. In particular, if $d(u, w) = 2$ then $9 = (A^4)_{uw} = 7 + h_{uw}$ so that $h_{uw} = 2$: each 2-path uvw lies in two hexagons. If no 3-path uvw lies in two hexagons then the graph is distance-regular with intersection array $\{3, 2, 2, 1, 1; 1, 1, 2, 2, 3\}$ (cf. Chapter 11) and hence is the Desargues graph. This is case (v) of the theorem. Now assume that the 3-path uvw lies in two hexagons, so that there are three paths $u \sim v_i \sim w_i \sim x$ ($i = 1, 2, 3$). The v_i and w_i need one more neighbor, say $v_i \sim y_i$ and $w_i \sim z_i$ ($i = 1, 2, 3$). The vertices y_i are distinct since there are no quadrangles, and similarly the z_i are distinct. The vertices y_i and z_j are nonadjacent, otherwise they would be a quadrangle (if $i = j$) or uv_jw_j would be in three hexagons (if $i \neq j$). Remain 6 more vertices, 3 each adjacent to two vertices y_i and 3 each adjacent to two vertices z_i . Call them s_i and t_i , where $s_i \sim y_j$ and $t_i \sim z_j$ whenever $i \neq j$. The final part is a matching between the s_i and the t_i . Now the 2-path $v_iw_iz_i$ is in two hexagons, and these must be of the form $t_jz_iw_iv_is_k$ with $j \neq i \neq k$, and necessarily $j = k$, that is, the graph is uniquely determined. This is case (vi) of the theorem.

In case g) we have $v = 20, q = 3, h = 6$. For an edge uv we have $(A^5 - 5A^3 + 4A)_{uv} = 12$, so that $2q_u + 2q_v + q_{uv} + h_{uv} = 4$. But that means that the edge uv cannot be in a quadrangle, contradiction.

In case h) we have $v = 24, q = 0, h = 12$. For an edge uv we have $(A^5 - 5A^3 + 4A)_{uv} = 10$, so that $h_{uv} = 2$. It follows that each vertex is in 3 hexagons, and each 2-path vuw is in a unique hexagon. Now one straightforwardly constructs the unique cubic bipartite graph on 24 vertices without quadrangles and such that each 2-path is in a unique hexagon. Starting from a vertex u , call its neighbors v_i ($i = 1, 2, 3$), call the six vertices at distance two w_{ij} ($i, j = 1, 2, 3$ and $i \neq j$), and let x_i ($i = 1, 2, 3$) be the three vertices opposite u in a hexagon on u , so that the three hexagons on u are $uv_iw_{ij}x_kw_{ji}v_j$ (with distinct i, j, k). Let the third neighbor of w_{ij} be y_{ij} , and let the third neighbor of x_k be z_k . Necessarily $z_k \sim y_{kj}$. Now each vertex y_{ij} still needs a neighbor and there are two more vertices, say $s \sim y_{12}, y_{23}, y_{31}$ and $t \sim y_{13}, y_{21}, y_{32}$. This is case (vii).

In case i) we have $v = 30, q = h = 0$ and we have Tutte's 8-case. This is case (viii). \square

Proposition 3.5.2 *Let Γ be a connected nonbipartite cubic graph such that all of its eigenvalues are integral. Then Γ is one of 5 possible graphs, namely (ix) K_4 , (x) $K_3 \times K_2$, (xi) the Petersen graph, (xii) the graph on 10 vertices defined*

by $i \sim (i+1) \pmod{10}$, $0 \sim 5$, $1 \sim 3$, $2 \sim 6$, $4 \sim 8$, $7 \sim 9$ (or, equivalently, the graph obtained from $K_{3,3}$ by replacing two nonadjacent vertices by a triangle with a $Y - \Delta$ operation), (xiii) Σ .

Proof. Consider $\Gamma \otimes K_2$. It is cubic and has integral eigenvalues, hence is one of the 8 graphs Δ found in the previous proposition. There is an involution σ of $\Delta = \Gamma \otimes K_2$ without fixed edges, that interchanges the two vertices x' and x'' for each vertex x of Γ . Now Γ can be retrieved as Δ/σ .

In cases (i), (iii), (viii) the graph Γ would be cubic on an odd number of vertices, impossible.

In case (ii), σ must interchange antipodes, and the quotient $2^3/\sigma$ is the complete graph K_4 . This is case (ix).

In case (iv), $C_6 \times K_2$, σ must interchange antipodes in the same copy of C_6 , and the quotient is $K_3 \times K_2$. This is case (x).

In case (v), $\Pi \otimes K_2$, we get the Petersen graph for a σ that interchanges antipodal vertices. This is case (xi). The group is $\text{Sym}(5).2$ and has two conjugacy classes of suitable involutions σ . The second one interchanges x' with $(12)x''$, and its quotient is obtained from Π by replacing the hexagon $13 \sim 24 \sim 15 \sim 23 \sim 14 \sim 25 \sim 13$ by the two triangles $13, 14, 15$ and $23, 24, 25$. This is case (xii).

In case (vi) there is no suitable σ . (An automorphism σ must interchange the two vertices u, x found in the previous proof, since this is the only pair of vertices joined by three 3-paths. But any shortest ux -path is mapped by σ into a different xu -path (since the path has odd length, and σ cannot preserve the middle edge) so that the number of such paths, which is 3, must be even.)

In case (vii) we get Σ . This is case (xiii). (The group of $\Sigma \otimes K_2$ has order 144, six times the order of the group $\text{Sym}(4)$ of Σ , and all possible choices of σ are equivalent.) \square

3.6 The largest Laplace eigenvalue

If $\mu_1 \leq \dots \leq \mu_n$ are the Laplace eigenvalues of a simple graph Γ , then $0 \leq n - \mu_n \leq \dots \leq n - \mu_2$ are the Laplace eigenvalues of the complement of Γ (see §1.3.2). Therefore $\mu_n \leq n$ with equality if and only if the complement of Γ is disconnected. If Γ is regular with valency k we know (by Proposition 3.4.1) that $\mu_n \leq 2k$, with equality if and only if Γ is bipartite. More generally:

Proposition 3.6.1 *Let Γ be a graph with adjacency matrix A (with eigenvalues $\theta_1 \geq \dots \geq \theta_n$), Laplacian L (with eigenvalues $\mu_1 \leq \dots \leq \mu_n$), and signless Laplacian Q (with eigenvalues $\rho_1 \geq \dots \geq \rho_n$). Then*

(i) (Zhang & Luo [313])

$$\mu_n \leq \rho_1.$$

If Γ is connected, then equality holds if and only if Γ is bipartite.

(ii) *Let d_x be the degree of the vertex x . If Γ has at least one edge then*

$$\rho_1 \leq \max_{x \sim y} (d_x + d_y).$$

Equality holds if and only if Γ is regular or bipartite semiregular.

(iii) (Yan [309])

$$2\theta_i \leq \rho_i \quad (1 \leq i \leq n).$$

Proof. (i) Apply Theorem 2.2.1 (vi).

(ii) Using Proposition 3.1.2 to bound the largest eigenvalue $\theta_1(L(\Gamma))$ of $L(\Gamma)$ by its maximum degree $\max_{x \sim y} (d_x + d_y - 2)$ we find $\rho_1 = \theta_1(L(\Gamma)) + 2 \leq \max_{x \sim y} (d_x + d_y)$, with equality if and only if $L(\Gamma)$ is regular so that Γ is regular or bipartite semiregular.

(iii) Since $Q = L + 2A$ and L is positive semidefinite, this follows from the Courant-Weyl inequalities (Theorem 2.8.1 (iii)). \square

Corollary 3.6.2 ([8]) *Let Γ be a graph on n vertices with at least one edge. Then*

$$\mu_n \leq \max_{x \sim y} (d_x + d_y).$$

If Γ is connected, then equality holds if and only if Γ is bipartite regular or semiregular. \square

For bipartite graphs, L and Q have the same spectrum (see Proposition 1.3.10). It follows by Perron-Frobenius that the largest Laplace eigenvalue of a connected bipartite graph decreases strictly when an edge is removed.

Interlacing provides a lower bound for μ_n :

Proposition 3.6.3 ([166]) *Let Γ be a graph on n vertices with at least one edge, and let d_x be the degree of the vertex x . Then*

$$\mu_n \geq 1 + \max_x d_x.$$

If Γ is connected, then equality holds if and only if $\max_x d_x = n - 1$.

Proof. If Γ has a vertex of degree d , then it has a subgraph $K_{1,d}$ (not necessarily induced), and $\mu_n \geq d + 1$. If equality holds, then Γ does not have a strictly larger bipartite subgraph. If Γ is moreover connected, then $d = n - 1$. \square

Deriving bounds on μ_n has become an industry—there are many papers, cf. [36, 124, 170, 223, 224, 231, 246, 312].

3.7 Laplace eigenvalues and degrees

The Schur inequality (Theorem 2.6.1) immediately yields an inequality between the sum of the largest m Laplace eigenvalues and the sum of the largest m vertex degrees. Grone [165] gave a slightly stronger result.

Proposition 3.7.1 *If Γ is connected, with Laplace eigenvalues $\nu_1 \geq \nu_2 \geq \dots \geq \nu_n = 0$ and vertex degrees $d_1 \geq d_2 \geq \dots \geq d_n > 0$, then for $1 \leq m \leq n - 1$ we have $1 + \sum_{i=1}^m d_i \leq \sum_{i=1}^m \nu_i$.*

Proof. Let x_i have degree d_i and put $Z = \{x_1, \dots, x_m\}$. Let $N(Z)$ be the set of vertices outside Z with a neighbor in Z . Instead of assuming that Γ is connected we just use that $N(Z)$ is nonempty. If we delete the vertices outside $Z \cup N(Z)$ then $\sum_{z \in Z} d_z$ does not change, while $\sum_{i=1}^m \nu_i$ does not increase, so we may assume $X = Z \cup N(Z)$. Let R be the quotient matrix of L for the partition $\{\{z\} \mid z \in Z\} \cup \{N(Z)\}$ of X , and let $\lambda_1 \geq \dots \geq \lambda_{m+1}$ be the eigenvalues of R . The matrix R has row sums 0, so $\lambda_{m+1} = 0$. By interlacing (Corollary 2.5.4) we have $\sum_{i=1}^m \nu_i \geq \sum_{i=1}^m \lambda_i = \sum_{i=1}^{m+1} \lambda_i = \text{tr } R = \sum_{z \in Z} d_z + R_{m+1, m+1}$ and the desired result follows since $R_{m+1, m+1} \geq 1$. \square

Second proof. We prove the following stronger statement:

For any graph Γ (not necessarily connected) and any subset Z of the vertex set X of Γ one has $h + \sum_{z \in Z} d_z \leq \sum_{i=1}^m \nu_i$, where d_z denotes the degree of the vertex z in Γ , and $m = |Z|$, and h is the number of connected components of the graph Γ_Z induced on Z that are not connected components of Γ .

We may assume that Γ is connected, and that Z and $X \setminus Z$ are nonempty. Now h is the number of connected components of Γ_Z .

The partition $\{Z, X \setminus Z\}$ of X induces a partition $L = \begin{bmatrix} B & -C \\ -C^\top & E \end{bmatrix}$. Since Γ is connected, B is nonsingular by Lemma 2.10.1(iii). All entries of B^{-1} are nonnegative. (Write $B = n(I - T)$ where $T \geq 0$, then $B^{-1} = \frac{1}{n}(I + T + T^2 + \dots) \geq 0$. If $h = 1$, then $B^{-1} > 0$.)

Since L is positive semidefinite, we can write $L = MM^\top$, where $M = \begin{bmatrix} P & 0 \\ Q & R \end{bmatrix}$ is a square matrix. Now $B = PP^\top$ and $-C = PQ^\top$. The eigenvalues of MM^\top are the same as those of $M^\top M$, and that latter matrix has submatrix $P^\top P + Q^\top Q$ of order m . By Schur's inequality we get $\sum_{i=1}^m \nu_i \geq \text{tr}(P^\top P + Q^\top Q) = \sum_{z \in Z} d_z + \text{tr } Q^\top Q$, and it remains to show that $\text{tr } Q^\top Q \geq h$.

Now $Q^\top Q = P^{-1}CC^\top P^{-1}$, so $\text{tr } Q^\top Q = \text{tr } B^{-1}CC^\top$. We have $B = L_Z + D$, where L_Z is the Laplacian of Γ_Z , and D is the diagonal matrix of the row sums of C . Since $CC^\top \geq D$ and $B^{-1} \geq 0$, we have $\text{tr } Q^\top Q \geq \text{tr } B^{-1}D$. If $L_Z u = 0$ then $(L_Z + D)^{-1}Du = u$. Since L_Z has eigenvalue 0 with multiplicity h , $B^{-1}D$ has eigenvalue 1 with multiplicity h . Since this matrix is positive semidefinite (since $D^{1/2}B^{-1}D^{1/2}$ is p.s.d.), its trace is at least h . \square

A lower bound for the individual ν_j was conjectured by Guo [171] and proved in Brouwer & Haemers [54].

Proposition 3.7.2 *Let Γ be a graph with Laplace eigenvalues $\nu_1 \geq \nu_2 \geq \dots \geq \nu_n = 0$ and with vertex degrees $d_1 \geq d_2 \geq \dots \geq d_n$. Let $1 \leq m \leq n$. If Γ is not $K_m + (n - m)K_1$, then $\nu_m \geq d_m - m + 2$.* \square

We saw the special case $m = 1$ in Proposition 3.6.3. The cases $m = 2$ and $m = 3$ were proved earlier in [222] and [171].

Examples with equality are given by complete graphs K_m with a pending edges at each vertex (where $a > 0$), with Laplace spectrum consisting of 0, $1^{m(a-1)}$, $a + 1$ and $\frac{1}{2}(m + a + 1 \pm \sqrt{(m + a + 1)^2 - 4m})$ with multiplicity $m - 1$ each, so that $\nu_m = a + 1 = d_m - m + 2$.

Further examples are complete graphs K_m with a pending edges attached at a single vertex. Here $n = m + a$, the Laplace spectrum consists of $m + a$, m^{m-2} , 1^a and 0, so that $\nu_m = 1 = d_m - m + 2$.

Any graph contained in $K_{a,a}$ and containing $K_{2,a}$ has $\nu_2 = a = d_2$, with equality for $m = 2$.

Any graph on n vertices with $d_1 = n - 1$ has equality for $m = 1$.

More generally, whenever one has an eigenvector u and vertices x, y with $u_x = u_y$, then u remains eigenvector, with the same eigenvalue, if we add or remove an edge between x and y . Many of the above examples can be modified by adding edges. This leads to many further cases of equality.

3.8 The Grone-Merris Conjecture

3.8.1 Threshold graphs

A *threshold graph* is a graph obtained from the graph K_0 by a sequence of operations of the form (i) add an isolated vertex, or (ii) take the complement.

Proposition 3.8.1 *Let Γ be a threshold graph with Laplace eigenvalues (in nonincreasing order) $\nu_1 \geq \nu_2 \geq \dots \geq \nu_n = 0$. Let d_x be the degree of the vertex x . Then*

$$\nu_j = \#\{x \mid d_x \geq j\}.$$

Proof. Induction on the number of construction steps of type (i) or (ii). \square

Grone & Merris [166] conjectured that this is the extreme case, and that for all undirected graphs and all t one has

$$\sum_{j=1}^t \nu_j \leq \sum_{j=1}^t \#\{x \mid d_x \geq j\}.$$

For $t = 1$ this is immediate from $\nu_1 \leq n$. For $t = n$ equality holds. This conjecture was proved in Hua Bai [13], see §3.8.2 below. There is a generalization to higher-dimensional simplicial complexes, see §3.9 below.

A variation on the Grone-Merris conjecture is the following.

Conjecture *Let Γ be a graph with e edges and Laplace eigenvalues $\nu_1 \geq \nu_2 \geq \dots \geq \nu_n = 0$. Then for each t we have $\sum_{i=1}^t \nu_i \leq e + \binom{t+1}{2}$.*

It is easy to see (by induction) that this inequality holds for threshold graphs. In [180] it is proved for trees, and in case $t = 2$. In [23] it is shown that there is a t such that the t -th inequality of this conjecture is sharper than the t -th Grone-Merris inequality if and only if the graph is non-split. In particular, this conjecture holds for split graphs. It also holds for regular graphs.

3.8.2 Proof of the Grone-Merris Conjecture

Very recently, Hua Bai [13] proved the Grone-Merris Conjecture. We repeat the statement of the theorem.

Theorem 3.8.2 *Let Γ be an undirected graph with Laplace eigenvalues (in non-increasing order) $\nu_1 \geq \nu_2 \geq \dots \geq \nu_n = 0$. Let d_x be the degree of the vertex x . Then for all t , $0 \leq t \leq n$, we have*

$$\sum_{i=1}^t \nu_i \leq \sum_{i=1}^t \#\{x \mid d_x \geq i\}. \quad (3.1)$$

The proof is by reducing to the case of a *semi-bipartite graph* (or *split graph*), that is a graph where the vertex set is the disjoint union of a nonempty subset inducing a clique (complete graph), and a nonempty subset inducing a coclique (edgeless graph). Then for semi-bipartite graphs a continuity argument proves the crucial inequality stated in the following lemma.

Lemma 3.8.3 *Let Γ be a semi-bipartite graph with clique of size c and Laplace eigenvalues be $\nu_1 \geq \nu_2 \geq \dots \geq \nu_n = 0$. Let δ be the maximum degree among the vertices in the coclique, so that $\delta \leq c$. If $\nu_c > c$ or $\nu_c = c > \delta$ then we have $\sum_{i=1}^c \nu_i \leq \sum_{i=1}^c \#\{x \mid d_x \geq i\}$.*

Proof of Theorem 3.8.2 (assuming Lemma 3.8.3). Consider counterexamples to (3.1) with minimal possible t .

Step 1 *If Γ is such a counterexample with minimal number of edges, and x, y are vertices in Γ of degree at most t , then they are nonadjacent.*

Indeed, if $x \sim y$ then let Γ' be the graph obtained from Γ by removing the edge xy . Then $\sum_{i=1}^t \#\{x \mid d'_x \geq i\} + 2 = \sum_{i=1}^t \#\{x \mid d_x \geq i\}$. The Laplace matrices L and L' of Γ and Γ' satisfy $L = L' + H$ where H has eigenvalues $2, 0^{n-1}$. By Theorem 2.8.2 we have $\sum_{i=1}^t \nu_i \leq \sum_{i=1}^t \nu'_i + 2$, and since Γ' has fewer edges than Γ we find $\sum_{i=1}^t \nu_i \leq \sum_{i=1}^t \nu'_i + 2 \leq \sum_{i=1}^t \#\{x \mid d'_x \geq i\} + 2 = \sum_{i=1}^t \#\{x \mid d_x \geq i\}$, contradiction. \square

Step 2 *There is a semi-bipartite counterexample Γ for the same t , with clique size $c := \#\{x \mid d_x \geq t\}$.*

Indeed, we can form a new graph Γ from the Γ of Step 1 by adding edges xy for every pair of nonadjacent vertices x, y , both of degree at least t . Now $\sum_{i=1}^t \#\{x \mid d_x \geq i\}$ does not change, and $\sum_{i=1}^t \nu_i$ does not decrease, and the new graph is semi-bipartite with the stated clique size. \square

This will be our graph Γ for the rest of the proof.

Step 3 *A semi-bipartite graph Δ of clique size c satisfies $\nu_{c+1} \leq c \leq \nu_{c-1}$.*

Indeed, since Δ contains the complete graph K_c with Laplace spectrum $c^{c-1}, 0$, we see by the Courant-Weyl inequalities (Theorem 2.8.1 (iii)) that $\nu_{c-1} \geq c$. And since Δ is contained in the complete semi-bipartite graph with clique of size c and coclique of size $n - c$ and all edges in-between, with Laplace spectrum $n^c, c^{n-c-1}, 0$, we have $\nu_{c+1} \leq c$. \square

Since t was chosen minimal, we have $\nu_t > \#\{x \mid d_x \geq t\} = c$. The previous step then implies $c \geq t$. If $c = t$ then $\nu_c > c$ and Lemma 3.8.3 gives a contradiction. So $c > t$.

All vertices in the coclique of Γ have degree at most $t - 1$ and all vertices in the clique have degree at least $c - 1$. So $\#\{x \mid d_x \geq i\} = c$ for $t \leq i \leq c - 1$. From Step 3 we have $\nu_i \geq \nu_{c-1} \geq c$ for $t \leq i \leq c - 1$. Since $\sum_{i=1}^t \nu_i > \sum_{i=1}^t \#\{x \mid d_x \geq i\}$ we also have $\sum_{i=1}^{c-1} \nu_i > \sum_{i=1}^{c-1} \#\{x \mid d_x \geq i\}$. Now if $\nu_c \geq c$ we contradict Lemma 3.8.3 (since $\#\{x \mid d_x \geq c\} \leq c$). So $\nu_c < c$.

Step 4 *The m -th Grone-Merris inequality for a graph Γ is equivalent to the $(n - 1 - m)$ -th Grone-Merris inequality for its complement $\bar{\Gamma}$ ($1 \leq m \leq n - 1$).*

Indeed, $\bar{\Gamma}$ has Laplace eigenvalues $\bar{\nu}_i = n - \nu_{n-i}$ ($1 \leq i \leq n-1$) and dual degrees $\#\{x \mid \bar{d}_x \geq i\} = n - \#\{x \mid d_x \geq n-i\}$, and $\sum_{i=1}^n \nu_i = \sum_{i=1}^n \#\{x \mid d_x \geq i\}$. \square

In our case $\bar{\Gamma}$ is semibipartite with clique size $n - c$, and by the above we have $\bar{\nu}_{n-c} = n - \nu_c > n - c$ and $\sum_{i=1}^{n-c} \bar{\nu}_i > \sum_{i=1}^{n-c} \#\{x \mid \bar{d}_x \geq i\}$. This contradicts Lemma 3.8.3.

This contradiction completes the proof of the Grone-Merris conjecture, except that Lemma 3.8.3 still has to be proved.

Proof of Lemma 3.8.3

Let Γ be a semi-bipartite graph with clique of size c and coclique of size $n - c$. The partition of the vertex set induces a partition of the Laplace matrix $L = \begin{bmatrix} K + D & -A \\ -A^\top & E \end{bmatrix}$, where K is the Laplacian of the complete graph K_c and A is the $c \times (n - c)$ adjacency matrix between vertices in the clique and the coclique, and D and E are diagonal matrices with the row and column sums of A .

Step 5 If $\nu_c \geq c$, then $\sum_{i=1}^c \#\{x \mid d_x \geq i\} = c^2 + \text{tr } D$.

Indeed, all vertices in the clique have degree at least c , for if some vertex x in the clique had degree $c - 1$ then we could move it to the coclique and find $\nu_c \leq c - 1$ from Step 3, contrary to the assumption. It follows that $\sum_{i=1}^c \#\{x \mid d_x \geq i\} = \sum_x \min(c, d_x) = c^2 + \text{tr } E = c^2 + \text{tr } D$. \square

Step 6 Suppose that the subspace W spanned by the L -eigenvectors belonging to ν_1, \dots, ν_c is spanned by the columns of $\begin{bmatrix} I \\ X \end{bmatrix}$. Then $L \begin{bmatrix} I \\ X \end{bmatrix} = \begin{bmatrix} I \\ X \end{bmatrix} Z$ for some matrix Z , and $\sum_{i=1}^c \nu_i = \text{tr } Z$.

Indeed, if $\begin{bmatrix} U \\ V \end{bmatrix}$ has these eigenvectors as columns, then $L \begin{bmatrix} U \\ V \end{bmatrix} = \begin{bmatrix} U \\ V \end{bmatrix} T$ where T is the diagonal matrix with the eigenvalues. Now $\begin{bmatrix} U \\ V \end{bmatrix} = \begin{bmatrix} I \\ X \end{bmatrix} U$, so that $L \begin{bmatrix} I \\ X \end{bmatrix} = \begin{bmatrix} I \\ X \end{bmatrix} Z$ where $Z = UTU^{-1}$ and $\text{tr } Z = \text{tr } T = \sum_{i=1}^c \nu_i$. \square

Suppose we are in the situation of the previous step, and that moreover X is nonpositive. Let δ be the maximum degree among the vertices in the coclique, so that $\delta \leq c$. We have to show that if $\nu_c > c$ or $\nu_c = c > \delta$ then $\text{tr } Z \leq c^2 + \text{tr } D$.

Now $L \begin{bmatrix} I \\ X \end{bmatrix} = \begin{bmatrix} K + D - AX \\ -A^\top + EX \end{bmatrix}$, so $Z = K + D - AX$, and $\text{tr } Z = \text{tr}(K + D - AX) = c(c - 1) + \text{tr } D - \text{tr}(AX)$, and we need $\text{tr}(AX) \geq -c$. But since $c < n$, the eigenvectors are orthogonal to $\mathbf{1}$ so that X has column sums -1 . Since X is nonpositive, $\text{tr}(AX) \geq -c$ follows, and we are done.

By interlacing ν_{c+1} is at most the largest eigenvalue of E , that is δ , which by hypothesis is smaller than ν_c . Hence the subspace of vectors $\begin{bmatrix} 0 \\ * \end{bmatrix}$ meets W trivially, so that W has a basis of the required form. Only nonpositivity of X remains, and the following lemma completes the proof.

Lemma 3.8.4 If $\nu_c > \delta$, then the invariant subspace W spanned by the L -eigenvectors for ν_i , $1 \leq i \leq c$, is spanned by the columns of $\begin{bmatrix} I \\ X \end{bmatrix}$ where X is

nonpositive.

Proof. We argue by continuity, viewing $L = L(A)$ and $X = X(A)$ as functions of the real-valued matrix A , where $0 \leq A \leq J$. (Now D has the row sums of A , and E has the column sums, and δ is the largest element of the diagonal matrix E .) We write J for the $c \times (n - c)$ all-1 matrix, and J_c for the all-1 matrix of order c , so that $JX = -J_c$.

Our hypothesis $\nu_c > \delta$ holds for all matrices $L^{(\alpha)} := L(\alpha A + (1 - \alpha)J) = \alpha L + (1 - \alpha)L^{(0)}$, for $0 \leq \alpha \leq 1$. Indeed, let $L^{(\alpha)}$ have eigenvalues $\nu_i^{(\alpha)}$, so that $\nu_c^{(0)} = n$ and $\nu_{c+1}^{(0)} = \nu_{n-1}^{(0)} = c$. The matrix $L^{(\alpha)}$ has lower left-hand corner $\alpha E + (1 - \alpha)cI$ so that $\delta^{(\alpha)} = \alpha\delta + (1 - \alpha)c$. The c -space W is orthogonal to $\mathbf{1}$, so that $\nu_c^{(\alpha)} \geq \alpha\nu_c + (1 - \alpha)c$ (by Theorem 2.4.1), and hence $\nu_c^{(\alpha)} > \delta^{(\alpha)}$ for $0 < \alpha \leq 1$, and also for $\alpha = 0$, since $\nu_c^{(0)} = n$ and $\delta^{(0)} = c$. It follows that $\nu_c^{(\alpha)} > \nu_{c+1}^{(\alpha)}$ for $0 \leq \alpha \leq 1$.

As we used already, $L(J)$ has spectrum $n^c, c^{n-c-1}, 0$, and one checks that $X(J) = -\frac{1}{n-c}J^\top < 0$, as desired. Above we found the condition $XZ = -A^\top + EX$ on X , that is, $X(K + D - AX) - EX + A^\top = 0$, that is, $X(K + J_c + D) = XJ_c + XAX + EX - A^\top = -X(J - A)X + EX - A^\top$. It follows, since $K + J_c + D$ is a positive diagonal matrix, that if $X \leq 0$ and $A > 0$, then $X < 0$. The matrix $X(A)$ depends continuously on A (in the region where $\nu_{c+1} < \nu_c$) and is strictly negative when $A > 0$. Then it is nonpositive when $A \geq 0$. \square

3.9 The Laplacian for hypergraphs

Let a *simplicial complex* on a finite set S be a collection \mathcal{C} of subsets of S (called simplices) that is an order ideal for inclusion, that is, such that if $A \in \mathcal{C}$ and $B \subseteq A$ then also $B \in \mathcal{C}$. Let the *dimension* of a simplex A be one less than its cardinality, and let the dimension of a simplicial complex be the maximum of the dimensions of its simplices. Given a simplicial complex \mathcal{C} , let \mathcal{C}_i (for $i \geq -1$) be the vector space (over any field) that has the simplices of dimension i as basis. Order the simplices arbitrarily (say, using some order on S) and define $\partial_i : \mathcal{C}_i \rightarrow \mathcal{C}_{i-1}$ by $\partial_i s_0 \dots s_i = \sum_j (-1)^j s_0 \dots \hat{s}_j \dots s_i$. Then $\partial_{i-1}\partial_i = 0$ for all $i \geq 0$.

Let N_i be the matrix of ∂_i on the standard basis, and put $L_i = N_{i+1}N_{i+1}^\top$ and $L'_i = N_i^\top N_i$. The matrices L_i generalize the Laplacian. Indeed, in the case of a 1-dimensional simplicial complex (that is, a graph) the ordinary Laplace matrix is just L_0 , and L'_0 is the all-1 matrix J .

Since $\partial_i\partial_{i+1} = 0$ we have $L_i L'_i = L'_i L_i = 0$, generalizing $LJ = JL = 0$.

We have $\text{tr } L_{i-1} = \text{tr } L'_i = (i + 1)|\mathcal{C}_i|$. This generalizes the facts that $\text{tr } L$ is twice the number of edges, and $\text{tr } J$ the number of vertices.

In case the underlying field is \mathbb{R} , we have the direct sum decomposition $\mathcal{C}_i = \text{im } N_{i+1} \oplus \ker(L_i + L'_i) \oplus \text{im } N_i^\top$. (Because then $M^\top Mx = 0$ if and only if $Mx = 0$.) Now $\ker N_i = \text{im } N_{i+1} \oplus \ker(L_i + L'_i)$ so that the i -th reduced homology group is $H_i(\mathcal{C}) := \ker N_i / \text{im } N_{i+1} \cong \ker(L_i + L'_i)$.

Example The spectrum of L_{m-2} for a simplicial complex containing all m -subsets of an n -set (the complete m -uniform hypergraph) consists of the eigenvalue n with multiplicity $\binom{n-1}{m-1}$ and all further eigenvalues are 0.

Indeed, we may regard simplices $s_0 \dots s_{m-1}$ as elements $s_0 \wedge \dots \wedge s_{m-1}$ of an exterior algebra. Then the expression $s_0 \dots s_{m-1}$ is defined regardless of the order of the factors, and also when factors are repeated. Now $N_i t_0 \dots t_i = \sum_j (-1)^j t_0 \dots \widehat{t_j} \dots t_i$ and for the complete $(i+2)$ -uniform hypergraph we have $N_i^\top t_0 \dots t_i = \sum_t t t_0 \dots t_i$, so that $L_i + L'_i = nI$. It follows that $N_{i+1}^\top N_{i+1} N_{i+1}^\top = n N_{i+1}^\top$, and L_i has eigenvalues 0 and n . The multiplicities follow by taking the trace.

Duval & Reiner [136] generalized the Grone-Merris conjecture. Given an m -uniform hypergraph \mathcal{H} , let d_x be the number of edges containing the vertex x . Let the spectrum of \mathcal{H} be that of the matrix L_{m-2} for the simplicial complex consisting of all subsets of edges of \mathcal{H} .

Conjecture *Let the m -uniform hypergraph \mathcal{H} have degrees d_x , and Laplace eigenvalues ν_i , ordered such that $\nu_1 \geq \nu_2 \geq \dots \geq 0$. Then for all t we have*

$$\sum_{j=1}^t \nu_j \leq \sum_{j=1}^t \#\{x \mid d_x \geq j\}.$$

Equality for all t holds if and only if \mathcal{H} is invariant under downshifting.

The part about ‘downshifting’ means the following: Put a total order on the vertices of \mathcal{H} in such a way that if $x \leq y$ then $d_x \geq d_y$. Now \mathcal{H} is said to be invariant under downshifting if whenever $\{x_1, \dots, x_m\}$ is an edge of \mathcal{H} , and $\{y_1, \dots, y_m\}$ is an m -set with $y_i \leq x_i$ for all i , then also $\{y_1, \dots, y_m\}$ is an edge of \mathcal{H} . If this holds for one total order, then it holds for any total order that is compatible with the degrees.

For $m = 2$ this is precisely the Grone-Merris conjecture. (And the graphs that are invariant for downshifting are precisely the threshold graphs.) The ‘if’ part of the equality case is a theorem:

Theorem 3.9.1 (Duval & Reiner [136]) *If \mathcal{H} is an m -uniform hypergraph with degrees d_x and Laplace eigenvalues ν_i with $\nu_1 \geq \nu_2 \geq \dots \geq 0$, and \mathcal{H} is invariant for downshifting, then $\nu_j = \#\{x \mid d_x \geq j\}$ for all t .*

In particular it follows that hypergraphs invariant for downshifting have integral Laplace spectrum.

For example, the complete m -uniform hypergraph on an underlying set of size n has degrees $\binom{n-1}{m-1}$ so that $\nu_j = n$ for $1 \leq j \leq \binom{n-1}{m-1}$ and $\nu_j = 0$ for $\binom{n-1}{m-1} < j \leq \binom{n}{m}$, as we already found earlier.

Dominance order

The conjecture and theorem can be formulated more elegantly in terms of *dominance order*. Let $\mathbf{a} = (a_i)$ and $\mathbf{b} = (b_i)$ be two finite nonincreasing sequences of nonnegative real numbers. We say that \mathbf{b} *dominates* \mathbf{a} , and write $\mathbf{a} \leq \mathbf{b}$, when $\sum_{i=1}^t a_i \leq \sum_{i=1}^t b_i$ for all t , and $\sum_{i=1}^\infty a_i = \sum_{i=1}^\infty b_i$, where missing elements are taken to be zero.

For example, in this notation Schur’s inequality (Theorem 2.6.1) says that $\mathbf{d} \leq \boldsymbol{\theta}$ if \mathbf{d} is the sequence of diagonal elements and $\boldsymbol{\theta}$ the sequence of eigenvalues of a real symmetric matrix.

If $\mathbf{a} = (a_j)$ is a finite nonincreasing sequences of nonnegative integers, then \mathbf{a}^\top denotes the sequence (a_j^\top) with $a_j^\top = \#\{i \mid a_i \geq j\}$. If \mathbf{a} is represented by a Ferrers diagram, then \mathbf{a}^\top is represented by the transposed diagram.

For example, the Duval-Reiner conjecture says that $\boldsymbol{\mu}' \trianglelefteq \mathbf{d}^\top$.

If \mathbf{a} and \mathbf{b} are two nonincreasing sequences, then let $\mathbf{a} \cup \mathbf{b}$ denote the (multiset) union of both sequences, with elements sorted in nonincreasing order.

Lemma 3.9.2

- (i) $\mathbf{a}^{\top\top} = \mathbf{a}$,
- (ii) $(\mathbf{a} \cup \mathbf{b})^\top = \mathbf{a}^\top + \mathbf{b}^\top$ and $(\mathbf{a} + \mathbf{b})^\top = \mathbf{a}^\top \cup \mathbf{b}^\top$,
- (iii) $\mathbf{a} \trianglelefteq \mathbf{b}$ if and only if $\mathbf{b}^\top \trianglelefteq \mathbf{a}^\top$.

□

3.10 Applications of eigenvectors

Sometimes it is not the eigenvalue but the eigenvector that is needed. We sketch very briefly some of the applications.

3.10.1 Ranking




Figure 3.1: Graph with Perron-Frobenius eigenvector

In a network, important people have many connections. One would like to pick out the vertices of highest degree and call them the most important. But it is not just the number of neighbors. Important people have connections to many other important people. If one models this and says that up to some constant of proportionality one's importance is the sum of the importances of one's neighbors in the graph, then the vector giving the importance of each vertex becomes an eigenvector of the graph, necessarily the Perron-Frobenius eigenvector if importance cannot be negative. The constant of proportionality is then the largest eigenvalue.

3.10.2 Google Page rank

Google uses a similar scheme to compute the Page Rank [40] of web pages. The authors described (in 1998) the algorithm as follows:

Suppose pages x_1, \dots, x_m are the pages that link to a page y . Let page x_i have d_i outgoing links. Then the PageRank of y is given by

$$PR(y) = 1 - \alpha + \alpha \sum_i \frac{PR(x_i)}{d_i}.$$

The PageRanks form a probability distribution: $\sum_x PR(x) = 1$. The vector of PageRanks can be calculated using a simple iterative algorithm, and corresponds to the principal eigenvector of the normalized link matrix of the web. A PageRank for 26 million web pages can be computed in a few hours on a medium size workstation. A suitable value for α is $\alpha = 0.85$.

In other words, let Γ be the directed graph on n vertices consisting of all web pages found, with an arrow from x to y when page x contains a hyperlink to page y . Let A be the adjacency matrix of Γ (with $A_{xy} = 1$ if there is a link from x to y). Let D be the diagonal matrix of outdegrees, so that the scaled matrix $S = D^{-1}A$ has row sums 1, and construct the positive linear combination $M = \frac{1-\alpha}{n}J + \alpha S$ with $0 < \alpha < 1$. Since $M > 0$ the matrix M has a unique positive left eigenvector u , normed so that $\sum u_x = 1$. Now $M\mathbf{1} = \mathbf{1}$ and hence $uM = u$. The PageRank of the web page x is the value u_x .

A small detail is the question what to do when page x does not have outgoing edges, so that row x in A is zero. One possibility is to do nothing (and take $D_{xx} = 1$). Then u will have eigenvalue less than 1.

The vector u is found by starting with an approximation (or just any positive vector) u_0 and then computing the limit of the sequence $u_i = u_0 M^i$. That is easy: the matrix M is enormous, but A is sparse: on average a web page does not have more than a dozen links. The constant α regulates the speed of convergence: convergence is determined by the 2nd largest eigenvalue, which is bounded by α ([211]). It is reported that 50 to 100 iterations suffice. A nonzero α guarantees that the matrix is irreducible. An α much less than 1 guarantees quick convergence. But an α close to 1 is better at preserving the information in A . Intuitively, u_x represents the expectation of finding oneself at page x after many steps, where each step consists of either (with probability α) clicking on a random link on the current page, or (with probability $1-\alpha$) picking a random internet page. Note that the precise value of u_x is unimportant—only the ordering among the values u_x is used.

There are many papers (and even books) discussing Google's PageRank. See e.g. [66], [24].

3.10.3 Cutting

Often the cheapest way to cut a connected graph into two pieces is by partitioning it into a single vertex (of minimal valency) and the rest. But in the area of clustering (see also below) one typically wants relatively large pieces. Here the second Laplace eigenvector helps. Without going into any detail, let us try the same example as above.

We see that cutting the edges where the 2nd Laplace eigenvector changes sign is fairly successful in this case.

3.10.4 Graph drawing

Often, a reasonable way to draw a connected graph is to take Laplace eigenvectors u and v for the 2nd and 3rd smallest Laplace eigenvalues, and draw the vertex x at the point with coordinates (u_x, v_x) . See, e.g. [218].




Figure 3.2: Graph with 2nd Laplace eigenvector

One can justify this as follows. Let the *energy* of an embedding $\rho : \Gamma \rightarrow \mathbb{R}^m$ be the sum of the squared edge lengths $\sum_e \|\rho(x) - \rho(y)\|^2$ where the sum is over all edges $e = xy$. Let R be the $m \times n$ matrix of which the columns are the vertex images $\rho(x)$. Then the energy of ρ equals RLR^\top . For graph drawing one would like to minimize the energy, given some normalization so that not all vertices are mapped close to the origin or close to some lower-dimensional subspace of \mathbb{R}^m . Pisanski & Shawe-Taylor [260] propose to require $R\mathbf{1} = 0$ and $RR^\top = I$, so that the origin is the center of mass, and $\|R^\top v\|^2 = \|v\|^2$ for all vectors $v \in \mathbb{R}^m$: no vector is almost perpendicular to the entire drawing. In this situation the minimum energy is $\sum_{i=2}^{m+1} \mu_i$, and this minimum is achieved when the row space of R contains the Laplace eigenvectors of μ_2, \dots, μ_{m+1} . The authors also discuss variations of this setup.

3.10.5 Clustering

Given a large data set, one often wants to cluster it. If the data is given as a set of vectors in some Euclidean space \mathbb{R}^m , then a popular clustering algorithm is *k-means*:

Given a set X of N vectors in \mathbb{R}^m and a number k , find a partition of X into k subsets X_1, \dots, X_k such that $\sum_{i=1}^k \sum_{x \in X_i} \|x - c_i\|^2$ is as small as possible, where $c_i = (1/|X_i|) \sum_{x \in X_i} x$ is the centroid of X_i .

The usual algorithm uses an iterative approach. First choose the k vectors c_i in some way, arbitrary or not. Then take X_i to be the subset of X consisting of the vectors closer to c_i than to the other c_j (breaking ties arbitrarily). Then compute new vectors c_i as the centroids of the sets X_i , and repeat. In common practical situations this algorithm converges quickly, but one can construct examples where this takes exponential time. The final partition found need not be optimal, but since the algorithm is fast, it can be repeated a number of times with different starting points c_i .

Now if the data is given as a graph, one can compute eigenvectors u_1, \dots, u_m for the m smallest eigenvalues μ_1, \dots, μ_m of the Laplace matrix L , and assign to the vertex x the vector $(u_i(x))_i$ and apply a vector space clustering algorithm such as *k-means* to the resulting vectors.

This is reasonable. For example, if the graph is disconnected with c connected components, then the first c eigenvalues of L are zero, and the first c eigenvectors are (linear combinations of) the characteristic functions of the connected components.

This approach also works when one has more detailed information—not adjacent/nonadjacent but a (nonnegative) similarity or closeness measure. (One uses an edge-weighted graph, with $A_{xy} = w(x, y)$ and $d_x = \sum_y w(x, y)$ and D the diagonal matrix with $D_{xx} = d_x$, and $L = D - A$. Again L is positive semidefinite, with $u^\top Lu = \sum w(x, y)(u(x) - u(y))^2$. The multiplicity of the eigenvalue 0 is the number of connected components of the underlying graph where points x, y are adjacent when $w(x, y) > 0$.)

Especially important is the special case where one searches for the cheapest cut of the graph into two relatively large pieces. If the graph is connected, then map the vertices into \mathbb{R}^1 using $x \mapsto u(x)$, where u is the eigenvector for the second smallest eigenvalue of L , and then use 2-means to cluster the resulting points. Compare §1.7 on the algebraic connectivity of a graph.

Several matrices related to the Laplacian have been used in this context. It seems useful to normalize the matrix, and to retain the property that if the graph is disconnected the characteristic functions of components are eigenvectors. A suitable matrix is $L_{\text{norm}} = D^{-1}L = I - D^{-1}A$.

There is a large body of literature on clustering in general and spectral clustering in particular. A few references are [169, 235, 288].

3.10.6 Searching an eigenspace

There exists a unique strongly regular graph¹ with parameters $(v, k, \lambda, \mu) = (162, 56, 10, 24)$ found as the second subconstituent of the McLaughlin graph. Its vertex set can be split into two halves such that each half induces a strongly regular graph with parameters $(v, k, \lambda, \mu) = (81, 20, 1, 6)$. How many such splits are there? Can we find them all?

In this and many similar situations one can search an eigenspace. The first graph has spectrum $56^1 \ 2^{140} \ (-16)^{21}$ and a split gives an eigenvector with eigenvalue -16 if we take the vector that is 1 on the subgraph and -1 on the rest.

It is easy to construct an explicit basis (u_i) for the 21-dimensional eigenspace, where the j -th coordinate of u_i is δ_{ij} . Construct the 2^{21} eigenvectors that are ± 1 on the first 21 coordinates and inspect the remaining coordinates. If all are ± 1 one has found a split into two regular graphs of valency 20. In this particular case there are 224 such subgraphs, 112 splits, and all subgraphs occurring are strongly regular with the abovementioned parameters.

3.11 Exercises

Exercise 1 Consider a graph with largest eigenvalue θ_1 and maximum valency k_{max} . Use interlacing to show that $\theta_1 \geq \sqrt{k_{\text{max}}}$. When does equality hold?

Exercise 2 Let Γ be a k -regular graph with n vertices and eigenvalues $k = \theta_1 \geq \dots \geq \theta_n$. Let Γ' be an induced subgraph of Γ with n' vertices and average degree k' .

- (i) Prove that $\theta_2 \geq \frac{nk' - n'k}{n - n'} \geq \theta_n$.

¹For strongly regular graphs, see Chapter 8. No properties are used except that the substructure of interest corresponds to an eigenvector of recognizable shape.

- (ii) What can be said in case of equality (on either side)?
- (iii) Deduce Hoffman's bound (Theorem 4.1.2) from the above inequality.

Exercise 3 Prove the conjecture from Section 3.8.1 for regular graphs. (Hint: use Cauchy-Schwarz.)

Chapter 4

Substructures

Interlacing yields information on subgraphs of a graph, and the way such subgraphs are embedded. In particular, one gets bounds on extremal substructures.

Probably the most important use of the spectrum is its use to imply connectivity, randomness, and expansion properties of a graph.

For the adjacency spectrum the important parameter is the second largest eigenvalue θ_2 , cf. §1.7, or rather the ratio θ_2/θ_1 . A small ratio implies high connectivity. For the Laplace spectrum it is the second smallest, μ_2 , or rather the ratio μ_2/μ_n . A large ratio implies high connectivity.

The Laplace eigenvector belonging to μ_2 has been used in graph partitioning problems.

4.1 Cliques and cocliques

A *clique* in a graph is a set of pairwise adjacent vertices. A *coclique* in a graph is a set of pairwise nonadjacent vertices. The *clique number* $\omega(\Gamma)$ is the size of the largest clique in Γ . The *independence number* $\alpha(\Gamma)$ is the size of the largest coclique in Γ .

Let Γ be a graph on n vertices (undirected, simple, and loopless) having an adjacency matrix A with eigenvalues $\theta_1 \geq \dots \geq \theta_n$. Both Corollaries 2.5.2 and 2.5.4 lead to a bound for $\alpha(\Gamma)$.

Theorem 4.1.1 $\alpha(\Gamma) \leq n - n_- = |\{i | \theta_i \geq 0\}|$ and $\alpha(\Gamma) \leq n - n_+ = |\{i | \theta_i \leq 0\}|$.

Proof. A has a principal submatrix $B = 0$ of size $\alpha = \alpha(\Gamma)$. Corollary 2.5.2 gives $\theta_\alpha \geq \eta_\alpha = 0$ and $\theta_{n-\alpha-1} \leq \eta_1 = 0$. \square

Theorem 4.1.2 If Γ is regular of nonzero degree k , then

$$\alpha(\Gamma) \leq n \frac{-\theta_n}{k - \theta_n},$$

and if a coclique C meets this bound, then every vertex not in C is adjacent to precisely $-\theta_n$ vertices of C .

Proof. We apply Corollary 2.5.4. The coclique gives rise to a partition of A with quotient matrix

$$B = \begin{bmatrix} 0 & k \\ \frac{k\alpha}{n-\alpha} & k - \frac{k\alpha}{n-\alpha} \end{bmatrix},$$

where $\alpha = \alpha(\Gamma)$. B has eigenvalues $\eta_1 = k = \theta_1$ (the row sum) and $\eta_2 = -k\alpha/(n - \alpha)$ (since $\text{trace } B = k + \eta_2$) and so $\theta_n \leq \eta_2$ gives the required inequality. If equality holds, then $\eta_2 = \theta_n$, and since $\eta_1 = \theta_1$, the interlacing is tight and hence the partition is equitable. \square

The first bound is due to Cvetković [103]. The second bound is an unpublished result of Hoffman.

Example The Petersen graph has 10 vertices and spectrum $3^1 1^5 (-2)^4$, and its independence number is 4. So both bounds are tight.

4.1.1 Using weighted adjacency matrices

Let us call a real symmetric matrix B a weighted adjacency matrix of a graph Γ when B has rows and columns indexed by the vertex set of Γ , has zero diagonal, and satisfies $B_{xy} = 0$ whenever $x \not\sim y$.

The proof of Theorem 4.1.1 applies to B instead of A , and we get

Theorem 4.1.3 $\alpha(\Gamma) \leq n - n_-(B)$ and $\alpha(\Gamma) \leq n - n_+(B)$.

Similarly, the proof of Theorem 4.1.2 remains valid for weighted adjacency matrices B with constant row sums.

Theorem 4.1.4 Let B be a weighted adjacency matrix of Γ with constant row sums b and smallest eigenvalue s . Then $\alpha(\Gamma) \leq n(-s)/(b - s)$.

4.2 Chromatic number

A *proper vertex coloring* of a graph is an assignment of colors to the vertices so that adjacent vertices get different colors. (In other words, a partition of the vertex set into cocliques.) The *chromatic number* $\chi(\Gamma)$ is the minimum number of colors of a proper vertex coloring of Γ .

Proposition 4.2.1 (Wilf [307]) Let Γ be connected with largest eigenvalue θ_1 . Then $\chi(\Gamma) \leq 1 + \theta_1$ with equality if and only if Γ is complete or is an odd cycle.

Proof. Put $m = \chi(\Gamma)$. Since Γ cannot be colored with $m - 1$ colors, while coloring vertices of degree less than $m - 1$ is easy, there must be an induced subgraph Δ of Γ with minimum degree at least $m - 1$. Now $\theta_1 \geq \theta_1(\Delta) \geq d_{\min}(\Delta) \geq m - 1 = \chi(\Gamma) - 1$. If equality holds, then by Perron-Frobenius $\Gamma = \Delta$ and Δ is regular of degree $m - 1$ (by Proposition 3.1.2), and the conclusion follows by Brooks' theorem. \square

Since each coclique (color class) has size at most $\alpha(\Gamma)$, we have $\chi(\Gamma) \geq n/\alpha(\Gamma)$ for a graph Γ with n vertices. Thus upper bounds for $\alpha(\Gamma)$ give lower bounds for $\chi(\Gamma)$. For instance if Γ is regular of degree $k = \theta_1$ then Theorem 4.1.2 implies that $\chi(\Gamma) \geq 1 - \frac{\theta_1}{\theta_n}$. This bound remains however valid for non-regular graphs.

Theorem 4.2.2 (Hoffman [195]) If Γ is not edgeless then $\chi(\Gamma) \geq 1 - \frac{\theta_1}{\theta_n}$.

Proof. Put $m = \chi(\Gamma)$. Since Γ is not edgeless, $\theta_n < 0$. Now, by part (i) of the following proposition, $\theta_1 + (m - 1)\theta_n \leq \theta_1 + \theta_{n-m+2} + \dots + \theta_n \leq 0$. \square

Proposition 4.2.3 Put $m = \chi(\Gamma)$. Then

$$(i) \quad \theta_1 + \theta_{n-m+2} + \dots + \theta_n \leq 0.$$

$$(ii) \quad \text{If } n > m, \text{ then } \theta_2 + \dots + \theta_m + \theta_{n-m+1} \geq 0.$$

$$(iii) \quad \text{If } n > tm, \text{ then } \theta_{t+1} + \dots + \theta_{t+m-1} + \theta_{n-t(m-1)} \geq 0.$$

Proof. Let A have orthonormal eigenvectors u_j , so that $Au_j = \theta_j u_j$.

(i) Let $\{X_1, \dots, X_m\}$ be a partition of Γ into m cocliques, where $m = \chi(\Gamma)$. Let x_j be the pointwise product of u_1 with the characteristic vector of X_j , so that $\sum x_j = u_1$. Now apply Corollary 2.5.3 to the vectors x_j , after deleting those that are zero. The matrix C defined there satisfies $C\mathbf{1} = \theta_1 \mathbf{1}$, and has zero diagonal, and has eigenvalues η_j interlacing those of A . Hence

$$0 = \text{tr}(C) = \eta_1 + \dots + \eta_m \geq \theta_1 + \theta_{n-m+2} + \dots + \theta_n.$$

(ii) Put $A' = A - (\theta_1 - \theta_n)u_1 u_1^\top$, then A' has the same eigenvectors u_j as A , but with eigenvalues $\theta_n, \theta_2, \dots, \theta_n$. Pick a non-zero vector y in

$$\langle u_{n-m+1}, \dots, u_n \rangle \cap \langle x_1, \dots, x_m \rangle^\perp.$$

The two spaces have non-trivial intersection since the dimensions add up to n and u_1 is orthogonal to both. Let y_j be the pointwise product of y with the characteristic vector of X_j , so that $\sum y_j = y$ and $y_j^\top A' y_j = 0$. Now apply Corollary 2.5.3 to the matrix A' and the vectors y_j , after deleting those that are zero. The matrix C defined there has zero diagonal, and smallest eigenvalue smaller than the Rayleigh quotient $\frac{y^\top A y}{y^\top y}$, which by choice of y is at most θ_{n-m+1} . We find

$$0 = \text{tr}(C) = \eta_1 + \dots + \eta_m \leq \theta_2 + \theta_3 + \dots + \theta_m + \theta_{n-m+1}.$$

(iii) The proof is as under (ii), but this time we move t (instead of just one) eigenvalues away (by subtracting multiples of $u_j u_j^\top$ for $1 \leq j \leq t$). The vector y must be chosen orthogonal to tm vectors, which can be done inside the $(tm - t + 1)$ -space $\langle u_{n-tm+t}, \dots, u_n \rangle$, assuming that this space is already orthogonal to u_1, \dots, u_t , i.e., assuming that $n > tm$. \square

The above proof of Theorem 4.2.2 using (i) above appeared in [174].

A coloring that meets the bound of Theorem 4.2.2 is called a *Hoffman coloring*. For regular graphs, the color classes of a Hoffman coloring are cocliques that meet Hoffman's coclique bound. So in this case all the color classes have equal size and the corresponding matrix partition is equitable.

In [175] more inequalities of the above kind are given. But the ones mentioned here, especially (i) and (ii), are by far the most useful.

Example The complete multipartite graph $K_{m \times a}$ has chromatic number m and spectrum $(am - a)^1, 0^{m(a-1)}, (-a)^{m-1}$. It has equality in Hoffman's inequality (and hence in (i)), and also in (ii).

Example The graph obtained by removing an edge from K_n has chromatic number $n - 1$ and spectrum $\frac{1}{2}(n - 3 + \sqrt{D}), 0, (-1)^{n-3}, \frac{1}{2}(n - 3 - \sqrt{D})$ with $D = (n + 1)^2 - 8$, with equality in (i).

Example Consider the generalized octagon of order $(2, 4)$ on 1755 vertices. It has spectrum $10^1 \ 5^{351} \ 1^{650} \ (-3)^{675} \ (-5)^{78}$. It is not 3-chromatic, as one sees by removing the largest 352 eigenvalues, i.e., by applying (iii) with $t = 352$.

The inequality (ii) looks a bit awkward, but can be made more explicit if the smallest eigenvalue θ_n has a large multiplicity.

Corollary 4.2.4 *If the eigenvalue θ_n has multiplicity m and $\theta_2 > 0$, then*

$$\chi(\Gamma) \geq \min(1 + m, 1 - \frac{\theta_n}{\theta_2}).$$

Proof. If $\chi(\Gamma) \leq m$, then $\theta_n = \theta_{n-\chi(\Gamma)+1}$, hence $\chi(\Gamma) \geq 1 - \frac{\theta_n}{\theta_2}$. \square

4.2.1 Using weighted adjacency matrices

If Γ has an m -coloring, then $\Gamma \times K_m$ has an independent set of size n , the number of vertices of Γ . This means that one can use bounds on the size of an independent set to obtain bounds on the chromatic number.

Example Consider the generalized octagon of order $(2, 4)$ again. Call it Γ , and call its adjacency matrix A . Now consider the weighted adjacency matrix B of $K_3 \times \Gamma$, where the K_3 is weighted with some number r , where $1 < r < \frac{3}{2}$. For each eigenvalue θ of A , we find eigenvalues $\theta + 2r$ (once) and $\theta - r$ (twice) as eigenvalues of B . Applying Theorem 4.1.3 we see that $\alpha(K_3 \times \Gamma) \leq 3(1 + 351) + 650 = 1706$ while Γ has 1755 vertices, so Γ is not 3-chromatic.

4.2.2 Rank and chromatic number

The easiest way for A to have low rank, is when it has many repeated rows. But then Γ contains large cocliques. People have conjectured that it might be true that $\chi(\Gamma) \leq \text{rk } A$ when $A \neq 0$. A counterexample was given by Alon & Seymour [7] who observed that the complement of the folded 7-cube (on 64 vertices) has chromatic number $\chi = 32$ (indeed, $\alpha = 2$), and rank 29 (indeed, the spectrum of the folded 7-cube is $7^1 \ 3^{21} \ (-1)^{35} \ (-5)^7$).

4.3 Shannon capacity

Shannon [282] defined the *Shannon capacity* of a graph as the capacity of the zero-error channel where a transmission consists of sending a vertex of the graph, and two transmissions can be confused when the corresponding vertices are joined by an edge.

The maximum number of possible messages of length 1 is $\alpha(\Gamma)$, and $\log \alpha(\Gamma)$ bits are transmitted. The maximum number of possible messages of length ℓ is $\alpha(\Gamma^\ell)$ where Γ^ℓ is defined (here) as the graph on sequences of ℓ vertices from Γ , where two sequences are adjacent when on each coordinate position their elements are equal or adjacent. This explains the definition $c(\Gamma) = \sup_{\ell \rightarrow \infty} \alpha(\Gamma^\ell)^{1/\ell}$. Now the channel capacity is $\log c(\Gamma)$.

For example, for the pentagon we find $c(\Gamma) \geq \sqrt{5}$ as shown by the 5-coclique 00, 12, 24, 31, 43.

Computing $c(\Gamma)$ is a difficult unsolved problem, even for graphs as simple as C_7 , the 7-cycle.

Clearly, $\alpha(\Gamma) \leq c(\Gamma) \leq \chi(\bar{\Gamma})$. (Indeed, if $m = \chi(\bar{\Gamma})$ then Γ can be covered by m cliques, and Γ^ℓ can be covered by m^ℓ cliques, and $\alpha(\Gamma^\ell) \leq m^\ell$.) In a few cases this suffices to determine $c(\Gamma)$.

One can sharpen the upper bound to the fractional clique covering number. For example, the vertices of C_5 can be doubly covered by 5 cliques, so the vertices of C_5^ℓ can be covered 2^ℓ times by 5^ℓ cliques, and $\alpha(C_5^\ell) \leq (5/2)^\ell$ so that $c(C_5) \leq 5/2$.

If A is the adjacency matrix of Γ , then $\otimes^\ell(A + I) - I$ is the adjacency matrix of Γ^ℓ .

The Hoffman upper bound for the size of cocliques is also an upper bound for $c(\Gamma)$ (and therefore, when the Hoffman bound holds with equality, also the Shannon capacity is determined).

Proposition 4.3.1 (Lovász [232]) *Let Γ be regular of valency k . Then*

$$c(\Gamma) \leq n(-\theta_n)/(k - \theta_n).$$

Proof. Use the weighted Hoffman bound (Theorem 4.1.4). If $B = A - \theta_n I$, then $\otimes^\ell B - (-\theta_n)^\ell I$ has constant row sums $(k - \theta_n)^\ell - (-\theta_n)^\ell$ and smallest eigenvalue $-(-\theta_n)^\ell$, so that $\alpha(\Gamma^\ell) \leq (n(-\theta_n)/(k - \theta_n))^\ell$. \square

Using $n = 5$, $k = 2$, $\theta_n = (-1 - \sqrt{5})/2$ we find for the pentagon $c(\Gamma) \leq \sqrt{5}$. Hence equality holds.

Haemers [172, 173] observed that if B is a matrix indexed by the vertices of Γ and $B_{xx} \neq 0$ for all x , and $B_{xy} = 0$ whenever $x \not\sim y$, then $c(\Gamma) \leq \text{rk } B$. Indeed, for such a matrix $\alpha(\Gamma) \leq \text{rk } B$ since an independent set determines a submatrix that is zero outside a nonzero diagonal. Now $\text{rk } \otimes^\ell B = (\text{rk } B)^\ell$.

Note that in the above the rank may be taken over any field.

Example The collinearity graph Γ of the generalized quadrangle $GQ(2, 4)$ (the complement of the Schläfli graph, cf. §8.6) on 27 vertices has spectrum $10^1 1^{20} (-5)^6$. Taking $B = A - I$ shows that $c(\Gamma) \leq 7$. (And $c(\Gamma) \geq \alpha(\Gamma) = 6$.) The complement $\bar{\Gamma}$ has $\alpha(\bar{\Gamma}) = 3$, but this is also the Hoffman bound, so $c(\bar{\Gamma}) = 3$.

Alon [3] proves that $c(\Gamma + \bar{\Gamma}) \geq 2\sqrt{n}$ for all Γ . Combined with the above example, this shows that the Shannon capacity of the disjoint sum of two graphs can be larger than the sum of their Shannon capacities.

4.4 Stars and star complements

Consider a graph Γ with vertex set X . By interlacing, the multiplicity of any given eigenvalue changes by at most 1 if we remove a vertex. But there is always a vertex such that removing it actually decreases the multiplicity. And that means that if θ is an eigenvalue of multiplicity m we can find a *star subset* for θ , that is, a subset S of X of size m such that $\Gamma \setminus S$ does not have eigenvalue θ . Now $X \setminus S$ is called a *star complement*.

Why precisely can we decrease the multiplicity? Let u be a θ -eigenvector of A , so that $(\theta I - A)u = 0$, and let x be a vertex with $u_x \neq 0$. Then removing x from Γ decreases the multiplicity of θ .

Indeed, removing x is equivalent to the two actions: (i) forcing $u_x = 0$ for eigenvectors u , and (ii) omitting the condition $\sum_{y \sim x} u_y = \theta u_x$ (row x of the

matrix equation $(\theta I - A)u = 0$ for eigenvectors u . Since A is symmetric, the column dependency $(\theta I - A)u = 0$ given by u is also a row dependency, and row x is dependent on the remaining rows, so that (ii) doesn't make a difference. But (i) does, as the vector u shows. So the multiplicity goes down.

This argument shows that the star sets for θ are precisely the sets S of size m such that no θ -eigenvector vanishes on all of S . Also, that any subgraph without eigenvalue θ is contained in a star complement.

Proposition 4.4.1 ([137, 108]) *Let Γ be a graph with eigenvalue θ of multiplicity m . Let S be a subset of the vertex set X of Γ , and let the partition $\{S, X \setminus S\}$ of X induce a partition $A = \begin{bmatrix} B & C \\ C^\top & D \end{bmatrix}$ of the adjacency matrix A . If S is a star set for θ (i.e., if $|S| = m$ and D does not have eigenvalue θ), then $B - \theta I = C(D - \theta I)^{-1}C^\top$.*

Proof. The row space of $A - \theta I$ has rank $n - m$. If S is a star set, then this row space is spanned by the rows of $[C^\top \ D - \theta I]$. Alternatively, apply Corollary 2.7.2 to $A - \theta I$. \square

This proposition says that the edges inside a star set are determined by the rest of the graph (and the value θ). Especially when m is large, this may be useful.

Stars and star complements have been used to study exceptional graphs with smallest eigenvalue not less than -2 , see, e.g., [107, 109, 110]. (One starts with the observation that if θ is the smallest eigenvalue of a graph, then a star complement has smallest eigenvalue larger than θ . But all graphs with smallest eigenvalue larger than -2 are explicitly known.) Several graphs and classes of graphs have been characterized by graph complement. See, e.g., [207, 109].

A *star partition* is a partition of X into star sets S_θ for θ , where θ runs through the eigenvalues of Γ . It was shown in [108] that every graph has a star partition.

4.5 The second largest eigenvalue

There is a tremendous amount of literature about the second largest eigenvalue of a regular graph. If the gap between the largest and second largest eigenvalues is large, then the graph has good connectivity, expansion and randomness properties. (About connectivity, see also §1.7.)

4.5.1 Bounds for the second largest eigenvalue

In this connection it is of interest how large this gap can become. Theorems by Alon-Boppana and Serre say that for large graphs θ_2 cannot be much smaller than $2\sqrt{k-1}$.

Proposition 4.5.1 (Alon-Boppana [2]) *If $k \geq 3$ then for k -regular graphs on n vertices one has*

$$\theta_2 \geq 2\sqrt{k-1} \left(1 - O\left(\frac{\log(k-1)}{\log n}\right)\right).$$

Proposition 4.5.2 (Serre [281]) *For each $\epsilon > 0$, there exists a positive constant $c = c(\epsilon, k)$ such that for any k -regular graph Γ on n vertices, the number of eigenvalues of Γ larger than $(2 - \epsilon)\sqrt{k-1}$ is at least cn .*

Quenell gives (weaker) explicit bounds:

Proposition 4.5.3 ([262]) *Let Γ be a finite graph with diameter d and minimal degree $k \geq 3$. Then for $2 \leq m \leq 1 + d/4$, the m^{th} eigenvalue of the adjacency matrix A of Γ satisfies $\lambda_m > 2\sqrt{k-1} \cos(\frac{\pi}{r+1})$, where $r = \lfloor d/(2m-2) \rfloor$.*

Alon [2] conjectured, and Friedman [147] proved that large random k -regular graphs have second largest eigenvalue smaller than $2\sqrt{k-1} + \epsilon$ (for fixed k , $\epsilon > 0$ and n sufficiently large). Friedman remarks that numerical experiments seem to indicate that random k -regular graphs in fact satisfy $\theta_2 < 2\sqrt{k-1}$.

A connected k -regular graph is called a *Ramanujan graph* when $|\theta| \leq 2\sqrt{k-1}$ for all eigenvalues $\theta \neq k$. (This notion was introduced in [234].) It is not difficult to find such graphs. For example, complete graphs, or Paley graphs, will do. Highly nontrivial was the construction of infinite sequences of Ramanujan graphs with given, constant, valency k and size n tending to infinity. Lubotzky, Phillips & Sarnak [234] and Margulis [240] constructed for each prime $p \equiv 1 \pmod{4}$ an infinite series of Ramanujan graphs with valency $k = p + 1$.

4.5.2 Large regular subgraphs are connected

We note the following trivial but useful result,

Proposition 4.5.4 *Let Γ be a graph with second largest eigenvalue θ_2 . Let Δ be a nonempty regular induced subgraph with largest eigenvalue $\rho > \theta_2$. Then Δ is connected.*

Proof. The multiplicity of the eigenvalue ρ of Δ is the number of connected components of Δ , and by interlacing this is 1. \square

4.5.3 Randomness

Let Γ be a regular graph of valency k on n vertices, and assume that (for some real constant λ) we have $|\theta| \leq \lambda$ for all eigenvalues $\theta \neq k$. The ratio λ/k determines randomness and expansion properties of Γ : the smaller λ/k , the more random, and the better expander Γ is.

For example, the following proposition says that most points have approximately the expected number of neighbors in a given subset of the vertex set. Here $\Gamma(x)$ denotes the set of neighbours of the vertex x in the graph Γ .

Proposition 4.5.5 *Let R be a subset of size r of the vertex set X of Γ . Then*

$$\sum_{x \in X} (|\Gamma(x) \cap R| - \frac{kr}{n})^2 \leq \frac{r(n-r)}{n} \lambda^2.$$

Proof. Apply interlacing to A^2 and the partition $\{R, X \setminus R\}$ of X . The sum of all entries of the matrix A^2 in the (R, R) -block equals the number of paths $y \sim x \sim z$, with $y, z \in R$ and $x \in X$, that is, $\sum_x (|\Gamma(x) \cap R|)^2$. \square

Rather similarly, the following proposition, a version of the *expander mixing lemma* from Alon & Chung [5], says that there are about the expected number of edges between two subsets.

Proposition 4.5.6 *Let S and T be two subsets of the vertex set of Γ , of sizes s and t , respectively. Let $e(S, T)$ be the number of ordered edges xy with $x \in S$ and $y \in T$. Then*

$$|e(S, T) - \frac{kst}{n}| \leq \lambda \sqrt{st(1 - \frac{s}{n})(1 - \frac{t}{n})} \leq \lambda \sqrt{st}.$$

Proof. Write the characteristic vectors χ_S and χ_T of the sets S and T as a linear combination of a set of orthonormal eigenvectors of A : $\chi_S = \sum \alpha_i u_i$ and $\chi_T = \sum \beta_i u_i$ where $Au_i = \theta_i u_i$. Then $e(S, T) = \chi_S^\top A \chi_T = \sum \alpha_i \beta_i \theta_i$. We have $\alpha_1 = s/\sqrt{n}$ and $\beta_1 = t/\sqrt{n}$ and $\theta_1 = k$. Now $|e(S, T) - \frac{kst}{n}| = |\sum_{i>1} \alpha_i \beta_i \theta_i| \leq \lambda \sum_{i>1} |\alpha_i \beta_i|$ and $\sum_{i>1} \alpha_i^2 \leq (\chi_S, \chi_S) - s^2/n = s(n-s)/n$, and $\sum_{i>1} \beta_i^2 \leq t(n-t)/n$, so that $|e(S, T) - \frac{kst}{n}| \leq \lambda \sqrt{st(n-s)(n-t)/n}$. \square

If S and T are equal or complementary, this says that

$$|e(S, T) - \frac{kst}{n}| \leq \lambda \frac{s(n-s)}{n}.$$

In particular, the average valency k_S of an induced subgraph S of size s satisfies $|k_S - \frac{ks}{n}| \leq \lambda \frac{n-s}{n}$. For example, the Hoffman-Singleton graph has $\theta_2 = 2$, $\theta_n = -3$, so $\lambda = 3$ and we find equality for subgraphs $\overline{K_{15}}$ ($s = 15$, $k_S = 0$), $10K_2$ ($s = 20$, $k_S = 1$) and $5C_5$ ($s = 25$, $k_S = 2$).

4.5.4 Expansion

An *expander* is a graph with the property that the number of points at distance at most one from any given (not too large) set is at least a fixed constant (larger than one) times the size of the given set. Expanders became famous because of their rôle in sorting networks (cf. Ajtai-Komlós-Szemerédi [1]) and have since found many other applications. The above proposition already implies that there cannot be too many vertices without neighbors in R . A better bound was given by Tanner [292] (in order to show that generalized polygons are good expanders).

Proposition 4.5.7 (Tanner [292]) *Let R be a set of r vertices of Γ and let $\Gamma(R)$ be the set of vertices adjacent to some point of R . Then*

$$\frac{|\Gamma(R)|}{n} \geq \frac{\rho}{\rho + \frac{\lambda^2}{k^2}(1 - \rho)}$$

where $\rho = r/n$.

Proof. Let χ be the characteristic vector of R . Write it as a linear combination of a set of orthonormal eigenvectors of A : $\chi = \sum \alpha_i u_i$ where $Au_i = \theta_i u_i$. Then $A\chi = \sum \alpha_i \theta_i u_i$ and $(A\chi, A\chi) = \sum \alpha_i^2 \theta_i^2$, so that $\|A\chi\|^2 \leq \alpha_0^2(\theta_0^2 - \lambda^2) + \lambda^2 \sum \alpha_i^2 = (\chi, u_0)^2(k^2 - \lambda^2) + \lambda^2(\chi, \chi) = \frac{r^2}{n}(k^2 - \lambda^2) + r\lambda^2$. Now let ψ be the characteristic vector of $\Gamma(R)$. Then $k^2 r^2 = (A\chi, 1)^2 = (A\chi, \psi)^2 \leq \|A\chi\|^2 \|\psi\|^2 \leq |\Gamma(R)| \cdot (\frac{r^2}{n}(k^2 - \lambda^2) + r\lambda^2)$, proving our claim. \square

The above used two-sided bounds on the eigenvalues different from the valency. It suffices to bound θ_2 . Let the *edge expansion constant* $h(\Gamma)$ of a graph Γ be the minimum of $e(S, T)/|S|$ where the minimum is taken over all partitions $\{S, T\}$ of the vertex set with $|S| \leq |T|$, and where $e(S, T)$ is the number of edges meeting both S and T . Then

Proposition 4.5.8 $\frac{1}{2}(k - \theta_2) \leq h(\Gamma) \leq \sqrt{k^2 - \theta_2^2}$.

Proof. For the lower bound, apply interlacing to A and a partition $\{S, T\}$ of the vertex set, with $s = |S|$ and $t = |T|$. Put $e = e(S, T)$. One finds $ne/st \geq k - \theta_2$, so that $e/s \geq (t/n)(k - \theta_2) \geq \frac{1}{2}(k - \theta_2)$. For the upper bound, consider a nonnegative vector w indexed by the point set X of Γ , with support of size at most $\frac{1}{2}n$. If w_x takes t different nonzero values $a_1 > \dots > a_t > 0$, then let $S_i = \{x \mid w_x \geq a_i\}$ ($1 \leq i \leq t$), and let $m_i = |S_i \setminus S_{i-1}|$ (with $S_0 = \emptyset$). Let $h = h(\Gamma)$. Now

$$h \sum_x w_x \leq \sum_{x \sim y} |w_x - w_y|.$$

Indeed, all S_i have size at most $\frac{1}{2}n$, so at least $h|S_i|$ edges stick out of S_i , and these contribute at least $h(m_1 + \dots + m_i)(a_i - a_{i+1})$ to $\sum_{x \sim y} |w_x - w_y|$ (with $a_{t+1} = 0$). The total contribution is at least $h \sum_i m_i a_i = h \sum_x w_x$.

Let u be an eigenvector of A with $Au = \theta_2 u$. We may assume that $u_x > 0$ for at most $\frac{1}{2}n$ points x (otherwise replace u by $-u$). Define a vector v by $v_x = \max(u_x, 0)$. Since $(Av)_x = \sum_{y \sim x} v_y \geq \sum_{y \sim x} u_y = (Au)_x = \theta_2 u_x = \theta_2 v_x$ if $v_x > 0$, we have $v^\top Av = \sum_x v_x (Av)_x \geq \theta_2 \sum_x v_x^2$.

Note that $\sum_{x \sim y} (v_x \pm v_y)^2 = k \sum_x v_x^2 \pm v^\top Av$.

Apply the above to the nonnegative vector w given by $w_x = v_x^2$. We find $h \sum_x v_x^2 \leq \sum_{x \sim y} |v_x^2 - v_y^2| \leq (\sum_{x \sim y} (v_x - v_y)^2 \cdot \sum_{x \sim y} (v_x + v_y)^2)^{1/2} = ((k \sum_x v_x^2)^2 - (v^\top Av)^2)^{1/2} \leq (\sum_x v_x^2) \sqrt{k^2 - \theta_2^2}$. \square

For similar results for not necessarily regular graphs, see §4.6.

4.5.5 Toughness and Hamiltonicity

As application of the above ideas, one can give bounds for the toughness of a graph in terms of the eigenvalues.

A connected, noncomplete graph Γ is called *t-tough* if one has $|S| \geq tc$ for every disconnecting set of vertices S such that the graph induced on its complement has $c \geq 2$ connected components. The *toughness* $\tau(\Gamma)$ of a graph Γ is the largest t such that Γ is *t-tough*. For example, the Petersen graph has toughness $4/3$.

This concept was introduced by Chvátal [87], who hoped that *t-tough* graphs would be *Hamiltonian* (i.e., have a circuit passing through all vertices) for sufficiently large t . People tried to prove this for $t = 2$, the famous ‘2-tough conjecture’, but examples were given in [18] of *t-tough* nonhamiltonian graphs for all $t < 9/4$. Whether a larger bound on τ suffices is still open.

Still, being tough seems to help. In [19] it was shown that a *t-tough* graph Γ on $n \geq 3$ vertices with minimum degree δ is Hamiltonian when $(t+1)(\delta+1) > n$.

Proposition 4.5.9 ([43]) *Let Γ be a connected noncomplete regular graph of valency k and let $|\theta| \leq \lambda$ for all eigenvalues $\theta \neq k$. Then $\tau(\Gamma) > k/\lambda - 2$.*

This proposition gives the right bound, in the sense that there are infinitely many graphs with $\tau(\Gamma) \leq k/\lambda$. The constant 2 can be improved a little. The result can be refined by separating out the smallest and the second largest eigenvalue. The main tool in the proof is Proposition 4.5.5.

See also the remarks following Theorem 8.3.2.

Krivelevich & Sudakov [219] show that, when n is large enough, a graph on n vertices, regular of degree $k = \theta_1$, and with second largest eigenvalue θ_2 satisfying

$$\frac{\theta_2}{\theta_1} < \frac{(\log \log n)^2}{1000 \log n \log \log \log n}$$

is Hamiltonian. Pyber [261] shows that it follows that every sufficiently large strongly regular graph is Hamiltonian.

4.5.6 The Petersen graph is not Hamiltonian

An amusing application of interlacing shows that the Petersen graph is not Hamiltonian. Indeed, a Hamilton circuit in the Petersen graph would give an induced C_{10} in its line graph. Now the line graph of the Petersen graph has spectrum $4^1 2^5 (-1)^4 (-2)^5$ and by interlacing the seventh eigenvalue $2 \cos \frac{3}{5}\pi = (1 - \sqrt{5})/2$ of C_{10} should be at most -1 , contradiction. (Cf. [247, 190].)

4.5.7 Diameter bound

Chung [85] gave the following diameter bound.

Proposition 4.5.10 *Let Γ be a graph on $n \geq 2$ vertices, regular of valency k , with diameter d , and with eigenvalues $k = \theta_1 \geq \dots \geq \theta_n$. Let $\lambda = \max_{i>1} |\theta_i|$. Then*

$$d \leq \left\lceil \frac{\log(n-1)}{\log(k/\lambda)} \right\rceil.$$

Proof. The graph Γ has diameter at most m when $A^m > 0$. Let A have orthonormal eigenvectors u_i with $Au_i = \theta_i u_i$. Then $A = \sum_i \theta_i u_i u_i^\top$. Take $u_1 = \frac{1}{\sqrt{n}} \mathbf{1}$. Now $(A^m)_{xy} = \sum_i \theta_i^m (u_i^\top u_i)_{xy} \geq \frac{k^m}{n} - \lambda^m \sum_{i>1} |(u_i)_x| \cdot |(u_i)_y|$ and $\sum_{i>1} |(u_i)_x| \cdot |(u_i)_y| \leq (\sum_{i>1} |(u_i)_x|^2)^{1/2} (\sum_{i>1} |(u_i)_y|^2)^{1/2} = (1 - |(u_1)_x|^2)^{1/2} (1 - |(u_1)_y|^2)^{1/2} = 1 - \frac{1}{n}$, so that $(A^m)_{xy} > 0$ if $k^m > (n-1)\lambda^m$. \square

4.6 Separation

Let Γ be a graph with Laplace matrix L and Laplace eigenvalues $0 = \mu_1 \leq \dots \leq \mu_n$. The Laplace matrix of a subgraph Γ' of Γ is not a submatrix of L , unless Γ' is a component. So the interlacing techniques of §2.5 do not work in such a straightforward manner here. But we can obtain results if we consider off-diagonal submatrices of L .

Proposition 4.6.1 *Let X and Y be disjoint sets of vertices of Γ , such that there is no edge between X and Y . Then*

$$\frac{|X||Y|}{(n-|X|)(n-|Y|)} \leq \left(\frac{\mu_n - \mu_2}{\mu_n + \mu_2} \right)^2.$$

Proof. Put $\mu = \frac{1}{2}(\mu_n + \mu_2)$ and define a matrix A of order $2n$ by

$$A = \begin{bmatrix} 0 & L - \mu I \\ L - \mu I & 0 \end{bmatrix}.$$

Let A have eigenvalues $\theta_1 \geq \dots \geq \theta_{2n}$. Then $\theta_{2n+1-i} = -\theta_i$ ($1 \leq i \leq 2n$) and $\theta_1 = \mu$ and $\theta_2 = \frac{1}{2}(\mu_n - \mu_2)$. The sets X and Y give rise to a partitioning of A (with rows and columns indexed by Y, \bar{Y}, \bar{X}, X) with quotient matrix

$$B = \begin{bmatrix} 0 & 0 & -\mu & 0 \\ 0 & 0 & -\mu + \mu \frac{|X|}{n-|Y|} & -\mu \frac{|X|}{n-|Y|} \\ -\mu \frac{|Y|}{n-|X|} & -\mu + \mu \frac{|Y|}{n-|X|} & 0 & 0 \\ 0 & -\mu & 0 & 0 \end{bmatrix}.$$

Let B have eigenvalues $\eta_1 \geq \dots \geq \eta_4$. Then $\eta_1 = \theta_1 = \mu$ and $\eta_4 = \theta_{2n} = -\mu$, and $\eta_1 \eta_2 \eta_3 \eta_4 = \det B = \mu^4 \frac{|X||Y|}{(n-|X|)(n-|Y|)} > 0$. Using interlacing we find

$$\mu^2 \frac{|X||Y|}{(n-|X|)(n-|Y|)} = -\eta_2 \eta_3 \leq -\theta_2 \theta_{2n-1} = \left(\frac{1}{2}(\mu_n - \mu_2)\right)^2,$$

which gives the required inequality. \square

One can rewrite Tanner's inequality (applied with $R = X$, $\Gamma(R) = V\Gamma \setminus Y$) in the form $|X||Y|/(n-|X|)(n-|Y|) \leq (\lambda/k)^2$ where $\lambda = \max(\theta_2, -\theta_n)$, and this is slightly weaker than the above, equivalent only when $\theta_n = -\theta_2$.

The vertex sets X and Y with the above property are sometimes called *disconnected vertex sets*. In the complement X and Y become sets such that all edges between X and Y are present. Such a pair is called a *biclique*.

For applications another form is sometimes handy:

Corollary 4.6.2 *Let Γ be a connected graph on n vertices, and let X and Y be disjoint sets of vertices, such that there is no edge between X and Y . Then*

$$\frac{|X||Y|}{n(n-|X|-|Y|)} \leq \frac{(\mu_n - \mu_2)^2}{4\mu_2\mu_n}.$$

Proof. Let K be the constant for which Proposition 4.6.1 says $|X||Y| \leq K(n-|X|)(n-|Y|)$. Then $|X||Y|(1-K) \leq n(n-|X|-|Y|)K$. \square

The above proposition gives bounds on vertex connectivity. For edge connectivity one has

Proposition 4.6.3 (Alon & Milman [6]) *Let A and B be subsets of $V\Gamma$ such that each point of A has distance at least ρ to each point of B . Let F be the set of edges which do not have both ends in A or both in B . Then*

$$|F| \geq \rho^2 \mu_2 \frac{|A||B|}{|A| + |B|}.$$

For $\rho = 1$ this yields:

Corollary 4.6.4 *Let Γ be a graph on n vertices, A a subset of $V\Gamma$, and F the set of edges with one end in A and one end outside A . Then*

$$|F| \geq \mu_2 |A| \left(1 - \frac{|A|}{n}\right).$$

Let χ be the characteristic vector of A . Then equality holds if and only if $\chi - \frac{|A|}{n} \mathbf{1}$ is a Laplace eigenvector with eigenvalue μ_2 .

Proof. Let u_i be an orthonormal system of Laplace eigenvectors, so that $Lu_i = \mu_i u_i$. Take $u_1 = \frac{1}{\sqrt{n}} \mathbf{1}$. Let $\chi = \sum \alpha_i u_i$. Now $|A| = (\chi, \chi) = \sum \alpha_i^2$ and $\alpha_1 = (\chi, u_1) = \frac{1}{\sqrt{n}} |A|$. We find $|F| = \sum_{a \in A, b \notin A, a \sim b} 1 = \sum_{x \sim y} (\chi_x - \chi_y)^2 = \chi^\top L \chi = \sum \alpha_i^2 \mu_i \geq (\sum_{i>1} \alpha_i^2) \mu_2$. \square

This is best possible in many situations.

Example The Hoffman-Singleton graph has Laplace spectrum $0^1 5^{28} 10^{21}$ and we find $|F| \geq |A||B|/10$ and this holds with equality for the 10-40 split into a Petersen subgraph and its complement.

4.6.1 Bandwidth

A direct consequence of Proposition 4.6.1 is an inequality of Helmberg, Mohar, Poljak and Rendl [189], concerning the bandwidth of a graph. A symmetric matrix M is said to have bandwidth w if $(M)_{i,j} = 0$ for all i, j satisfying $|i - j| > w$. The bandwidth $w(\Gamma)$ of a graph Γ is the smallest possible bandwidth for its adjacency matrix (or Laplace matrix). This number (and the vertex order realizing it) is of interest for some combinatorial optimization problems.

Theorem 4.6.5 *Suppose Γ is not edgeless and define $b = \left\lceil n \frac{\mu_2}{\mu_n} \right\rceil$, then*

$$w(\Gamma) \geq \begin{cases} b & \text{if } n - b \text{ is even,} \\ b - 1 & \text{if } n - b \text{ is odd.} \end{cases}$$

Proof. Order the vertices of Γ such that L has bandwidth $w = w(\Gamma)$. If $n - w$ is even, let X be the first $\frac{1}{2}(n - w)$ vertices and let Y be the last $\frac{1}{2}(n - w)$ vertices. Then Proposition 4.6.1 applies and thus we find the first inequality. If $n - w$ is odd, take for X and Y the first and last $\frac{1}{2}(n - w - 1)$ vertices and the second inequality follows. If b and w have different parity, then $w - b \geq 1$ and so the better inequality holds. \square

In case $n - w$ is odd, the bound can be improved a little by applying Proposition 4.6.1 with $|X| = \frac{1}{2}(n - w + 1)$ and $|Y| = \frac{1}{2}(n - w - 1)$. It is clear that the result remains valid if we consider graphs with weighted edges.

4.6.2 Perfect matchings

A more recent application of Proposition 4.6.1 is the following sufficient condition for existence of a perfect matching (a *perfect matching* in a graph is a subset of the edges, such that every vertex of the graph is incident with exactly one edge of the subset).

Theorem 4.6.6 ([53]) *Let Γ be a graph with n vertices, and Laplacian eigenvalues $0 = \mu_1 \leq \mu_2 \leq \dots \leq \mu_n$. If n is even and $\mu_n \leq 2\mu_2$, then Γ has a perfect matching.*

Except for Proposition 4.6.1, we need two more tools. The first one is Tutte's famous characterization of graphs with a perfect matching. The second one is an elementary observation.

Theorem 4.6.7 (Tutte [296]) *A graph $\Gamma = (V, E)$ has no perfect matching if and only if there exists a subset $S \subset V$, such that the subgraph of Γ induced by $V \setminus S$ has more than $|S|$ odd components.*

Lemma 4.6.8 *Let $x_1 \dots x_n$ be n positive integers such that $\sum_{i=1}^n x_i = k \leq 2n - 1$. Then for every integer ℓ , satisfying $0 \leq \ell \leq k$, there exists an $I \subset \{1, \dots, n\}$ such that $\sum_{i \in I} x_i = \ell$.*

Proof. Induction on n . The case $n = 1$ is trivial. If $n \geq 2$, assume $x_1 \geq \dots \geq x_n$. Then $n - 1 \leq k - x_1 \leq 2(n - 1) - 1$ and we apply the induction hypothesis to $\sum_{i=2}^n x_i = k - x_1$ with the same ℓ if $\ell \leq n - 1$, and $\ell - x_1$ otherwise. \square

Proof of Theorem 4.6.6. Assume $\Gamma = (V, E)$ has no perfect matching. By Tutte's theorem there exists a set $S \subset V$ of size s (say), such that the subgraph Γ' of Γ induced by $V \setminus S$ has $q > s$ odd components. But since n is even, $s + q$ is even, hence $q \geq s + 2$.

First assume $n \leq 3s + 3$. Then Γ' has at most $2s + 3$ vertices and at least $s + 2$ components. By Lemma 4.6.8, Γ' and hence Γ , has a pair of disconnected vertex sets X and Y with $|X| = \lfloor \frac{1}{2}(n - s) \rfloor$ and $|Y| = \lceil \frac{1}{2}(n - s) \rceil$. Now Proposition 4.6.1 implies

$$\left(\frac{\mu_n - \mu_2}{\mu_n + \mu_2} \right)^2 \geq \frac{|X| \cdot |Y|}{ns + |X| \cdot |Y|} = \frac{(n - s)^2 - \epsilon}{(n + s)^2 - \epsilon},$$

where $\epsilon = 0$ if $n - s$ is even and $\epsilon = 1$ if $n - s$ is odd. Using $n \geq 2s + 2$ we obtain

$$\frac{\mu_n - \mu_2}{\mu_n + \mu_2} > \frac{n - s - 1}{n + s} \geq \frac{s + 1}{3s + 2} > \frac{1}{3}.$$

Hence $2\mu_2 < \mu_n$.

Next assume $n \geq 3s + 4$. Now Γ' , and hence Γ , has a pair of disconnected vertex sets X and Y with $|X| + |Y| = n - s$ and $\min\{|X|, |Y|\} \geq s + 1$, so $|X| \cdot |Y| \geq (s + 1)(n - 2s - 1) > ns - 2s^2$. Now Proposition 4.6.1 implies

$$\left(\frac{\mu_n - \mu_2}{\mu_n + \mu_2} \right)^2 \geq \frac{|X| \cdot |Y|}{ns + |X| \cdot |Y|} \geq \frac{ns - 2s^2}{2ns - 2s^2} = \frac{1}{2} - \frac{s}{2n - 2s} > \frac{1}{4},$$

by use of $n \geq 3s + 4$. So

$$\frac{\mu_n - \mu_2}{\mu_n + \mu_2} > \frac{1}{2} > \frac{1}{3},$$

hence $2\mu_2 < \mu_n$. \square

The complete bipartite graphs $K_{l,m}$ with $l \leq m$ have Laplacian eigenvalues $\mu_2 = m$ and $\mu_n = n = l + m$. This shows that $2\mu_2$ can get arbitrarily close to μ_n for graphs with n even and no perfect matching.

If the graph is regular, the result can be improved considerably.

Theorem 4.6.9 [53, 89] *A connected k -regular graph on n vertices, where n is even, with (ordinary) eigenvalues $k = \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$, which satisfies*

$$\lambda_3 \leq \begin{cases} k - 1 + \frac{3}{k+1} & \text{if } k \text{ is even,} \\ k - 1 + \frac{4}{k+2} & \text{if } k \text{ is odd,} \end{cases}$$

has a perfect matching.

Proof. Let $\Gamma = (V, E)$ be a k -regular graph with $n = |V|$ even and no perfect matching. By Tutte's Theorem 4.6.7 there exists a set $S \subset V$ of size s such that $V \setminus S$ induces a subgraph with $q \geq s + 2$ odd components $\Gamma_1, \Gamma_2, \dots, \Gamma_q$ (say). Let t_i denote the number of edges in Γ between S and Γ_i , and let n_i be the number of vertices of Γ_i . Then clearly $\sum_{i=1}^q t_i \leq ks$, $s \geq 1$, and $t_i \geq 1$ (since Γ is connected). Hence $t_i < k$ and $n_i > 1$ for at least three values of i , say $i = 1, 2$ and 3 . Let ℓ_i denote the largest eigenvalue of Γ_i , and assume $\ell_1 \geq \ell_2 \geq \ell_3$. Then eigenvalue interlacing applied to the subgraph induced by the union of Γ_1, Γ_2 and Γ_3 gives $\ell_i \leq \lambda_i$ for $i = 1, 2, 3$.

Consider Γ_3 with n_3 vertices and e_3 edges (say). Then $2e_3 = kn_3 - t_3 \leq n_3(n_3 - 1)$. We saw that $t_3 < k$ and $n_3 > 1$, hence $k < n_3$. Moreover, the average degree \bar{d}_3 of Γ_3 equals $2e_3/n_3 = k - t_3/n_3$. Because n_3 is odd and $kn_3 - t_3$ is even, k and t_3 have the same parity, therefore $t_3 < k$ implies $t_3 \leq k - 2$. Also $k < n_3$ implies $k \leq n_3 - 1$ if k is even, and $k \leq n_3 - 2$ if k is odd. Hence

$$\bar{d}_3 \geq \begin{cases} k - \frac{k-2}{k+1} & \text{if } k \text{ is even,} \\ k - \frac{k-2}{k+2} & \text{if } k \text{ is odd.} \end{cases}$$

Note that $t_3 < n_3$ implies that Γ_3 cannot be regular. Next we use the fact that the largest adjacency eigenvalue of a graph is bounded from below by the average degree with equality if and only if the graph is regular (Proposition 3.1.2). Thus $\bar{d}_3 < \ell_3$. We saw that $\ell_3 \leq \lambda_3$, which finishes the proof. \square

From the above it is clear that n even and $\lambda_2 \leq k - 1$ implies existence of a perfect matching. In terms of the Laplacian matrix this translates into:

Corollary 4.6.10 *A regular graph with an even number of vertices and algebraic connectivity at least 1 has a perfect matching.*

But we can say more. The Laplacian matrix of a disjoint union of $n/2$ edges has eigenvalues 0 and 2. This implies that deletion of the edges of a perfect matching of a graph Γ reduces the eigenvalues of the Laplacian matrix of Γ by at most 2 (by the Courant-Weyl inequalities 2.8.1). Hence:

Corollary 4.6.11 *A regular graph with an even number of vertices and algebraic connectivity μ_2 has at least $\lfloor (\mu_2 + 1)/2 \rfloor$ disjoint perfect matchings.*

Cioabă, Gregory and Haemers [90] have improved the sufficient condition for a perfect matching from Theorem 4.6.9 to $\lambda_3 < \vartheta_k$ where $\vartheta_3 = 2.85577\dots$ (the largest root of $x^3 - x^2 - 6x + 2$), $\vartheta_k = (k - 2 + \sqrt{k^2 + 12})/2$ if $k \geq 4$ and even, and $\vartheta_k = (k - 3 + \sqrt{(k + 1)^2 + 16})/2$ if $k \geq 5$ and odd. They also prove that this bound is best possible by giving examples of k -regular graphs with n even, and $\lambda_3 = \vartheta_k$ that have no perfect matching. The example for $k = 3$ is presented in Figure 4.1.




Figure 4.1: A 3-regular graph with no perfect matching

4.7 Block Designs

In case we have a non-symmetric matrix N (say) we can still use interlacing by considering the matrix

$$A = \begin{bmatrix} 0 & N \\ N^\top & 0 \end{bmatrix}$$

like we did in the previous section. Then we find results in terms of the eigenvalues of A , which now satisfy $\theta_i = -\theta_{n-i+1}$ for $i = 1, \dots, n$. The positive eigenvalues of A are the singular values of N , they are also the square roots of the non-zero eigenvalues of NN^\top (and of $N^\top N$).

Suppose N is the 0-1 incidence matrix of an incidence structure (P, B) with point set P (rows) and block set B (columns). Then we consider the so-called *incidence graph* Γ of (P, B) , which is the bipartite graph with vertex set $P \cup B$, where two vertices are adjacent if they correspond to an incident point-block pair. An edge of Γ is called a *flag* of (P, B) .

An incidence structure (P, B) is called a t -(v, k, λ) design if $|P| = v$, all blocks are incident with k points, and for every t -set of points there are exactly λ blocks incident with all t points. For example (P, B) is a 1-(v, k, r) design precisely when N has constant column sums k (i.e. $N^\top \mathbf{1} = k\mathbf{1}$), and constant row sums r (i.e. $N\mathbf{1} = r\mathbf{1}$), in other words Γ is biregular with degrees k and r . Moreover, (P, B) is a 2-(v, k, λ) design if and only if $N^\top \mathbf{1} = k\mathbf{1}$ and $NN^\top = \lambda J + (r - \lambda)I$. Note that for $t \geq 1$, a t -design is also a $(t-1)$ -design. In particular a 2-(v, k, λ) design is also a 1-(v, k, r) design with $r = \lambda(v-1)/(k-1)$.

Theorem 4.7.1 *Let (P, B) be a 1-(v, k, r) design with b blocks and let (P', B') be a substructure with m' flags. Define $b = |B|$, $v' = |P'|$ and $b' = |B'|$. Then*

$$(m' \frac{v}{v'} - b'k)(m' \frac{b}{b'} - v'r) \leq \theta_2^2(v - v')(b - b').$$

Equality implies that all four substructures induced by P' or $V \setminus V'$ and B' or $B \setminus B'$ form a 1-design (possibly degenerate).

Proof. We apply Corollary 2.5.4. The substructure (P', B') gives rise to a partition of A with the following quotient matrix

$$B = \begin{bmatrix} 0 & 0 & \frac{m'}{v'} & r - \frac{m'}{v'} \\ 0 & 0 & \frac{b'k - m'}{v - v'} & r - \frac{b'k - m'}{v - v'} \\ \frac{m'}{b'} & k - \frac{m'}{b'} & 0 & 0 \\ \frac{v'r - m'}{b - b'} & k - \frac{v'r - m'}{b - b'} & 0 & 0 \end{bmatrix}.$$

We easily have $\theta_1 = -\theta_n = \eta_1 = -\eta_4 = \sqrt{rk}$ and

$$\det(B) = rk \left(\frac{m' \frac{v}{v'} - b'k}{v - v'} \right) \left(\frac{m' \frac{b}{b'} - v'r}{b - b'} \right).$$

Interlacing gives

$$\frac{\det(B)}{rk} = -\eta_2\eta_3 \leq -\theta_2\theta_{n-1} = \theta_2^2,$$

which proves the first statement. If equality holds then $\theta_1 = \eta_1$, $\theta_2 = \eta_2$, $\theta_{n-1} = \eta_3$ and $\theta_n = \eta_4$, so we have tight interlacing, which implies the second statement. \square

The above result becomes especially useful if we can express θ_2 in terms of the design parameters. For instance if (P, B) is a $2-(v, k, \lambda)$ design, then $\theta_2^2 = r - \lambda = \lambda \frac{v-k}{k-1}$ (see exercises) and if (P, B) is a generalized quadrangle of order (s, t) , then $\lambda_2^2 = s + t$ (see §8.6). Let us consider two special cases. (A 2-design (P, B) with $|P| = |B|$ is called *symmetric*.)

Corollary 4.7.2 *If a symmetric $2-(v, k, \lambda)$ design (P, B) has a symmetric $2-(v', k', \lambda')$ subdesign (P', B') (possibly degenerate) then*

$$(k'v - kv')^2 \leq (k - \lambda)(v - v')^2.$$

If equality holds, then the subdesign $(P', B \setminus B')$ is a $2-(v', v'(k - k')/(v - v'), \lambda - \lambda')$ design (possibly degenerate).

Proof. In Theorem 4.7.1 take $b = v$, $r = k$, $b' = v'$, $m' = v'k'$ and $\theta_2^2 = k - \lambda$. \square

Corollary 4.7.3 *Let X be a subset of the points and let Y be a subset of the blocks of a $2-(v, k, \lambda)$ design (P, B) , such that no point of X is incident with a block of Y , then*

$$kr|X||Y| \leq (r - \lambda)(v - |X|)(b - |Y|).$$

If equality holds then the substructure $(X, B') = (X, B \setminus Y)$ is a 2-design.

Proof. Take $m' = 0$, $v' = |X|$, $b' = |Y|$ and $\theta_2^2 = r - \lambda$. Now 4.7.1 gives the inequality and that (X, B') is a 1-design. But then (X, B') is a 2-design, because (P, B) is. \square

An example of a subdesign of a symmetric design is the incidence structure formed by the absolute point and lines of a polarity in a projective plane of order q . This gives a (degenerate) $2-(v', 1, 0)$ design in a $2-(q^2 + q + 1, q + 1, 1)$ design. The bound gives $v' \leq q\sqrt{q} + 1$. (See also the following section.) The $2-(q\sqrt{q} + 1, q + 1, 1)$ design, which is obtained in case of equality is called a *unital*. Other examples of symmetric designs that meet the bound can be found in Haemers & Shrikhande [182] or Jungnickel [210]. Wilbrink used Theorem 4.7.1 to shorten the proof of Feit's result on the number of points and blocks fixed by an automorphism group of a symmetric design (see [61]). The inequality of the second corollary is for example tight for hyperovals and (more generally) maximal arcs in finite projective planes.

4.8 Polarities

A *projective plane* is a point-line geometry such that any two points are on a unique line, and any two lines meet in a unique point. It is said to be of order q when all lines have $q + 1$ points and all points are on $q + 1$ lines. A projective plane of order q has $q^2 + q + 1$ points and as many lines.

A *polarity* of a point-block incidence structure is a map of order 2 interchanging points and blocks and preserving incidence. An *absolute point* is a point incident with its image under the polarity.

Suppose we have a projective plane of order q with a polarity σ . The polarity enables us to write the point-line incidence matrix N as a symmetric matrix, and then the number of absolute points is $\text{tr } N$. By definition we have $N^2 = NN^\top = J + qI$, which has one eigenvalue equal to $(q + 1)^2$ and all other eigenvalues equal to q . That means that N has spectrum $(q + 1)^1, \sqrt{q}^m, -\sqrt{q}^n$, for certain integers m, n , where this time exponents indicate multiplicities. The number of absolute points equals $a = q + 1 + (m - n)\sqrt{q}$. It follows that if q is not a square then $m = n$ and there are precisely $q + 1$ absolute points. If q is a square, and p is a prime dividing q , then $a \equiv 1 \pmod{p}$ so that a is nonzero.

(This is false in the infinite case: the polarity sending the point (p, q, r) to the line $pX + qY + rZ = 0$ has no absolute points over \mathbb{R} .)

With slightly more effort one finds bounds for the number of absolute points:

Proposition 4.8.1 *A polarity of a projective plane of order q has at least $q + 1$ and at most $q\sqrt{q} + 1$ absolute points.*

Proof. Suppose z is a non-absolute point. Now σ induces a map τ on the line z^σ defined for $y \in z^\sigma$ by: y^τ is the common point of y^σ and z^σ . Now $\tau^2 = 1$, and $y^\tau = y$ precisely when y is absolute. This shows that the number of absolute points on a non-absolute line is $q + 1 \pmod{2}$.

Now if q is odd, then take an absolute point x . This observation says that each line on x different from x^σ contains another absolute point, for a total of at least $q + 1$. On the other hand, if q is even, then each non-absolute line contains an absolute point, so that $q^2 + q + 1 - a \leq aq$ and $a \geq q + 1$.

For the upper bound, use interlacing: partition the matrix N into absolute / non-absolute points/lines and find the matrix of average row sums

$\begin{bmatrix} 1 & q \\ \frac{aq}{v-a} & q + 1 - \frac{aq}{v-a} \end{bmatrix}$ where $v = q^2 + q + 1$, with eigenvalues $q + 1$ and $1 - \frac{aq}{v-a}$. Now interlacing yields $1 - \frac{aq}{v-a} \geq -\sqrt{q}$, that is, $a \leq q\sqrt{q} + 1$, just like we found in the previous section. \square

The essential part of the proof of the lower bound was to show that there is at least one absolute point, and this used an eigenvalue argument.

4.9 Exercises

Exercise 1 Deduce Proposition 4.2.3(iii) (the part that says $(m - 1)\theta_{t+1} + \theta_{n-t(m-1)} \geq 0$) from Theorem 4.1.3.

Exercise 2 An (ℓ, m) -biqule in a graph Γ is a complete bipartite subgraph $K_{\ell, m}$ of Γ (not necessarily induced). Let $0 = \mu_1 \leq \dots \leq \mu_n$ be the Laplace

eigenvalues of Γ . Show that $\ell m / (n - \ell)(n - m) \leq ((\mu_n - \mu_2) / (2n - \mu_2 - \mu_n))^2$ if Γ is non-complete and contains an (ℓ, m) -biclique.

Exercise 3 Let A be the incidence graph of a 2 -(v, k, λ) design with b blocks and r blocks incident with each point. Express the spectrum of A in the design parameters v, k, λ, b and r .

Exercise 4 Let (P, B) is a 2 -(v, k, λ) design, and suppose that some block is repeated ℓ times (i.e. ℓ blocks are incident with exactly the same set of k points). Prove that $b \geq \ell v$ (this is Mann's inequality).

Chapter 5

Trees

Trees have a simpler structure than general graphs, and we can prove stronger results. For example, interlacing tells us that the multiplicity of an eigenvalue decreases by at most one when a vertex is removed. For trees Godsil's Lemma gives the same conclusion also when a path is removed.

5.1 Characteristic polynomials of trees

For a graph Γ with adjacency matrix A , let $\phi_\Gamma(t) := \det(tI - A)$ be its characteristic polynomial.

Note that since the characteristic polynomial of the disjoint union of two graphs is the product of their characteristic polynomials, results for trees immediately yield results for forests as well.

It will be useful to agree that $\phi_{T \setminus x, y} = 0$ if $x = y$.

Proposition 5.1.1 *Let T be a tree, and for $x, y \in T$, let P_{xy} be the unique path joining x and y in T .*

(i) *Let $e = xy$ be an edge in T that separates T into two subtrees A and B , with $x \in A$ and $y \in B$. Then*

$$\phi_T = \phi_A \phi_B - \phi_{A \setminus x} \phi_{B \setminus y}.$$

(ii) *Let x be a vertex of T . Then*

$$\phi_T(t) = t\phi_{T \setminus x}(t) - \sum_{y \sim x} \phi_{T \setminus \{x, y\}}(t).$$

(iii) *Let x be a vertex of T . Then*

$$\phi_{T \setminus x}(t)\phi_T(s) - \phi_{T \setminus x}(s)\phi_T(t) = (s - t) \sum_{y \in T} \phi_{T \setminus P_{xy}}(s)\phi_{T \setminus P_{xy}}(t).$$

(iv) *Let x be a vertex of T . Then*

$$\phi_{T \setminus x} \phi'_T - \phi'_{T \setminus x} \phi_T = \sum_{y \in T} \phi_{T \setminus P_{xy}}^2.$$

(v) Let x, y be vertices of T . Then

$$\phi_{T \setminus x} \phi_{T \setminus y} - \phi_{T \setminus x, y} \phi_T = \phi_{T \setminus P_{xy}}^2.$$

(vi) Let x, y, z be vertices of T where $z \in P_{xy}$. Then

$$\phi_{T \setminus x, y, z} \phi_T = \phi_{T \setminus x} \phi_{T \setminus y, z} - \phi_{T \setminus z} \phi_{T \setminus x, y} + \phi_{T \setminus y} \phi_{T \setminus x, z}.$$

(vii) We have $\phi'_T = \sum_{x \in T} \phi_{T \setminus x}$.

(viii) Let T have n vertices and c_m matchings of size m . Then

$$\phi_T(t) = \sum_m (-1)^m c_m t^{n-2m}.$$

Proof. Part (i) follows by expansion of the defining determinant. It can also be phrased as $\phi_T = \phi_{T \setminus e} - \phi_{T \setminus \{x, y\}}$. Part (ii) follows by applying (i) to all edges on x . Note that $\phi_{\{x\}}(t) = t$. Part (iii) follows from (ii) by induction on the size of T : expand in the LHS $\phi_T(s)$ and $\phi_T(t)$ using (ii), and then use induction. Part (iv) is immediate from (iii). Part (vii) follows by taking the derivative of the defining determinant. Part (viii) is a reformulation of the description in §1.2.1. Note that the only directed cycles in a tree are those of length 2. Part (v) is true if $T = P_{xy}$, and the general case follows from part (vi) and induction: the statement remains true when a subtree S is attached via an edge e at a vertex $z \in P_{xy}$. Finally, part (vi) follows from: if $\Gamma \setminus z = A + B$, then $\phi_\Gamma = \phi_{A \cup z} \phi_B + \phi_A \phi_{B \cup z} - \phi_A \phi_{\{z\}} \phi_B$, where of course $\phi_{\{z\}}(t) = t$. \square

Theorem 5.1.2 ('Godsil's Lemma', [157]) *Let T be a tree and θ an eigenvalue of multiplicity $m > 1$. Let P be a path in T . Then θ is eigenvalue of $T \setminus P$ with multiplicity at least $m - 1$.*

Proof. By parts (iv) and (vii) of the above Proposition we have

$$\phi'_T(t)^2 - \phi''_T(t) \phi_T(t) = \sum_{x, y \in T} \phi_{T \setminus P_{xy}}(t)^2.$$

Now θ is a root of multiplicity at least $2m - 2$ of the left hand side, and hence also of each of the terms on the right hand side. \square

As an application of Godsil's Lemma, consider a tree T with e distinct eigenvalues and maximum possible diameter $e - 1$. Let P be a path of length $e - 1$ (that is, with e vertices) in T . Then $T \setminus P$ has a spectrum that is independent of the choice of P : for each eigenvalue θ with multiplicity m of T , the forest $T \setminus P$ has eigenvalue θ with multiplicity $m - 1$ (and it has no other eigenvalues).

In particular, all eigenvalues of a path have multiplicity 1.

Note that going from T to $T \setminus x$ changes multiplicities by at most 1: they go up or down by at most one. Godsil's Lemma is one-sided: going from T to $T \setminus P$, the multiplicities go down by at most one, but they may well go up by more. For example, if one joins the centers x, y of two copies of $K_{1, m}$ by an edge, one obtains a tree T that has 0 as an eigenvalue of multiplicity $2m - 2$. For $P = xy$ the forest $T \setminus P$ has 0 with multiplicity $2m$.

5.2 Eigenvectors and multiplicities

For trees we have rather precise information about eigenvectors and eigenvalue multiplicities (Fiedler [142]).

Lemma 5.2.1 *Let T be a tree with eigenvalue θ , and let $Z = Z_T(\theta)$ be the set of vertices in T where all θ -eigenvectors vanish. If for some vertex $t \in T$ some component S of $T \setminus t$ has eigenvalue θ (in particular, if some θ -eigenvector of T vanishes at t), then $Z \neq \emptyset$.*

Proof. Consider proper subtrees S of T with eigenvalue θ and with a single edge st joining some vertex $s \in S$ with some vertex $t \in T \setminus S$, and pick a minimal one. If $|S| = 1$, then $\theta = 0$, and $t \in Z$. Assume $|S| > 1$. If a θ -eigenvector u of S is the restriction to S of a θ -eigenvector v of T , then v vanishes in t . So, if some θ -eigenvector v of T does not vanish at t then u and $v|_S$ are not dependent, and some linear combination vanishes in s and is a θ -eigenvector of $S \setminus s$, contradicting minimality of S . This shows that $t \in Z$. \square

Note that it is not true that the hypothesis of the lemma implies that $t \in Z$. For example, consider the tree T of type D_6 given by $1 \sim 2 \sim 3 \sim 4 \sim 5, 6$. It has $Z(0) = \{2, 4\}$, and the component $S = \{4, 5, 6\}$ of $T \setminus 3$ has eigenvalue 0, but $3 \notin Z(0)$.

Proposition 5.2.2 *Consider a tree T with eigenvalue θ , and let $Z = Z(\theta)$ be the set of vertices in T where all θ -eigenvectors vanish. Let $Z_0 = Z_0(\theta)$ be the set of vertices in Z that have a neighbor in $T \setminus Z$.*

(i) *Let S be a connected component of $T \setminus Z$. Then S has eigenvalue θ with multiplicity 1. If u is a θ -eigenvector of S , then u is nowhere zero.*

(ii) *Let $T \setminus Z$ have c connected components, and let $d = |Z_0|$. Then θ has multiplicity $c - d$.*

The components of $T \setminus Z(\theta)$ are called the *eigenvalue components* of T for θ .

Proof. (i) Suppose θ is eigenvalue of T with multiplicity greater than 1. Then some eigenvector has a zero coordinate and hence induces a θ -eigenvector on a proper subtree.

By Lemma 5.2.1, Z is nonempty.

If S is a connected component of $T \setminus Z$ then it has eigenvalue θ (otherwise $S \subset Z$, contradiction). Apply Lemma 5.2.1 to S instead of T to find that if some θ -eigenvector of S vanishes on a point of S , then there is a point $s \in S$ where all of its θ -eigenvectors vanish. But the restriction to S of a θ -eigenvector of T is a θ -eigenvector of S , so $s \in Z$, contradiction.

(ii) Each point of Z_0 imposes a linear condition, and since T is a tree, these conditions are independent. \square

We see that if the multiplicity of θ is not 1, then Z contains a vertex of degree at least three. In particular, $Z \neq \emptyset$, and hence $Z_0 \neq \emptyset$. Deleting a vertex in Z_0 from T increases the multiplicity of θ .

As an application we see that all eigenvalues of a path have multiplicity 1.

5.3 Sign patterns of eigenvectors of graphs

For a path, the i -th largest eigenvalue has multiplicity 1 and an eigenvector with $i - 1$ sign changes, that is, i areas of constant sign. It is possible to generalize this observation to more general graphs.

Given a real vector u , let the *support* $\text{supp } u$ be the set $\{i | u_i \neq 0\}$. For $*$ one of $<, >, \leq, \geq$ we also write $\text{supp}^* u$ for $\{i | u_i * 0\}$. Let $N(u)$ (resp. $N^*(u)$) be the number of connected components C of the subgraph induced by $\text{supp } u$ (resp. $\text{supp}^* u$) such that u does not vanish identically on C . Let $N_\theta(u)$ be the number of connected components C of the subgraph induced by $\text{supp } u$ such that u (does not vanish identically on C and) induces an eigenvector with eigenvalue θ on C .

Proposition 5.3.1 *Let Γ be a graph with eigenvalues $\theta_1 \geq \dots \geq \theta_n$, and let $\theta = \theta_j = \theta_{j+m-1}$ be an eigenvalue with multiplicity m . Let u be a vector with $Au \geq \theta u$. Let Δ be the subgraph of Γ induced by $\text{supp } u$, with eigenvalues $\eta_1 \geq \dots \geq \eta_t$. Then*

(i)

$$N^>(u) + N^<(u) \leq \#\{i \mid \eta_i \geq \theta\} \leq j + m - 1$$

(ii)

$$N^>(u) + N^<(u) - N_\theta(u) \leq \#\{i \mid \eta_i > \theta\} \leq j - 1$$

(iii) if Γ has c connected components, then

$$N^\geq(u) + N^\leq(u) \leq j + c - 1.$$

Proof. For a subset S of the vertex set of Γ , let I_S be the diagonal matrix with ones on the positions indexed by elements of S and zeros elsewhere.

Let C run through the connected components of $\text{supp}^> u$ and $\text{supp}^< u$ (resp. $\text{supp}^\geq u$ and $\text{supp}^\leq u$). Put $u_C = I_C u$. Then the space $U := \langle u_C | C \rangle$ has dimension $N^>(u) + N^<(u)$ (resp. $N^\geq(u) + N^\leq(u)$).

Let A be the adjacency matrix of Δ (resp. Γ). Define a real symmetric matrix B by $B_{CD} = u_C^\top (A - \theta I) u_D$. Then B has nonnegative row sums and nonpositive off-diagonal entries, so B is positive semidefinite. It follows that for $y \in U$ we have $y^\top (A - \theta I) y \geq 0$. This means that U intersects the space spanned by the eigenvectors of $A - \theta I$ with negative eigenvalue in 0.

For (i) $N^>(u) + N^<(u) \leq \#\{i \mid \eta_i \geq \theta\}$ follows.

The vectors $y \in U$ with $y^\top (A - \theta I) y = 0$ correspond to eigenvectors with eigenvalue 0 of B , and by Lemma 2.10.1 there are at most $N_\theta(u)$ (resp. c) independent such. This proves (ii) (resp. (iii)). \square

Remark For $j = 1$ the results follow from Perron-Frobenius. (If Γ is connected, then the eigenvector for θ_1 is nowhere zero and has constant sign.)

Examples a) Let Γ be connected and bipartite, and let θ be the smallest eigenvalue of Γ . The corresponding eigenvector u has different signs on the two sides of the bipartition, so $\text{supp}^> u$ and $\text{supp}^< u$ are the two sides of the bipartition, $N^>(u) + N^<(u) = n$ and $N(u) = 1$. We have equality in (i)–(iii).

b) Let Γ be the star $K_{1,s}$. The spectrum is $\sqrt{s}^1, 0^{s-1}, (-\sqrt{s})^1$. Let u be an eigenvector with eigenvalue $\theta = 0$ that has t nonzero coordinates. (Then $2 \leq t \leq s$.) Now $N^>(u) + N^<(u) = N(u) = t$ and $N^\geq(u) + N^\leq(u) = 2$, and for $t = s$ equality holds in (i)–(iii).

c) Let Γ be the Petersen graph. It has spectrum $3^1, 1^5, (-2)^4$. Let u be an eigenvector with eigenvalue $\theta = 1$ that vanishes on 4 points, so that $\text{supp } u$ induces $3K_2$ with spectrum $1^3, (-1)^3$. We find $N^>(u) + N^<(u) = N(u) = 3$ and $N^{\geq}(u) + N^{\leq}(u) = 2$, again equality in (i)–(iii).

d) Let Γ be the path P_n . The eigenvalues are $\theta_k = 2 \cos(k\pi/(n+1))$ for $k = 1, \dots, n$. The eigenvector u corresponding to θ_k has $k-1$ sign changes, so that $N^>(u) + N^<(u) = k$. If $\gcd(k, n+1) = 1$ then u has no zero entries, so that $N(u) = 1$. Now we have equality in (i)–(iii). If $\gcd(k, n+1) = r$, then u has $r-1$ zero entries, so that $N(u) = r$. Also, the eigenvalue θ_k is the k/r -th of each component of $\text{supp } u$, so that $\#\{i \mid \eta_i \geq \theta\} = k$ and $\#\{i \mid \eta_i > \theta\} = k-r$, with equality in (i) and the first inequality of (ii).

Remark It is not true that $N(u) \leq m$ if m is the multiplicity of θ for Γ . For example, in case b) above we have $N(u) = s$ and $m = s-1$. (And in case c) the opposite happens: $N(u) = 3$ and $m = 5$.)

5.4 Sign patterns of eigenvectors of trees

Proposition 5.4.1 *Let T be a tree with eigenvalue θ , and put $Z = Z(\theta)$. Let $T \setminus Z$ have eigenvalues $\eta_1 \geq \dots \geq \eta_m$. Let $g = \#\{i \mid \eta_i \geq \theta\}$ and $h = \#\{i \mid \eta_i > \theta\}$. Let u be a θ -eigenvector of T . Then $N^>(u) + N^<(u) = g$ and $N^>(u) + N^<(u) - N(u) = h$.*


Proof. Since $N(\cdot)$ and g and h are additive over connected components, we may assume that Z is empty. Now by Proposition 5.2.2(i), θ has multiplicity 1 and u is nowhere 0. Let T have n vertices, and let there be p edges xy with $u_x u_y > 0$ and q edges xy with $u_x u_y < 0$. Then $p + q = n - 1$. Since T is bipartite, also $-\theta$ is an eigenvalue, and an eigenvector v for $-\theta$ is obtained by switching the sign of u on one bipartite class. By Proposition 5.3.1 we have $q = N^>(u) + N^<(u) - 1 \leq h$ and $p = N^>(v) + N^<(v) - 1 \leq n - h - 1$, that is $q \geq h$, and hence equality holds everywhere. \square

Let a *sign change* for an eigenvector u of T be an edge $e = xy$ such that $u_x u_y < 0$.

Proposition 5.4.2 *Let T be a tree with j -th eigenvalue θ . If u is an eigenvector for θ with s sign changes, and $d = |Z_0(\theta)|$, then $d + s \leq j - 1$*

Proof. Let $T \setminus Z$ have c connected components, and let u be identically zero on c_0 of these. Then $s + c - c_0 = N^>(u) + N^<(u)$. Let $\theta = \theta_j = \theta_{j+m-1}$, where $m = c - d$ is the multiplicity of θ . By Proposition 5.3.1(i) we have $s + c - c_0 \leq j + m - 1$, that is, $d + s - c_0 \leq j - 1$. But we can make c_0 zero by adding a small multiple of some eigenvector that is nonzero on all of $T \setminus Z$. \square

Example For $T = E_6$ all eigenvalues have multiplicity 1, and $N^>(u) + N^<(u)$ takes the values 1, 2, 3, 4, 4, 6 for the six eigenvectors u . The sign patterns are:





We see that a small perturbation that would make u nonzero everywhere would give the two zeros in the second eigenvector the same sign, but the two zeros in the fifth eigenvector different sign (because $\theta_2 > 0$ and $\theta_5 < 0$) and for the perturbed vector u' we would find 0, 1, 2, 3, 4, 5 sign changes.

5.5 The spectral center of a tree

There are various combinatorial concepts ‘center’ for trees. One has the center/bicenter and the centroid/bicentroid. Here we define a concept of center using spectral methods. Closely related results can be found in Neumaier [251].

Proposition 5.5.1 *Let T be a tree (with at least two vertices) with second largest eigenvalue λ . Then there is a unique minimal subtree Y of T such that no connected component of $T \setminus Y$ has largest eigenvalue larger than λ . If $Z(\lambda) \neq \emptyset$ (and in particular if λ has multiplicity larger than 1) then $Y = Z_0(\lambda)$ and $|Y| = 1$. Otherwise $|Y| = 2$, and Y contains the endpoints of the edge on which the unique λ -eigenvector changes sign. In this latter case all connected components of $T \setminus Y$ have largest eigenvalue strictly smaller than λ .*

We call the set Y the *spectral center* of T .

Proof. If for some vertex y all connected components of $T \setminus y$ have largest eigenvalue at most λ , then pick $Y = \{y\}$. Otherwise, for each vertex y of T there is a unique neighbor y' in the unique component of $T \setminus y$ that has largest eigenvalue more than λ . Since T is finite, we must have $y'' = y$ for some vertex y . Now pick $Y = \{y, y'\}$. Clearly Y has the stated property and is minimal.

If $Z = \emptyset$ then λ has multiplicity 1 and by Proposition 5.4.2 there is a unique edge $e = pq$ such that the unique λ -eigenvector has different signs on p and q , and both components of $T \setminus e$ have largest eigenvalue strictly larger than λ , so that Y must contain both endpoints of e .

If $Z \neq \emptyset$, then all eigenvalue components for λ have eigenvalue λ , and any strictly larger subgraph has a strictly larger eigenvalue, so Y must contain Z_0 . By Proposition 5.4.2 we have $|Z_0| = 1$, say $Z_0 = \{y\}$. If Y is not equal to $\{y\}$, then Y also contains y' . This proves uniqueness.

Suppose that $Z_0 = \{y\}$ and $T \setminus y$ has a component with eigenvalue larger than λ . Let u be a vector that is 0 on y , and induces an eigenvector with the largest eigenvalue on each component of $T \setminus y$. Let c be the number of connected components of $T \setminus Z$. Proposition 5.3.1(i) now gives $c + 1 \leq 2 + (c - 1) - 1$, a contradiction. This shows that $|Y| = 1$ when Z is nonempty.

Finally, suppose that $Y = \{y, y'\}$ and that $T \setminus Y$ has largest eigenvalue λ . By Lemma 5.2.1 $Z \neq \emptyset$, contradiction. \square

Example If T is the path P_n with n vertices, then $\lambda = 2 \cos 2\pi/(n+1)$. If $n = 2m+1$ is odd, then Y consists of the middle vertex, and $T \setminus Y$ is the union of two paths P_m , with largest eigenvalue $\lambda = 2 \cos \pi/(m+1)$. If $n = 2m$ is even, then Y consists of the middle two vertices, and $T \setminus Y$ is the union of two paths P_{m-1} , with largest eigenvalue $2 \cos \pi/m < \lambda$.

5.6 Integral trees

An *integral tree* is a tree with only integral eigenvalues. Such trees are rare. A list of all integral trees on at most 50 vertices can be found in [46].

A funny result is

Proposition 5.6.1 (Watanabe [304]) *An integral tree cannot have a perfect matching, that is, must have an eigenvalue 0, unless it is K_2 .*

Proof. The constant term of the characteristic polynomial of a tree is, up to sign, the number of perfect matchings. It is also the product of all eigenvalues. If it is nonzero, then it is 1, since the union of two distinct perfect matchings contains a cycle. But then all eigenvalues are ± 1 and P_3 is not an induced subgraph, so we have K_2 . \square

This result can be extended a little. Let $SK_{1,m}$ be the tree on $2m + 1$ vertices obtained by subdividing all edges of $K_{1,m}$. The spectrum is $\pm\sqrt{m+1}$ $(\pm 1)^{m-1}$ 0.

Proposition 5.6.2 (Brouwer [46]) *If an integral tree has eigenvalue 0 with multiplicity 1, then it is $SK_{1,m}$, where $m = t^2 - 1$ for some integer $t \geq 1$.* \square

For a long time it has been an open question whether there exist integral trees of arbitrarily large diameter. Recently, this was settled in the affirmative by Csikvári. The construction is as follows. Define trees $T'(r_1, \dots, r_m)$ by induction: $T'()$ is the tree with a single vertex x_0 . $T'(r_1, \dots, r_m)$ is the tree obtained from $T'(r_1, \dots, r_{m-1})$ by adding r_m pendant edges to each vertex u with $d(u, x_0) = m - 1 \pmod{2}$. The diameter of this tree is $2m$ (assuming $r_1 > 1$) and it has $2m + 1$ distinct eigenvalues:

Proposition 5.6.3 (Csikvári [102]) *The tree $T'(r_1, \dots, r_m)$ has eigenvalues 0 and $\pm\sqrt{s_i}$ ($1 \leq i \leq m$), where $s_i = r_i + \dots + r_m$.*


Now all trees $T'(n_1^2 - n_2^2, \dots, n_{m-1}^2 - n_m^2, n_m^2)$ are integral of diameter $2m$ when $n_1 > n_2 > \dots > n_m$.

A short proof can be given using the following observation. If A and B are trees with fixed vertices x and y , respectively, then let $A \sim mB$ be the tree constructed on the union of A and m copies of B , where x is joined to the m copies of y . Now Proposition 5.1.1(i) and induction immediately yields that $T = A \sim mB$ has characteristic polynomial $\phi_T = \phi_B^{m-1}(\phi_A \phi_B - m\phi_{A \setminus x} \phi_{B \setminus y})$, where the last factor is symmetric in A and B .

Proof. Induction on m . The statement holds for $m \leq 1$. With $A = T'(r_3, \dots)$ and $B = T'(r_2, r_3, \dots)$ we have $T'(r_1, r_2, r_3, \dots) = A \sim r_1 B$ and $T'(r_1 + r_2, r_3, \dots) = B \sim r_1 A$. \square

5.7 Exercises

Exercise 1 Show that there are 6 integral trees on at most ten vertices, namely (i) K_1 , (ii) K_2 , (iii) $K_{1,4} = \hat{D}_4$, (iv) \hat{D}_5 , (v) \hat{E}_6 , (vi) $K_{1,9}$. (For notation, cf. §3.1.1.)



An integral tree on 31 vertices.
What is the spectrum?

Exercise 2 Show that the only trees that have integral Laplace spectrum are the stars $K_{1,m}$.

Chapter 6

Groups and graphs

6.1 $\Gamma(G, H, S)$

Let G be a finite group, and H a subgroup, and S a subset of G . We can define a graph $\Gamma(G, H, S)$ by taking as vertices the cosets gH ($g \in G$), and calling g_1H and g_2H adjacent when $Hg_2^{-1}g_1H \subseteq HSH$. The group G acts as a group of automorphisms on $\Gamma(G, H, S)$ via left multiplication, and this action is transitive. The stabilizer of the vertex H is the subgroup H .

A graph $\Gamma(G, H, S)$ with $H = 1$ is called a *Cayley graph*.

Conversely, let Γ be a graph with transitive group of automorphisms G . Let x be a vertex of Γ , and let $H := G_x$ be the stabilizer of x in G . Now Γ can be identified with $\Gamma(G, H, S)$, where $S = \{g \in G \mid x \sim gx\}$.

If Γ is moreover edge-transitive, then S can be chosen to have cardinality 1.

Instead of representing each vertex as a coset, one can represent each vertex y by the subgroup G_y fixing it. If $H = G_x$ and $y = gx$, then $G_y = gHg^{-1}$, so that now G acts by conjugation.

6.2 Spectrum

Let Γ be a graph and G a group of automorphisms. Let M be a matrix with rows and columns indexed by the vertex set of Γ , and suppose that M commutes with all elements of G (so that $gM = Mg$, or, equivalently, $M_{xy} = M_{gx,gy}$). Now $\text{tr } gM$ only depends on the conjugacy class of g in G , so the map $g \mapsto \text{tr } gM$ defines a class function on G .

(Also the spectrum of gM only depends on the conjugacy class of g in G , but it is not clear how the spectrum should be ordered. Having the trace, however, suffices: one can retrieve the spectrum of a matrix M from the traces of the powers M^i . People also introduce the zeta function of a graph Γ by $\zeta_\Gamma(-s) = \sum \lambda^s = \text{tr } L^s$, where the sum is over the eigenvalues λ of the Laplacian L , in order to have a single object the trace of which encodes the spectrum.)

If Γ has vertex set X , and $V = \mathbb{R}^X$ is the \mathbb{R} -vector space spanned by the vertices of Γ , then by Schur's Lemma M acts as a multiple of the identity on

each irreducible G -invariant subspace of V . In other words, the irreducible G -invariant subspaces are eigenspaces of M . If M acts like θI on the irreducible G -invariant subspace W with character χ , then $\text{tr } gM|_W = \theta\chi(g)$.

Example Let Γ be the Petersen graph, with as vertices the unordered pairs from a 5-set, adjacent when they are disjoint, and let $M = A$, the adjacency matrix. Now $f(g) := \text{tr } gA = \#\{x \mid x \sim gx\}$ defines a class function on $\text{Aut } \Gamma = \text{Sym}(5)$. Below we show f together with the character table of $\text{Sym}(5)$ (with top row indicating the cycle shape of the element):

	1	2	2 ²	3	4	5	2.3
χ_1	1	1	1	1	1	1	1
χ_2	1	-1	1	1	-1	1	-1
χ_3	4	2	0	1	0	-1	-1
χ_4	4	-2	0	1	0	-1	1
χ_5	5	1	1	-1	-1	0	1
χ_6	5	-1	1	-1	1	0	-1
χ_7	6	0	-2	0	0	1	0
f	0	0	4	0	2	5	6

We see that $f = 3\chi_1 - 2\chi_3 + \chi_5$. It follows that Γ has spectrum $3^1 (-2)^4 1^5$, where the eigenvalues are the coefficients of f written as linear combination of irreducible characters, and the multiplicities are the degrees of these characters. The permutation character is $\pi = \chi_1 + \chi_3 + \chi_5$ (obtained for $M = I$). It is *multiplicity free*, that is, no coefficients larger than 1 occur. In the general case the coefficient of an irreducible character χ in the expression for f will be the sum of the eigenvalues of M on the irreducible subspaces with character χ .

6.3 Covers

Let a *graph* $\Gamma = (X, E)$ consist of a set of vertices X and a set of edges E and an incidence relation between X and E (such that each edge is incident with one or two points). An edge incident with one point only is called a *loop*. A *homomorphism* $f : \Gamma \rightarrow \Delta$ of graphs is a map that sends vertices to vertices, edges to edges, loops to loops, and preserves incidence.

For example, the chromatic number of Γ is the smallest integer m such that there is a homomorphism from Γ to K_m .

The map f is called a *covering* when it is a surjective homomorphism, and for each vertex x of Γ and each edge e of Δ that is incident with $f(x)$, there is a unique edge \tilde{e} of Γ that is incident with x such that $f(\tilde{e}) = e$. Now Γ is called a *cover* of Δ .

If f is a covering, then paths in Δ starting at a vertex y of Δ lift uniquely to paths starting at a vertex x of Γ , for each $x \in f^{-1}(y)$.

The *universal cover* of a connected graph Δ is the unique tree T that is a cover. If a is a fixed vertex of Δ , then the vertices of T can be identified with the walks in Δ starting at a that never immediately retrace an edge, where two walks are adjacent when one is the extension of the other by one more edge. The tree T will be infinite when Δ contains at least one cycle. If f is the

covering map (that assigns to a walk its final vertex), then T has a group of automorphisms H acting regularly on the fibers of f .

Given an arbitrary collection of cycles \mathcal{C} in Δ , and a positive integer n_C for each $C \in \mathcal{C}$, one may consider the most general cover satisfying the restriction that the inverse image of the walk traversing C n_C times is closed. (For example, the ‘universal cover modulo triangles’ is obtained by requiring that the preimage of each triangle is a triangle.) There is a unique such graph, quotient of the universal cover. Again the covering group (the group preserving the fibers) acts regularly on the fibers.

Conversely, let Γ be a graph, and H a group of automorphisms. The *quotient graph* Γ/H has as vertices the H -orbits on the vertices of Γ , as edges the H -orbits on the edges of Γ , and a vertex x^H is incident with an edge e^H when some element of x^H is incident with some element of e^H .

The natural projection $\pi : \Gamma \rightarrow \Gamma/H$ is a homomorphism. It will be a covering when no vertex x of Γ is on two edges in an orbit e^H . In this case we also say that Γ is a cover of Γ/H .

Now let Γ be finite, and $f : \Gamma \rightarrow \Delta$ a covering. Let A_Γ and A_Δ be the adjacency matrices of Γ and Δ . Then $(A_\Delta)_{f(x),z} = \sum_{y \in f^{-1}(z)} (A_\Gamma)_{xy}$. If we view A_Γ and A_Δ as linear transformations on the vector spaces V_Γ and V_Δ spanned by the vertices of Γ and Δ , and extend f to a linear map, then this equation becomes $A_\Delta \circ f = f \circ A_\Gamma$. If u is an eigenvector of Δ with eigenvalue θ , then $u \circ f$ (defined by $(u \circ f)_y = u_{f(y)}$) is an eigenvector of Γ with the same eigenvalue, and the same holds for Laplacian eigenvectors and eigenvalues.

(This is immediately clear, but also follows from the fact that the partition of $V\Gamma$ into fibers $f^{-1}(z)$ is an equitable partition.)

For example, let Γ be the path on 6 vertices with a loop added on both sides and Δ the path on 2 vertices with a loop added on both sides. Then the map sending vertices 1, 4, 5 of Γ to one vertex of Δ and 2, 3, 6 to the other, is a covering. The ordinary spectrum of Δ is 2, 0, and hence also Γ has these eigenvalues. (It has spectrum 2, $\sqrt{3}$, 1, 0, -1 , $-\sqrt{3}$.)

Thus, the spectrum of Δ is a subset of the spectrum of Γ . We can be more precise and indicate which subset.

Let $V = \mathbb{R}^X$ be the vector space spanned by the vertices of Γ . Let G be a group of automorphisms of Γ . We can view the elements $g \in G$ as linear transformations of V (permuting the basis vectors). Let H be a subgroup of G , and let W be the subspace of V fixed by H .

Lemma 6.3.1 *Let M be a linear transformation of V that commutes with all $g \in G$. Then M preserves W and $\text{tr } M|_W = (1_H, \phi_M|_H) = (1_H^G, \phi_M)$ where ϕ_M is the class function on G defined by $\phi_M(g) = \text{tr } gM$.*

Proof. The orthogonal projection P from V onto W is given by

$$P = \frac{1}{|H|} \sum_{h \in H} h.$$

If M commutes with all $h \in H$ then $MPu = PMu$, so M preserves the fixed space W , and its restriction $M|_W$ has trace $\text{tr } PM$. Expanding P we find $\text{tr } M|_W = \text{tr } PM = \frac{1}{|H|} \sum_{h \in H} \text{tr } hM = (1_H, \phi_M|_H)$. The second equality

follows by Frobenius reciprocity. \square

Now assume that the map $\pi : \Gamma \rightarrow \Gamma/H$ is a covering. Then $\pi \circ A_\Gamma = A_{\Gamma/H} \circ \pi$. One can identify the vector space $V_{\Gamma/H}$ spanned by the vertices of Γ/H with the vector space W : the vertex x^H corresponds to $\frac{1}{\sqrt{|H|}} \sum_{h \in H} x^h \in W$. This identification identifies $A_{\Gamma/H}$ with $A|_W$. This means that the above lemma (applied with $M = A$) gives the spectrum of Γ/H . In precisely the same way, for $M = L$, it gives the Laplacian spectrum of Γ/H .

We see that for a covering the spectrum of the quotient Γ/H does not depend on the choice of H , but only on the permutation character 1_H^G . This is Sunada's observation, and has been used to construct cospectral graphs, see §13.2.4.

6.4 Cayley sum graphs

In §1.4.8 we discussed Cayley graphs for an abelian group G . A variation is the concept of *Cayley sum graph* with *sum set* S in an abelian group G . It has vertex set G , and two elements $g, h \in G$ are adjacent when $g + h \in S$. (Other terms are *addition Cayley graphs* or just *sum graphs*.)

It is easy to determine the spectrum of a Cayley sum graph.

Proposition 6.4.1 ([131]) *Let Γ be the Cayley sum graph with sum set S in the finite abelian group G . Let χ run through the $n = |G|$ characters of G . The spectrum of Γ consists of the numbers $\chi(S)$ for each real χ , and $\pm|\chi(S)|$ for each pair $\chi, \bar{\chi}$ of conjugate non-real characters, where $\chi(S) = \sum_{s \in S} \chi(s)$.*


Proof. If $\chi : G \rightarrow \mathbb{C}^*$ is a character of G , then $\sum_{y \sim x} \chi(y) = \sum_{s \in S} \chi(s - x) = (\sum_{s \in S} \chi(s))\chi(-x) = \chi(S)\overline{\chi(x)}$. Now Γ is undirected, so the spectrum is real. If χ is a real character, then we found an eigenvector χ , with eigenvalue $\chi(S)$. If χ is non-real, then pick a constant α so that $|\chi(S)| = \alpha^2 \chi(S)$. Then $\text{Re}(\alpha\chi)$ and $\text{Im}(\alpha\chi)$ are eigenvectors with eigenvalues $|\chi(S)|$ and $-|\chi(S)|$, respectively. \square

Chung [85] constructs Cayley sum graphs that are good expanders. For further material on Cayley sum graphs, see [4], [83], [164], [168].

6.4.1 (3,6)-fullerenes


An amusing application was given by DeVos et al. [131]. A *(3,6)-fullerene* is a cubic plane graph whose faces (including the outer face) have sizes 3 or 6. Fowler conjectured (cf. [146]) that such graphs have spectrum $\Phi \cup \{3, -1, -1, -1\}$ (as multiset), where $\Phi = -\Phi$, and this was proved in [131].

For example, the graph



has spectrum $3, \sqrt{5}, 1, (-1)^4, -\sqrt{5}$ with eigenvalues $3, -1, -1, -1$ together with the symmetric part $\pm\sqrt{5}, \pm 1$.

The proof goes as follows. Construct the bipartite double $\Gamma \otimes K_2$ of Γ . This is a cover of Γ , and both triangles and hexagons lift to hexagons, three at each vertex, so that $\Gamma \otimes K_2$ is a quotient of \mathcal{H} , the regular tessellation of the plane with hexagons.



Let \mathcal{H} have set of vertices H , and let $\Gamma \otimes K_2$ have vertex set U , and let Γ have vertex set V . Let $\pi : H \rightarrow U$ and $\rho : U \rightarrow V$ be the quotient maps. The graph $\Gamma \otimes K_2$ is bipartite with bipartite halves U_1 and U_2 , say. Fix a vertex $a_1 \in U_1$ and call it 0. Now $\pi^{-1}(U_1)$ is a lattice in \mathbb{R}^2 , and $\pi^{-1}(a_1)$ is a sublattice (because the concatenation of two walks of even length in Γ starting and ending in a again is such a walk), so the quotient $G = \pi^{-1}(U_1)/\pi^{-1}(a_1)$ is an abelian group, and G can be naturally identified with V . The automorphism of $\Gamma \otimes K_2$ that for each $u \in V$ interchanges the two vertices u_1, u_2 of $\rho^{-1}(u)$, lifts (for each choice of $\bar{a} \in \pi^{-1}(a_2)$) to an isometry of \mathcal{H} with itself that is a point reflection $x \mapsto v - x$ (where $v = \bar{a}$). It follows that if two edges x_1y_2 and z_1w_2 in \mathcal{H} are parallel, then $x + y = z + w$. Hence Γ is the Cayley sum graph for G where the sum set S is the set of three neighbors of a in Γ .

Now the spectrum follows. By the foregoing, the spectrum consists of the values $\pm|\chi(S)|$ for non-real characters χ of G , and $\chi(S)$ for real characters. Since $\text{tr } A = 0$ and Γ is cubic and not bipartite (it has four triangles) it suffices to show that there are precisely four real characters (then the corresponding eigenvalues must be $3, -1, -1, -1$). But this is clear since the number of real characters is the number of elements of order 2 in G , an abelian group with (at most) two generators, hence at most four, and fewer than four would force nonzero $\text{tr } A$. This proves Fowler's conjecture.

6.5 Exercises

Exercise 1 Show that a (3,6)-fullerene has precisely four triangles.

Chapter 7

Euclidean representations

The main goal of this chapter is the famous result by Cameron, Goethals, Seidel and Shult [77] characterizing graphs with smallest eigenvalue not less than -2 .

7.1 Examples

We have seen examples of graphs with smallest eigenvalue $\theta_{\min} \geq -2$. The most important example is formed by the line graphs (see §1.4.5), and people wanted to characterize line graphs by this condition and possibly some additional hypotheses.

Another series of examples are the so-called *cocktailparty graphs*, that is, the graphs $K_{m \times 2}$, i.e., $\overline{mK_2}$, with spectrum $2m - 2, 0^m, (-2)^{m-1}$. For $m \geq 4$ these are not line graphs.

And there are exceptional examples like the Petersen graph (with spectrum $3, 1^5, (-2)^4$), lots of them. It is easy to see that the Petersen graph is not a line graph. More generally, no line graph can have a *3-claw*, that is, an induced $K_{1,3}$ subgraph, as is immediately clear from the definition.

7.2 Euclidean representation

Now suppose Γ has smallest eigenvalue $\theta_{\min} \geq -2$. Then $A + 2I$ is positive semidefinite, so that $A + 2I$ is the Gram matrix of a collection of vectors in some Euclidean space \mathbb{R}^m (where $m = \text{rk}(A + 2I)$), cf. §2.9.

In this way we obtain a map $x \mapsto \bar{x}$ from vertices of Γ to vectors in \mathbb{R}^m , where

$$(\bar{x}, \bar{y}) = \begin{cases} 2 & \text{if } x = y \\ 1 & \text{if } x \sim y \\ 0 & \text{if } x \not\sim y. \end{cases}$$

The additive subgroup of \mathbb{R}^m generated by the vectors \bar{x} , for x in the vertex set X of Γ , is a *root lattice*: an integral lattice generated by *roots*: vectors with squared length 2. Root lattices have been classified. That classification is the subject of the next section.

7.3 Root lattices

We start with an extremely short introduction into lattices.

Lattice

A *lattice* Λ is a discrete additive subgroup of \mathbb{R}^n . Equivalently, it is a finitely generated free \mathbb{Z} -module with positive definite symmetric bilinear form.

Basis

Assume that our lattice Λ has dimension n , i.e., spans \mathbb{R}^n . Let $\{a_1, \dots, a_n\}$ be a \mathbb{Z} -basis of Λ . Let A be the matrix with the vectors a_i as rows. If we choose a different \mathbb{Z} -basis $\{b_1, \dots, b_n\}$, so that $b_i = \sum s_{ij} a_j$, and B is the matrix with the vectors b_i as rows, then $B = SA$, with $S = (s_{ij})$. Since S is integral and invertible, it has determinant ± 1 . It follows that $|\det A|$ is uniquely determined by Λ , independent of the choice of basis.

Volume

\mathbb{R}^n/Λ is an n -dimensional torus, compact with finite volume. Its volume is the volume of the fundamental domain, which equals $|\det A|$.

If Λ' is a sublattice of Λ , then $\text{vol}(\mathbb{R}^n/\Lambda') = \text{vol}(\mathbb{R}^n/\Lambda) \cdot |\Lambda/\Lambda'|$.

Gram matrix

Let G be the matrix (a_i, a_j) of inner products of basis vectors for a given basis. Then $G = AA^\top$, so $\text{vol}(\mathbb{R}^n/\Lambda) = \sqrt{\det G}$.

Dual lattice

The *dual* Λ^* of a lattice Λ is the lattice of vectors having integral inner products with all vectors in Λ : $\Lambda^* = \{x \in \mathbb{R}^n \mid (x, r) \in \mathbb{Z} \text{ for all } r \in \Lambda\}$.

It has a basis $\{a_1^*, \dots, a_n^*\}$ defined by $(a_i^*, a_j) = \delta_{ij}$.

Now $A^* A^\top = I$, so $A^* = (A^{-1})^\top$ and Λ^* has Gram matrix $G^* = G^{-1}$.

It follows that $\text{vol}(\mathbb{R}^n/\Lambda^*) = \text{vol}(\mathbb{R}^n/\Lambda)^{-1}$.

We have $\Lambda^{**} = \Lambda$.

Integral lattice

The lattice Λ is called *integral* when every two lattice vectors have an integral inner product.

For an integral lattice Λ one has $\Lambda \subseteq \Lambda^*$.

The lattice Λ is called *even* when (x, x) is an even integer for each $x \in \Lambda$. An even lattice is integral.

Discriminant

The *determinant*, or *discriminant*, $\text{disc } \Lambda$ of a lattice Λ is defined by $\text{disc } \Lambda = \det G$. When Λ is integral, we have $\text{disc } \Lambda = |\Lambda^*/\Lambda|$.

A lattice is called *self-dual* or *unimodular* when $\Lambda = \Lambda^*$, i.e., when it is integral with discriminant 1. An even unimodular lattice is called *Type II*, the remaining unimodular lattices are called *Type I*.

It can be shown that if there is an even unimodular lattice in \mathbb{R}^n , then n is divisible by 8.

Direct sums

If Λ and Λ' are lattices in \mathbb{R}^m and \mathbb{R}^n , respectively, then $\Lambda \perp \Lambda'$, the *orthogonal direct sum* of Λ and Λ' , is the lattice $\{(x, y) \in \mathbb{R}^{m+n} \mid x \in \Lambda \text{ and } y \in \Lambda'\}$. A lattice is called *irreducible* when it is not the orthogonal direct sum of two nonzero lattices.


Examples

\mathbb{Z}^n



The lattice \mathbb{Z}^n is unimodular, type I.

A_2



The triangular lattice in the plane \mathbb{R}^2 has basis $\{r, s\}$. Choose the scale such that r has length $\sqrt{2}$. Then the Gram matrix is $G = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$, so that $\det G = 3$ and $p, q \in A_2^*$. A fundamental region for A_2 is the parallelogram on $0, r, s$. A fundamental region for A_2^* is the parallelogram on $0, p, q$. Note that the area of the former is thrice that of the latter.

The representation of this lattice in \mathbb{R}^2 has nonintegral coordinates. It is easier to work in \mathbb{R}^3 , on the hyperplane $\sum x_i = 0$, and choose $r = (1, -1, 0)$, $s = (0, 1, -1)$. Then A_2 consists of the points (x_1, x_2, x_3) with $x_i \in \mathbb{Z}$ and $\sum x_i = 0$. The dual lattice A_2^* consists of the points (x_1, x_2, x_3) with $x_1 \equiv x_2 \equiv x_3 \equiv 0 \pmod{1}$ and $\sum x_i = 0$ (so that $3x_1 \in \mathbb{Z}$). It contains for example $p = \frac{1}{3}(2r + s) = (\frac{2}{3}, -\frac{1}{3}, -\frac{1}{3})$.

E_8

Let $\rho : \mathbb{Z}^n \rightarrow 2^n$ be coordinatewise reduction mod 2. Given a binary linear code C , the lattice $\rho^{-1}(C)$ is integral, since it is contained in \mathbb{Z}^n , but never unimodular, unless it is all of \mathbb{Z}^n , a boring situation.

Now suppose that C is self-orthogonal, so that any two code words have an even inner product. Then $\frac{1}{\sqrt{2}}\rho^{-1}(C)$ is an integral lattice. If $\dim C = k$ then $\text{vol}(\mathbb{R}^n/\rho^{-1}(C)) = 2^{n-k}$ and hence $\text{vol}(\mathbb{R}^n/\frac{1}{\sqrt{2}}\rho^{-1}(C)) = 2^{\frac{1}{2}n-k}$. In particular, $\frac{1}{\sqrt{2}}\rho^{-1}(C)$ will be unimodular when C is self-dual, and even when C is ‘doubly even’, i.e., has weights divisible by 4.

Let C be the $[8,4,4]$ extended Hamming code. Then $\frac{1}{\sqrt{2}}\rho^{-1}(C)$ is an even unimodular 8-dimensional lattice known as E_8 .

The code C has weight enumerator $1 + 14X^4 + X^8$ (that is, has one word of weight 0, 14 words of weight 4, and one word of weight 8). It follows that the *roots* (vectors r with $(r, r) = 2$) in this incarnation of E_8 are the 16 vectors $\pm \frac{1}{\sqrt{2}}(2, 0, 0, 0, 0, 0, 0, 0)$ (with 2 in any position), and the $16 \cdot 14 = 224$ vectors $\frac{1}{\sqrt{2}}(\pm 1, \pm 1, \pm 1, \pm 1, 0, 0, 0, 0)$ with ± 1 in the nonzero positions of a weight 4 vector. Thus, there are 240 roots.

Root lattices

A *root lattice* is an integral lattice generated by *roots* (vectors r with $(r, r) = 2$).

For example, A_2 and E_8 are root lattices.

The set of roots in a root lattice is a (reduced) *root system* Φ , i.e., satisfies

- (i) If $r \in \Phi$ and $\lambda r \in \Phi$, then $\lambda = \pm 1$.
- (ii) Φ is closed under the reflection w_r that sends s to $s - 2\frac{(r,s)}{(r,r)}r$ for each $r \in \Phi$.
- (iii) $2\frac{(r,s)}{(r,r)} \in \mathbb{Z}$.

Since Φ generates Λ and Φ is invariant under $W = \langle w_r \mid r \in \Phi \rangle$, the same holds for Λ , so root lattices have a large group of automorphisms.

A *fundamental system* of roots Π in a root lattice Λ is a set of roots generating Λ and such that $(r, s) \leq 0$ for distinct $r, s \in \Pi$. A *reduced fundamental system* of roots is a fundamental system that is linearly independent. A non-reduced fundamental system is called *extended*.

For example, in A_2 the set $\{r, s\}$ is a reduced fundamental system, and $\{r, s, -r - s\}$ is an extended fundamental system.

The *Dynkin diagram* of a fundamental system Π such that $(r, s) \neq -2$ for $r, s \in \Pi$, is the graph with vertex set Π where r and s are joined by an edge when $(r, s) = -1$. (The case $(r, s) = -2$ happens only for a non-reduced system with A_1 component. In that case we do not define the Dynkin diagram.)

Every root lattice has a reduced fundamental system: Fix some vector u , not orthogonal to any root. Put $\Phi^+(u) = \{r \in \Phi \mid (r, u) > 0\}$ and $\Pi(u) = \{r \in \Phi^+(u) \mid r \text{ cannot be written as } s + t \text{ with } s, t \in \Phi^+(u)\}$. Then $\Pi(u)$ is a reduced fundamental system of roots, and written on this basis each root has only positive or only negative coefficients.

(Indeed, if $r, s \in \Pi(u)$ and $(r, s) = 1$, then say $r - s \in \Phi^+(u)$ and $r = (r - s) + s$, contradiction. This shows that $\Pi(u)$ is a fundamental system. If

$\sum \gamma_r r = 0$, then separate the γ_r into positive and negative ones to get $\sum \alpha_r r = \sum \beta_s s = x \neq 0$ where all coefficients α_r, β_s are positive. Now $0 < (x, x) = \sum \alpha_r \beta_s (r, s) < 0$, contradiction. This shows that $\Pi(u)$ is reduced. Now also the last claim follows.)

Proposition 7.3.1 *Let Π be a reduced fundamental system.*

- (i) *For all $x \in \mathbb{R}^n$ there is a $w \in W$ such that $(w(x), r) \geq 0$ for all $r \in \Pi$.*
- (ii) *$\Pi = \Pi(u)$ for some u . (That is, W is transitive on reduced fundamental systems.)*
- (iii) *If Λ is irreducible, then there is a unique $\tilde{r} \in \Phi$ such that $\Pi \cup \{\tilde{r}\}$ is an extended fundamental system.*

Proof. (i) Let G be the Gram matrix of Π , and write $A = 2I - G$. Since G is positive definite, A has largest eigenvalue less than 2. Using Perron-Frobenius, let $\gamma = (\gamma_r)_{r \in \Pi}$ be a positive eigenvector of A . If $(x, s) < 0$ for some $s \in \Pi$, then put $x' = w_s(x) = x - (x, s)s$. Now

$$(x', \sum_r \gamma_r r) = (x, \sum_r \gamma_r r) - (G\gamma)_s(x, s) > (x, \sum_r \gamma_r r).$$

But W is finite, so after finitely many steps we reach the desired conclusion.


(ii) Induction on $|\Pi|$. Fix x with $(x, r) \geq 0$ for all $r \in \Pi$. Then $\Pi_0 = \Pi \cap x^\perp$ is a fundamental system of a lattice in a lower-dimensional space, so of the form $\Pi_0 = \Pi_0(u_0)$. Take $u = x + \varepsilon u_0$ for small $\varepsilon > 0$. Then $\Pi = \Pi(u)$.

(iii) If $r \in \Phi^+(u)$ has maximal (r, u) , then $\tilde{r} = -r$ is the unique root that can be added. It can be added, since $(\tilde{r}, s) \geq 0$ means $(r, s) < 0$, so that $r+s$ is a root, contradicting maximality of r . And it is unique because linear dependencies of an extended system correspond to an eigenvector with eigenvalue 2 of the extended Dynkin diagram, and by Perron-Frobenius up to a constant there is a unique such eigenvector when the diagram is connected, that is, when Λ is irreducible. \square


Classification

The irreducible root lattices one finds are A_n ($n \geq 0$), D_n ($n \geq 4$), E_6 , E_7 , E_8 . Each is defined by its Dynkin diagram.


(1) A_n : The lattice vectors are: $x \in \mathbb{Z}^{n+1}$ with $\sum x_i = 0$. There are $n(n+1)$ roots: $e_i - e_j$ ($i \neq j$). The discriminant is $n+1$, and $A_n^*/A_n \cong \mathbb{Z}_{n+1}$, with the quotient generated by $\frac{1}{n+1}(e_1 + \dots + e_n - ne_{n+1}) \in A_n^*$.




(2) D_n : The lattice vectors are: $x \in \mathbb{Z}^n$ with $\sum x_i \equiv 0 \pmod{2}$. There are $2n(n-1)$ roots $\pm e_i \pm e_j$ ($i \neq j$). The discriminant is 4, and D_n^*/D_n is isomorphic to \mathbb{Z}_4 when n is odd, and to $\mathbb{Z}_2 \times \mathbb{Z}_2$ when n is even. D_n^* contains e_1 and $\frac{1}{2}(e_1 + \dots + e_n)$. Note that $D_3 \cong A_3$.




(3) E_8 : (Recall that we already gave a construction of E_8 from the Hamming code.) The lattice is the span of D_8 and $c := \frac{1}{2}(e_1 + \dots + e_8)$. There are $240 = 112 + 128$ roots, of the forms $\pm e_i \pm e_j$ ($i \neq j$) and $\frac{1}{2}(\pm e_1 \pm \dots \pm e_8)$ with an even number of minus signs. The discriminant is 1, and $E_8^* = E_8$.



(4) E_7 : Take $E_7 = E_8 \cap c^\perp$. There are $126 = 56 + 70$ roots. The discriminant is 2, and E_7^* contains $\frac{1}{4}(1, 1, 1, 1, 1, 1, -3, -3)$.



(5) E_6 : For the vector $d = -e_7 - e_8$, take $E_6 = E_8 \cap \{c, d\}^\perp$. There are $72 = 32 + 40$ roots. The discriminant is 3, and E_6^* contains the vector $\frac{1}{3}(1, 1, 1, 1, -2, -2, 0, 0)$.



That this is all, is an easy consequence of the Perron-Frobenius theorem: $A = 2I - G$ is the adjacency matrix of a graph, namely the Dynkin diagram, and this graph has largest eigenvalue at most 2. These graphs were determined in Theorem 3.1.3. The connected graphs with largest eigenvalue less than 2 are the Dynkin diagrams of reduced fundamental systems of irreducible root systems and the connected graphs with largest eigenvalue 2 are the Dynkin diagrams of extended root systems.

In the pictures above, the reduced fundamental systems were drawn with black dots, and the additional element of the extended system with an open dot (and a name given in parentheses).

7.4 Cameron-Goethals-Seidel-Shult

Now return to the discussion of connected graphs Γ with smallest eigenvalue $\theta_{\min} \geq -2$. In §7.2 we found a map $x \mapsto \bar{x}$ from the vertex set X of Γ to some Euclidean space \mathbb{R}^m such that the inner product (\bar{x}, \bar{y}) takes the value 2, 1, 0 when $x = y$, $x \sim y$ and $x \not\sim y$, respectively.

Let Σ be the image of X under this map. Then Σ generates a root lattice Λ . Since Γ is connected, the root lattice is irreducible.

By the classification of root lattices, it follows that the root lattice is one of A_n , D_n , E_6 , E_7 or E_8 . Note that the graph is determined by Σ , so that the classification of graphs with $\theta_{\min} \geq -2$ is equivalent to the classification of subsets Φ of the root system with the property that all inner products are 2, 1, or 0, i.e., nonnegative.

Now A_n and D_n can be chosen to have integral coordinates, and $E_6 \subset E_7 \subset E_8$, so we have the two cases (i) $\Sigma \subset \mathbb{Z}^{m+1}$, and (ii) $\Sigma \subset E_8$. A graph is called *exceptional* in case (ii). Since E_8 has a finite number of roots, there are only finitely many exceptional graphs.

In case (i) one quickly sees what the structure of Γ has to be. Something like a line graph with attached cocktailparty graphs. This structure has been baptised *generalized line graph*. The precise definition will be clear from the proof of the theorem below.

Theorem 7.4.1 (i) *Let Γ be a connected graph with smallest eigenvalue $\theta_{\min} \geq -2$. Then either Γ is a generalized line graph, or Γ is one of finitely many exceptions, represented by roots in the E_8 lattice.*

(ii) *A regular generalized line graph is either a line graph or a cocktailparty graph.*

(iii) *A graph represented by roots in the E_8 lattice has at most 36 vertices, and every vertex has valency at most 28.*

Proof. (i) Consider the case $\Sigma \subset \mathbb{Z}^{m+1}$. Roots in \mathbb{Z}^{m+1} have shape $\pm e_i \pm e_j$. If some e_i has the same sign in all $\sigma \in \Sigma$ in which it occurs, then choose the basis such that this sign is +. Let I be the set of all such indices i . Then $\{x \mid \bar{x} = e_i + e_j \text{ for some } i, j \in I\}$ induced a line graph in Γ , with x corresponding to the edge ij on I . If $j \notin I$, then e_j occurs with both signs, and there are $\sigma, \tau \in \Sigma$ with $\sigma = \pm e_i + e_j$ and $\tau = \pm e_{i'} - e_j$. Since all inner products in Σ are nonnegative, $i = i'$ with $i \in I$, and $\sigma = e_i + e_j$, $\tau = e_i - e_j$. Thus, i is determined by j and we have a map $\phi : j \mapsto i$ from indices outside I to indices in I . Now for each $i \in I$ the set $\{x \mid \bar{x} = e_i \pm e_j \text{ for some } j \text{ with } \phi(j) = i\}$ induces a cocktailparty graph. Altogether we see in what way Γ is a line graph with attached cocktailparty graphs.

(ii) Now let Γ be regular. A vertex x with $\bar{x} = e_i - e_j$ is adjacent to all vertices with image $e_i \pm e_k$ different from $e_i + e_j$. But a vertex y with $\bar{y} = e_i + e_k$ where $i, k \in I$ is adjacent to all vertices with image $e_i \pm e_k$ without exception (and also to vertices with image $e_k \pm e_l$). Since Γ is regular both types of vertices cannot occur together, so that Γ is either a line graph or a cocktailparty graph.

(iii) Suppose $\Sigma \subset E_8$. Consider the 36-dimensional space of symmetric 8×8 matrices, equipped with the positive definite inner product $(P, Q) = \text{tr } PQ$. Associated with the 240 roots r of E_8 are 120 rank 1 matrices $P_r = rr^\top$ with mutual inner products $(P_r, P_s) = \text{tr } rr^\top ss^\top = (r, s)^2$. The Gram matrix of the

set of P_r for $r \in \Sigma$ is $G = 4I + A$. Since G is positive definite (it has smallest eigenvalue ≥ 2), the vectors P_r are linearly independent, and hence $|\Sigma| \leq 36$.

Finally, let r be a root of E_8 . The 56 roots s of E_8 that satisfy $(r, s) = 1$ fall into 28 pairs s, s' where $(s, s') = -1$. So, Σ can contain at most one member from each of these pairs, and each vertex of Γ has valency at most 28. \square

The bounds in (ii) are best possible: Take the graph $K_8 + L(K_8)$ and add edges joining $i \in K_8$ with $jk \in L(K_8)$ whenever i, j, k are distinct. This graph has 36 vertices, the vertices in K_8 have 28 neighbours, and the smallest eigenvalue is -2 . A representation in E_8 is given by $i \mapsto \frac{1}{2}(e_1 + \dots + e_8) - e_i$ and $jk \mapsto e_j + e_k$.

There is a large amount of literature on exceptional graphs.

7.5 Exercises

Exercise 1 Show that the following describes a root system of type E_6 . Take the following 72 vectors in \mathbb{R}^9 : 18 vectors $\pm(u, 0, 0), \pm(0, u, 0), \pm(0, 0, u)$ with $u \in \{(1, -1, 0), (0, 1, -1), (-1, 0, 1)\}$, and 54 vectors $\pm(u, v, w)$ with $u, v, w \in \{(\frac{2}{3}, -\frac{1}{3}, -\frac{1}{3}), (-\frac{1}{3}, \frac{2}{3}, -\frac{1}{3}), (-\frac{1}{3}, -\frac{1}{3}, \frac{2}{3})\}$.

Exercise 2 Show that the following describes a root system of type E_7 . Take the following 126 vectors in \mathbb{R}^7 : 60 vectors $\pm e_i \pm e_j$ with $1 \leq i < j \leq 6$, and 64 vectors $\pm(x_1, \dots, x_6, \frac{1}{\sqrt{2}})$ with $x_i = \pm \frac{1}{2}$ where an even number of x_i has $+$ sign, and 2 vectors $\pm(0, \dots, 0, \sqrt{2})$.

Chapter 8

Strongly regular graphs

8.1 Strongly regular graphs

A graph (simple, undirected and loopless) of order v is called *strongly regular* with parameters v, k, λ, μ whenever it is not complete or edgeless and

- (i) each vertex is adjacent to k vertices,
- (ii) for each pair of adjacent vertices there are λ vertices adjacent to both,
- (iii) for each pair of non-adjacent vertices there are μ vertices adjacent to both.

We require that both edges and non-edges occur, so that the parameters are well-defined.

In association scheme terminology (cf. §10.1), a strongly regular graph is a symmetric association scheme with two (nonidentity) classes, in which one relation is singled out to be the adjacency relation.

8.1.1 Simple examples

Easy examples of strongly regular graphs:

- (i) A quadrangle is strongly regular with parameters $(4, 2, 0, 2)$.
 - (ii) A pentagon is strongly regular with parameters $(5, 2, 0, 1)$.
 - (iii) The 3×3 grid, the Cartesian product of two triangles, is strongly regular with parameters $(9, 4, 1, 2)$.
 - (iv) The Petersen graph is strongly regular with parameters $(10, 3, 0, 1)$.
- (Each of these graphs is uniquely determined by its parameters, so if you do not know what a pentagon is, or what the Petersen graph is, this defines it.)

Each of these examples can be generalized in numerous ways. For example,

- (v) Let $q = 4t + 1$ be a prime power. The *Paley graph* $\text{Paley}(q)$ is the graph with the finite field \mathbb{F}_q as vertex set, where two vertices are adjacent when they differ by a (nonzero) square. It is strongly regular with parameters $(4t + 1, 2t, t - 1, t)$, as we shall see below. Doing this for $q = 5$ and $q = 9$, we find the examples (ii) and (iii) again. For $q = 13$ we find a graph that is locally a hexagon. For $q = 17$ we find a graph that is locally an 8-gon + diagonals.

(vi) The $m \times m$ grid, the Cartesian product of two complete graphs on m vertices, is strongly regular with parameters $(m^2, 2(m-1), m-2, 2)$ (for $m > 1$). For $m = 2$ and $m = 3$ we find the examples (i) and (iii) again.

(vii) The complete multipartite graph $K_{m \times a}$, with vertex set partitioned into m groups of size a , where two points are adjacent when they are from different groups, is strongly regular with parameters $(ma, (m-1)a, (m-2)a, (m-1)a)$ (for $m > 1$ and $a > 1$). For $m = a = 2$ we find Example (i) again.

The *complement* of a graph Γ is the graph $\bar{\Gamma}$ with the same vertex set as Γ , where two vertices are adjacent if and only if they are nonadjacent in Γ . The complement of a strongly regular graph with parameters (v, k, λ, μ) is again strongly regular, and has parameters $(v, v-k-1, v-2k+\mu-2, v-2k+\lambda)$. (Indeed, we keep the same association scheme, but now single out the other nonidentity relation.)

(viii) The Paley graph $\text{Paley}(q)$ is isomorphic to its complement. (Indeed, an isomorphism is given by multiplication by a nonsquare.) In particular we see that the pentagon and the 3×3 grid are (isomorphic to) their own complements.

(ix) The disjoint union mK_a of m complete graphs of size a is strongly regular with parameters $(ma, a-1, a-2, 0)$ (for $m > 1$ and $a > 1$). These graphs are the complements of those in Example (vii).

(x) The *triangular graph* on the pairs in an m -set, denoted by $T(m)$, or by $\binom{m}{2}$, has these pairs as vertices, where two pairs are adjacent whenever they meet in one point. These graphs are strongly regular, with parameters $(\binom{m}{2}, 2(m-2), m-2, 4)$, if $m \geq 4$. For $m = 4$ we find $K_{3 \times 2}$. For $m = 5$ we find the complement of the Petersen graph.

The four parameters are not independent. Indeed, if $\mu \neq 0$ we find the relation

$$v = 1 + k + \frac{k(k-1-\lambda)}{\mu}$$

by counting vertices at distance 0, 1 and 2 from a given vertex.

8.1.2 The Paley graphs

Above we claimed that the Paley graphs (with vertex set \mathbb{F}_q , where q is a prime power congruent 1 mod 4, and where two vertices are adjacent when their difference is a nonzero square) are strongly regular. Let us verify this.

Proposition 8.1.1 *The Paley graph $\text{Paley}(q)$ with $q = 4t+1$ is strongly regular with parameters $(v, k, \lambda, \mu) = (4t+1, 2t, t-1, t)$. It has eigenvalues $k, (-1 \pm \sqrt{q})/2$ with multiplicities 1, $2t, 2t$, respectively.*

Proof. The values for v and k are clear. Let $\chi : \mathbb{F}_q \rightarrow \{-1, 0, 1\}$ be the quadratic residue character defined by $\chi(0) = 0$, $\chi(x) = 1$ when x is a (nonzero) square, and $\chi(x) = -1$ otherwise. Note that $\sum_x \chi(x) = 0$, and that for nonzero a we have $\sum_z \chi(z^2 - az) = \sum_{z \neq 0} \chi(1 - \frac{a}{z}) = -1$. Now λ and μ follow from

$$4 \sum_{\substack{z \\ x \sim z \sim y}} 1 = \sum_{z \neq x, y} (\chi(z-x) + 1)(\chi(z-y) + 1) = -1 - 2\chi(x-y) + (q-2).$$

For the spectrum, see Theorem 8.1.3 below. \square

8.1.3 Adjacency matrix

For convenience we call an eigenvalue *restricted* if it has an eigenvector perpendicular to the all-ones vector $\mathbf{1}$.

Theorem 8.1.2 *For a simple graph Γ of order v , not complete or edgeless, with adjacency matrix A , the following are equivalent:*

- (i) Γ is strongly regular with parameters (v, k, λ, μ) for certain integers k, λ, μ ,
- (ii) $A^2 = (\lambda - \mu)A + (k - \mu)I + \mu J$ for certain real numbers k, λ, μ ,
- (iii) A has precisely two distinct restricted eigenvalues.

Proof. The equation in (ii) can be rewritten as

$$A^2 = kI + \lambda A + \mu(J - I - A).$$

Now (i) \iff (ii) is obvious.

(ii) \Rightarrow (iii): Let ρ be a restricted eigenvalue, and u a corresponding eigenvector perpendicular to $\mathbf{1}$. Then $Ju = 0$. Multiplying the equation in (ii) on the right by u yields $\rho^2 = (\lambda - \mu)\rho + (k - \mu)$. This quadratic equation in ρ has two distinct solutions. (Indeed, $(\lambda - \mu)^2 = 4(\mu - k)$ is impossible since $\mu \leq k$ and $\lambda \leq k - 1$.)

(iii) \Rightarrow (ii): Let r and s be the restricted eigenvalues. Then $(A - rI)(A - sI) = \alpha J$ for some real number α . So A^2 is a linear combination of A , I and J . \square

8.1.4 Imprimitve graphs

A strongly regular graph is called *imprimitve* if it, or its complement, is disconnected, and *primitive* otherwise. Imprimitve strongly regular graphs are boring.

If a strongly regular graph is not connected, then $\mu = 0$ and $k = \lambda + 1$. And conversely, if $\mu = 0$ or $k = \lambda + 1$ then the graph is a disjoint union aK_m of some number a of complete graphs K_m . In this case $v = am$, $k = m - 1$, $\lambda = m - 2$, $\mu = 0$ and the spectrum is $(m - 1)^a, (-1)^{a(m-1)}$.

If the complement of a strongly regular graph is not connected, then $k = \mu$. And conversely, if $k = \mu$ then the graph is the complete multipartite graph $K_{a \times m}$, the complement of aK_m , with parameters $v = am$, $k = \mu = (a - 1)m$, $\lambda = (a - 2)m$ and spectrum $(a - 1)m^1, 0^{a(m-1)}, (-m)^{a-1}$.

Let r and s ($r > s$) be the restricted eigenvalues of A . For a primitive strongly regular graph one has $k > r > 0$ and $s < -1$.

8.1.5 Parameters

Theorem 8.1.3 *Let Γ be a strongly regular graph with adjacency matrix A and parameters (v, k, λ, μ) . Let r and s ($r > s$) be the restricted eigenvalues of A and let f, g be their respective multiplicities. Then*

- (i) $k(k - 1 - \lambda) = \mu(v - k - 1)$,
- (ii) $rs = \mu - k, r + s = \lambda - \mu$,
- (iii) $f, g = \frac{1}{2}(v - 1 \mp \frac{(r+s)(v-1)+2k}{r-s})$.

(iv) If r and s are non-integral, then $f = g$ and $(v, k, \lambda, \mu) = (4t + 1, 2t, t - 1, t)$ for some integer t .

Proof. (i) Fix a vertex x of Γ . Let $\Gamma(x)$ and $\Delta(x)$ be the sets of vertices adjacent and non-adjacent to x , respectively. Counting in two ways the number of edges between $\Gamma(x)$ and $\Delta(x)$ yields (i). The equations (ii) are direct consequences of Theorem 8.1.2(ii), as we saw in the proof. Formula (iii) follows from $f + g = v - 1$ and $0 = \text{trace } A = k + fr + gs = k + \frac{1}{2}(r + s)(f + g) + \frac{1}{2}(r - s)(f - g)$. Finally, when $f \neq g$ then one can solve for r and s in (iii) (using (ii)) and find that r and s are rational, and hence integral. But $f = g$ implies $(\mu - \lambda)(v - 1) = 2k$, which is possible only for $\mu - \lambda = 1$, $v = 2k + 1$. \square

These relations imply restrictions for the possible values of the parameters. Clearly, the right hand sides of (iii) must be positive integers. These are the so-called *rationality conditions*.

8.1.6 The half case and cyclic strongly regular graphs

The case of a strongly regular graph with parameters $(v, k, \lambda, \mu) = (4t + 1, 2t, t - 1, t)$ for some integer t is called the *half case*. Such graphs are also called *conference graphs*. If such a graph exists, then v is the sum of two squares, see Theorem 9.4.2 below. The Paley graphs (§8.1.2, §9.4, §12.6) belong to this case, but there are many further examples.

A characterization of the Paley graphs of prime order is given by

Proposition 8.1.4 (Kelly [214], Bridges & Mena [38]) *A strongly regular graph with a regular cyclic group of automorphisms is a Paley graph with a prime number of vertices.*

(See the discussion of translation association schemes in BCN [48], §2.10. This result has been rediscovered several times.)

8.1.7 Strongly regular graphs without triangles

As an example of the application of the rationality conditions we classify the strongly regular graphs of girth 5.

Theorem 8.1.5 (Hoffman & Singleton [196]) *Suppose $(v, k, 0, 1)$ is the parameter set of a strongly regular graph. Then $(v, k) = (5, 2), (10, 3), (50, 7)$ or $(3250, 57)$.*

Proof. The rationality conditions imply that either $f = g$, which leads to $(v, k) = (5, 2)$, or $r - s$ is an integer dividing $(r + s)(v - 1) + 2k$. By Theorem 8.1.3(i)–(ii) we have

$$s = -r - 1, \quad k = r^2 + r + 1, \quad v = r^4 + 2r^3 + 3r^2 + 2r + 2,$$

and thus we obtain $r = 1, 2$ or 7 . \square

The first three possibilities are uniquely realized by the pentagon, the Petersen graph and the Hoffman-Singleton graph. For the last case existence is unknown (but see §10.5.1).

More generally we can look at strongly regular graphs of girth at least 4. Seven examples are known.

- (i) The *pentagon*, with parameters $(5, 2, 0, 1)$.
- (ii) The *Petersen graph*, with parameters $(10, 3, 0, 1)$. This is the complement of the triangular graph $T(5)$.
- (iii) The *folded 5-cube*, with parameters $(16, 5, 0, 2)$. This graph is obtained from the 5-cube 2^5 on 32 vertices by identifying antipodal vertices. (The complement of this graph is known as the Clebsch graph.)
- (iv) The *Hoffman-Singleton graph*, with parameters $(50, 7, 0, 1)$. There are many constructions for this graph, cf., e.g., [48], §13.1. A short one, due to N. Robertson, is the following. Take 25 vertices (i, j) and 25 vertices $(i, j)'$ with $i, j \in \mathbb{Z}_5$, and join (i, j) with $(i, j + 1)$, $(i, j)'$ with $(i, j + 2)'$, and (i, k) with $(j, i + k)'$ for all $i, j, k \in \mathbb{Z}_5$. Now the subsets $(i, *)$ become pentagons, the $(i, *)'$ become pentagons (drawn as pentagrams), and each of the 25 unions of $(i, *)$ with $(j, *)'$ induces a Petersen subgraph.
- (v) The *Gewirtz graph*, with parameters $(56, 10, 0, 2)$. This is the graph with as vertices the $77 - 21 = 56$ blocks of the unique Steiner system $S(3, 6, 22)$ not containing a given symbol, where two blocks are adjacent when they are disjoint. It is a subgraph of the following.
- (vi) The M_{22} *graph*, with parameters $(77, 16, 0, 4)$. This is the graph with as vertices the 77 blocks of the unique Steiner system $S(3, 6, 22)$, adjacent when they are disjoint. It is a subgraph of the following.
- (vii) The *Higman-Sims graph*, with parameters $(100, 22, 0, 6)$. This is the graph with as $1 + 22 + 77$ vertices an element ∞ , the 22 symbols of $S(3, 6, 22)$, and the 77 blocks of $S(3, 6, 22)$. The element ∞ is adjacent to the 22 symbols, each symbol is adjacent to the 21 blocks containing it, and blocks are adjacent when disjoint. The (rank 3) automorphism group of this graph is $HS.2$, where HS is the sporadic simple group of Higman and Sims. This graph can be partitioned into two halves, each inducing a Hoffman-Singleton graph, cf. [48], §13.1.

Each of these seven graphs is uniquely determined by its parameters. It is unknown whether there are any further examples. There are infinitely many feasible parameter sets. For the parameters $(324, 57, 0, 12)$ nonexistence was shown in Gavrilyuk & Makhnev [150] and in Kaski & Östergård [213].

8.1.8 Further parameter restrictions

Except for the rationality conditions, a few other restrictions on the parameters are known. We mention two of them. The *Krein conditions*, due to Scott [270], can be stated as follows:

$$(r + 1)(k + r + 2rs) \leq (k + r)(s + 1)^2,$$

$$(s + 1)(k + s + 2rs) \leq (k + s)(r + 1)^2.$$

When equality holds in one of these, the subconstituents of the graph (the induced subgraphs on the neighbours and on the nonneighbours of a given point) are both strongly regular (in the wide sense) again. For example, in the Higman-Sims graph with parameters $(v, k, \lambda, \mu) = (100, 22, 0, 6)$ and $k, r, s = 22, 2, -8$ the second subconstituent of any point has parameters $(77, 16, 0, 4)$.

Seidel's *absolute bound* for the number of vertices of a primitive strongly regular graph (see Corollary 9.6.8 below) reads

$$v \leq f(f+3)/2, \quad v \leq g(g+3)/2.$$

For example, the parameter set (28,9,0,4) (spectrum $9^1 \ 1^{21} \ (-5)^6$) is ruled out both by the second Krein condition and by the absolute bound.

A useful identity is an expression for the *Frame quotient* (cf. [48], 2.2.4 and 2.7.2). One has

$$fg(r-s)^2 = vk(v-1-k).$$

(as is easy to check directly from the expressions for f and g given in Theorem 8.1.3(iii)). From this one immediately concludes that if v is prime, then $r-s = \sqrt{v}$ and we are in the 'half case' $(v, k, \lambda, \mu) = (4t+1, 2t, t-1, t)$.

The Frame quotient, Krein conditions and absolute bound are special cases of general (in)equalities for association schemes—see also §10.4 below. In Brouwer & van Lint [56] one may find a list of known restrictions and constructions. It is a sequel to Hubaut's [202] earlier survey of constructions.

Using the above parameter conditions, Neumaier [249] derives the μ -bound:

Theorem 8.1.6 *For a primitive strongly regular graph $\mu \leq s^3(2s+3)$. If equality holds, then $r = -s^2(2s+3)$.*

Examples of equality in the μ -bound are known for $s = -2$ (the Schläfli graph, with $(v, k, \lambda, \mu) = (27, 16, 10, 8)$) and $s = -3$ (the McLaughlin graph, with $(v, k, \lambda, \mu) = (275, 162, 105, 81)$).

Brouwer & Neumaier [58] showed that a connected partial linear space with girth at least 5 and more than one line, in which every point is collinear with m other points, contains at least $\frac{1}{2}m(m+3)$ points. It follows that a strongly regular graph with $\mu = 2$ either has $k \geq \frac{1}{2}\lambda(\lambda+3)$ or has $(\lambda+1)|k$.

Bagchi [12] showed that any $K_{1,1,2}$ -free strongly regular graph is either the collinearity graph of a generalized quadrangle or satisfies $k \geq (\lambda+1)(\lambda+2)$. (It follows that in the above condition on $\mu = 2$ the $(\lambda+1)|k$ alternative only occurs for the $m \times m$ grid, where $m = \lambda+2$.)

8.1.9 Strongly regular graphs from permutation groups

Suppose G is a permutation group, acting on a set Ω . The *rank* of the action is the number of orbits of G on $\Omega \times \Omega$. (These latter orbits are called *orbitals*.) If R is an orbital, or a union of orbitals, then (Ω, R) is a directed graph that admits G as group of automorphisms.

If G is transitive of rank 3 and its orbitals are symmetric (for all $x, y \in \Omega$ the pairs (x, y) and (y, x) belong to the same orbital), say with orbitals I, R, S , where $I = \{(x, x) \mid x \in \Omega\}$, then (Ω, R) and (Ω, S) is a pair of complementary strongly regular graphs.

For example, let G be $\text{Sym}(n)$ acting on a set Σ of size 5. This action induces an action on the set Ω of unordered pairs of elements in Σ , and this latter action is rank 3, and gives the pair of graphs $T(5)$ and $\overline{T(5)}$, where this latter graph is the Petersen graph.

The rank 3 groups have been classified by the combined effort of many people, including Foulser, Kantor, Liebler, Liebeck and Saxl, see [212, 225, 226].

8.1.10 Strongly regular graphs from quasisymmetric designs

As an application of Theorem 8.1.2, we show that quasisymmetric block designs give rise to strongly regular graphs. A *quasisymmetric design* is a $2-(v, k, \lambda)$ design (see §4.7) such that any two blocks meet in either x or y points, for certain fixed distinct x, y . Given this situation, we may define a graph Γ on the set of blocks, and call two blocks adjacent when they meet in x points. Let N be the point-block matrix of the design and A the adjacency matrix of Γ . Then $N^\top N = kI + xA + y(J - I - A)$. Since each of NN^\top , NJ , and JN is a linear combination of I and J , we see that A^2 can be expressed in terms of A, I, J , so that Γ is strongly regular by part (ii) of Theorem 8.1.2. (For an application, see §9.3.2.)

A large class of quasisymmetric block designs is provided by the $2-(v, k, \lambda)$ designs with $\lambda = 1$ (also known as Steiner systems $S(2, k, v)$). Such designs have only two intersection numbers since no two blocks can meet in more than one point. This leads to a substantial family of strongly regular graphs, including the triangular graphs $T(m)$ (derived from the trivial design consisting of all pairs from an m -set).

8.1.11 Symmetric 2-designs from strongly regular graphs

Conversely, some families of strongly regular graphs lead to designs. Let A be the adjacency matrix of a strongly regular graph with parameters (v, k, λ, λ) (i.e., with $\lambda = \mu$; such a graph is sometimes called a (v, k, λ) graph). Then, by Theorem 8.1.2

$$AA^\top = A^2 = (k - \lambda)I + \lambda J,$$

which reflects that A is the incidence matrix of a symmetric $2-(v, k, \lambda)$ design. (And in this way one obtains precisely all symmetric 2-designs possessing a polarity without absolute points.) For instance, the triangular graph $T(6)$ provides a symmetric $2-(15, 8, 4)$ design, the complementary design of the design of points and planes in the projective space $PG(3, 2)$. Similarly, if A is the adjacency matrix of a strongly regular graph with parameters $(v, k, \lambda, \lambda + 2)$, then $A + I$ is the incidence matrix of a symmetric $2-(v, k + 1, \lambda + 2)$ design (and in this way one obtains precisely all symmetric 2-designs possessing a polarity with all points absolute). For instance, the Gewirtz graph with parameters $(56, 10, 0, 2)$ provides a biplane $2-(56, 11, 2)$.

8.1.12 Latin square graphs

A *transversal design* of *strength* t and *index* λ is a triple $(X, \mathcal{G}, \mathcal{B})$, where X is a set of points, \mathcal{G} is a partition of X into *groups*, and \mathcal{B} is a collection of subsets of X called *blocks* such that (i) $t \leq |\mathcal{G}|$, (ii) every block meets every group in precisely one point, and (iii) every t -subset of X that meets each group in at most one point is contained in precisely λ blocks.

Suppose X is finite and $t < |\mathcal{G}|$. Then all groups $G \in \mathcal{G}$ have the same size m , and the number of blocks is λm^t . Given a point $x_0 \in X$, the groups not on x_0 together with the blocks $B \setminus \{x_0\}$ for $x_0 \in B \in \mathcal{B}$ form a transversal design of strength $t - 1$ with the same index λ .

Equivalent to the concept of transversal design is that of *orthogonal array*. An orthogonal array with strength t and index λ over an alphabet of size m is a $k \times N$ array (with $N = \lambda m^t$) such that for any choice of t rows and prescribed symbols on these rows there are precisely λ columns that satisfy the demands.

When $t = 2$ the strength is usually not mentioned, and one talks about transversal designs $TD_\lambda(k, m)$ or orthogonal arrays $OA_\lambda(m, k)$, where k is the block size and m the group size.

When $\lambda = 1$ the index is suppressed from the notation. Now a $TD(k, m)$ or $OA(m, k)$ is equivalent to a set of $k - 2$ mutually orthogonal Latin squares of order m . (The k rows of the orthogonal array correspond to row index, column index, and Latin square number; the columns correspond to the m^2 positions.)

The dual of a transversal design is a *net*. An (m, k) -net is a set of m^2 points together with km lines, partitioned into k parallel classes, where two lines from different parallel classes meet in precisely one point.

Given a point-line incidence structure, the *point graph* or *collinearity graph* is the graph with the points as vertices, adjacent when they are collinear. Dually, the *block graph* is the graph with the lines as vertices, adjacent when they have a point in common.

The collinearity graph of an (m, t) -net, that is, the block graph of a transversal design $TD(t, m)$ (note the new use of t here!), is strongly regular with parameters $v = m^2$, $k = t(m - 1)$, $\lambda = m - 2 + (t - 1)(t - 2)$, $\mu = t(t - 1)$ and eigenvalues $r = m - t$, $s = -t$. One says that a strongly regular graph 'is a pseudo Latin square graph', or 'has Latin square parameters' when there are t and m such that (v, k, λ, μ) have the above values. One also says that it has ' $OA(m, t)$ parameters'.

There is extensive literature on nets and transversal designs.

Proposition 8.1.7 *Suppose Γ is a strongly regular graph with $OA(m, t)$ parameters with a partition into cocliques of size m . Then the graph Δ obtained from Γ by adding edges so that these cocliques become cliques is again strongly regular and has $OA(m, t + 1)$ parameters.*

Proof. More generally, let Γ be a strongly regular graph with a partition into cocliques that meet the Hoffman bound. Then the graph Δ obtained from Γ by adding edges so that these cocliques become cliques has spectrum $k + m - 1$, $(r - 1)^f$, $(s + m - 1)^h$, $(s - 1)^{g-h}$, where m is the size of the cocliques, and $h = v/m - 1$. The proposition is the special case $m = r - s$. \square

For example, from the Hall-Janko graph with $OA(10, 4)$ parameters (100, 36, 12, 14) and a partition into ten 10-cocliques (which exists) one obtains a strongly regular graph with $OA(10, 5)$ parameters (100, 45, 20, 20), and hence also a symmetric design 2-(100, 45, 20). But an $OA(10, 5)$ (three mutually orthogonal Latin squares of order 10) is unknown.

8.1.13 Partial Geometries

A *partial geometry* with parameters (s, t, α) is a point-line geometry (any two points are on at most one line) such that all lines have size $s + 1$, there are $t + 1$ lines on each point, and given a line and a point outside, the point is collinear

with α points on the given line. One calls this structure a $pg(s, t, \alpha)$. Note that the dual of a $pg(s, t, \alpha)$ is a $pg(t, s, \alpha)$ (where ‘dual’ means that the names ‘point’ and ‘line’ are swapped).

Partial geometries were introduced by Bose [31].

One immediately computes the number of points $v = (s + 1)(st + \alpha)/\alpha$ and lines $b = (t + 1)(st + \alpha)/\alpha$.

The extreme examples of partial geometries are *generalized quadrangles* (partial geometries with $\alpha = 1$) and Steiner systems $S(2, K, V)$ (partial geometries with $\alpha = s + 1$). Many examples are also provided by nets (with $t = \alpha$) or their duals, the transversal designs (with $s = \alpha$).

The collinearity graph of a $pg(s, t, \alpha)$ is complete if $\alpha = s + 1$, and otherwise strongly regular with parameters $v = (s + 1)(st + \alpha)/\alpha$, $k = s(t + 1)$, $\lambda = s - 1 + t(\alpha - 1)$, $\mu = \alpha(t + 1)$, and eigenvalues $\theta_1 = s - \alpha$, $\theta_2 = -t - 1$. (Note: earlier we used s for the smallest eigenvalue, but here s has a different meaning!)

A strongly regular graph is called *geometric* when it is the collinearity graph of a partial geometry. It is called *pseudo-geometric* when there are integers s, t, α such that the parameters (v, k, λ, μ) have the above-given values.

Bose [31] showed that a pseudo-geometric graph with given t and sufficiently large s must be geometric. Neumaier [249] showed that the same conclusion works in all cases, and hence derives a contradiction in the non-pseudo-geometric case.

Theorem 8.1.8 (Bose-Neumaier) *A strongly regular graph with $s < -1$ and $r > \frac{1}{2}s(s + 1)(\mu + 1) - 1$ is the block graph of a linear space or transversal design.*

It follows immediately (from this and the μ -bound) that

Theorem 8.1.9 *For any fixed $s = -m$ there are only finitely many primitive strongly regular graphs with smallest eigenvalue s , that are not the block graph of a linear space or transversal design with $t + 1 = -m$.* \square

8.2 Strongly regular graphs with eigenvalue -2

For later use we give Seidel’s classification [274] of the strongly regular graphs with $s = -2$.

Theorem 8.2.1 *Let Γ be a strongly regular graph with smallest eigenvalue -2 . Then Γ is one of*

- (i) *the complete n -partite graph $K_{n \times 2}$, with parameters $(v, k, \lambda, \mu) = (2n, 2n - 2, 2n - 4, 2n - 2)$, $n \geq 2$,*
- (ii) *the lattice graph $L_2(n) = K_n \times K_n$, with parameters $(v, k, \lambda, \mu) = (n^2, 2(n - 1), n - 2, 2)$, $n \geq 3$,*
- (iii) *the Shrikhande graph, with parameters $(v, k, \lambda, \mu) = (16, 6, 2, 2)$,*
- (iv) *the triangular graph $T(n)$ with parameters $(v, k, \lambda, \mu) = (\binom{n}{2}, 2(n - 2), n - 2, 4)$, $n \geq 5$,*
- (v) *one of the three Chang graphs, with parameters $(v, k, \lambda, \mu) = (28, 12, 6, 4)$,*

(vi) the Petersen graph, with parameters $(v, k, \lambda, \mu) = (10, 3, 0, 1)$,


(vii) the Clebsch graph, with parameters $(v, k, \lambda, \mu) = (16, 10, 6, 6)$,

(viii) the Schläfli graph, with parameters $(v, k, \lambda, \mu) = (27, 16, 10, 8)$.

Proof. If Γ is imprimitive, then we have case (i). Otherwise, the μ -bound gives $\mu \leq 8$, and the rationality conditions give $(r+2) | (\mu-2)(\mu-4)$ and integrality of v gives $\mu | 2r(r+1)$. For $\mu = 2$ we find the parameters of $L_2(n)$, for $\mu = 4$ those of $T(n)$, and for the remaining values for μ only the parameter sets $(v, k, \lambda, \mu) = (10, 3, 0, 1)$, $(16, 10, 6, 6)$, and $(27, 16, 10, 8)$ survive the parameter conditions and the absolute bound. It remains to show that the graph is uniquely determined by its parameters in each case. Now Shrikhande [283] proved uniqueness of the graph with $L_2(n)$ parameters, with the single exception of $n = 4$, where there is one more graph, now known as the Shrikhande graph, and Chang [79, 80] proved uniqueness of the graph with $T(n)$ parameters, with the single exception of $n = 8$, where there are three more graphs, now known as the Chang graphs. In the remaining three cases uniqueness is easy to see. \square

Let us give definitions for the graphs involved.

The *Shrikhande graph* is the result of Seidel switching the lattice graph $L_2(4)$ with respect to an induced circuit of length 8. It is the complement of the Latin square graph for the cyclic Latin square of order 4. It is locally a hexagon.



The three *Chang graphs* are the result of switching $T(8)$ (the line graph of K_8) with respect to (a) a 4-coclique $\overline{K_4}$, that is, 4 pairwise disjoint edges in K_8 ; (b) $K_3 + K_5$, that is, 8 edges forming a triangle and a (disjoint) pentagon in K_8 ; (c) the line graph of the cubic graph formed by an 8-circuit plus edges between opposite vertices.

The *Clebsch graph* is the complement of the folded 5-cube.

The *Schläfli graph* is the collinearity graph of $GQ(2, 4)$ (cf. §8.6).

8.3 Connectivity

For a graph Γ , let $\Gamma_i(x)$ denote the set of vertices at distance i from x in Γ . Instead of $\Gamma_1(x)$ we write $\Gamma(x)$.

Proposition 8.3.1 *If Γ is a primitive strongly regular graph, then for each vertex x the subgraph $\Gamma_2(x)$ is connected.*

Proof. Note that $\Gamma_2(x)$ is regular of valency $k - \mu$. If it is not connected, then its eigenvalue $k - \mu$ would have multiplicity at least two, and hence would be not larger than the second largest eigenvalue r of Γ . Then $x^2 + (\mu - \lambda)x + \mu - k \leq 0$ for $x = k - \mu$, i.e., $(k - \mu)(k - \lambda - 1) \leq 0$, contradiction. \square

The *vertex connectivity* $\kappa(\Gamma)$ of a connected non-complete graph Γ is the smallest integer m such that Γ can be disconnected by removing m vertices.

Theorem 8.3.2 ([57]). *Let Γ be a connected strongly regular graph of valency k . Then $\kappa(\Gamma) = k$, and the only disconnecting sets of size k are the sets of all neighbors of some vertex x .*

Proof. Clearly, $\kappa(\Gamma) \leq k$. Let S be a disconnecting set of vertices not containing all neighbors of some vertex. Let $\Gamma \setminus S = A + B$ be a separation of $\Gamma \setminus S$. Since the eigenvalues of $A \cup B$ interlace those of Γ , it follows that at least one of A and B , say B , has largest eigenvalue at most r . It follows that the average valency of B is at most r . Since B has an edge, $r > 0$.

Now let $|S| \leq k$. Since B has average valency at most r , we can find two points x, y in B such that $|S \cap \Gamma(x)| + |S \cap \Gamma(y)| \geq 2(k - r)$, so that these points have at least $k - 2r$ common neighbors in S .

If Γ has nonintegral eigenvalues, then we have $(v, k, \lambda, \mu) = (4t + 1, 2t, t - 1, t)$ for some t , and $r = (-1 + \sqrt{v})/2$. The inequality $\max(\lambda, \mu) \geq k - 2r$ gives $t \leq 2$, but for $t = 2$ the eigenvalues are integral, so we have $t = 1$ and Γ is the pentagon. But the claim is true in that case.

Now let r, s be integral. If $s \leq -3$, then $\mu = k + rs \leq k - 3r$ and $\lambda = \mu + r + s \leq k - 2r - 3$, so that no two points can have $k - 2r$ common neighbors.

Therefore $s = -2$, and we have one of the eight cases in Seidel's classification. But not case (i), since $r > 0$.

Since both A and B contain an edge, both B and A have size at most $\bar{\mu} = v - 2k + \lambda$, so that both A and B have size at least $k - \lambda$, and $v \geq 3k - 2\lambda$. This eliminates cases (vii) and (viii).

If B is a clique, then $|B| \leq r + 1 = k - \lambda - 1$, contradiction. So, B contains two nonadjacent vertices, and their neighbors must be in $B \cup S$, so $2k - \mu \leq |B| + |S| - 2$ and $k - \mu + 2 \leq |B| \leq \bar{\mu}$.

In cases (iii), (v), (vi) we have $\bar{\mu} = k - \mu + 2$, so equality holds and $|B| = \bar{\mu}$ and $|S| = k$. Since $v < 2\bar{\mu} + k$, we have $|A| < \bar{\mu}$ and A must be a clique (of size $v - k - \bar{\mu} = k - \lambda$). But the Petersen graph does not contain a 3-clique, and the Shrikhande graph does not contain 4-cliques; also, if A is a 6-clique in a Chang graph, and $a, b, c \in A$, then $\Gamma(a) \cap S$, $\Gamma(b) \cap S$, and $\Gamma(c) \cap S$ are three 7-sets in the 12-set S that pairwise meet in precisely two points, impossible. This eliminates cases (iii), (v), (vi).

We are left with the two infinite families of lattice graphs and triangular graphs. In both cases it is easy to see that if x, y are nonadjacent, then there exist k paths joining x and y , vertex disjoint apart from x, y , and entirely contained in $\{x, y\} \cup \Gamma(x) \cup \Gamma(y)$. Hence $|S| = k$, and if S separates x, y then $S \subseteq \Gamma(x) \cup \Gamma(y)$.

The subgraph $\Delta := \Gamma \setminus (\{x, y\} \cup \Gamma(x) \cup \Gamma(y))$ is connected (if one removes a point and its neighbors from a lattice graph, the result is a smaller lattice graph, and the same holds for a triangular graph), except in the case of the triangular graph $\binom{5}{2}$ where Δ is empty.

Each vertex of $(\Gamma(x) \cup \Gamma(y)) \setminus S$ has a neighbor in Δ and we find a path of length 4 disjoint from S joining x and y , except in the case of the triangular graph $\binom{5}{2}$, where each vertex of $\Gamma(x) \setminus S$ is adjacent to each vertex of $\Gamma(y) \setminus S$, and we find a path of length 3 disjoint from S joining x and y . \square

We remark that it is not true that for every strongly regular graph Γ with vertex x the vertex connectivity of the subgraph $\Gamma_2(x)$ equals its valency $k - \mu$. A counterexample is given by the graph Γ that is the complement of the strongly regular graph Δ with parameters $(96, 19, 2, 4)$ constructed by Haemers for $q = 4$,

see [175], p. 76 or [56], §8A. Indeed, we have $\Delta(x) \cong K_3 + 4C_4$, so that $\Gamma_2(x)$ has degree 16 and vertex connectivity 15.

Erdős & Chvátal [88] showed that if a graph Γ on at least 3 vertices has vertex connectivity κ and largest independent set of size α , and $\alpha \leq \kappa$ then Γ has a Hamiltonian circuit. Bigalke & Jung [25] showed that if Γ is 1-tough, with $\alpha \leq \kappa + 1$ and $\kappa \geq 3$, and Γ is not the Petersen graph, then Γ is Hamiltonian. Such results imply for example that if Γ is strongly regular with smallest eigenvalue s , and s is not integral, or $-s \leq \mu + 1$, then Γ is Hamiltonian. This, together with explicit inspection of the Hoffman-Singleton graph, the Gewirtz graph, and the M_{22} graph, shows that all connected strongly regular graphs on fewer than 99 vertices are Hamiltonian, except for the Petersen graph.

8.4 Cocliques and colorings

In §2.5 we have derived some bounds the size of a coclique in terms of eigenvalues. These bounds are especially useful for strongly regular graphs. Moreover, strongly regular graphs for which the bounds of Hoffman and Cvetković are tight have a very special structure:

Theorem 8.4.1 *Let Γ be a strongly regular graph with eigenvalues k (degree), r and s ($r > s$) and multiplicities 1, f and g , respectively. Suppose that Γ is not complete multi-partite (i.e. $r \neq 0$) and let C be a coclique in Γ . Then*

- (i) $|C| \leq g$,
- (ii) $|C| \leq ns/(s - k)$,
- (iii) *if $|C| = g = ns/(s - k)$, then the subgraph Γ' of Γ induced by the vertices which are not in C , is strongly regular with eigenvalues $k' = k + s$ (degree), $r' = r$ and $s' = r + s$ and respective multiplicities 1, $f - g + 1$ and $g - 1$.*

Proof. Parts (i) and (ii) follow from Theorems 4.1.1 and 4.1.2. Assume $|C| = g = ns/(s - k)$. By Theorem 2.5.4, Γ' is regular of degree $k + s$. Apply Lemma 2.11.1 to $P = A - \frac{k-r}{n}J$, where A is the adjacency matrix of Γ . Since Γ is regular, A and J commute and therefore P has eigenvalues r and s with multiplicities $f + 1$ and g , respectively. We take $Q = -\frac{k-r}{n}J$ of size $|C| = g$ and $R = A' - \frac{k-r}{n}J$, where A' is the adjacency matrix of Γ' . Lemma 2.11.1 gives the eigenvalues of R : r ($f + 1 - g$ times), s (0 times), $r + s$ ($g - 1$ times) and $r + s + g(k - r)/n$ (1 time). Since Γ' is regular of degree $k + s$ and A' commutes with J we obtain the required eigenvalues for A' . By Theorem 8.1.2 Γ' is strongly regular. \square

For instance, an $(m - 1)$ -coclique in $\overline{T(m)}$ is tight for both bounds and the graph on the remaining vertices is $T(m - 1)$.

Also for the chromatic number we can say more in case of a strongly regular graph.

Theorem 8.4.2 *If Γ is a primitive strongly regular graph, not the pentagon, then*

$$\chi(\Gamma) \geq 1 - \frac{s}{r}.$$

Proof. Since Γ is primitive, $r > 0$ and by Corollary 4.2.4, it suffices to show that the multiplicity g of s satisfies $g \geq -s/r$ for all primitive strongly regular graphs but the pentagon. First we check this claim for all feasible parameter sets with at most 23 vertices. Next we consider strongly regular graphs with $v \geq 24$ and $r < 2$. The complements of these graphs have $s > -3$, and by Theorem 8.1.3 (iv), $s = -2$. By use of Theorem 8.2 we easily find that all these graphs satisfy the claim.

Assume that Γ is primitive, that $r \geq 2$, and that the claim does not hold (that is $g < -s/r$). Now $(v - 1 - g)r + gs + k = 0$ gives

$$g^2 < -sg/r = v - 1 - g + k/r \leq v - 1 - g + k/2 < 3v/2 - g.$$

This implies $g(g + 3) \leq 3v/2 = 2\sqrt{3v/2}$. By use of the absolute bound $v \leq g(g + 3)/2$, we get $v/2 < 2\sqrt{3v/2}$, so $v < 24$. Contradiction. \square

For example if Γ is the complement of the triangular graph $T(m)$ then Γ is strongly regular with eigenvalues $k = \frac{1}{2}(m-2)(m-3)$, $r = 1$ and $s = 3 - m$ (for $m \geq 4$). The above bound gives $\chi(\Gamma) \geq m - 2$, which is tight, whilst Hoffman's lower bound (Theorem 4.2.2) equals $\frac{1}{2}m$. On the other hand, if m is even, Hoffman's bound is tight for the complement of Γ whilst the above bound is much less. We saw (see §4.2) that a Hoffman coloring (i.e. a coloring with $1 - k/s$ classes) corresponds to an equitable partition of the adjacency matrix. For the complement this gives an equitable partition into maximal cliques, which is called a spread of the strongly regular graph. For more application of eigenvalues to the chromatic number we refer to [140] and [158]. See also §8.7.

8.5 Automorphisms

Let A be the adjacency matrix of a graph Γ , and P the permutation matrix that describes an automorphism ϕ of Γ . Then $AP = PA$. If ϕ has order m , then $P^m = I$, so that the eigenvalues of AP are m -th roots of unity times eigenvalues of A .

Apply this in the special case of strongly regular graphs. Suppose ϕ has f fixed points, and moves g points to a neighbor. Then $f = \text{tr } P$ and $g = \text{tr } AP$. Now consider $M = A - sI$. It has spectrum $(k - s)^1, (r - s)^f, 0^g$ with multiplicities written as exponents. Hence MP has eigenvalues $k - s, (r - s)\zeta$ for certain m -th roots of unity ζ , and 0. It follows that $g - sf = \text{tr } MP \equiv k - s \pmod{r - s}$.

For example, for the Petersen graph every automorphism satisfies $f \equiv g + 1 \pmod{3}$.

For example, for a hypothetical Moore graph on 3250 vertices (cf. §10.5.1), every automorphism satisfies $8f + g \equiv 5 \pmod{15}$.

In some cases, where a structure is given locally, it must either be a universal object, or a quotient, where the quotient map preserves local structure, that is, only identifies points that are far apart. In the finite case arguments like those in this section can be used to show that $f = g = 0$ is impossible, so that nontrivial quotients do not exist. For an example, see [47].

8.6 Generalized quadrangles

A *generalized n -gon* is a connected bipartite graph of diameter n and girth $2n$. (The girth of a graph is the length of a shortest circuit.)

It is common to call the vertices in one color class of the unique 2-coloring *points*, and the other vertices *lines*. For example, a generalized 3-gon is the same thing as a projective plane: any two points have an even distance at most 3, hence are joined by a line, and similarly any two lines meet in a point; finally two lines cannot meet in two points since that would yield a quadrangle, but the girth is 6.

A *generalized quadrangle* is a generalized 4-gon. In terms of points and lines, the definition becomes: a *generalized quadrangle* is an incidence structure (P, L) with set of points P and set of lines L , such that two lines meet in at most one point, and if p is a point not on the line m , then there is a unique point q on m and a unique line n on p such that q is on n .

8.6.1 Parameters

A generalized n -gon is called *firm* (*thick*) when each vertex has at least 2 (resp. 3) neighbors, that is, when each point is on at least two (three) lines, and each line is on at least two (three) points.

An example of a non-firm generalized quadrangle is a pencil of lines on one common point x_0 . Each point different from x_0 is on a unique line, and $\Gamma_3(x_0) = \emptyset$.

Proposition 8.6.1 (i) *If a generalized n -gon Γ has a pair of opposite vertices x, y where x has degree at least two, then every vertex has an opposite, and Γ is firm.*

(ii) *A thick generalized n -gon has parameters: each line has the same number of points, and each point is on the same number of lines. When moreover n is odd then the number of points on each line equals the number of lines through each point.*

Proof. For a vertex x of a generalized n -gon, let $k(x)$ be its degree. Call two vertices of a generalized n -gon *opposite* when they have distance n . If x and y are opposite then each neighbor of one is on a unique shortest path to the other, and we find $k(x) = k(y)$.

(i) Being non-opposite gives a bijection between $\Gamma(x)$ and $\Gamma(y)$, and hence if $k(x) > 1$ then also each neighbor z of x has an opposite and satisfies $k(z) > 1$. Since Γ is connected, it is firm.

(ii) Let x, z be two points joined by the line y . Let w be opposite to y . Since $k(w) > 2$ there is a neighbor u of w opposite to both x and z . Now $k(x) = k(u) = k(z)$. Since Γ is connected and bipartite this shows that $k(p)$ is independent of the point p . If n is odd, then a vertex opposite a point is a line. \square

A firm, non-thick generalized quadrangle is the vertex-edge incidence graph of a complete bipartite graph.

The *halved graph* of a bipartite graph Γ , is the graph on the same vertex set, where two vertices are adjacent when they have distance 2 in Γ . The *point graph* and *line graph* of a generalized n -gon are the two components of its halved graph containing the points and lines, respectively.

The point graph and line graph of a finite thick generalized n -gon are distance-regular of diameter $\lfloor n/2 \rfloor$ (see Chapter 11). In particular, the point graph and line graph of a thick generalized quadrangle are strongly regular (see Theorem 8.6.2).

It is customary to let $GQ(s, t)$ denote a finite generalized quadrangle with $s + 1$ points on each line and $t + 1$ lines on each point. Note that it is also customary to use s to denote the smallest eigenvalue of a strongly regular graph, so in this context one has to be careful to avoid confusion.

It is a famous open problem whether a thick generalized n -gon can have finite s and infinite t . In the special case of generalized quadrangles a little is known: Cameron, Kantor, Brouwer, and Cherlin [74, 44, 81] show that this cannot happen for $s + 1 \leq 5$.

8.6.2 Constructions of generalized quadrangles

Suppose V is a vector space provided with a nondegenerate quadratic form f of Witt index 2 (that is, such that the maximal totally singular subspaces have vector space dimension 2). Consider in the projective space PV the singular projective points and the totally singular projective lines. These will form a generalized quadrangle.

Indeed, f defines a bilinear form B on V via $B(x, y) = f(x+y) - f(x) - f(y)$. Call x and y *orthogonal* when $B(x, y) = 0$. When two singular vectors are orthogonal, the subspace spanned by them is totally singular. And conversely, in a totally singular subspace any two vectors are orthogonal. The collection of all vectors orthogonal to a given vector is a hyperplane. We have to check that if $P = \langle x \rangle$ is a singular projective point, and L is a totally singular projective line not containing P , then P has a unique neighbor on L . But the hyperplane of vectors orthogonal to x meets L , and cannot contain L otherwise f would have larger Witt index.

This construction produces generalized quadrangles over arbitrary fields. If V is a vector space over a finite field \mathbb{F}_q , then a nondegenerate quadratic form can have Witt index 2 in dimensions 4, 5, and 6. A hyperbolic quadric in 4 dimensions yields a generalized quadrangle with parameters $GQ(q, 1)$, a parabolic quadric in 5 dimensions yields a generalized quadrangle with parameters $GQ(q, q)$, and an elliptic quadric in 6 dimensions yields a generalized quadrangle with parameters $GQ(q, q^2)$.

Other constructions, and other parameters occur.

In the below we'll meet $GQ(2, t)$ for $t = 1, 2, 4$ and $GQ(3, 9)$. Let us give simple direct descriptions for $GQ(2, 1)$ and $GQ(2, 2)$.

The unique $GQ(2, 1)$ is the 3-by-3 grid: 9 points, 6 lines. Its point graph is $K_3 \times K_3$.

The unique $GQ(2, 2)$ is obtained by taking as points the 15 pairs from a 6-set, and as lines the 15 partitions of that 6-set into three pairs. Now collinearity is being disjoint. Given a point ac , and a line $\{ab, cd, ef\}$, the two points ab and cd on this line are not disjoint from ac , so that ef is the unique point on this line collinear with ac , and the line joining ac and ef is $\{ac, bd, ef\}$.

8.6.3 Strongly regular graphs from generalized quadrangles

As mentioned before, the point graph (collinearity graph) of a finite thick generalized quadrangle is strongly regular. The parameters and eigenvalue can be obtained in a straightforward way (see exercises).

Theorem 8.6.2 *The collinearity graph of a finite generalized quadrangle with parameters $GQ(s, t)$ is strongly regular with parameters*

$$v = (s + 1)(st + 1), \quad k = s(t + 1), \quad \lambda = s - 1, \quad \mu = t + 1$$

and spectrum

$$\begin{array}{ll} s(t + 1) & \text{with multiplicity } 1, \\ s - 1 & \text{with multiplicity } st(s + 1)(t + 1)/(s + t), \\ -t - 1 & \text{with multiplicity } s^2(st + 1)/(s + t). \end{array}$$

In particular, if a $GQ(s, t)$ exists, then $(s + t) | s^2(st + 1)$.

8.6.4 Generalized quadrangles with lines of size 3

Let a *weak generalized quadrangle* be a point-line geometry with the properties that two lines meet in at most one point, and given a line m and a point p outside there is a unique pair (q, n) such that $p \sim n \sim q \sim m$, where \sim denotes incidence. The difference with the definition of a generalized quadrangle is that connectedness is not required. (But of course, as soon as there is a point and a line then the geometry is connected.)

Theorem 8.6.3 *A weak generalized quadrangle where all lines have size 3 is one of the following:*

- (i) a coclique (no lines),
- (ii) a pencil (all lines passing through a fixed point),
- (iii) the unique $GQ(2, 1)$,
- (iv) the unique $GQ(2, 2)$,
- (v) the unique $GQ(2, 4)$.

Proof. After reducing to the case of $GQ(2, t)$ one finds $(t + 2) | (8t + 4)$, i.e., $(t + 2) | 12$, i.e., $t \in \{1, 2, 4, 10\}$, and $t = 10$ is ruled out by the Krein conditions. Alternatively, or afterwards, notice that the point graphs have eigenvalue 1, so that their complements have smallest eigenvalue -2 , and apply Seidel's classification. Cases (iii), (iv), (v) here have point graphs that are the complements of the lattice graph $K_3 \times K_3$, the triangular graph $T(6)$, and the Schläfli graph, respectively. \square

This theorem can be used in the classification of root lattices, where the five cases correspond to A_n , D_n , E_6 , E_7 , E_8 , respectively.

And the classification of root lattices can be used in the classification of graphs with smallest eigenvalue -2 . Indeed, for such graphs $A + 2I$ is positive semidefinite, and one can represent these graphs by vectors in a Euclidean space such that $(x, y) = 2, 1, 0$ when x and y are equal, adjacent, nonadjacent, respectively. The lattice spanned by these vectors is a root lattice.

8.7 The (81,20,1,6) strongly regular graph

Large parts of this section are taken from [51]. Sometimes the graph of this section is called the *Brouwer-Haemers graph*.

Let $\Gamma = (X, E)$ be a strongly regular graph with parameters $(v, k, \lambda, \mu) = (81, 20, 1, 6)$. Then Γ has spectrum $\{20^1, 2^{60}, -7^{20}\}$, where the exponents denote multiplicities. We will show that up to isomorphism there is a unique such graph Γ . More generally we give a short proof for the fact (due to Ivanov & Shpectorov [206]) that a strongly regular graph with parameters $(v, k, \lambda, \mu) = (q^4, (q^2 + 1)(q - 1), q - 2, q(q - 1))$ that is the collinearity graph of a partial quadrangle (that is, in which all maximal cliques have size q) is the second subconstituent of the collinearity graph of a generalized quadrangle $GQ(q, q^2)$. In the special case $q = 3$ this will imply our previous claim, since $\lambda = 1$ implies that all maximal cliques have size 3, and it is known (see Cameron, Goethals & Seidel [76]) that there is a unique generalized quadrangle $GQ(3, 9)$ (and this generalized quadrangle has an automorphism group transitive on the points).

8.7.1 Descriptions

Let us first give a few descriptions of our graph on 81 vertices. Note that the uniqueness shows that all constructions below give isomorphic graphs, something, which is not immediately obvious from the description in all cases.

A. Let X be the point set of $AG(4, 3)$, the 4-dimensional affine space over \mathbf{F}_3 , and join two points when the line connecting them hits the hyperplane at infinity (a $PG(3, 3)$) in a fixed elliptic quadric Q . This description shows immediately that $v = 81$ and $k = 20$ (since $|Q| = 10$). Also $\lambda = 1$ since no line meets Q in more than two points, so that the affine lines are the only triangles. Finally $\mu = 6$, since a point outside Q in $PG(3, 3)$ lies on 4 tangents, 3 secants and 6 exterior lines with respect to Q , and each secant contributes 2 to μ . We find that the group of automorphisms contains $G = 3^4 \cdot PGO_4^- \cdot 2$, where the last factor 2 accounts for the linear transformations that do not preserve the quadratic form Q , but multiply it by a constant. In fact this is the full group, as will be clear from the uniqueness proof.

B. A more symmetric form of this construction is found by starting with $X = \mathbf{1}^\perp / \langle \mathbf{1} \rangle$ in \mathbf{F}_3^6 provided with the standard bilinear form. The corresponding quadratic form $(Q(x) = \text{wt}(x))$, the number of nonzero coordinates of x is elliptic, and if we join two vertices $x + \langle \mathbf{1} \rangle, y + \langle \mathbf{1} \rangle$ of X when $Q(x - y) = 0$, i.e., when their difference has weight 3, we find the same graph as under A. This construction shows that the automorphism group contains $G = 3^4 \cdot (2 \times \text{Sym}(6)) \cdot 2$, and again this is the full group.

C. There is a unique strongly regular graph Γ with parameters $(112, 30, 2, 10)$, the collinearity graph of the unique generalized quadrangle with parameters $GQ(3, 9)$. Its second subconstituent is an $(81, 20, 1, 6)$ strongly regular graph, and hence isomorphic to our graph Γ . (See Cameron, Goethals & Seidel [76].) We find that $\text{Aut } \Gamma$ contains (and in fact it equals) the point stabilizer in $U_4(3) \cdot D_8$ acting on $GQ(3, 9)$.

D. The graph Γ is the coset graph of the truncated ternary Golay code C : take the 3^4 cosets of C and join two cosets when they contain vectors differing in only one place.

E. The graph Γ is the Hermitean forms graph on \mathbf{F}_9^2 ; more generally, take the q^4 matrices M over \mathbf{F}_{q^2} satisfying $M^\top = \overline{M}$, where $\overline{}$ denotes the field automorphism $x \rightarrow x^q$ (applied entrywise), and join two matrices when their difference has rank 1. This will give us a strongly regular graph with parameters $(v, k, \lambda, \mu) = (q^4, (q^2 + 1)(q - 1), q - 2, q(q - 1))$.

F. The graph Γ is the graph with vertex set \mathbf{F}_{81} , where two vertices are joined when their difference is a fourth power. (This construction was given by Van Lint & Schrijver [228].)

8.7.2 Uniqueness

Now let us embark upon the uniqueness proof. Let $\Gamma = (X, E)$ be a strongly regular graph with parameters $(v, k, \lambda, \mu) = (q^4, (q^2 + 1)(q - 1), q - 2, q(q - 1))$ and assume that all maximal cliques (we shall just call them lines) of Γ have size q . Let Γ have adjacency matrix A . Using the spectrum of A - it is $\{k^1, (q - 1)^f, (q - 1 - q^2)^g\}$, where $f = q(q - 1)(q^2 + 1)$ and $g = (q - 1)(q^2 + 1)$ - we can obtain some structure information. Let \mathbf{T} be the collection of subsets of X of cardinality q^3 inducing a subgraph that is regular of degree $q - 1$.

1. **Claim.** *If $T \in \mathbf{T}$, then each point of $X \setminus T$ is adjacent to q^2 points of T .*
Look at the matrix B of average row sums of A , with sets of rows and columns partitioned according to $\{T, X \setminus T\}$. We have

$$B = \begin{bmatrix} q - 1 & q^2(q - 1) \\ q^2 & k - q^2 \end{bmatrix}$$

with eigenvalues k and $q - 1 - q^2$, so interlacing is tight, and by Corollary 2.5.4(ii) it follows that the row sums are constant in each block of A .

2. **Claim.** *Given a line L , there is a unique $T_L \in \mathbf{T}$ containing L .*
Let Z be the set of vertices in $X \setminus L$ without a neighbor in L . Then $|Z| = q^4 - q - q(k - q + 1) = q^3 - q$. Let $T = L \cup Z$. Each vertex of Z is adjacent to $q\mu = q^2(q - 1)$ vertices with a neighbor in L , so T induces a subgraph that is regular of degree $q - 1$.
3. **Claim.** *If $T \in \mathbf{T}$ and $x \in X \setminus T$, then x is on at least one line L disjoint from T , and T_L is disjoint from T for any such line L .*
The point x is on $q^2 + 1$ lines, but has only q^2 neighbors in T . Each point of L has q^2 neighbors in T , so each point of T has a neighbor on L and hence is not in T_L .

4. **Claim.** *Any $T \in \mathbf{T}$ induces a subgraph Δ isomorphic to $q^2 K_q$.*
It suffices to show that the multiplicity m of the eigenvalue $q - 1$ of Δ is (at least) q^2 (it cannot be more). By interlacing we find $m \geq q^2 - q$, so we need some additional work. Let $M := A - (q - 1/q^2)J$. Then M has spectrum $\{(q - 1)^{f+1}, (q - 1 - q^2)^g\}$, and we want that M_T , the submatrix of M with rows and columns indexed by T , has eigenvalue $q - 1$ with multiplicity (at

least) $q^2 - 1$, or, equivalently (by Lemma 2.11.1), that $M_{X \setminus T}$ has eigenvalue $q - 1 - q^2$ with multiplicity (at least) $q - 2$. But for each $U \in \mathbf{T}$ with $U \cap T = \emptyset$ we find an eigenvector $x_U = (2 - q)\chi_U + \chi_{X \setminus (T \cup U)}$ of $M_{X \setminus T}$ with eigenvalue $q - 1 - q^2$. A collection $\{x_U | U \in \mathbf{U}\}$ of such eigenvectors cannot be linearly dependent when $\mathbf{U} = \{U_1, U_2, \dots\}$ can be ordered such that $U_i \not\subset \bigcup_{j < i} U_j$ and $\bigcup \mathbf{U} \neq X \setminus T$, so we can find (using Claim 3) at least $q - 2$ linearly independent such eigenvectors, and we are done.

5. **Claim.** Any $T \in \mathbf{T}$ determines a unique partition of X into members of \mathbf{T} .

Indeed, we saw this in the proof of the previous step.

Let Π be the collection of partitions of X into members of \mathbf{T} . We have $|\mathbf{T}| = q(q^2 + 1)$ and $|\Pi| = q^2 + 1$. Construct a generalized quadrangle $GQ(q, q^2)$ with point set $\{\infty\} \cup \mathbf{T} \cup X$ as follows: The $q^2 + 1$ lines on ∞ are $\{\infty\} \cup \pi$ for $\pi \in \Pi$. The q^2 remaining lines on each $T \in \mathbf{T}$ are $\{T\} \cup L$ for $L \subset T$. It is completely straightforward to check that we really have a generalized quadrangle $GQ(q, q^2)$.

8.7.3 Independence and chromatic numbers

We have $\alpha(\Gamma) = 15$ and $\chi(\Gamma) = 7$.

Clearly, the independence number of our graph is one less than the independence number of the unique $GQ(3, 9)$ of which it is the second subconstituent. So it suffices to show that $\alpha(\Delta) = 16$, where Δ is the collinearity graph of $GQ(3, 9)$.

It is easy to indicate a 16-coclique: define $GQ(3, 9)$ in $PG(5, 3)$ provided with the nondegenerate elliptic quadratic form $\sum_{i=1}^6 x_i^2$. There are 112 isotropic points, 80 of weight 3 and 32 of weight 6. Among the 32 of weight 6, 16 have coordinate product 1, and 16 have coordinate product -1 , and these two 16-sets are cocliques.

That there is no larger coclique can be seen by cubic counting.

Let C be a 16-coclique in Δ . Let there be n_i vertices outside that have i neighbors inside. Then

$$\sum n_i = 96, \quad \sum i n_i = 480, \quad \sum \binom{i}{2} n_i = 1200, \quad \sum \binom{i}{3} n_i = 2240,$$

so that

$$\sum (i - 4)^2 (i - 10) n_i = 0.$$

(Here the quadratic counting is always possible in a strongly regular graph, and the last equation can be written because the second subconstituent is itself strongly regular.) Now each point is on 10 lines, and hence cannot have more than 10 neighbors in C . It follows that each point has either 4 or 10 neighbors in C . In particular, C is maximal.

As an aside: Solving these equations gives $n_4 = 80$, $n_{10} = 16$. Let D be the set of 16 vertices with 10 neighbors in C . If two vertices $d_1, d_2 \in D$ are adjacent then they can have only 2 common neighbors in C , but each has 10 neighbors in C , contradiction. So, also D is a 16-coclique, which means that 16-cocliques in Δ come in pairs.

Since $81/15 > 5$, we have $\chi(\Gamma) \geq 6$. Since Δ has a split into two Gewirtz graphs, and the Gewirtz graph has chromatic number 4, it follows that $\chi(\Delta) \leq 8$. (And

in fact equality holds.) This shows that for our graph $6 \leq \chi(\Gamma) \leq 8$. In fact $\chi(\Gamma) = 7$ can be seen by computer (Edwin van Dam, pers. comm.).

Since $\lambda = 1$, the maximum clique size equals 3. And from the uniqueness proof it is clear that Γ admits a partition into 27 triangles. So the complement of Γ has chromatic number 27.

8.8 Strongly regular graphs and 2-weight codes

8.8.1 Codes, graphs and projective sets

In this section we show the equivalence of three kinds of objects:

- (i) projective two-weight codes,
- (ii) subsets X of a projective space such that $|X \cap H|$ takes two values when H ranges through the hyperplanes of the projective space,
- (iii) strongly regular graphs defined by a difference set that is a cone in a vector space.

This equivalence is due to Delsarte [126]. An extensive survey of this material was given by Calderbank & Kantor [73].

A *linear code* is a linear subspace of some finite vector space with fixed basis. For basic terminology and results on codes, see MacWilliams & Sloane [236] and Van Lint [227]. A linear code C is called *projective* when its dual C^\perp has minimum weight at least three, that is, when no two coordinate positions of C are linearly dependent. The *weight* of a vector is its number of nonzero coordinates. A *two-weight code* is a linear code in which precisely two nonzero weights occur.

Let us first discuss the correspondence between linear codes and subsets of projective spaces.

8.8.2 The correspondence between linear codes and subsets of a projective space

A linear code C of word length n over the alphabet \mathbb{F}_q is a linear subspace of the vector space \mathbb{F}_q^n . The *weight* of a vector is its number of nonzero coordinates. We call C an $[n, m, w]$ -code if C has dimension m and minimum nonzero weight w . We say that C has *effective length* (or *support*) $n - z$ when there are precisely z coordinate positions j such that $c_j = 0$ for all $c \in C$. The *dual* C^\perp of a code C is the linear code $\{d \in \mathbb{F}_q^n \mid \langle c, d \rangle = 0 \text{ for all } c \in C\}$, where $\langle c, d \rangle = \sum c_i d_i$ is the standard inner product (bilinear form).

Let us call two linear codes of length n over \mathbb{F}_q *equivalent* when one arises from the other by permutation of coordinates or multiplication of coordinates by a nonzero constant. E.g., the \mathbb{F}_3 -codes generated by $\begin{pmatrix} 1111 \\ 0012 \end{pmatrix}$ and $\begin{pmatrix} 1212 \\ 1100 \end{pmatrix}$ are equivalent. If we study codes up to equivalence, and assume that n is chosen minimal, i.e., that the generator matrix has no zero columns, we may identify the set of columns in a $m \times n$ generator matrix with points of a projective space

$PG(m-1, q)$. In this way, we find a subset X of $PG(m-1, q)$, possibly with repeated points, or, if you prefer, a weight function $w : PG(m-1, q) \rightarrow \mathbb{N}$.

Choosing one code in an equivalence class means choosing a representative in \mathbb{F}_q^m for each $x \in X$, and fixing an order on X . Now the code words can be identified with the linear functionals f , and the x -coordinate position is $f(x)$.

Clearly, the code has word length $n = |X|$. Note that the code will have dimension m if and only if X spans $PG(m-1, q)$, i.e., if and only if X is not contained in a hyperplane.

The weight of the code word f equals the number of x such that $f(x) \neq 0$. But a nonzero f vanishes on a hyperplane of $PG(m-1, q)$. Consequently, the number of words of nonzero weight w in the code equals $q-1$ times the number of hyperplanes H that meet X in $n-w$ points. In particular the minimum distance of the code is n minus the maximum size of $H \cap X$ for a hyperplane H .

The minimum weight of the dual code equals the minimum number of points of X that are dependent. So, it is 2 if and only if X has repeated points, and 3 when X has no repeated points but has three collinear points.

Example Take for X the entire projective space $PG(m-1, q)$, so that $n = |X| = (q^m-1)/(q-1)$. We find the so-called simplex code: all words have weight q^{m-1} , and we have an $[n, m, q^{m-1}]$ -code over \mathbb{F}_q . Its dual is the $[n, n-m, 3]$ Hamming code. It is perfect!

8.8.3 The correspondence between projective two-weight codes, subsets of a projective space with two intersection numbers, and affine strongly regular graphs

Given a subset X of size n of $PG(m-1, q)$, let us define a graph Γ with vertex set \mathbb{F}_q^m , with $x \sim y$ if and only if $\langle y-x \rangle \in X$. Then clearly Γ is regular of valency $k = (q-1)n$. We show below that this graph has eigenvalues $k - qw_i$ when the linear code has weights w_i . Hence if a linear code has only two nonzero weights, and its dual has minimum weight at least 3, then we have a strongly regular graph.

Let us look at the details.

Let $F = \mathbb{F}_q$ and $K = \mathbb{F}_{q^k}$ and let $\text{tr} : K \rightarrow F$ be the trace map defined by $\text{tr}(x) = x + x^q + \dots + x^{q^{k-1}}$. Then the F -linear maps $f : F \rightarrow K$ are precisely the maps f_a defined by $f_a(x) = \text{tr}(ax)$, for $a \in K$. If $\text{tr}(ax) = 0$ for all x , then $a = 0$.

(Indeed, first of all, these maps are indeed F -linear. If $a \neq 0$, then $\text{tr}(ax)$ is a polynomial of degree q^{k-1} and cannot have q^k zeros. It follows that we find q^k distinct maps f_a . But this is the total number of F -linear maps from K to F (since K is a vector space of dimension k over F , and such a map is determined by its values on a basis).)

Let G be a finite abelian group. If $a : G \rightarrow \mathbb{C}$ is any function, and we define the matrix A by $A_{xy} = a(y-x)$, then the eigenspaces of A have a basis consisting of characters of G .

(Indeed, if $\chi : G \rightarrow \mathbb{C}^*$ is a character (a homomorphism from the additively written group G into the multiplicative group of nonzero complex numbers),

then

$$(A\chi)_x = \sum_{y \in G} a(y-x)\chi(y) = \left(\sum_{z \in G} a(z)\chi(z) \right) \chi(x)$$

so that χ (regarded as column vector) is an eigenvector of A with eigenvalue $\sum_{z \in G} a(z)\chi(z)$. But G has $|G|$ characters, and these are linearly independent, so this gives us the full spectrum of A .)

Example. The matrix $A = \begin{pmatrix} a & b & c \\ c & a & b \\ b & c & a \end{pmatrix}$ has eigenvectors $\begin{pmatrix} 1 \\ \omega \\ \omega^2 \end{pmatrix}$ with eigenvalues $a + b\omega + c\omega^2$, where ω runs through the cube roots of unity.

Now apply this to the adjacency matrix A of the graph Γ . Let $D := \{d \in \mathbb{F}_q^m \mid \langle d \rangle \in X\}$, so that $|D| = (q-1)|X|$. Then the neighbors of the vertex x of Γ are the points $x+d$ for $d \in D$, and we see that Γ has valency $k = |D| = (q-1)n$. The eigenvalues of A are the sums $\sum_{d \in D} \chi(d)$, where χ is a character of the additive group of \mathbb{F}_q^m . Let $\zeta = e^{2\pi i/p}$ be a primitive p -th root of unity, and let $\text{tr} : \mathbb{F}_q \rightarrow \mathbb{F}_p$ be the trace function. Then the characters χ are of the form

$$\chi_a(x) = \zeta^{\text{tr}(\langle a, x \rangle)}.$$

Now

$$\sum_{\lambda \in \mathbb{F}_q} \chi_a(\lambda x) = \begin{cases} q & \text{if } \langle a, x \rangle = 0 \\ 0 & \text{otherwise.} \end{cases}$$

(Indeed, if S denotes this sum, then $\chi_a(\mu x)S = S$ for all μ , so if $S \neq 0$, then $\text{tr}(\langle a, \mu x \rangle) = \text{tr}(\mu \langle a, x \rangle) = 0$ for all μ , and by the above $\langle a, x \rangle = 0$.)

Thus, we find, if D_0 is a set of representatives for X ,

$$\sum_{d \in D} \chi_a(d) = \sum_{d \in D_0} \sum_{\lambda \in \mathbb{F}_q \setminus \{0\}} \chi_a(\lambda d) = q \cdot |H_a \cap X| - |X|$$

where H_a is the hyperplane $\{\langle x \rangle \mid \langle a, x \rangle = 0\}$ in $PG(m-1, q)$. This shows that if H_a meets X in m_a points, so that the corresponding $q-1$ code words have weight $w_a = n - m_a$, then the corresponding eigenvalue is $qm_a - n = (q-1)n - qw_a = k - qw_a$.

We have proved:

Theorem 8.8.1 *There is a 1-1-1 correspondence between*

- (i) *linear codes C of effective word length n and dimension m and $(q-1)f_i$ words of weight w_i , and*
- (ii) *weighted subsets X of total size n of the projective space $PG(m-1, q)$ such that for f_i hyperplanes H we have $|X \setminus H| = w_i$, and*
- (iii) *graphs Γ , without loops but possibly with multiple edges, with vertex set \mathbb{F}_q^m , invariant under translation and dilatation, and with eigenvalues $k - qw_i$ of multiplicity $(q-1)f_i$, where $k = n(q-1)$.*

If the code C is *projective*, that is, no two coordinate positions are dependent (i.e., the dual code has minimum weight at least 3), then X has no repeated points, and we find an ordinary subset under (ii), and a simple graph under (iii) (that is, without multiple edges).

Corollary 8.8.2 *There is a 1-1-1 correspondence between*

- (i) *projective linear codes C of effective word length n and dimension m with precisely two nonzero weights w_1 and w_2 , and*
- (ii) *subsets X of size n of the projective space $PG(m-1, q)$ such that for each hyperplane H we have $|X \cap H| = w_i$, $i \in \{1, 2\}$, and*
- (iii) *strongly regular graphs Γ , with vertex set \mathbb{F}_q^m , invariant under translation and dilatation, and with eigenvalues $k - qw_i$, where $k = n(q-1)$.*

For example, if we take a hyperoval in $PG(2, q)$, q even, we find a two-weight $[q+2, 3, q]$ -code over \mathbb{F}_q . If we take the curve $\{(1, t, t^2, \dots, t^{m-1}) \mid t \in \mathbb{F}_q\} \cup \{(0, 0, \dots, 0, 1)\}$ in $PG(m-1, q)$, q arbitrary, we find a $[q+1, m, q-m+2]$ -code over \mathbb{F}_q . (These codes are optimal: they reach the Singleton bound.)

A 1-1 correspondence between projective codes and 2-weight codes was shown in Brouwer & van Eupen [50].

8.8.4 Duality for affine strongly regular graphs

Let X be a subset of $PG(m-1, q)$ such that all hyperplanes meet it in either m_1 or m_2 points. In the dual projective space (where the rôles of points and hyperplanes have been interchanged), the collection Y of hyperplanes that meet X in m_1 points, is a set with the same property: there are numbers n_1 and n_2 such that each point is in either n_1 or in n_2 hyperplanes from Y .

Indeed, let $x \in X$ be in n_1 hyperplanes from Y . We can find n_1 (independent of the choice of x) by counting hyperplanes on pairs x, y of distinct points in X :

$$n_1 \cdot (m_1 - 1) + \left(\frac{q^{m-1} - 1}{q - 1} - n_1\right) \cdot (m_2 - 1) = (|X| - 1) \cdot \frac{q^{m-2} - 1}{q - 1}.$$

In a similar way we find n_2 , the number of hyperplanes from Y on a point outside X . Computation yields $(m_1 - m_2)(n_1 - n_2) = q^{k-2}$. This proves:

Proposition 8.8.3 *The difference of the weights in a projective 2-weight code, and the difference of the nontrivial eigenvalues of an affine strongly regular graph, is a power of p , where p is the characteristic of the field involved.*

Let Γ and Δ be the strongly regular graphs corresponding to X and Y , respectively. We see that Γ and Δ both have q^k vertices; Γ has valency $k = (q-1)|X|$ and multiplicity $f = (q-1)|Y|$, and for Δ these values have interchanged rôles. We call Δ the *dual* of Γ . (More generally it is possible to define the dual of an association scheme with a regular abelian group of automorphisms, cf. [48], p. 68.)

Example. The ternary Golay code is a perfect $[11, 6, 5]$ code over \mathbb{F}_3 , and its dual C is a $[11, 5, 6]$ code with weights 6 and 9. The corresponding strongly regular graph Γ has parameters $(v, k, v-k-1, \lambda, \mu, r, s, f, g) = (243, 22, 220, 1, 2, 4, -5, 132, 110)$ (it is the *Berlekamp-van Lint-Seidel graph*) and its dual has parameters $(243, 110, 132, 37, 60, 2, -25, 220, 22)$, and we see that $k, v-k-1$ interchange place with g, f . The code corresponding to Δ is a $[55, 5, 36]$ ternary code.

Example. The quaternary Hill code ([193]) is a $[78, 6, 56]$ code over \mathbb{F}_4 with weights 56 and 64. The corresponding strongly regular graph has parameters $(4096, 234, 3861, 2, 14, 10, -22, 2808, 1287)$. Its dual has parameters $(4096, 1287, 2808, 326, 440, 7, -121, 3861, 234)$, corresponding to a quaternary $[429, 6, 320]$ code with weights 320 and 352. This code lies outside the range of the tables, but its residue is a world record $[109, 5, 80]$ code. The binary $[234, 12, 112]$ code derived from the Hill code has a $[122, 11, 56]$ code as residue - also this is a world record.

8.8.5 Cyclotomy

In this section we take D to be a union of cosets of a subgroup of the multiplicative group of a field \mathbb{F}_q . (I.e., the q here corresponds to the q^k of the previous sections.)

Let $q = p^e$, p prime and $e|(q-1)$, say $q = em + 1$. Let $K \subseteq \mathbb{F}_q^*$ be the subgroup of the e -th powers (so that $|K| = m$). Let α be a primitive element of \mathbb{F}_q . For $J \subseteq \{0, 1, \dots, e-1\}$ put $u := |J|$ and $D := D_J := \bigcup \{\alpha^j K \mid j \in J\} = \{\alpha^{ie+j} \mid j \in J, 0 \leq i < m\}$. Define a (directed) graph $\Gamma = \Gamma_J$ with vertex set \mathbb{F}_q and edges (x, y) whenever $y - x \in D$. Note that Γ will be undirected iff either -1 is an e -th power (i.e., q is even or $e|(q-1)/2$) or $J + (q-1)/2 = J$ (arithmetic in \mathbb{Z}_e).

Let $A = A_J$ be the adjacency matrix of Γ defined by $A(x, y) = 1$ if (x, y) is an edge of Γ and $= 0$ otherwise. Let us compute the eigenvalues of A . For each (additive) character χ of \mathbb{F}_q we have

$$(A\chi)(x) = \sum_{y \sim x} \chi(y) = \left(\sum_{u \in D} \chi(u) \right) \chi(x).$$

So each character gives us an eigenvector, and since these are all independent we know all eigenvalues. Their explicit determination requires some theory of Gauss sums. Let us write $A\chi = \theta(\chi)\chi$. Clearly, $\theta(1) = mu$, the valency of Γ . Now assume $\chi \neq 1$. Then $\chi = \chi_g$ for some g , where

$$\chi_g(\alpha^j) = \exp\left(\frac{2\pi i}{p} \text{tr}(\alpha^{j+g})\right)$$

and $\text{tr} : \mathbb{F}_q \rightarrow \mathbb{F}_p$ is the trace function.

If μ is any multiplicative character of order e (say, $\mu(\alpha^j) = \zeta^j$, where $\zeta = \exp(\frac{2\pi i}{e})$), then

$$\sum_{i=0}^{e-1} \mu^i(x) = \begin{cases} e & \text{if } \mu(x) = 1 \\ 0 & \text{otherwise.} \end{cases}$$

Hence,

$$\begin{aligned} \theta(\chi_g) &= \sum_{u \in D} \chi_g(u) = \sum_{j \in J} \sum_{u \in K} \chi_{j+g}(u) = \frac{1}{e} \sum_{j \in J} \sum_{x \in \mathbb{F}_q^*} \chi_{j+g}(x) \sum_{i=0}^{e-1} \mu^i(x) = \\ &= \frac{1}{e} \sum_{j \in J} \left(-1 + \sum_{i=1}^{e-1} \sum_{x \neq 0} \chi_{j+g}(x) \mu^i(x) \right) = \frac{1}{e} \sum_{j \in J} \left(-1 + \sum_{i=1}^{e-1} \mu^{-i}(\alpha^{j+g}) G_i \right) \end{aligned}$$

where G_i is the Gauss sum $\sum_{x \neq 0} \chi_0(x) \mu^i(x)$.

In general, determination of Gauss sums seems to be complicated, but there are a few explicit results. For our purposes the most interesting is the following:

Proposition 8.8.4 (Stickelberger and Davenport & Hasse, see McEliece & Rumsey [242]) *Suppose $e > 2$ and p is semiprimitive mod e , i.e., there exists an l such that $p^l \equiv -1 \pmod{e}$. Choose l minimal and write $\kappa = 2lt$. Then*

$$G_i = (-1)^{t+1} \varepsilon^{it} \sqrt{q},$$

where

$$\varepsilon = \begin{cases} -1 & \text{if } e \text{ is even and } (p^l + 1)/e \text{ is odd} \\ +1 & \text{otherwise.} \end{cases}$$

Under the hypotheses of this proposition, we have

$$\sum_{i=1}^{e-1} \mu^{-i}(\alpha^{j+g}) G_i = \sum_{i=1}^{e-1} \zeta^{-i(j+g)} (-1)^{t+1} \varepsilon^{it} \sqrt{q} = \begin{cases} (-1)^t \sqrt{q} & \text{if } r \neq 1, \\ (-1)^{t+1} \sqrt{q}(e-1) & \text{if } r = 1, \end{cases}$$

where $\zeta = \exp(2\pi i/e)$ and $r = r_{g,j} = \zeta^{-j-g} \varepsilon^t$ (so that $r^e = \varepsilon^{et} = 1$), and hence

$$\theta(\chi_g) = \frac{u}{e} (-1 + (-1)^t \sqrt{q}) + (-1)^{t+1} \sqrt{q} \cdot \#\{j \in J \mid r_{g,j} = 1\}.$$

If we abbreviate the cardinality in this formula with $\#$ then: If $\varepsilon^t = 1$ then $\# = 1$ if $g \in -J \pmod{e}$, and $= 0$ otherwise. If $\varepsilon^t = -1$ (then e is even and p is odd) then $\# = 1$ if $g \in \frac{1}{2}e - J \pmod{e}$, and $= 0$ otherwise. We proved:

Theorem 8.8.5 *Let $q = p^\kappa$, p prime and $e \mid (q-1)$, where p is semiprimitive mod e , i.e., there is an $l > 0$ such that $p^l \equiv -1 \pmod{e}$. Choose l minimal with this property and write $\kappa = 2lt$. Choose u , $1 \leq u \leq e-1$ and assume that q is even or u is even or $e \mid (q-1)/2$. Then the graphs Γ_J (where J is arbitrary for q even or $e \mid (q-1)/2$ and satisfies $J + (q-1)/2 = J \pmod{e}$ otherwise) are strongly regular with eigenvalues*

$$\begin{aligned} k &= \frac{q-1}{e} u && \text{with multiplicity } 1, \\ \theta_1 &= \frac{u}{e} (-1 + (-1)^t \sqrt{q}) && \text{with multiplicity } q-1-k, \\ \theta_2 &= \frac{u}{e} (-1 + (-1)^t \sqrt{q}) + (-1)^{t+1} \sqrt{q} && \text{with multiplicity } k. \end{aligned}$$

(Obviously, when t is even we have $r = \theta_1$, $s = \theta_2$, and otherwise $r = \theta_2$, $s = \theta_1$.)

Clearly, if $e \mid e'(q-1)$, then the set of e -th powers is a union of cosets of the set of e' -th powers, so when applying the above theorem we may assume that e has been chosen as large as possible, i.e., $e = p^l + 1$. Then the restriction ' q is even or u is even or $e \mid (q-1)/2$ ' is empty, and J can always be chosen arbitrarily.

The above construction can be generalized. Pick several values e_i ($i \in I$) with $e_i \mid (q-1)$. Let K_i be the subgroup of \mathbb{F}_q^* of the e_i -th powers. Let J_i be a subset of $\{0, 1, \dots, e_i - 1\}$. Let $D_i := D_{J_i} := \bigcup \{\alpha^j K_i \mid j \in J_i\}$. Put $D := \bigcup D_i$. If the D_i are mutually disjoint, then D defines a graph of which we can compute the spectrum.

For example, let p be odd, and take $e_i = p^{l_i} + 1$ ($i = 1, 2$) and $q = p^\kappa$ where $\kappa = 4l_i s_i$ ($i = 1, 2$). Pick J_1 to consist of even numbers only, and J_2 to consist

of odd numbers only. Then $D_1 \cap D_2 = \emptyset$ and $g \in -J_i \pmod{e_i}$ cannot happen for $i = 1, 2$ simultaneously. This means that the resulting graph will be strongly regular with eigenvalues

$$\theta(\chi_g) = \left(\frac{|J_1|}{e_1} + \frac{|J_2|}{e_2}\right)(-1 + \sqrt{q}) - \sqrt{q} \cdot \delta(g \in -J_i \pmod{e_i} \text{ for } i = 1 \text{ or } i = 2)$$

(where $\delta(P) = 1$ if P holds, and $\delta(P) = 0$ otherwise). See also [62]. In the special case $p = 3$, $l_1 = 1$, $l_2 = 2$, $e_1 = 4$, $e_2 = 10$, $J_1 = \{0\}$, $J_2 = \{1\}$, the difference set consists of the powers α^i with $i \equiv 0 \pmod{4}$ or $i \equiv 1 \pmod{10}$, i.e., is the set $\{1, \alpha, \alpha^4, \alpha^8, \alpha^{11}, \alpha^{12}, \alpha^{16}\} \langle \alpha^{20} \rangle$, and we found the first graph from De Lange [220] again. (It has parameters $(v, k, \lambda, \mu) = (6561, 2296, 787, 812)$ and spectrum $2296^1 \ 28^{4264} \ (-53)^{2296}$.)

8.9 Table

Below a table with the feasible parameters for strongly regular graphs on at most 100 vertices. Here *feasible* means that the parameters v, k, λ, μ and multiplicities f, g are integers, with $0 \leq \lambda < k - 1$ and $0 < \mu < k < v$. In some cases a feasible parameter set is ruled out by the absolute bound or the Krein conditions, or the restriction that the order of a conference graph must be the sum of two squares. For some explanation of the comments, see after the table.

\exists	v	k	λ	μ	r^f	s^g	comments
!	5	2	0	1	0.618^2	-1.618^2	pentagon; Paley(5); Seidel 2-graph-*
!	9	4	1	2	1^4	-2^4	Paley(9); 3^2 ; 2-graph-*
!	10	3	0	1	1^5	-2^4	Petersen graph [259]; $NO_4^-(2)$; $NO_3^{-\perp}(5)$; 2-graph
		6	3	4	1^4	-2^5	$\binom{5}{2}$; 2-graph
!	13	6	2	3	1.303^6	-2.303^6	Paley(13); 2-graph-*
!	15	6	1	3	1^9	-3^5	$O_5(2)$ polar graph; $Sp_4(2)$ polar graph; $NO_4^-(3)$; 2-graph-*
			8	4	2^5	-2^9	$\binom{6}{2}$; 2-graph-*
!	16	5	0	2	1^{10}	-3^5	$q_{22}^2 = 0$; vanLint-Schrijver(1); $VO_4^-(2)$ affine polar graph; projective binary [5,4] code with weights 2, 4; $RSHCD^-$; 2-graph
			10	6	2^5	-2^{10}	Clebsch graph [93, 99, 92, 274]; $q_{11}^1 = 0$; vanLint-Schrijver(2); 2-graph
2!	16	6	2	2	2^6	-2^9	Shrikhande graph [283]; 4^2 ; from a partial spread; projective binary [6,4] code with weights 2, 4; $RSHCD^+$; 2-graph
		9	4	6	1^9	-3^6	$OA(4,3)$; $Bilin_{2 \times 2}(2)$; Goethals-Seidel(2,3); $VO_4^+(2)$ affine polar graph; 2-graph
!	17	8	3	4	1.562^8	-2.562^8	Paley(17); 2-graph-*
!	21	10	3	6	1^{14}	-4^6	
		10	5	4	3^6	-2^{14}	$\binom{7}{2}$
-	21	10	4	5	1.791^{10}	-2.791^{10}	Conf
!	25	8	3	2	3^8	-2^{16}	5^2
		16	9	12	1^{16}	-4^8	$OA(5,4)$
15!	25	12	5	6	2^{12}	-3^{12}	complete enumeration by Paulus [255]; Paley(25); $OA(5,3)$; 2-graph-*
10!	26	10	3	4	2^{13}	-3^{12}	complete enumeration by Paulus [255]; 2-graph
		15	8	9	2^{12}	-3^{13}	$S(2,3,13)$; 2-graph
!	27	10	1	5	1^{20}	-5^6	$q_{22}^2 = 0$; $O_6^-(2)$ polar graph; $GQ(2,4)$; 2-graph-*
		16	10	8	4^6	-2^{20}	Schläfli graph; unique by Seidel [274]; $q_{11}^1 = 0$; 2-graph-*
-	28	9	0	4	1^{21}	-5^6	Krein ₂ ; Absolute bound

\exists	v	k	λ	μ	r^f	s^g	comments
		18	12	10	4^6	-2^{21}	Krein ₁ ; Absolute bound
4!	28	12	6	4	4^7	-2^{20}	Chang graphs [80]; $\binom{8}{2}$; 2-graph
		15	6	10	1^{20}	-5^7	$NO_6^+(2)$; Goethals-Seidel(3,3); Taylor 2-graph for $U_3(3)$
41!	29	14	6	7	2.193^{14}	-3.193^{14}	complete enumeration by Bussemaker & Spence [pers.comm.]; Paley(29); 2-graph-*
—	33	16	7	8	2.372^{16}	-3.372^{16}	Conf
3854!	35	16	6	8	2^{20}	-4^{14}	complete enumeration by McKay & Spence [244]; 2-graph-*
		18	9	9	3^{14}	-3^{20}	S(2,3,15); lines in $PG(3,2)$; $O_6^+(2)$ polar graph; 2-graph-*
!	36	10	4	2	4^{10}	-2^{25}	6^2
		25	16	20	1^{25}	-5^{10}	
180!	36	14	4	6	2^{21}	-4^{14}	$U_3(3).2/L_2(7).2$ - subconstituent of the Hall-Janko graph; complete enumeration by McKay & Spence [244]; $RSHCD^-$; 2-graph
		21	12	12	3^{14}	-3^{21}	2-graph
!	36	14	7	4	5^8	-2^{27}	$\binom{9}{2}$
		21	10	15	1^{27}	-6^8	
32548!	36	15	6	6	3^{15}	-3^{20}	complete enumeration by McKay & Spence [244]; $OA(6,3)$; $NO_6^-(2)$; $RSHCD^+$; 2-graph
		20	10	12	2^{20}	-4^{15}	$NO_5^-(3)$; 2-graph
+	37	18	8	9	2.541^{18}	-3.541^{18}	Paley(37); 2-graph-*
28!	40	12	2	4	2^{24}	-4^{15}	complete enumeration by Spence [286]; $O_5(3)$ polar graph; $Sp_4(3)$ polar graph
		27	18	18	3^{15}	-3^{24}	$NU(4,2)$
+	41	20	9	10	2.702^{20}	-3.702^{20}	Paley(41); 2-graph-*
78!	45	12	3	3	3^{20}	-3^{24}	complete enumeration by Coolsaet, Degraer & Spence [95]; $U_4(2)$ polar graph
		32	22	24	2^{24}	-4^{20}	$NO_5^+(3)$
!	45	16	8	4	6^9	-2^{35}	$\binom{10}{2}$
		28	15	21	1^{35}	-7^9	
+	45	22	10	11	2.854^{22}	-3.854^{22}	Mathon [241]; 2-graph-*
!	49	12	5	2	5^{12}	-2^{36}	7^2
		36	25	30	1^{36}	-6^{12}	$OA(7,6)$
—	49	16	3	6	2^{32}	-5^{16}	Bussemaker-Haemers-Mathon-Wilbrink [70]
		32	21	20	4^{16}	-3^{32}	
+	49	18	7	6	4^{18}	-3^{30}	$OA(7,3)$; Pasechnik(7)
		30	17	20	2^{30}	-5^{18}	$OA(7,5)$
+	49	24	11	12	3^{24}	-4^{24}	Paley(49); $OA(7,4)$; 2-graph-*
!	50	7	0	1	2^{28}	-3^{21}	Hoffman-Singleton graph [196]; $U_3(5^2).2/Sym(7)$
		42	35	36	2^{21}	-3^{28}	
—	50	21	4	12	1^{42}	-9^7	Absolute bound
		28	18	12	8^7	-2^{42}	Absolute bound
+	50	21	8	9	3^{25}	-4^{24}	switch skewhad ² + *; 2-graph
		28	15	16	3^{24}	-4^{25}	S(2,4,25); 2-graph
+	53	26	12	13	3.140^{26}	-4.140^{26}	Paley(53); 2-graph-*
!	55	18	9	4	7^{10}	-2^{44}	$\binom{11}{2}$
		36	21	28	1^{44}	-8^{10}	
!	56	10	0	2	2^{35}	-4^{20}	Sims-Gewirtz graph [152, 153, 52]; $L_3(4).2^2/Alt(6).2^2$
		45	36	36	3^{20}	-3^{35}	Witt: intersection-2 graph of a 2-(21,6,4) design with block intersections 0, 2
—	56	22	3	12	1^{48}	-10^7	Krein ₂ ; Absolute bound
		33	22	15	9^7	-2^{48}	Krein ₁ ; Absolute bound
—	57	14	1	4	2^{38}	-5^{18}	Wilbrink-Brouwer [306]
		42	31	30	4^{18}	-3^{38}	
+	57	24	11	9	5^{18}	-3^{38}	S(2,3,19)
		32	16	20	2^{38}	-6^{18}	
—	57	28	13	14	3.275^{28}	-4.275^{28}	Conf
+	61	30	14	15	3.405^{30}	-4.405^{30}	Paley(61); 2-graph-*
—	63	22	1	11	1^{55}	-11^7	Krein ₂ ; Absolute bound
		40	28	20	10^7	-2^{55}	Krein ₁ ; Absolute bound
+	63	30	13	15	3^{35}	-5^{27}	intersection-8 graph of a 2-(36,16,12) design with block intersections 6, 8; $O_7(2)$ polar graph; $Sp_6(2)$ polar graph; 2-graph-*

\exists	v	k	λ	μ	r^f	s^g	comments
		32	16	16	4^{27}	-4^{35}	S(2,4,28); intersection-6 graph of a 2-(28,12,11) design with block intersections 4, 6; $NU(3,3)$; 2-graph-*
!	64	14	6	2	6^{14}	-2^{49}	8^2 ; from a partial spread of 3-spaces: projective binary [14,6] code with weights 4, 8
		49	36	42	1^{49}	-7^{14}	OA(8,7)
167!	64	18	2	6	2^{45}	-6^{18}	complete enumeration by Haemers & Spence [184]; $GQ(3,5)$; from a hyperoval: projective 4-ary [6,3] code with weights 4, 6
		45	32	30	5^{18}	-3^{45}	
-	64	21	0	10	1^{56}	-11^7	Krein ₂ ; Absolute bound
		42	30	22	10^7	-2^{56}	Krein ₁ ; Absolute bound
+	64	21	8	6	5^{21}	-3^{42}	OA(8,3); $Bilin_{2 \times 3}(2)$; from a Baer subplane: projective 4-ary [7,3] code with weights 4, 6; from a partial spread of 3-spaces: projective binary [21,6] code with weights 8, 12
		42	26	30	2^{42}	-6^{21}	OA(8,6)
+	64	27	10	12	3^{36}	-5^{27}	from a unital: projective 4-ary [9,3] code with weights 6, 8; $VO_6^-(2)$ affine polar graph; $RSHCD^-$; 2-graph
		36	20	20	4^{27}	-4^{36}	2-graph
+	64	28	12	12	4^{28}	-4^{35}	OA(8,4); from a partial spread of 3-spaces: projective binary [28,6] code with weights 12, 16; $RSHCD^+$; 2-graph
		35	18	20	3^{35}	-5^{28}	OA(8,5); Goethals-Seidel(2,7); $VO_6^+(2)$ affine polar graph; 2-graph
-	64	30	18	10	10^8	-2^{55}	Absolute bound
		33	12	22	1^{55}	-11^8	Absolute bound
?	65	32	15	16	3.531^{32}	-4.531^{32}	2-graph-*
!	66	20	10	4	8^{11}	-2^{54}	$\binom{12}{2}$
		45	28	36	1^{54}	-9^{11}	
?	69	20	7	5	5^{23}	-3^{45}	
		48	32	36	2^{45}	-6^{23}	S(2,6,46) does not exist
-	69	34	16	17	3.653^{34}	-4.653^{34}	Conf
+	70	27	12	9	6^{20}	-3^{49}	S(2,3,21)
		42	23	28	2^{49}	-7^{20}	
+	73	36	17	18	3.772^{36}	-4.772^{36}	Paley(73); 2-graph-*
?	75	32	10	16	2^{56}	-8^{18}	2-graph-*
		42	25	21	7^{18}	-3^{56}	2-graph-*
-	76	21	2	7	2^{56}	-7^{19}	Haemers [176]
		54	39	36	6^{19}	-3^{56}	
?	76	30	8	14	2^{57}	-8^{18}	2-graph?
		45	28	24	7^{18}	-3^{57}	2-graph?
?	76	35	18	14	7^{19}	-3^{56}	2-graph?
		40	18	24	2^{56}	-8^{19}	2-graph?
!	77	16	0	4	2^{55}	-6^{21}	S(3,6,22); $M_{22}.2/2^4$: Sym(6); unique by Brouwer [42]; subconstituent of Higman-Sims graph; intersection-6 graph of a 2-(56,16,6) design with block intersections 4, 6
		60	47	45	5^{21}	-3^{55}	Witt 3-(22,6,1): intersection-2 graph of a 2-(22,6,5) design with block intersections 0, 2
-	77	38	18	19	3.887^{38}	-4.887^{38}	Conf
!	78	22	11	4	9^{12}	-2^{65}	$\binom{13}{2}$
		55	36	45	1^{65}	-10^{12}	
!	81	16	7	2	7^{16}	-2^{64}	9^2 ; from a partial spread: projective ternary [8,4] code with weights 3, 6
		64	49	56	1^{64}	-8^{16}	OA(9,8)
!	81	20	1	6	2^{60}	-7^{20}	unique by Brouwer & Haemers [51]; $VO_4^-(3)$ affine polar graph; projective ternary [10,4] code with weights 6, 9
		60	45	42	6^{20}	-3^{60}	
+	81	24	9	6	6^{24}	-3^{56}	OA(9,3); $VNO_4^+(3)$ affine polar graph; from a partial spread: projective ternary [12,4] code with weights 6, 9

\exists	v	k	λ	μ	r^f	s^g	comments
		56	37	42	2^{56}	-7^{24}	OA(9,7)
+	81	30	9	12	3^{50}	-6^{30}	$VNO_4^-(3)$ affine polar graph; Hamada-Helleseth [188]: projective ternary [15,4] code with weights 9, 12
		50	31	30	5^{30}	-4^{50}	
+	81	32	13	12	5^{32}	-4^{48}	OA(9,4); $\text{Bilin}_{2 \times 2}(3)$; $VO_4^+(3)$ affine polar graph; from a partial spread: projective ternary [16,4] code with weights 9, 12
		48	27	30	3^{48}	-6^{32}	OA(9,6)
-	81	40	13	26	1^{72}	-14^8	Absolute bound
		40	25	14	13^8	-2^{72}	Absolute bound
+	81	40	19	20	4^{40}	-5^{40}	Paley(81); OA(9,5); projective ternary [20,4] code with weights 12, 15; 2-graph-*
+	82	36	15	16	4^{41}	-5^{40}	2-graph
		45	24	25	4^{40}	-5^{41}	S(2,5,41); 2-graph
?	85	14	3	2	4^{34}	-3^{50}	
		70	57	60	2^{50}	-5^{34}	
+	85	20	3	5	3^{50}	-5^{34}	$O_5(4)$ polar graph; $Sp_4(4)$ polar graph
		64	48	48	4^{34}	-4^{50}	
?	85	30	11	10	5^{34}	-4^{50}	
		54	33	36	3^{50}	-6^{34}	S(2,6,51)?
?	85	42	20	21	4.110^{42}	-5.110^{42}	2-graph-*
?	88	27	6	9	3^{55}	-6^{32}	
		60	41	40	5^{32}	-4^{55}	
+	89	44	21	22	4.217^{44}	-5.217^{44}	Paley(89); 2-graph-*
!	91	24	12	4	10^{13}	-2^{77}	$\binom{14}{2}$
		66	45	55	1^{77}	-11^{13}	
-	93	46	22	23	4.322^{46}	-5.322^{46}	Conf
?	95	40	12	20	2^{75}	-10^{19}	2-graph-*
		54	33	27	9^{19}	-3^{75}	2-graph-*
+	96	19	2	4	3^{57}	-5^{38}	Haemers(4)
		76	60	60	4^{38}	-4^{57}	
+	96	20	4	4	4^{45}	-4^{50}	$GQ(5,3)$
		75	58	60	3^{50}	-5^{45}	
?	96	35	10	14	3^{63}	-7^{32}	
		60	38	36	6^{32}	-4^{63}	
-	96	38	10	18	2^{76}	-10^{19}	Degraer
		57	36	30	9^{19}	-3^{76}	
?	96	45	24	18	9^{20}	-3^{75}	2-graph?
		50	22	30	2^{75}	-10^{20}	2-graph?
+	97	48	23	24	4.424^{48}	-5.424^{48}	Paley(97); 2-graph-*
?	99	14	1	2	3^{54}	-4^{44}	
		84	71	72	3^{44}	-4^{54}	
?	99	42	21	15	9^{21}	-3^{77}	
		56	28	36	2^{77}	-10^{21}	
+	99	48	22	24	4^{54}	-6^{44}	2-graph-*
		50	25	25	5^{44}	-5^{54}	S(2,5,45); 2-graph-*
!	100	18	8	2	8^{18}	-2^{81}	10^2
		81	64	72	1^{81}	-9^{18}	
!	100	22	0	6	2^{77}	-8^{22}	Higman-Sims graph [192]; $HS01.2/M_{22}.2$; unique by Gewirtz [152]; $q_{11}^2 = 0$
		77	60	56	7^{22}	-3^{77}	$q_{11}^1 = 0$
+	100	27	10	6	7^{27}	-3^{72}	OA(10,3)
		72	50	56	2^{72}	-8^{27}	OA(10,8)?
?	100	33	8	12	3^{66}	-7^{33}	
		66	44	42	6^{33}	-4^{66}	
+	100	33	14	9	8^{24}	-3^{75}	S(2,3,25)
		66	41	48	2^{75}	-9^{24}	
-	100	33	18	7	13^{11}	-2^{88}	Absolute bound
		66	39	52	1^{88}	-14^{11}	Absolute bound
+	100	36	14	12	6^{36}	-4^{63}	Hall-Janko graph; $J_2.2/U_3(3).2$; subconstituent of $G_2(4)$ graph; OA(10,4)
		63	38	42	3^{63}	-7^{36}	OA(10,7)?
+	100	44	18	20	4^{55}	-6^{44}	Jørgensen-Klin graph [209]; $RSHCD^-$; 2-graph

\exists	v	k	λ	μ	r^f	s^g	comments
		55	30	30	5^{44}	-5^{55}	2-graph
+	100	45	20	20	5^{45}	-5^{54}	OA(10,5)?; $RSHCD^+$; 2-graph
		54	28	30	4^{54}	-6^{45}	OA(10,6)?; 2-graph

Comments

Comment	Explanation
$q_{11}^1 = 0, q_{22}^2 = 0$ m^2	zero Krein parameter, see §10.4 Hamming graph $H(2, m)$, i.e., lattice graph $L_2(m)$, i.e., grid graph $m \times m$, i.e., $K_m \times K_m$, see §11.3.1, §1.4.5
$\binom{m}{2}$	Johnson graph $J(m, 2)$, i.e., triangular graph $T(m)$, see §11.3.2, §1.4.5
$OA(n, t)$ ($t \geq 3$)	block graph of an orthogonal array $OA(n, t)$ ($t - 2$ mutually orthogonal Latin squares of order n)
$S(2, k, v)$	block graph of a Steiner system $S(2, k, v)$, i.e., of a $2-(v, k, 1)$ design
Goethals-Seidel(k, r)	graph constructed from a Steiner system $S(2, k, v)$ (with $r = (v - 1)/(k - 1)$) and a Hadamard matrix of order $r + 1$ as in [160]
2-graph	graph in the switching class of a regular 2-graph, see §9.2
2-graph- $*$ $RSHCD^\pm$	descendant of a regular 2-graph, see §9.2 Regular 2-graph derived from a regular symmetric Hadamard matrix with constant diagonal (cf. §9.5, [56], [160])
Taylor 2-graph for $U_3(q)$	graph derived from Taylor's regular 2-graph (cf. [56], [294], [295])
Paley(q)	Paley graph on \mathbb{F}_q , see §9.4, §12.6
vanLint-Schrijver(u)	graph constructed by the cyclotomic construction of [228], taking the union of u classes
$\text{Bilin}_{2 \times d}(q)$	graph on the $2 \times d$ matrices over \mathbb{F}_q , adjacent when their difference has rank 1
$GQ(s, t)$	collinearity graph of a generalized quadrangle with pa- rameters $GQ(s, t)$, see §8.6.3
$O_{2d}^\varepsilon(q), O_{2d+1}(q)$	isotropic points on a nondegenerate quadric in the pro- jective space $PG(2d - 1, q)$ or $PG(2d, q)$, joined when the connecting line is totally singular
$Sp_{2d}(q)$	points of $PG(2d - 1, q)$ provided with a nondegenerate symplectic form, joined when the connecting line is totally isotropic
$U_d(q)$	isotropic points of $PG(d - 1, q^2)$ provided with a non- degenerate Hermitean form, joined when the connect- ing line is totally isotropic

continued...

Comment	Explanation
$NO_{2d}^\varepsilon(2)$	nonisotropic points of $PG(2d-1, 2)$ provided with a nondegenerate quadratic form, joined when they are orthogonal, i.e., when the connecting line is a tangent
$NO_{2d}^\varepsilon(3)$	one class of nonisotropic points of $PG(2d-1, 3)$ provided with a nondegenerate quadratic form, joined when they are orthogonal, i.e., when the connecting line is elliptic
$NO_{2d+1}^\varepsilon(q)$	one class of nondegenerate hyperplanes of $PG(2d, q)$ provided with a nondegenerate quadratic form, joined when their intersection is degenerate
$NO_{2d+1}^{\perp}(5)$	one class of nonisotropic points of $PG(2d, 5)$ provided with a nondegenerate quadratic form, joined when they are orthogonal
$NU_n(q)$	nonisotropic points of $PG(n-1, q)$ provided with a nondegenerate Hermitean form, joined when the connecting line is a tangent
$VO_{2d}^\varepsilon(q)$	vectors of a $2d$ -dimensional vector space over \mathbb{F}_q provided with a nondegenerate quadratic form Q , where two vectors u and v are joined when $Q(v-u) = 0$
$VNO_{2d}^\varepsilon(q)$ (q odd)	vectors of a $2d$ -dimensional vector space over \mathbb{F}_q provided with a nondegenerate quadratic form Q , where two vectors u and v are joined when $Q(v-u)$ is a nonzero square

8.10 Exercises

Exercise 1 ([160]) Consider the graph on the set of flags (incident point-line pairs) of the projective plane $PG(2, 4)$ where (p, L) and (q, M) are adjacent when $p \neq q$ and $L \neq M$ and either $p \in M$ or $q \in L$. Show that this graph is strongly regular with parameters $(v, k, \lambda, \mu) = (105, 32, 4, 12)$.

Exercise 2 ([21]) Consider the graph on the cosets of the perfect ternary Golay code (an $[11, 6, 5]$ code over \mathbb{F}_3), where two cosets are adjacent when they differ by a vector of weight 1. Show that this graph is strongly regular with parameters $(v, k, \lambda, \mu) = (243, 22, 1, 2)$. It is known as the *Berlekamp-van Lint-Seidel graph*.

Exercise 3 For a strongly regular graph Γ and a vertex x of Γ , let Δ be the subgraph of Γ induced on the set of vertices different from x and nonadjacent to x . If Γ has no triangles and spectrum k^1, r^f, s^g , then show that Δ has spectrum $(k-\mu)^1, r^{f-k}, s^{g-k}, (-\mu)^{k-1}$. Conclude if Γ is primitive that $f \geq k$ and $g \geq k$, and that if $f = k$ or $g = k$ then Δ is itself complete or strongly regular. Determine all strongly regular graphs with $\lambda = 0$ and $f = k$.

Exercise 4 ([32]) Show that having a constant k almost follows from having constant λ, μ . More precisely: Consider a graph Γ with the property that any two adjacent (non-adjacent) vertices have λ (resp. μ) common neighbors. Show that if Γ is not regular, then either $\mu = 0$ and Γ is a disjoint union of $(\lambda+2)$ -cliques, or $\mu = 1$, and Γ is obtained from a disjoint union of $(\lambda+1)$ -cliques by adding a new vertex, adjacent to all old vertices.

Exercise 5 Prove Theorem 8.6.2.

Exercise 6 A spread in a generalized quadrangle is a subset S of the lines such that every point is on exactly one line of S . Prove that a $GQ(q^2, q)$ has no spread. Hint: A spread is a coclique in the line graph.

Exercise 7 ([209]) Show that the strongly regular graph with parameters $(v, k, \lambda, \mu) = (100, 45, 20, 20)$ obtained from the Hall-Janko graph in §8.1.12 can be switched into a strongly regular graph with parameters $(100, 55, 30, 30)$.

Exercise 8 There exist strongly regular graphs in \mathbb{F}_3^4 , invariant for translation and dilatation, with parameters $(v, k, \lambda, \mu) = (81, 20, 1, 6)$ and $(81, 30, 9, 12)$. Determine the corresponding ternary codes and their weight enumerators.

Exercise 9 With C and D as in §8.7, show that $C \cup D$ induces a distance-regular graph of diameter three with intersection array $\{10, 9, 4; 1, 6, 10\}$.

Exercise 10 With Γ as in §8.7, show that $\chi(\Gamma) \geq 6$ also follows from Corollary 4.2.4 applied to the induced subgraph of Γ , obtained by deleting all vertices of one color class.

Exercise 11 Under what conditions is the Hamming code cyclic? Negacyclic? Constacyclic?

Exercise 12 A *cap* in a projective space is a collection of points, no three on a line. Show that a $[n, n-m, 4]$ code over \mathbb{F}_q exists if and only if there is a cap of size n in $PG(m-1, q)$. Construct for $m > 0$ a $[2^{m-1}, 2^{m-1}-m, 4]$ binary code.

Exercise 13 Given a two-weight code over \mathbb{F}_q of word length n , dimension m and weights w_1 and w_2 . Express the parameters $v, k, \lambda, \mu, r, s, f, g$ of the corresponding strongly regular graph in terms of q, n, k, w_1 and w_2 .

Chapter 9

Regular two-graphs

9.1 Strong graphs

Let us call a graph (*possibly improper*) *strongly regular* when it is strongly regular or complete or edgeless. Above (Theorem 8.1.2) we saw that a graph Γ is (*possibly improper*) strongly regular if and only if its adjacency matrix A satisfies $A^2 \in \langle A, I, J \rangle$, where $\langle \dots \rangle$ denotes the \mathbb{R} -span. In particular, this condition implies that Γ is regular, so that $AJ = JA$.

Consider the Seidel matrix $S = J - I - 2A$ (see §1.8.2). We have $\langle A, I, J \rangle = \langle S, I, J \rangle$. If $A^2 \in \langle A, I, J \rangle$ then also $S^2 \in \langle S, I, J \rangle$, but the converse does not hold. For example, consider the path P_3 of length 2. We have $S^2 = S + 2I$, but A only satisfies the cubic equation $A^3 = 2A$.

We call a graph *strong* whenever its Seidel matrix S satisfies $S^2 \in \langle S, I, J \rangle$. Thus a (*possibly improper*) strongly regular graph is strong, and conversely a regular strong graph is (*possibly improper*) strongly regular. As we saw, a strong graph need not be regular. Another example is given by $C_5 + K_1$, where the Seidel matrix satisfies $S^2 = 5I$. But the following properties are satisfied (recall that an eigenvalue is called *restricted* if it has an eigenvector orthogonal to the all-ones vector $\mathbf{1}$):

Proposition 9.1.1 *For a graph Γ with v vertices and Seidel matrix S the following holds:*

- (i) Γ is strong if and only if S has at most two restricted eigenvalues. In this case $(S - \rho_1 I)(S - \rho_2 I) = (v - 1 + \rho_1 \rho_2)J$, where ρ_1 and ρ_2 are restricted eigenvalues of S .
- (ii) Γ is strong and regular if and only if Γ is (*possibly improper*) strongly regular. In this case the eigenvalue ρ_0 of S for $\mathbf{1}$ satisfies $(\rho_0 - \rho_1)(\rho_0 - \rho_2) = v(v - 1 + \rho_1 \rho_2)$.
- (iii) If Γ is strong with restricted eigenvalues ρ_1 and ρ_2 , and $v - 1 + \rho_1 \rho_2 \neq 0$, then Γ is regular, and hence (*possibly improper*) strongly regular.
- (iv) S has a single restricted eigenvalue if and only if $S = \pm(J - I)$, that is, if and only if Γ is complete or edgeless.

Proof. (i) If Γ is strong then $S^2 + \alpha S + \beta I = \gamma J$ for some constants α, β and γ . If ρ is a restricted eigenvalue of S with eigenvector \mathbf{v} orthogonal to $\mathbf{1}$, then $(\rho^2 + \alpha\rho + \beta)\mathbf{v} = \gamma J\mathbf{v} = \mathbf{0}$, so $\rho^2 + \alpha\rho + \beta = 0$. Therefore S has at most two restricted eigenvalues. Conversely, if S has just two restricted eigenvalues ρ_1 and ρ_2 , then $(S - \rho_1 I)(S - \rho_2 I) \in \langle J \rangle$, so Γ is strong. And if $(S - \rho_1 I)(S - \rho_2 I) = \gamma J$, then the diagonal entries show that $\gamma = v - 1 + \rho_1\rho_2$.

(ii) We know that (possibly improper) strongly regular implies strong and regular. Suppose Γ is strong and regular, then $S^2 \in \langle S, I, J \rangle$ and $SJ \in \langle J \rangle$, this implies that the adjacency matrix $A = (J - S - I)/2$ of Γ satisfies $A^2 \in \langle A, I, J \rangle$, so Γ is (possibly improper) strongly regular by Theorem 8.1.2.

(iii) If Γ is not regular, then J is not a polynomial in S , so $v - 1 + \rho_1\rho_2 = 0$ follows from part (i). \square

We see that $v - 1 + \rho_1\rho_2 = 0$ if and only if S has exactly two distinct eigenvalues ρ_1 and ρ_2 . Recall that two graphs Γ and $\tilde{\Gamma}$ are switching equivalent (see Section 1.8.2) if their Seidel matrices S and \tilde{S} are similar by some diagonal matrix $D = \text{diag}(\pm 1, \dots, \pm 1)$ (i.e. $\tilde{S} = DSD$). So switching equivalent graphs have the same Seidel spectrum, and therefore the property of being strong with two Seidel eigenvalues is invariant under Seidel switching.

Suppose Γ is a strong graph on v vertices with two Seidel eigenvalues ρ_1 and ρ_2 (so $v - 1 + \rho_1\rho_2 = 0$). Clearly, Γ is regular of degree k if and only if its Seidel matrix has constant row sum $v - 1 - 2k$. Therefore $v - 1 - 2k = \rho_0$ is an eigenvalue of S , so either $\rho_0 = \rho_1$, or $\rho_0 = \rho_2$. Switching in Γ produces another strong graph, which may or may not be regular. If it is regular, then it is regular of degree either $(v - 1 - \rho_1)/2$ or $(v - 1 - \rho_2)/2$.

Examples (i) If Γ is P_3 , then the Seidel eigenvalues are -1 and 2 , so a regular graph that is switching equivalent must have degree either $3/2$ or 0 . The former is impossible, but the latter happens: P_3 is switching equivalent to $3K_1$.

(ii) If Γ is $C_5 + K_1$, then the eigenvalues are $\pm\sqrt{5}$, and so can never be equal to the row sum. So this graph cannot be switched into a regular one.

(iii) If Γ is the 4×4 grid (the lattice graph $L_2(4)$), then $v = 16$ and $\rho_0 = \rho_1 = 3$, $\rho_2 = -5$. So Γ is strong with two eigenvalues. Switching in Γ with respect to a coclique of size 4 gives again a regular graph with the same parameters as Γ , but which is not isomorphic to Γ . This is the Shrikhande graph (see Section 8.2). Switching with respect to the union of two parallel lines in the grid (that is, two disjoint 4-cliques in Γ) gives a regular graph of degree 10, the Clebsch graph (see Section 8.2).

Strong graphs were introduced by Seidel [274].

9.2 Two-graphs

A *two-graph* $\Omega = (V, \Delta)$ consists of a finite set V , together with a collection Δ of unordered triples from V , such that every 4-subset of V contains an even number of triples from Δ . The triples from Δ are called *coherent*.

From a graph $\Gamma = (V, E)$, one can construct a two-graph $\Omega = (V, \Delta)$ by defining a triple from V to be coherent if the three vertices induce a subgraph in Γ with an odd number of edges. It is easily checked that out of the four triples in any graph on four vertices, 0, 2, or 4 are coherent. So Ω is a two-graph. We call Ω the two-graph *associated* to Γ .

Observe that Seidel switching does not change the parity of the number of edges in any 3-vertex subgraph of Γ . Therefore switching equivalent graphs have the same associated two-graph. Conversely, from any two-graph $\Omega = (V, \Delta)$ one can construct a graph Γ as follows. Take $\omega \in V$. Define two vertices $x, y \in V \setminus \{\omega\}$ to be adjacent in Γ if $\{\omega, x, y\} \in \Delta$, and define ω to be an isolated vertex of Γ . We claim that every triple $\{x, y, z\} \in \Delta$ has an odd number of edges in Γ , which makes Ω the two-graph associated to Γ . If $\omega \in \{x, y, z\}$ this is clear. If $\omega \notin \{x, y, z\}$, the 4-subgraph condition implies that $\{x, y, z\} \in \Delta$ whenever from the triples $\{\omega, y, z\}$, $\{\omega, x, y\}$, $\{\omega, x, z\}$ just one, or all three are coherent. Hence $\{x, y, x\}$ has one or three edges in Γ . Thus we have established a one-to-one correspondence between two-graphs and switching classes of graphs.

Small two-graphs were enumerated in [71]. The number of nonisomorphic two-graphs on n vertices for small n is

n	0	1	2	3	4	5	6	7	8	9	10
#	1	1	1	2	3	7	16	54	243	2038	33120


There is an explicit formula for arbitrary n . See, e.g., [237].

For the graph Γ with an isolated vertex ω , obtained from Ω as indicated above, the graph $\Gamma \setminus \omega$ plays an important role. It is called the *descendant* of Ω with respect to ω , and will be denoted by Γ_ω .

Since switching equivalent graphs have the same Seidel spectrum, we can define the eigenvalues of a two-graph to be the Seidel eigenvalues of any graph in the corresponding switching class.

Seidel & Tsaranov [278] classified the two-graphs with smallest Seidel eigenvalue not less than -3 :

Theorem 9.2.1 (i) *A graph Γ with smallest Seidel eigenvalue larger than -3 is switching equivalent to the void graph on n vertices, to the one-edge graph on n vertices, or to one of the following $2 + 3 + 5$ graphs on $5, 6, 7$ vertices, respectively:*



(ii) *A graph Γ with smallest Seidel eigenvalue not less than -3 is switching equivalent to a subgraph of mK_2 or of $\overline{T(8)}$, the complement of the line graph of K_8 . \square*

9.3 Regular two-graphs

A two-graph (V, Δ) is called *regular* (of degree a) if every unordered pair from V is contained in exactly a triples from Δ . Suppose $\Omega = (V, \Delta)$ is a two-graph, and let ∇ be the set of non-coherent triples, then it easily follows that $\overline{\Omega} = (V, \nabla)$ is also a two-graph, called the *complement* of Ω . Moreover, Ω is regular of degree a if and only if the complement $\overline{\Omega}$ is regular of degree $\bar{a} = v - 2 - a$. The following result relates regular two-graphs with strong graphs and strongly regular graphs.

Theorem 9.3.1 *For a graph Γ with v vertices, its associated two-graph Ω , and any descendant Γ_ω of Ω the following are equivalent.*

- (i) Γ is strong with two Seidel eigenvalues ρ_1 and ρ_2 .
- (ii) Ω is regular of degree a .
- (iii) Γ_ω is (possibly improper) strongly regular with parameters $(v-1, k, \lambda, \mu)$ with $\mu = k/2$.

The parameters are related by $v = 1 - \rho_1\rho_2$, $a = k = 2\mu = -(\rho_1 + 1)(\rho_2 + 1)/2$, and $\lambda = (3k - v)/2 = 1 - (\rho_1 + 3)(\rho_2 + 3)/4$. The restricted Seidel eigenvalues of Γ_ω are ρ_1 and ρ_2 , and $\rho_1 + \rho_2 = v - 2a - 2 = \bar{a} - a$.

Proof. (ii) \Rightarrow (iii): Let x be a vertex of Γ_ω . The number of coherent triples containing ω and x equals the number of edges in Γ_ω containing x , so Γ_ω is regular of degree a . For two vertices x and y in Γ_ω , let $p(x, y)$ denote the number of vertices z ($z \neq x, y$) adjacent to x but not to y . If x and y are distinct non-adjacent, then $p(x, y) + p(y, x) = a$, and the number μ of common neighbors of x and y equals $k - p(x, y) = k - p(y, x)$. Therefore $\mu = k/2 = a/2$ is independent of x and y . Similarly, if x and y are adjacent, then $p(x, y) + p(y, x) = \bar{a}$ (the degree of the complement), and the number λ of common neighbors of x and y equals $k - 1 - p(x, y) = k - 1 - p(y, x)$, which implies $\lambda = (3k - v)/2$, which is independent of x and y .

(iii) \Rightarrow (ii): If Γ_ω is strongly regular and $k = 2\mu$, then Theorem 8.1.3 gives $\lambda = (3k - v)/2$. With the relations above this shows that Ω is regular of degree k .

(i) \Rightarrow (iii): Switch in Γ with respect to the neighbors of ω , then ω becomes isolated, and $\Gamma \setminus \omega = \Gamma_\omega$. If S_ω is the Seidel matrix of Γ_ω , then

$$S = \begin{bmatrix} 0 & \mathbf{1}^\top \\ \mathbf{1} & S_\omega \end{bmatrix}$$

is the Seidel matrix of Γ . We know $(S - \rho_1 I)(S - \rho_2 I) = 0$. This gives $(S_\omega - \rho_1 I)(S_\omega - \rho_2 I) = -J$. Therefore Γ_ω is strongly regular with restricted Seidel eigenvalues ρ_1 and ρ_2 and $v - 1 = -\rho_1\rho_2$ vertices. From $S = J - 2A - I$ we get the adjacency eigenvalues $r = -(\rho_1 + 1)/2$ and $s = -(\rho_2 + 1)/2$ of Γ_ω . Now the parameters of Γ_ω follow from Theorem 8.1.3.

(iii) \Rightarrow (i): Suppose Γ_ω is strongly regular with $k = 2\mu$ and Seidel matrix S_ω . Then it follows readily that $S_\omega \mathbf{1} = (\rho_1 + \rho_2)\mathbf{1}$ and $(S_\omega - \rho_1 I)(S_\omega - \rho_2 I) = -J$. This implies that S satisfies $(S - \rho_1 I)(S - \rho_2 I) = 0$. \square

Small regular two-graphs have been classified. The table below gives the numbers of nonisomorphic nontrivial regular two-graphs with $\rho_1 = -3$ or $\rho_1 = -5$ or $v \leq 50$.

v	6	10	14	16	18	26	28	30	36
ρ_1, ρ_2	$\pm\sqrt{5}$	± 3	$\pm\sqrt{13}$	$-3, 5$	$\pm\sqrt{17}$	± 5	$-3, 9$	$\pm\sqrt{29}$	$-5, 7$
#	1	1	1	1	1	4	1	6	227
v	38	42	46	50	76	96	126	176	276
ρ_1, ρ_2	$\pm\sqrt{37}$	$\pm\sqrt{41}$	$\pm\sqrt{45}$	± 7	$-5, 15$	$-5, 19$	$-5, 25$	$-5, 35$	$-5, 55$
#	≥ 191	≥ 18	≥ 97	≥ 54	?	?	1	1	1

9.3.1 Related strongly regular graphs

Given the parameters of a regular two-graph Ω , we find three parameter sets for strongly regular graphs that may be related, namely that of the descendants, and the two possible parameter sets for regular graphs in the switching class of Ω . The parameters are given by:

Proposition 9.3.2 (i) Let Γ be strongly regular with parameters (v, k, λ, μ) . The associated two-graph Ω is regular if and only if $v = 2(2k - \lambda - \mu)$. If this is the case, then it has degree $a = 2(k - \mu)$, and Γ_ω is strongly regular with parameters $(v - 1, 2(k - \mu), k + \lambda - 2\mu, k - \mu)$.

(ii) Conversely, if Γ is regular of valency k , and the associated two-graph Ω is regular of degree a , then Γ is strongly regular with parameters $\lambda = k - (v - a)/2$ and $\mu = k - a/2$, and k satisfies the quadratic $2k^2 - (v + 2a)k + (v - 1)a = 0$.

Proof. (i) By definition, Ω is regular of degree a if and only if $a = \lambda + (v - 2k + \lambda) = 2(k - \mu)$. The parameters follow immediately.

(ii) The quadratic expresses that $k - \frac{1}{2}v \in \{r, s\}$. \square

In the case of the regular two-graph on 6 vertices, the descendants are pentagons, and there are no regular graphs in the switching class.

In the case of the regular two-graph on 10 vertices, the descendants are grid graphs 3×3 . The switching class contains both the Petersen graph and its complement. Therefore Ω is isomorphic to its complement (and so are the descendants).

In the case of the regular two-graph on 16 vertices, the descendants are isomorphic to the triangular graph $T(6)$ (with parameters $(15, 8, 4, 4)$ and spectrum $8^1 2^5 (-2)^9$). The switching class contains the grid graph 4×4 and the Shrikhande graph (both with parameters $(16, 6, 2, 2)$ and spectrum $6^1 2^6 (-2)^9$), and the Clebsch graph (with parameters $(16, 10, 6, 6)$ and spectrum $10^1 2^5 (-2)^{10}$).

It remains to specify what switching sets are needed to switch between two strongly regular graphs associated to the same regular two-graph.

Proposition 9.3.3 Let Γ be strongly regular with parameters (v, k, λ, μ) , associated with a regular two-graph.

(i) The graph Γ is switched into a strongly regular graph with the same parameters if and only if every vertex outside the switching set S is adjacent to half of the vertices of S .

(ii) The graph Γ is switched into a strongly regular graph with parameters $(v, k + c, \lambda + c, \mu + c)$ where $c = \frac{1}{2}v - 2\mu$ if and only if the switching set S has size $\frac{1}{2}v$ and is regular of valency $k - \mu$. \square

For example, in order to switch the 4×4 grid graph into the Shrikhande graph, we can switch with respect to a 4-coclique. And in order to switch the 4×4 grid graph into the Clebsch graph, we need a split into two halves that are regular with valency 4, and the union of two disjoint K_4 's works.

Regular two-graphs were introduced by Graham Higman and further investigated by Taylor [293].

9.3.2 The regular two-graph on 276 points

If N is the point-block incidence matrix of the unique Steiner system $S(4, 7, 23)$, then $NN^\top = 56I + 21J$, $NJ = 77J$, $JN = 7J$. Since any two blocks in this Steiner system meet in 1 or 3 points, we have $N^\top N = 7I + A + 3(J - I - A)$ where A describes the relation of meeting in 1 point. As we already saw in §8.1.10, A is the adjacency matrix of a strongly regular graph—in this case one with parameters $(v, k, \lambda, \mu) = (253, 112, 36, 60)$ and spectrum $112^1 \ 2^{30} \ (-26)^{22}$. The Seidel matrix $S = J - I - 2A$ has spectrum $28^1 \ (-5)^{230} \ 51^{22}$ and satisfies $(S - 51I)(S + 5I) = -3J$. Now $S' = \begin{pmatrix} J - I & J - 2N \\ J - 2N^\top & S \end{pmatrix}$ satisfies $(S' - 55I)(S' + 5I) = 0$ and hence is the Seidel matrix of a regular two-graph on 276 vertices. This two-graph is unique (Goethals & Seidel [161]). Its group of automorphisms is Co_3 , acting 2-transitively.

9.3.3 Coherent subsets

A *clique*, or *coherent* subset in a two-graph $\Omega = (V, \Delta)$ is a subset C of V such that all triples in C are coherent. If $x \notin C$, then x determines a partition $\{C_x, C'_x\}$ of C into two possibly empty parts such that a triple xyz with $y, z \in C$ is coherent precisely when y and z belong to the same part of the partition.

Proposition 9.3.4 (Taylor [295]) *Let C be a nonempty coherent subset of the regular two-graph Ω with eigenvalues ρ_1, ρ_2 , where $\rho_2 < 0$. Then*

- (i) $|C| \leq 1 - \rho_2$, with equality iff for each $x \notin C$ we have $|C_x| = |C'_x|$,
and
- (ii) $|C| \leq m(\rho_2)$.

Proof. (i) Let $c = |C|$. Counting incoherent triples that meet C in two points, we find $\frac{1}{2}c(c-1)\bar{a} = \sum_{x \notin C} |C_x| \cdot |C'_x| \leq \sum_{x \notin C} (c/2)^2 = \frac{1}{4}c^2(v-c)$. It follows that $c^2 - (v-2\bar{a})c - 2\bar{a} \leq 0$. But the two roots of $x^2 - (v-2\bar{a})x - 2\bar{a} = 0$ are $1 - \rho_1$ and $1 - \rho_2$, hence $1 - \rho_1 \leq c \leq 1 - \rho_2$.

(ii) This follows by making a system of equiangular lines in \mathbb{R}^m as in §9.6.1 corresponding to the complement of Ω . We can choose unit vectors for the points in C such that their images form a simplex (any two have the same inner product) and hence $|C|$ is bounded by the dimension $m = v - m(\rho_1) = m(\rho_2)$. \square

9.3.4 Completely regular two-graphs

In a regular two-graph each pair is in $a_2 = a$ coherent triples, that is, in a_2 3-cliques, and each coherent triple is in a_3 4-cliques, where a_3 is the number of common neighbours of two adjacent vertices in any strongly regular graph Γ_ω , so that $a_3 = -\frac{1}{4}(\rho_1 + 3)(\rho_2 + 3) + 1$ by Theorem 9.3.1.

Let a *t-regular two-graph* be a regular two-graph in which every i -clique is contained in a nonzero constant number a_i of $(i+1)$ -cliques, for $2 \leq i \leq t$. By Proposition 9.3.4 we must have $t \leq -\rho_2$. A *completely regular two-graph* is a t -regular two-graph with $t = -\rho_2$. For example, the regular two-graph on 276 points (§9.3.2) is completely regular. Neumaier [252] introduced this concept and gave parameter restrictions strong enough to leave only a finite list of feasible parameters. There are five examples, and two open cases.

#	ρ_1	ρ_2	v	a_2	a_3	a_4	a_5	a_6	a_7	existence
1	3	-3	10	4	1					unique [273]
2	5	-3	16	6	1					unique [273]
3	9	-3	28	10	1					unique [273]
4	7	-5	36	16	6	2	1			unique (BH)
5	19	-5	96	40	12	2	1			none (NP)
6	25	-5	126	52	15	2	1			none [252]
7	55	-5	276	112	30	2	1			unique [161]
8	21	-7	148	66	25	8	3	2	1	none [252]
9	41	-7	288	126	45	12	3	2	1	none [28]
10	161	-7	1128	486	165	36	3	2	1	?
11	71	-9	640	288	112	36	10	4	3	none (BH)
12	351	-9	3160	1408	532	156	30	4	3	?
13	253	-11	2784	1270	513	176	49	12	5	none [252]

Here (BH) refers to an unpublished manuscript by Blokhuis and Haemers, while (NP) is the combination of Neumaier [252] who showed that a derived graph on 95 vertices must be locally $GQ(3, 3)$, and Pasechnik [254] who classified such graphs and found none on 95 vertices.

9.4 Conference matrices

The Seidel matrix of $C_5 + K_1$ is an example of a so called *conference matrix*. An $n \times n$ matrix S is a *conference matrix* if all diagonal entries are 0, the off-diagonal entries are ± 1 , and $SS^\top = (n-1)I$.

Multiplying a row or column by -1 (switching) does not affect the conference matrix property. It was shown in [128] that any conference matrix can be switched into a form where it is either symmetric or skew symmetric:

Lemma 9.4.1 *Let S be a conference matrix of order n with $n > 2$. Then n is even and one can find diagonal matrices D and E with diagonal entries ± 1 such that $(DSE)^\top = DSE$ if and only if $n \equiv 2 \pmod{4}$. One can find such D and E with $(DSE)^\top = -DSE$ if and only if $n \equiv 0 \pmod{4}$.*

Proof. Switch rows and columns so as to make all non-diagonal entries of the first row and column equal to 1. The second row now has $n/2$ entries 1 and equally many entries -1 (since it has inner product zero with the first row). So, n is even, say $n = 2m + 2$. Let there be a, b, c, d entries 1, $-1, 1, -1$ in the third row below the entries 1, 1, $-1, -1$ of the second row, respectively. We may assume (by switching the first column and all rows except the first if required) that $S_{23} = 1$. If $S_{32} = 1$ then $a + b = m - 1$, $c + d = m$, $a + c + 1 = m$, $a - b - c + d + 1 = 0$ imply $a + 1 = b = c = d = \frac{1}{2}m$ so that m is even. If $S_{32} = -1$ then $a + b = m - 1$, $c + d = m$, $a + c = m$, $a - b - c + d + 1 = 0$ imply $a = b = c - 1 = d = \frac{1}{2}(m - 1)$ so that m is odd. This proves that after switching the first row and column to 1, the matrix S has become symmetric in case $n \equiv 2 \pmod{4}$, while after switching the first row to 1 and the first column to -1 , the matrix S has become skew symmetric in case $n \equiv 0 \pmod{4}$. \square

Thus, if $n \equiv 2 \pmod{4}$, S gives rise to a strong graph with two eigenvalues and its associated two-graph is regular of degree $(n-2)/2$. The descendants are strongly regular with parameters $(n-1, (n-2)/2, (n-6)/4, (n-2)/4)$. We call

these graphs *conference graphs*. Conference graphs are characterized among the strongly regular graphs by $f = g$ (f and g are the multiplicities of the restricted eigenvalues), and are the only cases in which non-integral eigenvalues can occur.

The following condition is due to Belevitch [20].

Theorem 9.4.2 *If n is the order of a symmetric conference matrix, then $n - 1$ is the sum of two integral squares.*

Proof. $CC^\top = (n - 1)I$ implies that I and $(n - 1)I$ are rationally congruent (two matrices A and B are rationally congruent if there exist a rational matrix R such that $RAR^\top = B$). A well-known property (essentially Lagrange's four squares theorem) states that for every positive rational number α , the 4×4 matrix αI_4 is rationally congruent to I_4 . This implies that the $n \times n$ matrix αI_n is rationally congruent to $\text{diag}(1, \dots, 1, \alpha, \dots, \alpha)$ where the number of ones is divisible by 4. Since $n \equiv 2 \pmod{4}$, I must be rationally congruent to $\text{diag}(1, \dots, 1, n - 1, n - 1)$. This implies that $n - 1$ is the sum of two squares. \square

Note that this theorem also gives a necessary condition for the existence of conference graphs. For example, 21 is not the sum of two squares, therefore there exists no conference matrix of order 22, and no strongly regular graph with parameters $(21, 10, 4, 5)$.

For many values of n conference matrices are known to exist, see for example [159]. The following construction, where $n - 1$ is an odd prime power, is due to Paley [253]. Let S_ω be a matrix whose rows and columns are indexed by the elements of a finite field \mathbf{F}_q of order q , q odd, by $(S_\omega)_{i,j} = \chi(i - j)$, where χ is the quadratic residue character (that is, $\chi(0) = 0$ and $\chi(x) = 1$ if x is a square, and -1 if x is not a square). It follows that S is symmetric if $q \equiv 1 \pmod{4}$, and S is skew symmetric if $q \equiv 3 \pmod{4}$. In both cases

$$S = \begin{bmatrix} 0 & \mathbf{1}^\top \\ \mathbf{1} & S_\omega \end{bmatrix}$$

is a conference matrix. If $n \equiv 2 \pmod{4}$, S represents a regular two-graph and all its descendants are isomorphic. They are the *Paley graphs*, that we already encountered in §8.1.2.

9.5 Hadamard matrices

Closely related to conference matrices are Hadamard matrices. A matrix H of order n is called a *Hadamard matrix* if every entry is 1 or -1 , and $HH^\top = nI$. If H is a Hadamard matrix, then so is H^\top . If a row or a column of a Hadamard matrix is multiplied by -1 , the matrix remains a Hadamard matrix. The *core* of a Hadamard matrix H (with respect to the first row and column) is the matrix C of order $n - 1$ obtained by first multiplying rows and columns of H by ± 1 so as to obtain a Hadamard matrix of which the first row and column consist of ones only, and then deleting the first row and column. Now all entries of C are ± 1 , and we have $CC^\top = C^\top C = nI - J$, and $C\mathbf{1} = C^\top\mathbf{1} = -\mathbf{1}$. This implies that the $(0, 1)$ matrix $N = \frac{1}{2}(C + J)$ satisfies $N^\top\mathbf{1} = (\frac{1}{2}n - 1)\mathbf{1}$ and $NN^\top = \frac{1}{4}nI + (\frac{1}{4}n - 1)J$, so that, for $n > 2$, N is the incidence matrix of a symmetric 2 -($n - 1, \frac{1}{2}n - 1, \frac{1}{4}n - 1$) design. Conversely, if N is the incidence matrix of a 2-design with these parameters, then $2N - J$ is the core of a Hadamard matrix.

Note that the design parameters imply that n is divisible by 4 if $n > 2$. The famous Hadamard conjecture states that this condition is sufficient for existence of a Hadamard matrix of order n . Many constructions are known (see below), but the conjecture is still far from being solved.

A Hadamard matrix H is *regular* if H has constant row and column sum (ℓ say). Now $-H$ is a regular Hadamard matrix with row sum $-\ell$. From $HH^\top = nI$ we get that $\ell^2 = n$, so $\ell = \pm\sqrt{n}$, and n is a square. If H is a regular Hadamard matrix with row sum ℓ , then $N = \frac{1}{2}(H + J)$ is the incidence matrix of a symmetric 2 -($n, (n+\ell)/2, (n+2\ell)/4$) design. Conversely, if N is the incidence matrix of a 2 -design with these parameters (a *Menon design*), then $2N - J$ is a regular Hadamard matrix.

A Hadamard matrix H is *graphical* if it is symmetric with constant diagonal. Without loss of generality we assume that the diagonal elements are 1 (otherwise we replace H by $-H$). If H is a graphical Hadamard matrix of order n , then $S = H - I$ is the Seidel matrix of a strong graph Γ with two Seidel eigenvalues: $-1 \pm \sqrt{n}$. In other words, Γ is in the switching class of a regular two-graph. The descendent of Γ with respect to some vertex has Seidel matrix $C - I$, where C is the corresponding core of H . It is a strongly regular graph with parameters $(v, k, \lambda, \mu) = (n-1, \frac{1}{2}n-1, \frac{1}{4}n-1, \frac{1}{4}n-1)$. From $\text{tr } S = 0$ it follows that also for a graphical Hadamard matrix n is a square. If, in addition, H is regular with row sum $\ell = \pm\sqrt{n}$, then Γ is a strongly regular graph with parameters $(n, (n-\ell)/2, (n-2\ell)/4, (n-2\ell)/4)$. And conversely, a strongly regular graph with one of the above parameter sets gives rise to a Hadamard matrix of order n .

There is an extensive literature on Hadamard matrices. See, e.g., [271, 272, 100].

Constructions

There is a straightforward construction of Hadamard matrices from conference matrices. If S is a skew symmetric conference matrix, then $H = S + I$ is a Hadamard matrix, and if S is a symmetric conference matrix, then

$$H = \begin{bmatrix} S+I & S-I \\ S-I & -S-I \end{bmatrix}$$

is a Hadamard matrix. Thus the conference matrices constructed in the previous section give Hadamard matrices of order $n = 4m$ if $4m-1$ is a prime power, and if m is odd and $2m-1$ is a prime power. Some small Hadamard matrices are:

$$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}, \quad \begin{bmatrix} 1 & 1 & 1 & -1 \\ 1 & 1 & -1 & 1 \\ 1 & -1 & 1 & 1 \\ -1 & 1 & 1 & 1 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 1 & -1 & -1 & -1 \\ -1 & 1 & -1 & -1 \\ -1 & -1 & 1 & -1 \\ -1 & -1 & -1 & 1 \end{bmatrix}.$$

Observe that the two Hadamard matrices of order 4 are regular and graphical. One easily verifies that, if H_1 and H_2 are Hadamard matrices, then so is the Kronecker product $H_1 \otimes H_2$. Moreover, if H_1 and H_2 are regular with row sums ℓ_1 and ℓ_2 , respectively, then $H_1 \otimes H_2$ is regular with row sum $\ell_1 \ell_2$. Similarly, the Kronecker product of two graphical Hadamard matrices is graphical again. With

the small Hadamard matrices given above, we can make Hadamard matrices of order $n = 2^t$ and regular graphical Hadamard matrices of order $n = 4^t$ with row sum $\ell = \pm 2^t$.

Let $RSHCD$ be the set of pairs (n, ε) such that there exists a regular symmetric Hadamard matrix H with row sums $\ell = \varepsilon\sqrt{n}$ and constant diagonal, with diagonal entries 1. If $(m, \delta), (n, \varepsilon) \in RSHCD$, then $(mn, \delta\varepsilon) \in RSHCD$.

We mention some direct constructions:

- (i) $(4, \pm 1), (36, \pm 1), (100, \pm 1), (196, \pm 1) \in RSHCD$.
- (ii) If there exists a Hadamard matrix of order m , then $(m^2, \pm 1) \in RSHCD$.
- (iii) If both $a - 1$ and $a + 1$ are odd prime powers, then $(a^2, 1) \in RSHCD$.
- (iv) If $a + 1$ is a prime power and there exists a symmetric conference matrix of order a , then $(a^2, 1) \in RSHCD$.
- (v) If there is a set of $t - 2$ mutually orthogonal Latin squares of order $2t$, then $(4t^2, 1) \in RSHCD$.
- (vi) $(4t^4, \pm 1) \in RSHCD$.

See [160], [56] and [271], §5.3. For the third part of (i), see [209]. For the fourth part of (i), cf. [160], Theorem 4.5 (for $k = 7$) and [204]. For (ii), cf. [160], Theorem 4.4, and [179]. For (iii), cf. [271], Corollary 5.12. For (iv), cf. [271], Corollary 5.16. For (v), consider the corresponding Latin square graph. For (vi), see [186].

9.6 Equiangular lines

9.6.1 Equiangular lines in \mathbb{R}^d and two-graphs

Seidel (cf. [221, 229, 129]) studied systems of lines in Euclidean space \mathbb{R}^d , all passing through the origin 0, with the property that any two make the same angle φ . The cases $\varphi = 0$ (only one line) and $\varphi = \frac{\pi}{2}$ (at most d lines, mutually orthogonal) being trivial, we assume $0 < \varphi < \frac{\pi}{2}$. Let $\alpha = \cos \varphi$, so that $0 < \alpha < 1$. Choose for each line ℓ_i a unit vector x_i on ℓ_i (determined up to sign). Then $x_i^\top x_i = 1$ for each i , and $x_i^\top x_j = \pm \cos \varphi = \pm \alpha$ for $i \neq j$.

For the Gram matrix G of the vectors x_i this means that $G = I + \alpha S$, where S is the *Seidel adjacency matrix* of a graph Γ . (That is, S is symmetric with zero diagonal, and has entries -1 and 1 for adjacent and nonadjacent vertices, respectively.) Note that changing the signs of some of the x_i corresponds to *Seidel switching* of Γ .

Conversely, let S be the Seidel adjacency matrix of a graph on at least two vertices, and let θ be the smallest eigenvalue of S . (Then $\theta < 0$ since $S \neq 0$ and $\text{tr } S = 0$.) Now $S - \theta I$ is positive semi-definite, and $G = I - \frac{1}{\theta} S$ is the Gram matrix of a set of vectors in \mathbb{R}^d , where $d = \text{rk}(S - \theta I) = n - m(\theta)$ where n is the number of vertices of the graph, and $m(\theta)$ the multiplicity of θ as eigenvalue of S .

We see that there is a 1-1 correspondence between dependent equiangular systems of n lines and two-graphs on n vertices, and more precisely between equiangular systems of n lines spanning \mathbb{R}^d (with $d < n$) and two-graphs on n vertices such that the smallest eigenvalue has multiplicity $n - d$.

Thus, in order to find large sets of equiangular lines, one has to find large graphs where the smallest Seidel eigenvalue has large multiplicity (or, rather, small comultiplicity).

9.6.2 Bounds on equiangular sets of lines in \mathbb{R}^d or \mathbb{C}^d

An upper bound for the size of an equiangular system of lines (and hence an upper bound for the multiplicity of the smallest Seidel eigenvalue of a graph) is given by the so-called *Absolute bound* due to M. Gerzon (cf. [221]):

Theorem 9.6.1 (‘Absolute bound’) *The cardinality n of a system of equiangular lines in Euclidean space \mathbb{R}^d is bounded by $\frac{1}{2}d(d+1)$.*

Proof. Let $X_i = x_i x_i^\top$ be the rank 1 matrix that is the projection onto the line ℓ_i . Then $X_i^2 = X_i$ and

$$\operatorname{tr} X_i X_j = (x_i^\top x_j)^2 = \begin{cases} 1 & \text{if } i = j \\ \alpha^2 & \text{otherwise.} \end{cases}$$

We prove that the matrices X_i are linearly independent. Since they are symmetric, that will show that there are at most $\frac{1}{2}d(d+1)$. So, suppose that $\sum c_i X_i = 0$. Then $\sum_i c_i X_i X_j = 0$ for each j , so that $c_j(1 - \alpha^2) + \alpha^2 \sum c_i = 0$ for each j . This means that all c_j are equal, and since $\sum c_i = \operatorname{tr} \sum c_i X_i = 0$, they are all zero. \square

In \mathbb{C}^d one can study lines (1-spaces) in the same way, choosing a spanning unit vector in each and agreeing that $\langle x \rangle$ and $\langle y \rangle$ make angle $\phi = \arccos \alpha$ where $\alpha = |x^* y|$. (Here x^* stands for \bar{x}^\top .) The same argument now proves

Proposition 9.6.2 *The cardinality n of a system of equiangular lines in \mathbb{C}^d is bounded by d^2 .* \square

There are very few systems of lines in \mathbb{R}^d that meet the absolute bound, but it is conjectured that systems of d^2 equiangular lines in \mathbb{C}^d exist for all d . Such systems are known for $d = 1, 2, 3, 4, 5, 6, 7, 8, 19$ ([310, 198, 199, 163, 9]). In quantum information theory they are known as SICPOVMs.

The special bound gives an upper bound for n in terms of the angle ϕ , or an upper bound for ϕ (equivalently, a lower bound for $\alpha = \cos \phi$) in terms of n .

Proposition 9.6.3 (‘Special bound’) *If there is a system of $n > 1$ lines in \mathbb{R}^d or \mathbb{C}^d such that the cosine of the angle between any two lines is at most α , then $\alpha^2 \geq (n-d)/(n-1)d$, or, equivalently, $n \leq d(1-\alpha^2)/(1-\alpha^2 d)$ if $1-\alpha^2 d > 0$.*

Proof. Let x_i ($1 \leq i \leq n$) be unit vectors in \mathbb{R}^d or \mathbb{C}^d with $|x_i^* x_j| \leq \alpha$ for $i \neq j$. Put $X_i = x_i x_i^*$ and $Y = \sum_i X_i - \frac{n}{d} I$. Then $\operatorname{tr} X_i X_j = |x_i^* x_j|^2 \leq \alpha^2$ for $i \neq j$, and $\operatorname{tr} X_i = \operatorname{tr} X_i^2 = 1$. Now $0 \leq \operatorname{tr} Y Y^* \leq n(n-1)\alpha^2 + n - \frac{n^2}{d}$. \square

Complex systems of lines with equality in the special bound are known as *equiangular tight frames*. There is a lot of recent literature.

If equality holds in the absolute bound, then the X_i span the vector space of all symmetric matrices, and in particular I is a linear combination of the X_i . If equality holds in the special bound, the same conclusion follows. In both cases the following proposition shows (in the real case) that the graph Γ belongs to a regular two-graph.

Proposition 9.6.4 Suppose x_i ($1 \leq i \leq n$) are unit vectors in \mathbb{R}^d or \mathbb{C}^d with $|x_i^* x_j| = \alpha$ for $i \neq j$, where $0 < \alpha < 1$. Put $X_i = x_i x_i^*$ and suppose that there are constants c_i such that $I = \sum c_i X_i$. Then $c_i = d/n$ for all i and $n = d(1 - \alpha^2)/(1 - \alpha^2 d)$.

If the x_i are vectors in \mathbb{R}^d , and G is the Gram matrix of the x_i , and $G = I + \alpha S$, then S has eigenvalues $(n - d)/(\alpha d)$ and $-1/\alpha$ with multiplicities d and $n - d$, respectively. If $n > d + 1$ and $n \neq 2d$, then these eigenvalues are odd integers.

Proof. If $I = \sum c_i X_i$ then $X_j = \sum_i c_i X_i X_j$ for each j , so that $c_j(1 - \alpha^2) + \alpha^2 \sum c_i = 1$ for each j . This means that all c_j are equal, and since $\sum c_i = \text{tr} \sum c_i X_i = \text{tr} I = d$, they all equal d/n . Our equation now becomes $(d/n)(1 - \alpha^2) + \alpha^2 d = 1$, so that $n = d(1 - \alpha^2)/(1 - \alpha^2 d)$.

If F is the $d \times n$ matrix whose columns are the vectors x_i , then $G = F^\top F$, while $FF^\top = \sum x_i x_i^\top = \sum X_i = (n/d)I$. It follows that FF^\top has eigenvalue n/d with multiplicity d , and $G = F^\top F$ has the same eigenvalues, and in addition 0 with multiplicity $n - d$. The spectrum of S follows. If the two eigenvalues of the integral matrix S are not integers, they are conjugate algebraic integers, and then have the same multiplicity, so that $n = 2d$. Since $S = J - I - 2A$, the eigenvalues of S , when integral, are odd. \square

Graphs for which the Seidel adjacency matrix S has only two eigenvalues are strong (cf. §9.1, Proposition 9.1.1) and belong to the switching class of a regular two-graph (Theorem 9.3.1).

The known lower and upper bounds for the maximum number of equiangular lines in \mathbb{R}^d are given in the table below. For these bounds, see van Lint & Seidel [229], Lemmens & Seidel [221], Seidel [277] (p. 884).

d	1	2	3	4	5	6	7–14	15	16	17–18	
N_{\max}	1	3	6	6	10	16	28	36	40	48	
d	19		20		21		22		23–42		43
N_{\max}	72–76		90–96		126		176		276		344

Bounds for the size of systems of lines in \mathbb{R}^d or \mathbb{C}^d with only a few different, specified, angles, or just with a given total number of different angles, were given by Delsarte, Goethals & Seidel [129].

9.6.3 Bounds on sets of lines with few angles and sets of vectors with few distances

In the case of equiangular lines the absolute value of the inner product took only one value. Generalizing that, one has

Theorem 9.6.5 ([129]) For a set of n unit vectors in \mathbb{R}^d such that the absolute value of the inner product between distinct vectors takes s distinct values different from 1, one has $n \leq \binom{d+2s-1}{d-1}$. If one of the inner products is 0, then $n \leq \binom{d+2s-2}{d-1}$.

There are several examples of equality. For example, from the root system of E_8 one gets 120 lines in \mathbb{R}^8 with $|\alpha| \in \{0, \frac{1}{2}\}$.

Theorem 9.6.6 ([129]) *For a set of n unit vectors in \mathbb{C}^d such that the absolute value of the inner product between distinct vectors takes s distinct values different from 1, one has $n \leq \binom{d+s-1}{d-1}^2$. If one of the inner products is 0, then $n \leq \binom{d+s-1}{d-1} \binom{d+s-2}{d-1}$.*

For example, there are systems of 40 vectors in \mathbb{C}^4 with $|\alpha| \in \{0, \frac{1}{\sqrt{3}}\}$ and 126 vectors in \mathbb{C}^6 with $|\alpha| \in \{0, \frac{1}{2}\}$.

For sets of unit vectors instead of sets of lines it may be more natural to look at the inner product itself, instead of using the absolute value.

Theorem 9.6.7 ([130]) *For a set of n unit vectors in \mathbb{R}^d such that the inner product between distinct vectors takes s distinct values, one has $n \leq \binom{d+s-1}{d-1} + \binom{d+s-2}{d-1}$. If the set is antipodal, then $n \leq 2 \binom{d+s-2}{d-1}$.*

For example, in the antipodal case the upper bound is met with equality for $s = 1$ by a pair of vectors $\pm x$ (with $n = 2$), for $s = 2$ by the vectors $\pm e_i$ of a coordinate frame (with $n = 2d$), and for $s = 6$ by the set of shortest nonzero vectors in the Leech lattice in \mathbb{R}^{24} (with inner products $-1, 0, \pm \frac{1}{4}, \pm \frac{1}{2}$ and size $n = 2 \binom{28}{5}$).

In the general case the upper bound is met with equality for $s = 1$ by a simplex (with $n = d + 1$). For $s = 2$ one has

d	2	5	6	22	23	3, 4, 7–21, 24–39
N_{\max}	5	16	27	275	276–277	$\frac{1}{2}d(d+1)$

with examples of equality in the bound $n \leq \frac{1}{2}d(d+3)$ for $d = 2, 6, 22$. The upper bounds for $d > 6$, $d \neq 22$ are due to Musin [248].

Corollary 9.6.8 ([130]) *Let Γ be a regular graph on n vertices, with smallest eigenvalue $\theta_{\min} < -1$ of multiplicity $n - d$. Then $n \leq \frac{1}{2}d(d+1) - 1$.*

(Earlier we saw for strongly regular graphs that $n \leq \frac{1}{2}f(f+3)$. Here $d = f + 1$, so this gives the same bound, but applies to a larger class of graphs.)

Theorem 9.6.9 ([27]) *A set of vectors in \mathbb{R}^d such that the distance between distinct vectors takes s values has size at most $\binom{d+s}{d}$.*

For $d \leq 8$, the maximal size of a 2-distance set in \mathbb{R}^d was determined by Lisoněk [230]. The results are

d	1	2	3	4	5	6	7	8
N_{\max}	3	5	6	10	16	27	29	45

so that equality holds in the Blokhuis bound $\binom{d+2}{2}$ for $d = 1$ and $d = 8$.

The above gave generalizations of the absolute bound. There are also analogs of the special bound, see [129, 130].

Chapter 10

Association schemes

10.1 Definition

An *association scheme with d classes* is a finite set X together with $d+1$ relations R_i on X such that

- (i) $\{R_0, R_1, \dots, R_d\}$ is a partition of $X \times X$;
- (ii) $R_0 = \{(x, x) | x \in X\}$;
- (iii) if $(x, y) \in R_i$, then also $(y, x) \in R_i$, for all $x, y \in X$ and $i \in \{0, \dots, d\}$;
- (iv) for any $(x, y) \in R_k$ the number p_{ij}^k of $z \in X$ with $(x, z) \in R_i$ and $(z, y) \in R_j$ depends only on i, j and k .

The numbers p_{ij}^k are called the *intersection numbers* of the association scheme. The above definition is the original definition of Bose & Shimamoto [34]; it is what Delsarte [127] calls a symmetric association scheme. In Delsarte's more general definition, (iii) is replaced by:

- (iii') for each $i \in \{0, \dots, d\}$ there exists a $j \in \{0, \dots, d\}$ such that $(x, y) \in R_i$ implies $(y, x) \in R_j$,
- (iii'') $p_{ij}^k = p_{ji}^k$, for all $i, j, k \in \{0, \dots, d\}$.

It is also very common to require just (i), (ii), (iii'), (iv), and to call the scheme 'commutative' when it also satisfies (iii''). Define $n = |X|$, and $n_i = p_{ii}^0$. Clearly, for each $i \in \{1, \dots, d\}$, (X, R_i) is a simple graph which is regular of degree n_i .

Theorem 10.1.1 *The intersection numbers of an association scheme satisfy*

- (i) $p_{0j}^k = \delta_{jk}$, $p_{ij}^0 = \delta_{ij} n_j$, $p_{ij}^k = p_{ji}^k$,
- (ii) $\sum_i p_{ij}^k = n_j$, $\sum_j n_j = n$,
- (iii) $p_{ij}^k n_k = p_{ik}^j n_j$,
- (iv) $\sum_l p_{ij}^l p_{kl}^m = \sum_l p_{kj}^l p_{il}^m$.

Proof. Equations (i), (ii) and (iii) are straightforward. The expressions at both sides of (iv) count quadruples (w, x, y, z) with $(w, x) \in R_i$, $(x, y) \in R_j$, $(y, z) \in R_k$, for a fixed pair $(w, z) \in R_m$. \square

It is convenient to write the intersection numbers as entries of the so-called *intersection matrices* L_0, \dots, L_d :

$$(L_i)_{kj} = p_{ij}^k.$$

Note that $L_0 = I$ and $L_i L_j = \sum p_{ij}^k L_k$. From the definition it is clear that an association scheme with two classes is the same as a pair of complementary strongly regular graphs. If (X, R_1) is strongly regular with parameters (v, k, λ, μ) , then the intersection matrices of the scheme are

$$L_1 = \begin{bmatrix} 0 & k & 0 \\ 1 & \lambda & k - \lambda - 1 \\ 0 & \mu & k - \mu \end{bmatrix}, \quad L_2 = \begin{bmatrix} 0 & 0 & v - k - 1 \\ 0 & k - \lambda - 1 & v - 2k + \lambda \\ 1 & k - \mu & v - 2k + \mu - 2 \end{bmatrix}.$$

10.2 The Bose-Mesner algebra

The relations R_i of an association scheme are described by their adjacency matrices A_i of order n defined by

$$(A_i)_{xy} = \begin{cases} 1 & \text{whenever } (x, y) \in R_i, \\ 0 & \text{otherwise.} \end{cases}$$

In other words, A_i is the adjacency matrix of the graph (X, R_i) . In terms of the adjacency matrices, the axioms (i)–(iv) become

- (i) $\sum_{i=0}^d A_i = J$,
- (ii) $A_0 = I$,
- (iii) $A_i = A_i^\top$, for all $i \in \{0, \dots, d\}$,
- (iv) $A_i A_j = \sum_k p_{ij}^k A_k$, for all $i, j, k \in \{0, \dots, d\}$.

From (i) we see that the $(0, 1)$ matrices A_i are linearly independent, and by use of (ii)–(iv) we see that they generate a commutative $(d+1)$ -dimensional algebra \mathcal{A} of symmetric matrices with constant diagonal. This algebra was first studied by Bose & Mesner [33] and is called the *Bose-Mesner algebra* of the association scheme.

Since the matrices A_i commute, they can be diagonalized simultaneously (see Marcus & Minc [239]), that is, there exist a matrix S such that for each $A \in \mathcal{A}$, $S^{-1}AS$ is a diagonal matrix. Therefore \mathcal{A} is semisimple and has a unique basis of minimal idempotents E_0, \dots, E_d (see Burrow [67]). These are matrices satisfying

$$E_i E_j = \delta_{ij} E_i, \quad \sum_{i=0}^d E_i = I.$$

The matrix $\frac{1}{n}J$ is a minimal idempotent (idempotent is clear, and minimal follows since $\text{rk } J = 1$). We shall take $E_0 = \frac{1}{n}J$. Let P and $\frac{1}{n}Q$ be the matrices

relating our two bases for \mathcal{A} :

$$A_j = \sum_{i=0}^d P_{ij} E_i, \quad E_j = \frac{1}{n} \sum_{i=0}^d Q_{ij} A_i.$$

Then clearly

$$PQ = QP = nI.$$

It also follows that

$$A_j E_i = P_{ij} E_i,$$

which shows that the P_{ij} are the eigenvalues of A_j and that the columns of E_i are the corresponding eigenvectors. Thus $m_i = \text{rk } E_i$ is the multiplicity of the eigenvalue P_{ij} of A_j (provided that $P_{ij} \neq P_{kj}$ for $k \neq i$). We see that $m_0 = 1$, $\sum_i m_i = n$, and $m_i = \text{trace } E_i = n(E_i)_{jj}$ (indeed, E_i has only eigenvalues 0 and 1, so $\text{rk } E_k$ equals the sum of the eigenvalues).

Theorem 10.2.1 *The numbers P_{ij} and Q_{ij} satisfy*

- (i) $P_{i0} = Q_{i0} = 1$, $P_{0i} = n_i$, $Q_{0i} = m_i$,
- (ii) $P_{ij} P_{ik} = \sum_{l=0}^d p_{jk}^l P_{il}$,
- (iii) $m_i P_{ij} = n_j Q_{ji}$, $\sum_i m_i P_{ij} P_{ik} = nn_j \delta_{jk}$, $\sum_i n_i Q_{ij} Q_{ik} = nm_j \delta_{jk}$,
- (iv) $|P_{ij}| \leq n_j$, $|Q_{ij}| \leq m_j$.

Proof. Part (i) follows easily from $\sum_i E_i = I = A_0$, $\sum_i A_i = J = nE_0$, $A_i J = n_i J$, and $\text{trace } E_i = m_i$. Part (ii) follows from $A_j A_k = \sum_l p_{jk}^l A_l$. The first equality in (iii) follows from $m_i P_{ij} = \text{tr } A_j E_i = n_j Q_{ji}$, and the other two follow since $PQ = nI$. The first inequality of (iv) holds because the P_{ij} are eigenvalues of the n_j -regular graphs (X, R_j) . The second inequality then follows by use of (iii). \square

Relations (iii) are often referred to as the *orthogonality relations*, since they state that the rows (and columns) of P (and Q) are orthogonal with respect to a suitable weight function.

If $d = 2$, and (X, R_1) is strongly regular with parameters (v, k, λ, μ) and spectrum $k^1 r^f s^g$, the matrices P and Q are

$$P = \begin{bmatrix} 1 & k & v-k-1 \\ 1 & r & -r-1 \\ 1 & s & -s-1 \end{bmatrix}, \quad Q = \begin{bmatrix} 1 & f & g \\ 1 & fr/k & gs/k \\ 1 & -f \frac{r+1}{v-k-1} & -g \frac{s+1}{v-k-1} \end{bmatrix}.$$

In general the matrices P and Q can be computed from the intersection numbers of the scheme, as follows from the following

Theorem 10.2.2 *For $i = 0, \dots, d$, the intersection matrix L_j has eigenvalues P_{ij} ($0 \leq i \leq d$).*

Proof. Theorem 10.2.1(ii) yields

$$\sum_{k,l} P_{il} (L_j)_{lk} (P^{-1})_{km} = P_{ij} \sum_k P_{ik} (P^{-1})_{km} = \delta_{im} P_{ij},$$

hence $PL_jP^{-1} = \text{diag}(P_{0j}, \dots, P_{dj})$. \square

Thanks to this theorem, it is relatively easy to compute P , $Q (= \frac{1}{n}P^{-1})$ and $m_i (= Q_{0i})$. It is also possible to express P and Q in terms of the (common) eigenvectors of the L_j . Indeed, $PL_jP^{-1} = \text{diag}(P_{0j}, \dots, P_{dj})$ implies that the rows of P are left eigenvectors and the columns of Q are right eigenvectors. In particular, m_i can be computed from the right eigenvector u_i and the left eigenvector v_i^\top , normalized such that $(u_i)_0 = (v_i)_0 = 1$, by use of $m_i u_i^\top v_i = n$. Clearly, each m_i must be an integer. These are the *rationality conditions* for an association scheme. As we saw in the case of a strongly regular graph, these conditions can be very powerful.

10.3 The Linear Programming Bound

One of the main reasons association schemes have been studied is that they yield upper bounds for the size of substructures.

Let Y be a nonempty subset of X , and let its *inner distribution* be the vector a defined by $a_i = |(Y \times Y) \cap R_i|/|Y|$, the average number of elements of Y in relation R_i to a given one. Let χ be the characteristic vector of Y . Then $a_i = \frac{1}{|Y|} \chi^\top A_i \chi$.

Theorem 10.3.1 (Delsarte) $aQ \geq 0$.

Proof. We have $|Y|(aQ)_j = |Y| \sum a_i Q_{ij} = \chi^\top \sum Q_{ij} A_i \chi = n \chi^\top E_j \chi \geq 0$ since E_j is positive semidefinite. \square

Example Consider the schemes of the triples from a 7-set, where two triples are in relation R_i when they have $3 - i$ elements in common ($i = 0, 1, 2, 3$). We find

$$P = \begin{bmatrix} 1 & 12 & 18 & 4 \\ 1 & 5 & -3 & -3 \\ 1 & 0 & -3 & 2 \\ 1 & -3 & 3 & -1 \end{bmatrix} \quad \text{and} \quad Q = \begin{bmatrix} 1 & 6 & 14 & 14 \\ 1 & 5/2 & 0 & -7/2 \\ 1 & -1 & -7/3 & 7/3 \\ 1 & -9/2 & 7 & -7/2 \end{bmatrix}.$$

How many triples can we find such that any two meet in at most one point? For the inner distribution a of such a collection Y we have $a_1 = 0$, so $a = (1, 0, s, t)$, and $aQ \geq 0$ gives the three inequalities

$$6 - s - \frac{9}{2}t \geq 0, \quad 14 - \frac{7}{3}s + 7t \geq 0, \quad 14 + \frac{7}{3}s - \frac{7}{2}t \geq 0.$$

The linear programming problem is to maximize $|Y| = 1 + s + t$ given these inequalities, and the unique solution is $s = 6$, $t = 0$. This shows that one can have at most 7 triples that pairwise meet in at most one point in a 7-set, and if one has 7, then no two are disjoint. Of course an example is given by the Fano plane.

How many triples can we find such that any two meet in at least one point? Now $a = (1, r, s, 0)$ and the optimal solution of $aQ \geq 0$ is $(1, 8, 6, 0)$. An example of such a collection is given by the set of 15 triples containing a fixed point.

How many triples can we find such that no two meet in precisely one point? Now $a = (1, r, 0, t)$ and the maximum value of $1 + r + t$ is 5. An example is given by the set of 5 triples containing two fixed points.

10.4 The Krein parameters

The Bose-Mesner algebra \mathcal{A} is not only closed under ordinary matrix multiplication, but also under componentwise (Hadamard, Schur) multiplication (denoted \circ). Clearly $\{A_0, \dots, A_d\}$ is the basis of minimal idempotents with respect to this multiplication. Write

$$E_i \circ E_j = \frac{1}{n} \sum_{k=0}^d q_{ij}^k E_k.$$

The numbers q_{ij}^k thus defined are called the *Krein parameters*. (Our q_{ij}^k are those of Delsarte, but differ from Seidel's [276] by a factor n .) As expected, we now have the analogue of Theorems 10.1.1 and 10.2.1.

Theorem 10.4.1 *The Krein parameters of an association scheme satisfy*

- (i) $q_{0j}^k = \delta_{jk}$, $q_{ij}^0 = \delta_{ij} m_j$, $q_{ij}^k = q_{ji}^k$,
- (ii) $\sum_i q_{ij}^k = m_j$, $\sum_j m_j = n$,
- (iii) $q_{ij}^k m_k = q_{ik}^j m_j$,
- (iv) $\sum_l q_{ij}^l q_{kl}^m = \sum_l q_{kj}^l q_{il}^m$,
- (v) $Q_{ij} Q_{ik} = \sum_{l=0}^d q_{jk}^l Q_{il}$,
- (vi) $n m_k q_{ij}^k = \sum_l n_l Q_{li} Q_{lj} Q_{lk}$.

Proof. Let $\sum(A)$ denote the sum of all entries of the matrix A . Then $JAJ = \sum(A)J$, $\sum(A \circ B) = \text{trace } AB^\top$ and $\sum(E_i) = 0$ if $i \neq 0$, since then $E_i J = n E_i E_0 = 0$. Now (i) follows by use of $E_i \circ E_0 = \frac{1}{n} E_i$, $q_{ij}^0 = \sum(E_i \circ E_j) = \text{trace } E_i E_j = \delta_{ij} m_j$, and $E_i \circ E_j = E_j \circ E_i$, respectively. Equation (iv) follows by evaluating $E_i \circ E_j \circ E_k$ in two ways, and (iii) follows from (iv) by taking $m = 0$. Equation (v) follows from evaluating $A_i \circ E_j \circ E_k$ in two ways, and (vi) follows from (v), using the orthogonality relation $\sum_l n_l Q_{li} Q_{lj} Q_{lk} = \delta_{mk} m_k n$. Finally, by use of (iii) we have

$$m_k \sum_j q_{ij}^k = \sum_j q_{ik}^j m_j = n \cdot \text{trace } (E_i \circ E_k) = n \sum_l (E_i)_{ll} (E_k)_{ll} = m_i m_k,$$

proving (ii). □

The above results illustrate a dual behavior between ordinary multiplication, the numbers p_{ij}^k and the matrices A_i and P on the one hand, and Schur multiplication, the numbers q_{ij}^k and the matrices E_i and Q on the other hand. If two association schemes have the property that the intersection numbers of one are the Krein parameters of the other, then the converse is also true. Two such schemes are said to be (formally) dual to each other. One scheme may have several (formal) duals, or none at all (but when the scheme is invariant under a regular abelian group, there is a natural way to define a dual scheme, cf. Delsarte [127]). In fact usually the Krein parameters are not even integers. But they cannot be negative. These important restrictions, due to Scott [270] are the so-called *Krein conditions*.

Theorem 10.4.2 *The Krein parameters of an association scheme satisfy $q_{ij}^k \geq 0$ for all $i, j, k \in \{0, \dots, d\}$.*

Proof. The numbers $\frac{1}{n}q_{ij}^k$ ($0 \leq k \leq d$) are the eigenvalues of $E_i \circ E_j$ (since $(E_i \circ E_j)E_k = \frac{1}{n}q_{ij}^k E_k$). On the other hand, the Kronecker product $E_i \otimes E_j$ is positive semidefinite, since each E_i is. But $E_i \circ E_j$ is a principal submatrix of $E_i \otimes E_j$, and therefore is positive semidefinite as well, i.e., has no negative eigenvalue. \square

The Krein parameters can be computed by use of equation 10.4.1(vi). This equation also shows that the Krein condition is equivalent to

$$\sum_l n_l Q_{li} Q_{lj} Q_{lk} \geq 0 \quad \text{for all } i, j, k \in \{0, \dots, d\}.$$

In case of a strongly regular graph we obtain

$$q_{11}^1 = \frac{f^2}{v} \left(1 + \frac{r^3}{k^2} - \frac{(r+1)^3}{(v-k-1)^2} \right) \geq 0,$$

$$q_{22}^2 = \frac{g^2}{v} \left(1 + \frac{s^3}{k^2} - \frac{(s+1)^3}{(v-k-1)^2} \right) \geq 0$$

(the other Krein conditions are trivially satisfied in this case), which is equivalent to the result mentioned in section §8.1.5.

Neumaier [250] generalized Seidel's absolute bound to association schemes, and obtained the following.

Theorem 10.4.3 *The multiplicities m_i ($0 \leq i \leq d$) of an association scheme with d classes satisfy*

$$\sum_{q_{ij}^k \neq 0} m_k \leq \begin{cases} m_i m_j & \text{if } i \neq j, \\ \frac{1}{2} m_i (m_i + 1) & \text{if } i = j. \end{cases}$$

Proof. The left hand side equals $\text{rk}(E_i \circ E_j)$. But $\text{rk}(E_i \circ E_j) \leq \text{rk}(E_i \otimes E_j) = \text{rk } E_i \cdot \text{rk } E_j = m_i m_j$. And if $i = j$, then $\text{rk}(E_i \circ E_i) \leq \frac{1}{2} m_i (m_i + 1)$. Indeed, if the rows of E_i are linear combinations of m_i rows, then the rows of $E_i \circ E_i$ are linear combinations of the $m_i + \frac{1}{2} m_i (m_i - 1)$ rows that are the elementwise products of any two of these m_i rows. \square

For strongly regular graphs with $q_{11}^1 = 0$ we obtain Seidel's bound: $v \leq \frac{1}{2} f(f+3)$. But in case $q_{11}^1 > 0$, Neumaier's result states that the bound can be improved to $v \leq \frac{1}{2} f(f+1)$.

10.5 Automorphisms

Let π be an automorphism of an association scheme, and suppose there are N_i points x such that x and $\pi(x)$ are in relation R_i .

Theorem 10.5.1 (G. Higman) *For each j the number $\frac{1}{n} \sum_{i=0}^d N_i Q_{ij}$ is an algebraic integer.*

Proof. The automorphism is represented by a permutation matrix S where $SM = MS$ for each M in the Bose-Mesner algebra. Let $E = E_j$ be one of the idempotents. Then E has eigenvalues 0 and 1, and S has eigenvalues that are roots of unity, so ES has eigenvalues that are zero or a root of unity, and $\text{tr}ES$ is an algebraic integer. But $\text{tr}ES = \frac{1}{n} \sum_i N_i Q_{ij}$. \square

If one puts $a_j = \frac{1}{n} \sum_i N_i Q_{ij}$, then $N_h = \sum_j a_j P_{jh}$ for all h .

10.5.1 The Moore graph on 3250 vertices

Let Γ be a strongly regular graph with parameters $(v, k, \lambda, \mu) = (3250, 57, 0, 1)$ (an unknown Moore graph of diameter two, cf. Theorem 8.1.5).

$$\text{For such a graph } Q = \begin{bmatrix} 1 & 1729 & 1520 \\ 1 & \frac{637}{3} & -\frac{640}{3} \\ 1 & -\frac{13}{3} & \frac{10}{3} \end{bmatrix}.$$

Aschbacher [10] proved that there is no such graph with a rank three group. G. Higman (unpublished, cf. Cameron [75]) proved that there is no such graph with a vertex transitive group.

Proposition 10.5.2 (G. Higman) Γ is not vertex-transitive.

Proof. Consider any nontrivial group of automorphisms G of such a graph. The collection of points fixed by G has the properties $\lambda = 0$ and $\mu = 1$. Also, two nonadjacent fixed vertices are adjacent to the same number of fixed vertices, so the fixed subgraph is either a strongly regular Moore graph (and then has 5, 10 or 50 vertices), or all fixed vertices have distance at most 1 to some fixed vertex (so that there are at most $k + 1 = 58$ of them).

Consider an involution π . If π does not interchange the endpoints of some edge, then $N_1 = 0$, and $N_0 + N_2 = 3250$. But if $\{x, y\}$ is an orbit of π , then the unique common neighbor z of x and y is fixed, and z occurs for at most 28 pairs $\{x, y\}$, so $N_2 \leq 56N_0$, so that $N_0 = 58$, $N_1 = 0$, $N_2 = 3192$ and $\frac{1}{3250}(58 \times 1729 - 3192 \times \frac{13}{3}) = \frac{133}{5}$ is not an integer, contradiction.

So, π must interchange two adjacent points x and y , and hence interchanges the remaining 56 neighbors u of x with the remaining 56 neighbors v of y . If $\{u, v\}$ is such an orbit, then the unique common neighbor of u and v is fixed, and these are all the fixed points. So $N_0 = 56$, that is, π is an odd permutation, since it is the product of 1597 transpositions. Let N be the subgroup of G consisting of the even permutations. Then N does not have any involutions, so is not transitive, and if G was transitive N has two orbits interchanged by any element outside N . But α has fixed points and cannot interchange the two orbits of N . A contradiction, so G was not transitive. \square

10.6 P - and Q -polynomial association schemes

In many cases, the association scheme carries a distance function such that relation R_i is the relation of having distance i . Such schemes are called *metric*. They are characterized by the fact that p_{jk}^i is zero whenever one of i, j, k is larger than the sum of the other two, while p_{jk}^i is nonzero for $i = j + k$. Note that whether a scheme is metric depends on the ordering of the relations R_i . A scheme may be

metric for more than one ordering. Metric association schemes are essentially the same objects as distance-regular graphs (see Chapter 11 below).

Dually, a *cometric* scheme is defined by $q_{jk}^i = 0$ for $i > j + k$ and $q_{jk}^i > 0$ for $i = j + k$.

There are several equivalent formulations of the metric (cometric) property.

An association scheme is called *P-polynomial* if there exist polynomials f_k of degree k with real coefficients, and real numbers z_i such that $P_{ik} = f_k(z_i)$. Clearly we may always take $z_i = P_{i1}$. By the orthogonality relation 10.2.1(iii) we have

$$\sum_i m_i f_j(z_i) f_k(z_i) = \sum_i m_i P_{ij} P_{ik} = n n_j \delta_{jk},$$

which shows that the f_k are orthogonal polynomials.

Dually, a scheme is called *Q-polynomial* when the same holds with Q instead of P . The following result is due to Delsarte [127] (Theorem 5.6, p. 61).

Theorem 10.6.1 *An association scheme is metric (resp. cometric) if and only if it is P-polynomial (resp. Q-polynomial).*

Proof. Let the scheme be metric. Then

$$A_1 A_i = p_{1i}^{i-1} A_{i-1} + p_{1i}^i A_i + p_{1i}^{i+1} A_{i+1}.$$

Since $p_{1i}^{i+1} \neq 0$, A_{i+1} can be expressed in terms of A_1 , A_{i-1} and A_i . Hence for each j there exists a polynomial f_j of degree j such that $A_j = f_j(A_1)$, and it follows that $P_{ij} E_i = A_j E_i = f_j(A_1) E_i = f_j(P_{i1}) E_i$, hence $P_{ij} = f_j(P_{i1})$.

Now suppose that the scheme is *P-polynomial*. Then the f_j are orthogonal polynomials, and therefore they satisfy a 3-term recurrence relation (see Szegő [291] p. 42)

$$\alpha_{j+1} f_{j+1}(z) = (\beta_j - z) f_j(z) + \gamma_{j-1} f_{j-1}(z).$$

Hence

$$P_{i1} P_{ij} = -\alpha_{j+1} P_{i,j+1} + \beta_j P_{ij} + \gamma_{j-1} P_{i,j-1} \quad \text{for } i = 0, \dots, d.$$

Since $P_{i1} P_{ij} = \sum_l p_{1j}^l P_{il}$ and P is nonsingular, it follows that $p_{1j}^l = 0$ for $|l - j| > 1$. Now the full metric property easily follows by induction. The proof for the cometric case is similar. \square

Given a sequence of nonzero real numbers, let its number of *sign changes* be obtained by first removing all zeros from the sequence, and then counting the number of consecutive pairs of different sign. (Thus, the number of sign changes in 1, -1, 0, 1 is 2.)

Proposition 10.6.2 (i) *Let (X, \mathcal{R}) be a P-polynomial association scheme, with relations ordered according to the P-polynomial ordering and eigenspaces ordered according to descending real order on the $\theta_i := P_{i1}$. Then both row i and column i of both matrices P and Q have precisely i sign changes ($0 \leq i \leq d$).*

(ii) *Dually, if (X, \mathcal{R}) is a Q-polynomial association scheme, and the eigenspaces are ordered according to the Q-polynomial ordering and the relations are ordered according to descending real order on the $\sigma_i := Q_{i1}$, then row i and column i of the matrices P and Q have precisely i sign changes ($0 \leq i \leq d$).*

Proof. Since $m_i P_{ij} = n_j Q_{ji}$ the statements about P and Q are equivalent. Define polynomials p_j of degree j for $0 \leq j \leq d+1$ by $p_{-1}(x) = 0$, $p_0(x) = 1$, $(x - a_j)p_j(x) = b_{j-1}p_{j-1} + c_{j+1}p_{j+1}(x)$, taking $c_{d+1} = 1$. Then $A_j = p_j(A)$, and $p_{d+1}(x) = 0$ has as roots the eigenvalues of A . The numbers in row j of P are $p_i(\theta_j)$ ($0 \leq i \leq d$), and by the theory of Sturm sequences the number of sign changes is the number of roots of p_{d+1} larger than θ_j , which is j . The numbers in column i of P are the values of p_i evaluated at the roots of p_{d+1} . Since p_i has degree i , and there is at least one root of p_{d+1} between any two roots of p_i there are i sign changes. The proof in the Q -polynomial case is similar. \square

Example Consider the Hamming scheme $H(4, 2)$, the association scheme on the binary vectors of length 4, where the relation is their Hamming distance. Now

$$P = Q = \begin{bmatrix} 1 & 4 & 6 & 4 & 1 \\ 1 & 2 & 0 & -2 & -1 \\ 1 & 0 & -2 & 0 & 1 \\ 1 & -2 & 0 & 2 & -1 \\ 1 & -4 & 6 & -4 & 1 \end{bmatrix}.$$

10.7 Exercises

Exercise 1 Show that the number of relations of valency 1 in an association scheme is 2^m for some $m \geq 0$, and $2^m | n$. Hint: the relations of valency 1 form an elementary abelian 2-group with operation $i \oplus j = k$ when $A_i A_j = A_k$.

Exercise 2 Show that for the special case where Y is a coclique in a strongly regular graph, the linear programming bound is the Hoffman bound (Theorem 4.1.2).

Exercise 3 Show that if Γ is a relation of valency k in an association scheme, and θ is a negative eigenvalue of Γ , then $|S| \leq 1 - k/\theta$ for each clique S in Γ .

Exercise 4 Consider a primitive strongly regular graph Γ on v vertices with eigenvalues k^1, r^f, s^g ($k > r > s$) with a Hoffman coloring (that is a coloring with $1 - k/s$ colors). Consider the following relations on the vertex set of Γ :

R_0 : identity,

R_1 : adjacent in Γ ,

R_2 : nonadjacent in Γ with different colors,

R_3 : nonadjacent in Γ with the same color.

Prove that these relations define an association scheme on the vertex set of Γ , and determine the matrices P and Q .

Chapter 11

Distance regular graphs

Consider a connected simple graph with vertex set X of diameter d . Define $R_i \subset X^2$ by $(x, y) \in R_i$ whenever x and y have graph distance i . If this defines an association scheme, then the graph (X, R_1) is called *distance-regular*. By the triangle inequality, $p_{ij}^k = 0$ if $i + j < k$ or $|i - j| > k$. Moreover, $p_{ij}^{i+j} > 0$. Conversely, if the intersection numbers of an association scheme satisfy these conditions, then (X, R_1) is easily seen to be distance-regular.

Many of the association schemes that play a rôle in combinatorics are metric. Families of distance-regular graphs with unbounded diameter include the Hamming graphs, the Johnson graphs, the Grassmann graphs and graphs associated to dual polar spaces. Recently Van Dam & Koolen [121] constructed a new such family, the 15th, and the first without transitive group.

Many constructions and results for strongly regular graphs are the $d = 2$ special case of corresponding results for distance-regular graphs.

The monograph [48] is devoted to the theory of distance-regular graphs, and gives the state of the theory in 1989.

11.1 Parameters

Conventionally, the parameters are $b_i = p_{i+1,1}^i$ and $c_i = p_{i-1,1}^i$ (and $a_i = p_{i,1}^i$). The *intersection array* of a distance-regular graph of diameter d is $\{b_0, \dots, b_{d-1}; c_1, \dots, c_d\}$. The valencies $p_{i,i}^0$, that were called n_i above, are usually called k_i here. We have $c_i k_i = b_{i-1} k_{i-1}$. The total number of vertices is usually called v .

It is easy to see that one has $b_0 \geq b_1 \geq \dots \geq b_{d-1}$ and $c_1 \leq c_2 \leq \dots \leq c_d$ and $c_j \leq b_{d-j}$ ($1 \leq j \leq d$).

11.2 Spectrum

A distance-regular graph Γ of diameter d has $d + 1$ distinct eigenvalues, and the spectrum is determined by the parameters. (Indeed, the matrices P and Q of any association scheme are determined by the parameters p_{jk}^i , and for a distance-regular graph the p_{jk}^i are determined again in terms of the b_i and c_i .)

The eigenvalues of Γ are the eigenvalues of the tridiagonal matrix $L_1 = (p_{1k}^j)$

of order $d + 1$ that here gets the form

$$L_1 = \begin{pmatrix} 0 & b_0 & & & 0 \\ c_1 & a_1 & b_1 & & \\ & c_2 & a_2 & b_2 & \\ & & \dots & \dots & \\ 0 & & & c_d & a_d \end{pmatrix}.$$

If $L_1 u = \theta u$ and $u_0 = 1$, then the multiplicity of θ as eigenvalue of Γ equals $m(\theta) = v / (\sum k_i u_i^2)$.

11.3 Examples

11.3.1 Hamming graphs

Let Q be a set of size q . The *Hamming graph* $H(d, q)$ is the graph with vertex set Q^d , where two vertices are adjacent when they agree in $d - 1$ coordinates.

This graph is distance-regular, with parameters $c_i = i$, $b_i = (q - 1)(d - i)$, diameter d and eigenvalues $(q - 1)d - qi$ with multiplicity $\binom{d}{i}(q - 1)^i$ ($0 \leq i \leq d$). (Indeed, $H(d, q)$ is the Cartesian product of d copies of K_q , see §1.4.6.)

For $q = 2$ this graph is also known as the hypercube 2^d , often denoted Q_d . For $d = 2$ the graph $H(2, q)$ is also called $L_2(q)$.

Cospectral graphs

In §1.8.1 we saw that there are precisely two graphs with the spectrum of $H(4, 2)$.

In §8.2 we saw that there are precisely two graphs with the spectrum of $H(2, 4)$. Here we give a graph cospectral with $H(3, 3)$ (cf. [183]).

The graphs $H(d, q)$ have q^d vertices, and dq^{d-1} maximal cliques ('lines') of size




Figure 11.1: The geometry of the Hamming graph $H(3, 3)$

q . Let N be the point-line incidence matrix. Then $NN^\top - dI$ is the adjacency matrix of $\Gamma = H(d, q)$, and $N^\top N - qI$ is the adjacency matrix of the graph Δ on the lines, where two lines are adjacent when they have a vertex in common. It follows that for $d = q$ the graphs Γ and Δ are cospectral. In Γ any two vertices at distance two have $c_2 = 2$ common neighbors. If $q \geq 3$, then two vertices at distance two in Δ have 1 or q common neighbors (and both occur), so that Δ is not distance-regular, and in particular not isomorphic to Γ . For $q = 3$ the geometry is displayed in Figure 11.1. See also §13.2.2.

11.3.2 Johnson graphs

Let X be a set of size n . The *Johnson graph* $J(n, m)$ is the graph with vertex set $\binom{X}{m}$, the set of all m -subsets of X , where two m -subsets are adjacent when they have $m - 1$ elements in common. For example, $J(n, 0)$ has a single vertex; $J(n, 1)$ is the complete graph K_n ; $J(n, 2)$ is the triangular graph $T(n)$.

This graph is distance-regular, with parameters $c_i = i^2$, $b_i = (m - i)(n - m - i)$, diameter $d = \min(m, n - m)$ and eigenvalues $(m - i)(n - m - i) - i$ with multiplicity $\binom{n}{i} - \binom{n}{i-1}$.

The *Kneser graph* $K(n, m)$ is the graph with vertex set $\binom{X}{m}$, where two m -subsets are adjacent when they have maximal distance in $J(n, m)$ (i.e., are disjoint when $n \geq 2m$, and have $2m - n$ elements in common otherwise). These graphs are not distance-regular in general, but the *Odd graph* O_{m+1} , which equals $K(2m + 1, m)$, is.

Sending a vertex (m -set) to its complement in X is an isomorphism from $J(n, m)$ onto $J(n, n - m)$ and from $K(n, m)$ onto $K(n, n - m)$. Thus, we may always assume that $n \geq 2m$.

11.3.3 Grassmann graphs

Let V be a vector space of dimension n over the field \mathbb{F}_q . The *Grassmann graph* $Gr(n, m)$ is the graph with vertex set $\begin{bmatrix} V \\ m \end{bmatrix}$, the set of all m -subspaces of V , where two m -subspaces are adjacent when they intersect in an $(m - 1)$ -space. This graph is distance-regular, with parameters $c_i = \begin{bmatrix} i \\ 1 \end{bmatrix}^2$, $b_i = q^{2i+1} \begin{bmatrix} m-i \\ 1 \end{bmatrix} \begin{bmatrix} n-m-i \\ 1 \end{bmatrix}$, diameter $d = \min(m, n - m)$, and eigenvalues $q^{i+1} \begin{bmatrix} m-i \\ 1 \end{bmatrix} \begin{bmatrix} n-m-i \\ 1 \end{bmatrix} - \begin{bmatrix} i \\ 1 \end{bmatrix}$ with multiplicity $\begin{bmatrix} n \\ i \end{bmatrix} - \begin{bmatrix} n \\ i-1 \end{bmatrix}$. (Here $\begin{bmatrix} n \\ i \end{bmatrix} = (q^n - 1) \cdots (q^{n-i+1} - 1) / (q^i - 1) \cdots (q - 1)$ is the q -binomial coefficient, the number of m -subspaces of an n -space.)

11.3.4 Van Dam-Koolen graphs

Van Dam & Koolen [121] construct distance-regular graphs $vDK(m)$ with the same parameters as $Gr(2m + 1, m)$. (They call them the *twisted Grassmann graphs*.) These graphs are ugly, the group of automorphisms is not transitive. The existence of such examples reinforces the idea that the parameters of distance-regular graphs of large diameter are strongly restricted, while there is some freedom for the actual structure. The construction is as follows. Let V be a vector space of dimension $2m + 1$ over \mathbb{F}_q , and let H be a hyperplane of V . Take as vertices the $(m + 1)$ -subspaces of V not contained in H , and the $(m - 1)$ -subspaces contained in H , where two subspaces of the same dimension are adjacent when their intersection has codimension 1 in both, and two subspaces of different dimension are adjacent when one contains the other. This graph is the line graph (concurrence graph on the set of lines) of the partial linear space of which the points are the m -subspaces of V , with natural incidence, while the point graph (collinearity graph on the set of points) is $Gr(2m + 1, m)$. It follows that $vDK(m)$ and $Gr(2m + 1, m)$ are cospectral.

11.4 Bannai-Ito conjecture

The most famous problem about distance-regular graphs was the Bannai-Ito conjecture ([17], p. 237): show that there are only finitely many distance-regular graphs with fixed valency k larger than 2. After initial work by Bannai & Ito, the conjecture was attacked by Jack Koolen and coauthors in a long series of papers. After 25 years a complete proof was given by Sejeong Bang, Arturas Dubickas, Jack Koolen, and Vincent Moulton [15].

11.5 Connectedness

For strongly regular graphs we had Theorem 8.3.2 stating that the vertex connectivity $\kappa(\Gamma)$ equals the valency k . In [55] it was shown that the same holds for distance-regular graphs.

For strongly regular graphs we also had Proposition 8.3.1 that says that the induced subgraph on the vertices at maximal distance from a given vertex is connected. This is a very important property, but for distance-regular graphs additional hypotheses are needed. For example, there are two generalized hexagons with parameters $GH(2, 2)$ (duals of each other) and in one of them the subgraphs $\Gamma_3(x)$ are disconnected.

11.6 Degree of eigenvalues

For strongly regular graphs we saw that eigenvalues are integral, except in the ‘half case’ where they are quadratic. Something similar happens for distance-regular graphs.

Polygons have eigenvalues of high degree: for an n -gon the degree of the i -th eigenvalue is $\phi(m)$ where $m = \gcd(i, n)$, where ϕ is the Euler totient function. But elsewhere only integral and quadratic eigenvalues seem to occur.

For the case of a P - and Q -polynomial scheme of diameter at least 3, Bannai & Ito [17] Theorem 7.11 show that the eigenvalues are integers.

There is precisely one known distance-regular graph of valency larger than 2 with a cubic eigenvalue, namely the Biggs-Smith graph, the unique graph with intersection array $\{3, 2, 2, 2, 1, 1, 1; 1, 1, 1, 1, 1, 1, 3\}$. It has 102 vertices, and spectrum $3^1 2^{18} 0^{17} ((1 \pm \sqrt{17})/2)^9 \theta_j^{16}$ where the θ_j are the three roots of $\theta^3 + 3\theta^2 - 3 = 0$.

A result in this direction is

Proposition 11.6.1 *The only distance-regular graph of diameter 3 with a cubic eigenvalue is the heptagon.*

Proof. Let Γ be a distance-regular graph of diameter 3 on n vertices with a cubic eigenvalue. Since algebraically conjugate eigenvalues have the same multiplicity we have three eigenvalues θ_i with multiplicity $f = (n-1)/3$. Since $\text{tr}A = 0$ we find that $\theta_1 + \theta_2 + \theta_3 = -k/f$. Now k/f is rational and an algebraic integer, hence an integer, and $k \geq (n-1)/3$. The same reasoning applies to A_i for $i = 2, 3$ and hence $k_i \geq (n-1)/3$, and we must have equality. Since $k = k_2 = k_3$ we see that $b_1 = c_2 = b_2 = c_3$.

Write $\mu := c_2$. The distinct eigenvalues $k, \theta_1, \theta_2, \theta_3$ of A are the eigenvalues of the matrix L_1 (Theorem 10.2.2) and hence $k - 1 = k + \theta_1 + \theta_2 + \theta_3 = \text{tr} L_1 = a_1 + a_2 + a_3 = (k - \mu - 1) + (k - 2\mu) + (k - \mu)$, so that $k = 2\mu$ and $a_2 = 0$.

Let $d(x, y) = 3$ and put $A = \Gamma(x) \cap \Gamma_2(y)$, $B = \Gamma_2(x) \cap \Gamma(y)$, so that $|A| = |B| = c_3 = \mu$. Every vertex in B is adjacent to every vertex in A , and hence two vertices in B have at least $\mu + 1$ common neighbors, so must be adjacent. Thus B is a clique, and $\mu = |B| \leq a_2 + 1$, that is, $\mu = 1$, $k = 2$. \square

11.7 Moore graphs and generalized polygons

Any k -regular graph of diameter d has at most

$$1 + k + k(k - 1) + \dots + k(k - 1)^{d-1}$$

vertices, as is easily seen. A graph for which equality holds is called a *Moore graph*. Moore graphs are distance-regular, and those of diameter 2 were dealt with in Theorem 8.1.5. Using the rationality conditions Damerell [123] and Bannai & Ito [16] showed:

Theorem 11.7.1 *A Moore graph with diameter $d \geq 3$ is a $(2d + 1)$ -gon.*

A strong non-existence result of the same nature is the theorem of Feit & G. Higman [139] about finite generalized polygons. We recall that a *generalized m -gon* is a point-line incidence geometry such that the incidence graph is a connected, bipartite graph of diameter m and girth $2m$. It is called *regular* of order (s, t) for certain (finite or infinite) cardinal numbers s, t if each line is incident with $s + 1$ points and each point is incident with $t + 1$ lines. From such a regular generalized m -gon of order (s, t) , where s and t are finite and $m \geq 3$, we can construct a distance-regular graph with valency $s(t + 1)$ and diameter $d = \lfloor \frac{m}{2} \rfloor$ by taking the collinearity graph on the points.

Theorem 11.7.2 *A finite generalized m -gon of order (s, t) with $s > 1$ and $t > 1$ satisfies $m \in \{2, 3, 4, 6, 8\}$.*

Proofs of this theorem can be found in Feit & Higman [139], Brouwer, Cohen & Neumaier [48] and Van Maldeghem [298]; again the rationality conditions do the job. The Krein conditions yield some additional information:

Theorem 11.7.3 *A finite regular generalized m -gon with $s > 1$ and $t > 1$ satisfies $s \leq t^2$ and $t \leq s^2$ if $m = 4$ or 8 ; it satisfies $s \leq t^3$ and $t \leq s^3$ if $m = 6$.*

This result is due to Higman [191] and Haemers & Roos [181].

11.8 Primitivity

A distance-regular graph Γ of diameter d is called *imprimitive* when one of the relations (X, R_i) with $i \neq 0$ is disconnected. This can happen in three cases: either Γ is an n -gon, and $i|n$, or $i = 2$, and Γ is bipartite, or $i = d$, and Γ is *antipodal*, that is, having distance d is an equivalence relation. Graphs can be both bipartite and antipodal. The $2n$ -gons fall in all three cases.

11.9 Euclidean representations

Let Γ be distance regular, and let θ be a fixed eigenvalue. Let $E = E_j$ be the idempotent in the association scheme belonging to θ , so that $AE = \theta E$. Let $u_i = Q_{ij}/n$, so that $E = \sum u_i A_i$. Let $f = \text{rk} E$.

The map sending vertex x of Γ to the vector $\bar{x} = Ee_x$, column x of E , provides a representation of Γ by vectors in an f -dimensional Euclidean space, namely the column span of E , where graph distances are translated into inner products: if $d(x, y) = i$ then $(\bar{x}, \bar{y}) = E_{xy} = u_i$.

If this map is not injective, and $\bar{x} = \bar{y}$ for two vertices x, y at distance $i \neq 0$, then $u_i = u_0$ and any two vertices at distance i have the same image. For $i = 1$ this happens when $\theta = k$. Otherwise, Γ is imprimitive, and either $i = 2$ and Γ is bipartite and $\theta = -k$, or $i = d$ and Γ is antipodal, or $2 < i < d$ and Γ is a polygon.

This construction allows one to translate problems about graphs into problems in Euclidean geometry. Especially when f is small, this is a very useful tool.

As an example of the use of this representation, let us prove Terwilliger's Tree Bound. Call an induced subgraph T of Γ *geodetic* when distances measured in T equal distances measured in Γ .

Proposition 11.9.1 *Let Γ be distance regular, and let θ be an eigenvalue different from $\pm k$. Let T be a geodetic tree in Γ . Then the multiplicity f of the eigenvalue θ is at least the number of endpoints of T .*

Proof. We show that the span of the vectors \bar{x} for $x \in T$ has a dimension not less than the number e of endpoints of T . Induction on the size of T . If $T = \{x, y\}$ then $\bar{x} \neq \bar{y}$ since $k \neq \theta$. Assume $|T| > 2$. If $x \in T$, and S is the set of endpoints of T adjacent to x , then for $y, z \in S$ and $w \in T \setminus S$ we have $(\bar{w}, \bar{y} - \bar{z}) = 0$. Pick x such that S is nonempty, and x is an endpoint of $T' = T \setminus S$. By induction $\dim \langle \bar{w} \mid w \in T' \rangle \geq e - |S| + 1$. Since $\theta \neq \pm k$ we have $\dim \langle \bar{y} - \bar{z} \mid x, y \in S \rangle = |S| - 1$. \square

Example For a distance-regular graph without triangles, $f \geq k$. Equality can hold. For example, the Higman-Sims graph is strongly regular with parameters $(v, k, \lambda, \mu) = (100, 22, 0, 6)$ and spectrum $22^1 \ 2^{77} \ (-8)^{22}$.

11.10 Extremality

This section gives a simplified account of the theory developed by Fiol and Garriga and coauthors. The gist is that among the graphs with a given spectrum with $d + 1$ distinct eigenvalues the distance-regular graphs are extremal in the sense that they have a maximal number of pairs of vertices at mutual distance d .

Let Γ be a connected k -regular graph with adjacency matrix A with eigenvalues $k = \theta_1 \geq \dots \geq \theta_n$. Suppose that A has precisely $d + 1$ distinct eigenvalues (so that the diameter of Γ is at most d). Define an inner product on the $(d + 1)$ -dimensional vector space of real polynomials modulo the minimum polynomial of A by

$$\langle p, q \rangle = \frac{1}{n} \text{tr } p(A)q(A) = \frac{1}{n} \sum_{i=1}^n p(\theta_i)q(\theta_i).$$

Note that $\langle p, p \rangle \geq 0$ for all p , and $\langle p, p \rangle = 0$ if and only if $p(A) = 0$. By applying Gram-Schmidt to the sequence of polynomials x^i ($0 \leq i \leq d$) we find a sequence of orthogonal polynomials p_i of degree i ($0 \leq i \leq d$) satisfying $\langle p_i, p_j \rangle = 0$ for $i \neq j$ and $\langle p_i, p_i \rangle = p_i(k)$. This latter normalization is possible since $p_i(k) \neq 0$.

(Indeed, suppose that p_i changes sign at values α_j ($0 \leq j \leq h$) inside the interval (θ_n, k) . Put $q(x) = \prod_{j=1}^h (x - \alpha_j)$. Then all terms in $\langle p_i, q \rangle$ have the same sign, and not all are zero, so $\langle p_i, q \rangle \neq 0$, hence $h = i$, so that all zeros of p_i are in the interval (θ_n, k) , and $p_i(k) \neq 0$.)

The Hoffman polynomial (the polynomial p such that $p(A) = J$) equals $p_0 + \dots + p_d$. Indeed, $\langle p_i, p \rangle = \frac{1}{n} \text{tr } p_i(A)J = p_i(k) = \langle p_i, p_i \rangle$ for all i .

If Γ is distance-regular, then the p_i are the polynomials for which $A_i = p_i(A)$.

Theorem 11.10.1 *Let Γ be connected and regular of degree k , with $d+1$ distinct eigenvalues. Define the polynomials p_i as above. Let $\bar{k}_d := \frac{1}{n} \sum_x k_d(x)$ be the average number of vertices at distance d from a given vertex in Γ . Then $\bar{k}_d \leq p_d(k)$, and equality holds if and only if Γ is distance-regular.*

Proof. We follow Fiol, Gago & Garriga [144]. Use the inner product $\langle M, N \rangle = \frac{1}{n} \text{tr } M^\top N$ on the space $M_n(\mathbb{R})$ of real matrices of order n . If M, N are symmetric, then $\langle M, N \rangle = \frac{1}{n} \sum_{x,y} (M \circ N)_{xy}$. If $M = p(A)$ and $N = q(A)$ are polynomials in A , then $\langle M, N \rangle = \langle p, q \rangle$.

Since $\langle A_d, p_d(A) \rangle = \langle A_d, J \rangle = \bar{k}_d$, the orthogonal projection A'_d of A_d on the space $\langle I, A, \dots, A^d \rangle = \langle p_0(A), \dots, p_d(A) \rangle$ of polynomials in A equals

$$A'_d = \sum_j \frac{\langle A_d, p_j(A) \rangle}{\langle p_j, p_j \rangle} p_j(A) = \frac{\langle A_d, p_d(A) \rangle}{p_d(k)} p_d(A) = \frac{\bar{k}_d}{p_d(k)} p_d(A).$$

Now $\|A'_d\|^2 \leq \|A_d\|^2$ gives $\bar{k}_d^2 / p_d(k) \leq \bar{k}_d$, and the inequality follows since $p_d(k) > 0$. When equality holds, $A_d = p_d(A)$.

Now it follows by downward induction on h that $A_h = p_h(A)$ ($0 \leq h \leq d$). Indeed, from $\sum_j p_j(A) = J = \sum_j A_j$ it follows that $p_0(A) + \dots + p_h(A) = A_0 + \dots + A_h$. Hence $p_h(A)_{xy} = 0$ if $d(x, y) > h$, and $p_h(A)_{xy} = 1$ if $d(x, y) = h$. Since $\langle xp_{h+1}, p_j \rangle = \langle p_{h+1}, xp_j \rangle = 0$ for $j \neq h, h+1, h+2$, we have $xp_{h+1} = ap_h + bp_{h+1} + cp_{h+2}$ and hence $AA_{h+1} = ap_h(A) + bA_{h+1} + cA_{h+2}$ for certain a, b, c with $a \neq 0$. But then $p_h(A)_{xy} = 0$ if $d(x, y) < h$, so that $p_h(A) = A_h$.

Finally, the three-term recurrence for the p_h now becomes the three-term recurrence for the A_h that defines distance-regular graphs. \square

Noting that $p_d(k)$ depends on the spectrum only, we see that this provides a characterization of distance-regularity in terms of the spectrum and the number of pairs of vertices far apart (at mutual distance d). See [115], [143], [144] and Theorem 13.5.3 below.

11.11 Exercises

Exercise 1 Determine the spectrum of a strongly regular graph minus a vertex. (Hint: if the strongly regular graph has characteristic polynomial $p(x) = (x - k)(x - r)^f(x - s)^g$, then the graph obtained after removing one vertex has characteristic polynomial $((x - k)(x - \lambda + \mu) + \mu)(x - r)^{f-1}(x - s)^{g-1}$.)

Determine the spectrum of a strongly regular graph minus two (non)adjacent vertices.

Show that the spectrum of a distance-regular graph minus a vertex does not depend on the vertex chosen. Give an example of two nonisomorphic cospectral graphs both obtained by removing a vertex from the same distance-regular graph.

Chapter 12

p -ranks

Designs or graphs with the same parameters can sometimes be distinguished by considering the p -rank of associated matrices. For example, there are three nonisomorphic 2-(16,6,2) designs, with point-block incidence matrices of 2-rank 6, 7 and 8 respectively.

Tight bounds on the occurrence of certain configurations are sometimes obtained by computing a rank in some suitable field, since p -ranks of integral matrices may be smaller than their ranks over \mathbb{R} .

Our first aim is to show that given the parameters (say, the real spectrum), only finitely many primes p are of interest.

12.1 Reduction mod p

A technical difficulty is that one would like to talk about eigenvalues that are zero or nonzero mod p for some prime p , but it is not entirely clear what that might mean when the eigenvalues are nonintegral. Necessarily some arbitrariness will be involved. For example $(5 + \sqrt{2})(5 - \sqrt{2}) \equiv 0 \pmod{23}$ and one point of view is that this means that 23 is not a prime in $\mathbb{Q}(\sqrt{2})$, and one gets into algebraic number theory. But another point of view is that if one ‘reduces mod 23’, mapping to a field of characteristic 23, then at least one factor must become 0. However, the sum of $5 + \sqrt{2}$ and $5 - \sqrt{2}$ does not become 0 upon reduction mod 23, so not both factors become 0. Since these factors are conjugate, the ‘reduction mod 23’ cannot be defined canonically, it must involve some arbitrary choices. We follow Isaacs [205], who follows Brauer.

Let R be the ring of algebraic integers in \mathbb{C} , and let p be a prime. Let M be a maximal ideal in R containing the ideal pR . Put $F = R/M$. Then F is a field of characteristic p . Let $r \mapsto \bar{r}$ be the quotient map $R \rightarrow R/M = F$. This will be our ‘reduction mod p ’. (It is not canonical because M is not determined uniquely.)

Lemma 12.1.1 (Isaacs [205], 15.1) *Let $U = \{z \in \mathbb{C} \mid z^m = 1 \text{ for some integer } m \text{ not divisible by } p\}$. Then the quotient map $R \rightarrow R/M = F$ induces an isomorphism of groups $U \rightarrow F^*$ from U onto the multiplicative group F^* of F . Moreover, F is algebraically closed, and is algebraic over its prime field.*

One consequence is that on integers ‘reduction mod p ’ has the usual meaning: if m is an integer not divisible by p then some power is 1 (mod p) and it follows

that $\bar{m} \neq 0$. More generally, if $\bar{\theta} = 0$, then $p|N(\theta)$, where $N(\theta)$ is the *norm* of θ , the product of its conjugates, up to sign the constant term of its minimal polynomial.

12.2 The minimal polynomial

Let M be a matrix of order n over a field F . For each eigenvalue θ of M in F , let $m(\theta)$ be the geometric multiplicity of θ , so that $\text{rk}(M - \theta I) = n - m(\theta)$.

Let $e(\theta)$ be the algebraic multiplicity of the eigenvalue θ , so that the characteristic polynomial of M factors as $c(x) := \det(xI - M) = \prod (x - \theta)^{e(\theta)} c_0(x)$, where $c_0(x)$ has no roots in F . Then $m(\theta) \leq e(\theta)$.

The *minimal polynomial* $p(x)$ of M is the unique monic polynomial over F of minimal degree such that $p(M) = 0$. The numbers $\theta \in F$ for which $p(\theta) = 0$ are precisely the eigenvalues of M (in F). By Cayley-Hamilton, $p(x)$ divides $c(x)$. It follows that if $p(x) = \prod (x - \theta)^{h(\theta)} p_0(x)$, where $p_0(x)$ has no roots in F , then $1 \leq h(\theta) \leq e(\theta)$.

In terms of the Jordan decomposition of M , $m(\theta)$ is the number of Jordan blocks for θ , $h(\theta)$ is the size of the largest block, and $e(\theta)$ is the sum of the sizes of all Jordan blocks for θ .

We see that $n - e(\theta) + h(\theta) - 1 \leq \text{rk}(M - \theta I) \leq n - e(\theta)/h(\theta)$, and also that $1 \leq \text{rk}((M - \theta I)^i) - \text{rk}((M - \theta I)^{i+1}) \leq m(\theta)$ for $1 \leq i \leq h - 1$.

12.3 Bounds for the p -rank

Let M be a square matrix of order n , and let $\text{rk}_p(M)$ be its p -rank. Let R and F be as above in §12.1. Use a suffix F or p to denote rank or multiplicity over the field F or \mathbb{F}_p (instead of \mathbb{C}).

Proposition 12.3.1

Let M be an integral square matrix. Then $\text{rk}_p(M) \leq \text{rk}(M)$.

Let M be a square matrix with entries in R . Then $\text{rk}_F(\bar{M}) \leq \text{rk}(M)$.

Proof. The rank of a matrix is the size of the largest submatrix with nonzero determinant. □

Proposition 12.3.2 Let M be an integral square matrix. Then

$$\text{rk}_p(M) \geq \sum \{m(\theta) \mid \bar{\theta} \neq 0\}.$$

Proof. Let M have order n . Then $\text{rk}_p(M) = n - m_p(0) \geq n - e_p(0) = n - e_F(0) = \sum \{e_F(t) \mid t \neq 0\} = \sum \{e(\theta) \mid \bar{\theta} \neq 0\} \geq \sum \{m(\theta) \mid \bar{\theta} \neq 0\}$. □

Proposition 12.3.3 Let the integral square matrix M be diagonalizable. Then $\text{rk}_F(\bar{M} - \bar{\theta}I) \leq n - e(\theta)$ for each eigenvalue θ of M .

Proof. $\text{rk}_F(\bar{M} - \bar{\theta}I) \leq \text{rk}(M - \theta I) = n - m(\theta) = n - e(\theta)$. □

It follows that if $\bar{\theta} = 0$ for a unique θ , then $\text{rk}_p(M) = n - e(\theta)$. We can still say something when $\bar{\theta} = 0$ for two eigenvalues θ , when one has multiplicity 1:

Proposition 12.3.4 *Let the integral square matrix M be diagonalizable, and suppose that $\theta = 0$ for only two eigenvalues θ , say θ_0 and θ_1 , where $e(\theta_0) = 1$. Let M have minimal polynomial $p(x) = (x - \theta_0)f(x)$. Then $\text{rk}_F(\overline{M}) = n - e(\theta_1) - \varepsilon$, where $\varepsilon = 1$ if $f(M) = 0$ and $\varepsilon = 0$ otherwise.*

Proof. By the above $n - e(\theta_1) - 1 \leq \text{rk}_F(\overline{M}) \leq n - e(\theta_1)$. By the previous section $n - e_F(0) + h_F(0) - 1 \leq \text{rk}_F(\overline{M}) \leq n - e_F(0)/h_F(0)$. Since $e_F(0) = e(\theta_1) + 1$ we find $\text{rk}_F(\overline{M}) = n - e(\theta_1) - \varepsilon$, where $\varepsilon = 1$ if $h_F(0) = 1$ and $\varepsilon = 0$ otherwise. But $h_F(0) = 1$ iff $\overline{f(M)} = 0$. \square

If M is a matrix with integral entries, then the minimal polynomial $p(x)$ and its factor $f(x)$ have integral coefficients. In particular, if M is an integral symmetric matrix with constant row sums k , and the eigenvalue k of M has multiplicity 1, then $f(M) = (f(k)/n)J$ and the condition $\overline{f(M)} = 0$ becomes $\overline{c} = 0$, where $c = \frac{1}{n} \prod_{\theta \neq k} (k - \theta)$ is an integer.

12.4 Interesting primes p

Let A be an integral matrix of order n , and let $M = A - aI$ for some integer a . If θ is an eigenvalue of A , then $\theta - a$ is an eigenvalue of M .

If $\bar{\theta} = \bar{a}$ for no θ , then $\text{rk}_p(M) = n$.

If $\bar{\theta} = \bar{a}$ for a unique θ , then $\text{rk}_p(M) = \text{rk}_F(\overline{M}) = \text{rk}_F(M - (\bar{\theta} - a)I) \leq \text{rk}(A - \theta I) = n - m(\theta)$ by Proposition 12.3.1, but also $\text{rk}_p(M) \geq n - m(\theta)$ by Proposition 12.3.2, so that $\text{rk}_p(M) = n - m(\theta)$.

So, if the p -rank of M is interesting, if it gives information not derivable from the spectrum of A and the value a , then at least two eigenvalues of M become zero upon reduction mod p . But if $\bar{\theta} - \bar{a} = \overline{\eta - a} = 0$, then $\bar{\theta} - \bar{\eta} = 0$, and in particular $p | N(\theta - \eta)$, which happens for finitely many p only.

Example The unique distance-regular graph with intersection array $\{4, 3, 2; 1, 2, 4\}$ has 14 vertices and spectrum $4, \sqrt{2}^6, (-\sqrt{2})^6, -4$ (with multiplicities written as exponents).

Let A be the adjacency matrix of this graph, and consider the p -rank of $M = A - aI$ for integers a . The norms of $\theta - a$ are $4 - a, a^2 - 2, -4 - a$, and if these are all nonzero mod p then the p -rank of M is 14. If p is not 2 or 7, then at most one of these norms can be 0 mod p , and for $a \equiv 4 \pmod{p}$ or $a \equiv -4 \pmod{p}$ the p -rank of M is 13. If $a^2 \equiv 2 \pmod{p}$ then precisely one of the eigenvalues $\sqrt{2} - a$ and $-\sqrt{2} - a$ reduces to 0, and the p -rank of M is 8. Finally, for $p = 2$ and $p = 7$ we need to look at the matrix M itself, and find $\text{rk}_2(A) = 6$ and $\text{rk}_7(A \pm 3I) = 8$.

12.5 Adding a multiple of J

Let A be an integral matrix of order n with row and column sums k , and consider the rank and p -rank of $M = M_b = A + bJ$. Since J has rank 1, all these matrices differ in rank by at most 1, so either all have the same rank r , or two ranks $r, r + 1$ occur, and in the latter case rank $r + 1$ occurs whenever the row space of M contains the vector $\mathbf{1}$.

The matrix M has row sums $k + bn$.

If $p \nmid n$, then the row space of M over \mathbb{F}_p contains $\mathbf{1}$ when $k + bn \not\equiv 0 \pmod{p}$. On the other hand, if $k + bn \equiv 0 \pmod{p}$, then all rows have zero row sum \pmod{p} while $\mathbf{1}$ has not, so that $\mathbf{1}$ is not in the row space over \mathbb{F}_p . Thus, we are in the second case, where the smaller p -rank occurs for $b = -k/n$ only.

If $p|n$ and $p \nmid k$, then all row sums are nonzero \pmod{p} for all b , and we are in the former case: the rank is independent of b , and the row space over \mathbb{F}_p always contains $\mathbf{1}$.

Finally, if $p|n$ and also $p|k$, then further inspection is required.

Example (Cf. Peeters [258]). According to [183], there are precisely ten graphs with the spectrum $7^1 \sqrt{7}^8 (-1)^7 (-\sqrt{7})^8$, one of which is the Klein graph, the unique distance-regular graph with intersection array $\{7, 4, 1; 1, 2, 7\}$. It turns out that the p -ranks of $A - aI + bJ$ for these graphs depend on the graph only for $p = 2$ ([258]). Here $n = 24$ and $k = 7 - a + 24b$.

graph	$\text{rk}_2(A + I)$	$\text{rk}_2(A + I + J)$	$\text{rk}_3(A - aI + bJ)$				$\text{rk}_7(A - aI)$	
#1,2	14	14	$a \backslash b$				$a \backslash b$	0
#3,8,9	15	14	0	24	24	24	0	15
#4,7	13	12	1	16	15	16	1-5	24
#5	12	12	2	16	16	16	6	17
#6	11	10						
#10	9	8						

Interesting primes (dividing the norm of the difference of two eigenvalues) are 2, 3 and 7. All p -ranks follow from the parameters except possibly $\text{rk}_2(A + I + bJ)$, $\text{rk}_3(A - I + bJ)$, $\text{rk}_3(A + I)$, $\text{rk}_7(A)$.

The interesting 2-rank is $\text{rk}_2(A + I)$, and inspection of the graphs involved shows that this takes the values 9, 11, 12, 13, 14, 15 where 9 occurs only for the Klein graph. The value of $\text{rk}_2(A + I + J)$ follows, since a symmetric matrix with zero diagonal has even 2-rank, and the diagonal of a symmetric matrix lies in the \mathbb{F}_2 -space of its rows. Hence if $\text{rk}_2(A + I)$ is even, then $\text{rk}_2(A + I + J) = \text{rk}_2(A + I)$, and if $\text{rk}_2(A + I)$ is odd then $\text{rk}_2(A + I + J) = \text{rk}_2(A + I) - 1$.

The 3-rank of $A - I + bJ$ is given by Proposition 12.3.4. Here $f(x) = (x + 2)((x + 1)^2 - 7)$ and $k = 6 + 24b$, so that $f(k)/n \equiv 0 \pmod{3}$ is equivalent to $b \equiv 1 \pmod{3}$.

One has $\text{rk}_3(A + I) = 16$ in all ten cases.

The value of $\text{rk}_7(A)$ can be predicted: We have $\det(A + J) = -7^8 \cdot 31$, so the Smith Normal Form (§12.8) of $A + J$ has at most 8 entries divisible by 7 and $\text{rk}_7(A + J) \geq 16$. By Proposition 12.3.3, $\text{rk}_7(A + J) = 16$. Since $7 \nmid n$ and $\mathbf{1}$ is in the row space of $A + J$ but not in that of A , $\text{rk}_7(A) = 15$.

12.6 Paley graphs

Let q be a prime power, $q \equiv 1 \pmod{4}$, and let Γ be the graph with vertex set \mathbb{F}_q where two vertices are adjacent whenever their difference is a nonzero square. (Then Γ is called the *Paley graph* of order q .) In order to compute the p -rank of the Paley graphs, we first need a lemma.

Lemma 12.6.1 *Let $p(x, y) = \sum_{i=0}^{d-1} \sum_{j=0}^{e-1} c_{ij} x^i y^j$ be a polynomial with coefficients in a field F . Let $A, B \subseteq F$, with $m := |A| \geq d$ and $n := |B| \geq e$.*

Consider the $m \times n$ matrix $P = (p(a, b))_{a \in A, b \in B}$ and the $d \times e$ matrix $C = (c_{ij})$. Then $\text{rk}_F(P) = \text{rk}_F(C)$.

Proof. For any integer s and subset X of F , let $Z(s, X)$ be the $|X| \times s$ matrix $(x^i)_{x \in X, 0 \leq i \leq s-1}$. Note that if $|X| = s$ then this is a Vandermonde matrix and hence invertible. We have $P = Z(d, A) C Z(e, B)^\top$, so $\text{rk}_F(P) \leq \text{rk}_F(C)$, but P contains a submatrix $Z(d, A') C Z(e, B')$ with $A' \subseteq A$, $B' \subseteq B$, $|A'| = d$, $|B'| = e$, and this submatrix has the same rank as C . \square

For odd prime powers $q = p^e$, p prime, let Q be the $\{0, \pm 1\}$ -matrix of order q with entries $Q_{xy} = \chi(y - x)$ ($x, y \in \mathbb{F}_q$, χ the quadratic residue character, $\chi(0) = 0$).

Proposition 12.6.2 ([49]) $\text{rk}_p Q = ((p+1)/2)^e$.

Proof. Applying the above lemma with $p(x, y) = \chi(y - x) = (y - x)^{(q-1)/2} = \sum_i (-1)^i \binom{(q-1)/2}{i} x^i y^{(q-1)/2-i}$, we see that $\text{rk}_p Q$ equals the number of binomial coefficients $\binom{(q-1)/2}{i}$ with $0 \leq i \leq (q-1)/2$ not divisible by p . Now Lucas' Theorem says that if $l = \sum_i l_i p^i$ and $k = \sum_i k_i p^i$ are the p -ary expansions of l and k , then $\binom{l}{k} \equiv \prod_i \binom{l_i}{k_i} \pmod{p}$. Since $\frac{1}{2}(q-1) = \sum_i \frac{1}{2}(p-1)p^i$, this means that for each p -ary digit of i there are $(p+1)/2$ possibilities and the result follows. \square

For Lucas' Theorem, cf. MacWilliams & Sloane [236], §13.5, p. 404 (and references given there). Note that this proof shows that each submatrix of Q of order at least $(q+1)/2$ has the same rank as Q .

The relation between Q here and the adjacency matrix A of the Paley graph is $Q = 2A + I - J$. From $Q^2 = qI - J \equiv -J \pmod{p}$ and $(2A + I)^2 = qI + (q-1)J \equiv -J \pmod{p}$ it follows that both $\langle Q \rangle$ and $\langle 2A + I \rangle$ contain $\mathbf{1}$, so $\text{rk}_p(A + \frac{1}{2}I) = \text{rk}_p(2A + I) = \text{rk}_p(Q) = ((p+1)/2)^e$.

12.7 Strongly regular graphs

Let Γ be a strongly regular graph with adjacency matrix A , and assume that A has integral eigenvalues k, r, s with multiplicities $1, f, g$, respectively. We investigate the p -rank of a linear combination of A , I and J .

The following proposition shows that only the case $p|(r-s)$ is interesting.

Proposition 12.7.1 Let $M = A + bJ + cI$. Then M has eigenvalues $\theta_0 = k + bv + c$, $\theta_1 = r + c$, $\theta_2 = s + c$, with multiplicities $m_0 = 1$, $m_1 = f$, $m_2 = g$, respectively.

(i) If none of the θ_i vanishes \pmod{p} , then $\text{rk}_p M = v$.

(ii) If precisely one θ_i vanishes \pmod{p} , then M has p -rank $v - m_i$.

Put $e := \mu + b^2v + 2bk + b(\mu - \lambda)$.

(iii) If $\theta_0 \equiv \theta_1 \equiv 0 \pmod{p}$, $\theta_2 \not\equiv 0 \pmod{p}$, then $\text{rk}_p M = g$ if and only if $p|e$, and $\text{rk}_p M = g + 1$ otherwise.

(iii)' If $\theta_0 \equiv \theta_2 \equiv 0 \pmod{p}$, $\theta_1 \not\equiv 0 \pmod{p}$, then $\text{rk}_p M = f$ if and only if $p|e$, and $\text{rk}_p M = f + 1$ otherwise.

(iv) In particular, if $k \equiv r \equiv 0 \pmod{p}$ and $s \not\equiv 0 \pmod{p}$, then $\text{rk}_p A = g$. And if $k \equiv s \equiv 0 \pmod{p}$ and $r \not\equiv 0 \pmod{p}$, then $\text{rk}_p A = f$.

(v) If $\theta_1 \equiv \theta_2 \equiv 0 \pmod{p}$, then $\text{rk}_p M \leq \min(f+1, g+1)$.

Proof. Parts (i) and (ii) and (v) are immediate from Propositions 12.3.1, 12.3.2. Suppose $\theta_0 \equiv \theta_1 \equiv 0 \pmod{p}$, $\theta_2 \not\equiv 0 \pmod{p}$. Then we can apply Proposition 12.3.4 with the two eigenvalues 0 and θ_2 . Since $\text{rk}_p(M - \theta_2 I) = v - g$, and $g \leq \text{rk}_p M \leq g + 1$, it follows that $\text{rk}_p M = g$ if and only if $M(M - \theta_2 I) \equiv 0 \pmod{p}$. But using $(A - rI)(A - sI) = \mu J$ and $r + s = \lambda - \mu$, we find $M(M - \theta_2 I) \equiv (A + bJ - rI)(A + bJ - sI) = eJ$. Part (iii)' is similar. \square

Thus, the only interesting case (where the structure of Γ plays a rôle) is that where p divides both θ_1 and θ_2 , so that $p \mid (r - s)$. In particular, only finitely many primes are of interest. In this case we only have the upper bound (v).

Looking at the idempotents sometimes improves this bound by 1: We have $E_1 = (r - s)^{-1}(A - sI - (k - s)v^{-1}J)$ and $E_2 = (s - r)^{-1}(A - rI - (k - r)v^{-1}J)$. Thus, if $k - s$ and v are divisible by the same power of p (so that $(k - s)/v$ can be interpreted in \mathbb{F}_p), then $\text{rk}_p(A - sI - (k - s)v^{-1}J) \leq \text{rk } E_1 = f$, and, similarly, if $k - r$ and v are divisible by the same power of p then $\text{rk}_p(A - rI - (k - r)v^{-1}J) \leq \text{rk } E_2 = g$.

For $M = A + bJ + cI$ and $p \mid (r + c)$, $p \mid (s + c)$ we have $ME_1 = JE_1 = 0$ (over \mathbb{F}_p) so that $\text{rk}_p \langle M, \mathbf{1} \rangle \leq g + 1$, and hence $\text{rk}_p M \leq g$ (and similarly $\text{rk}_p M \leq f$) in case $\mathbf{1} \notin \langle M \rangle$.

Much more detail is given in [49] and [257].

In the table below we give for a few strongly regular graphs for each prime p dividing $r - s$ the p -rank of $A - sI$ and the unique b_0 such that $\text{rk}_p(A - sI - b_0 J) = \text{rk}_p(A - sI - bJ) - 1$ for all $b \neq b_0$, or ‘-’ in case $\text{rk}_p(A - sI - bJ)$ is independent of b . (When $p \nmid v$ we are in the former case, and b_0 follows from the parameters. When $p \mid v$ and $p \nmid \mu$, we are in the latter case.)

For a description of most of these graphs, see [56].

Name	v	k	λ	μ	r^f	s^g	p	$\text{rk}_p(A - sI)$	b_0
Folded 5-cube	16	5	0	2	1^{10}	$(-3)^5$	2	6	-
Schläfli	27	16	10	8	4^6	$(-2)^{20}$	2	6	0
							3	7	-
$T(8)$	28	12	6	4	4^7	$(-2)^{20}$	2	6	0
							3	8	2
3 Chang graphs	28	12	6	4	4^7	$(-2)^{20}$	2	8	-
							3	8	2
$G_2(2)$	36	14	4	6	2^{21}	$(-4)^{14}$	2	8	-
							3	14	-
$Sp_4(3)$	40	12	2	4	2^{24}	$(-4)^{15}$	2	16	-
							3	11	1
$O_5(3)$	40	12	2	4	2^{24}	$(-4)^{15}$	2	10	-
							3	15	1
Hoffman-Singleton	50	7	0	1	2^{28}	$(-3)^{21}$	5	21	-
Gewirtz	56	10	0	2	2^{35}	$(-4)^{20}$	2	20	-
							3	20	1
M_{22}	77	16	0	4	2^{55}	$(-6)^{21}$	2	20	0
Brouwer-Haemers	81	20	1	6	2^{60}	$(-7)^{20}$	3	19	-
Higman-Sims	100	22	0	6	2^{77}	$(-8)^{22}$	2	22	-
							5	23	-
Hall-Janko	100	36	14	12	6^{36}	$(-4)^{63}$	2	36	0
							5	23	-

continued...

Name	v	k	λ	μ	r^f	s^g	p	$\text{rk}_p(A - sI)$	b_0
$GQ(3, 9)$	112	30	2	10	2^{90}	$(-10)^{21}$	2	22	-
							3	20	1
001... in $S(5, 8, 24)$	120	42	8	18	2^{99}	$(-12)^{20}$	2	20	-
							7	20	5
$Sp_4(5)$	156	30	4	6	4^{90}	$(-6)^{65}$	2	66	-
							5	36	1
Sub McL	162	56	10	24	2^{140}	$(-16)^{21}$	2	20	0
							3	21	-
Edges of Ho-Si	175	72	20	36	2^{153}	$(-18)^{21}$	2	20	0
							5	21	-
01... in $S(5, 8, 24)$	176	70	18	34	2^{154}	$(-18)^{21}$	2	22	-
							5	22	3
switched version of previous	176	90	38	54	2^{153}	$(-18)^{22}$	2	22	-
							5	22	3
Cameron	231	30	9	3	9^{55}	$(-3)^{175}$	2	55	1
							3	56	1
Berlekamp-van Lint-Seidel	243	22	1	2	4^{132}	$(-5)^{110}$	3	67	-
Delsarte	243	110	37	60	2^{220}	$(-25)^{22}$	3	22	-
$S(4, 7, 23)$	253	112	36	60	2^{230}	$(-26)^{22}$	2	22	0
							7	23	5
McLaughlin	275	112	30	56	2^{252}	$(-28)^{22}$	2	22	0
							3	22	1
							5	23	-
switched version of previous plus isolated point	276	140	58	84	2^{252}	$(-28)^{23}$	2	24	-
							3	23	2
							5	24	3
$G_2(4)$	416	100	36	20	20^{65}	$(-4)^{350}$	2	38	-
							3	65	1
Dodecads mod 1	1288	792	476	504	8^{1035}	$(-36)^{252}$	2	22	0
							11	230	3

Table 12.1: p -ranks of some strongly regular graphs

12.8 Smith Normal Form

The *Smith Normal Form* $S(M)$ of an integral matrix M is a diagonal matrix $S(M) = PMQ = \text{diag}(s_1, \dots, s_n)$, where P and Q are integral with determinant ± 1 and $s_1 | s_2 | \dots | s_n$. It exists and is uniquely determined up to the signs of the s_i . The s_i are called the *elementary divisors* or *invariant factors*. For example, if $M = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 3 & 1 & 1 & 1 \end{bmatrix}$, then $S(M) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \end{bmatrix}$.

Let $\langle M \rangle$ denote the row space of M over \mathbb{Z} . By the fundamental theorem for finitely generated abelian groups, the group $\mathbb{Z}^n / \langle M \rangle$ is isomorphic to a direct sum $\mathbb{Z}_{s_1} \oplus \dots \oplus \mathbb{Z}_{s_m} \oplus \mathbb{Z}^s$ for certain s_1, \dots, s_m, s , where $s_1 | \dots | s_m$. Since $\mathbb{Z}^n / \langle M \rangle \cong \mathbb{Z}^n / \langle S(M) \rangle$, we see that $\text{diag}(s_1, \dots, s_m, 0^t)$ is the Smith Normal Form of M , when M has r rows and $n = m + s$ columns, and $t = \min(r, n) - m$.

If M is square then $\prod s_i = \det S(M) = \pm \det M$. More generally, $\prod_{i=1}^t s_i$ is the g.c.d. of all minors of M of order t .

The Smith Normal Form is a finer invariant than the p -rank: the p -rank is just the number of s_i not divisible by p . (It follows that if M is square and $p^e || \det M$, then $\text{rk}_p M \geq n - e$.)

We give some examples of graphs distinguished by Smith Normal Form or p -rank.

Example Let A and B be the adjacency matrices of the lattice graph $K_4 \times K_4$ and the Shrikhande graph. Then $S(A) = S(B) = \text{diag}(1^6, 2^4, 4^5, 12)$, but $S(A + 2I) = \text{diag}(1^6, 8^1, 0^9)$ and $S(B + 2I) = \text{diag}(1^6, 2^1, 0^9)$. All have 2-rank equal to 6.

Example An example where the p -rank suffices to distinguish, is given by the Chang graphs, strongly regular graphs with the same parameters as the triangular graph $T(8)$, with $(v, k, \lambda, \mu) = (28, 12, 6, 4)$ and spectrum $12^1 4^7 (-2)^{20}$. If A is the adjacency matrix of the triangular graph and B that of one of the Chang graphs then $S(A) = \text{diag}(1^6, 2^{15}, 8^6, 24^1)$ and $S(B) = \text{diag}(1^8, 2^{12}, 8^7, 24^1)$, so that A and B have different 2-rank.

Example Another example is given by the point graph and the line graph of the $GQ(3, 3)$ constructed in §8.6.2. The 2-ranks of the adjacency matrices are 10 and 16 respectively.




Figure 12.1: Graphs with same Laplacian SNF $(1^3, 5, 15, 0)$

Concerning the Smith Normal Form of the Laplacian, Grone, Merris & Watkins [167] gave the pair of graphs in Figure 12.1 that both have $S(L) = \text{diag}(1^3, 5, 15, 0)$. The Laplacian spectrum of the left one (which is $K_2 \times K_3$) is $0, 2, 3^2, 5^2$. That of the right one is $0, 0.914, 3.572, 5^2, 5.514$, where the three non-integers are roots of $\lambda^3 - 10\lambda^2 + 28\lambda - 18 = 0$.

12.8.1 Smith Normal Form and spectrum

There is no very direct connection between Smith Normal Form and spectrum. For example, the matrix $\begin{bmatrix} 3 & 1 \\ 1 & 3 \end{bmatrix}$ has eigenvalues 2 and 4, and invariant factors 1 and 8.

Proposition 12.8.1 *Let M be an integral matrix of order n , with invariant factors s_1, \dots, s_n .*

- (i) *If a is an integral eigenvalue of M , then $a|s_n$.*
- (ii) *If a is an integral eigenvalue of M with geometric multiplicity m , then $a|s_{n-m+1}$.*
- (iii) *If M is diagonalizable with distinct eigenvalues a_1, \dots, a_m , all integral, then we have $s_n|a_1 a_2 \cdots a_m$.*

Proof. Part (i) is a special case of (ii). Part (ii) is Proposition 12.8.4 below. For (iii) we may assume that all a_i are nonzero. It suffices to show that every element in $\mathbb{Z}^n / \langle M \rangle$ has an order dividing $a_1 a_2 \cdots a_m$. We show by induction on k that if $u = \sum u_i$ is integral and is sum of k left eigenvectors u_i of M , with $u_i M = a_i u_i$, then $a_1 \cdots a_k u \in \langle M \rangle$. Indeed, since $uM = \sum a_i u_i \in \langle M \rangle$ and $a_k u - uM = \sum (a_k - a_i) u_i$ is integral and sum of at most $k-1$ eigenvectors, we find by induction that $a_1 \cdots a_{k-1} (a_k u - uM) \in \langle M \rangle$, hence $a_1 \cdots a_k u \in \langle M \rangle$. \square

The invariant factors are determined when we know for each prime p and each $i \geq 0$ how many invariant factors are divisible by p^i , and the following proposition tells us.

Proposition 12.8.2 *Let A be an integral matrix of order n , p a prime number and i a nonnegative integer. Put $M_i := M_i(A) := \{x \in \mathbb{Z}^n \mid p^{-i} Ax \in \mathbb{Z}^n\}$. Let $\overline{M}_i \subseteq \mathbb{F}_p^n$ be the mod p reduction of M_i . Then \overline{M}_i is an \mathbb{F}_p -vector space, and the number of invariant factors of A divisible by p^i equals $\dim_p \overline{M}_i$.*

Proof. $\dim_p \overline{M}_i$ does not change when A is replaced by PAQ where P and Q are integral matrices of determinant 1. So we may assume that A is already in Smith Normal Form. Now the statement is obvious. \square

There is a dual statement:

Proposition 12.8.3 *Let A be an integral matrix of order n , p a prime number and i a nonnegative integer. Put $N_i := N_i(A) := \{p^{-i} Ax \mid x \in M_i\}$. Then the number of invariant factors of A not divisible by p^{i+1} equals $\dim_p \overline{N}_i$.*

Proof. $\dim_p \overline{N}_i$ does not change when A is replaced by PAQ where P and Q are integral matrices of determinant 1. So we may assume that A is already in Smith Normal Form. Now the statement is obvious. \square

Proposition 12.8.4 *Let A be a square integral matrix with integral eigenvalue a of (geometric) multiplicity m . Then the number of invariant factors of A divisible by a is at least m .*

Proof. Let $W = \{x \in \mathbb{Q}^n \mid Ax = ax\}$ be the a -eigenspace of A over \mathbb{Q} , so that $\dim_{\mathbb{Q}}(W) = m$. By the Proposition 12.8.2 it suffices to show that $\dim_p \overline{W} = m$ for all primes p , where \overline{W} is the mod p reduction of $W \cap \mathbb{Z}^n$. Pick a basis x_1, \dots, x_m of W consisting of m integral vectors, chosen in such a way that the $n \times m$ matrix X that has columns x_j has a (nonzero) minor of order m with the minimum possible number of factors p . If upon reduction mod p these vectors become dependent, that is, if $\sum c_j \overline{x_j} = 0$ where not all c_j vanish, then $\sum c_j x_j$ has coefficients divisible by p , so that $y := \frac{1}{p} \sum c_j x_j \in W \cap \mathbb{Z}^n$, and we can replace some x_j (with nonzero c_j) by y and get a matrix X' where the minors have fewer factors p , contrary to assumption. So, the x_i remain independent upon reduction mod p , and $\dim_p \overline{W} = m$. \square

Example Let $q = p^t$ for some prime p . Consider the adjacency matrix A of the graph Γ of which the vertices are the lines of $PG(3, q)$, where two lines are adjacent when they are disjoint. This graph is strongly regular, with eigenvalues $k = q^4$, $r = q$, $s = -q^2$ and multiplicities 1 , $f = q^4 + q^2$, $g = q^3 + q^2 + q$, respectively. Since $\det A$ is a power of p , all invariant factors are powers of p . Let p^i occur as invariant factor with multiplicity e_i .

Claim. We have $e_0 + e_1 + \cdots + e_t = f$ and $e_{2t} + \cdots + e_{3t} = g$ and $e_{4t} = 1$ and $e_i = 0$ for $t < i < 2t$ and $3t < i < 4t$ and $i > 4t$. Moreover, $e_{3t-i} = e_i$ for $0 \leq i < t$.

Proof. The total number of invariant factors is the size of the matrix, so $\sum_i e_i = f + g + 1$. The number of factors p in $\det A$ is $\sum_i i e_i = t(f + 2g + 4)$. Hence $\sum_i (i - t)e_i = t(g + 3)$.

Let $m_i := \sum_{j \geq i} e_j$. By (the proof of) Proposition 12.8.4 we have $m_{4t} \geq 1$ and $m_{2t} \geq g + 1$. (The $+1$ follows because $\mathbf{1}$ is orthogonal to eigenvectors with eigenvalue other than k , but has a nonzero (mod p) inner product with itself, so that $\mathbf{1} \notin W$ for an eigenspace W with $\mathbf{1} \notin W$.)

The matrix A satisfies the equation $(A - rI)(A - sI) = \mu J$, that is, $A(A + q(q - 1)I) = q^3 I + q^3(q - 1)J$, and the right-hand side is divisible by p^{3t} . If $x \in \mathbb{Z}^n$ and $p^{-i}(A + q(q - 1)I)x \in \mathbb{Z}^n$, then $p^{-i}(A + q(q - 1)I)x \in M_{3t-i}(A)$ for $0 \leq i \leq 3t$. If $0 \leq i < t$, then $p^{-i}q(q - 1)x = 0 \pmod{p}$, so that $\bar{N}_i \subseteq \bar{M}_{3t-i}$. Also $\mathbf{1} \in \bar{M}_{3t-i}$, while $\mathbf{1} \notin \bar{N}_i$ because $\mathbf{1}^\top p^{-i}Ax = p^{4t-i}Ax$ reduces to $0 \pmod{p}$ for integral x , unlike $\mathbf{1}^\top \mathbf{1}$. By Proposition 12.8.3 we find $m_{3t-i} \geq e_0 + \cdots + e_i + 1$ ($0 \leq i < t$).

Adding the inequalities $-\sum_{0 \leq i \leq h} e_i + \sum_{i \geq 3t-h} e_i \geq 1$ ($0 \leq h < t$), and $t \sum_{i \geq 2t} e_i \geq t(g + 1)$ and $t \sum_{i \geq 4t} e_i \geq t$ yields

$$\sum_{0 \leq i < t} (i - t)e_i + \sum_{2t \leq i \leq 3t} (i - t)e_i + 2t \sum_{3t+1 \leq i < 4t} e_i + 3t \sum_{i \geq 4t} e_i \geq t(g + 3)$$

and equality must hold everywhere since $\sum_i (i - t)e_i = t(g + 3)$. \square

Note that our conclusion also holds for any strongly regular graph with the same parameters as this graph on the lines of $PG(3, q)$.

In the particular case $q = p$, the invariant factors are $1, p, p^2, p^3, p^4$ with multiplicities $e, f - e, g - e, e, 1$, respectively, where $e = \frac{1}{3}p(2p^2 + 1)$ in the case of the lines of $PG(3, p)$ (cf. [135]). Indeed, the number e of invariant factors not divisible by p is the p -rank of A , determined in Sin [284].

For $p = 2$, there are 3854 strongly regular graphs with parameters $(35, 16, 6, 8)$ ([244]), and the 2-ranks occurring are 6, 8, 10, 12, 14 (with frequencies 1, 3, 44, 574, 3232, respectively)—they must be even because A is alternating (mod 2).

The invariant factors of the disjointness graph of the lines of $PG(3, 4)$ are $1^{36} 2^{16} 4^{220} 16^{32} 32^{16} 64^{36} 256^1$, with multiplicities written as exponents.

One can generalize the above observations, and show for example that if p is a prime, and A is the adjacency matrix of a strongly regular graph, and $p^a || k$, $p^b || r$, $p^c || s$, where $a \geq b + c$ and $p \nmid v$, and A has e_i invariant factors s_j with $p^i || s_j$, then $e_i = 0$ for $\min(b, c) < i < \max(b, c)$ and $b + c < i < a$ and $i > a$. Moreover, $e_{b+c-i} = e_i$ for $0 \leq i < \min(b, c)$.

Chapter 13

Spectral characterizations

In this chapter, we consider the question to what extent graphs are determined by their spectrum. First we give several constructions of families of cospectral graphs, and then give cases in which it has been shown that the graph is determined by its spectrum.

Let us abbreviate ‘determined by the spectrum’ to DS. Here, of course, ‘spectrum’ (and DS) depends on the type of adjacency matrix. If the matrix is not specified, we mean the ordinary adjacency matrix.

Large parts of this chapter were taken from Van Dam & Haemers [117, 118].

13.1 Generalized adjacency matrices

Let $A = A_\Gamma$ be the adjacency matrix of a graph Γ . The choice of 0, 1, 0 in A to represent equality, adjacency and non-adjacency was rather arbitrary, and one can more generally consider a matrix $xI + yA + z(J - I - A)$ that uses x, y, z instead. Any such matrix, with $y \neq z$, is called a *generalized adjacency matrix* of Γ . The spectrum of any such matrix is obtained by scaling and shifting from that of a matrix of the form $A + yJ$, so for matters of cospectrality we can restrict ourselves to this case.

Call two graphs Γ and Δ *y-cospectral* (for some real y) when $A_\Gamma - yJ$ and $A_\Delta - yJ$ have the same spectrum. Then 0-cospectral is what we called cospectral, and $\frac{1}{2}$ -cospectral is Seidel-cospectral, and 1-cospectrality is cospectrality for the complementary graphs. Call two graphs *just y-cospectral* when they are *y-cospectral* but not *z-cospectral* for any $z \neq y$.

The graphs $K_{1,4}$ and $K_1 + C_4$ are just 0-cospectral. The graphs $2K_3$ and $2K_1 + K_4$ are just $\frac{1}{3}$ -cospectral. The graphs $K_1 + C_6$ and \hat{E}_6 (cf. §1.3.7) are *y-cospectral* for all y .

Proposition 13.1.1

- (i) (Johnson & Newman [208]) *If two graphs are y-cospectral for two distinct values of y, then for all y.*
- (ii) (Van Dam, Haemers & Koolen [119]) *If two graphs are y-cospectral for an irrational value of y, then for all y.*

Proof. Define $p(x, y) = \det(A_\Gamma - xI - yJ)$. Thus for fixed y , $p(x, y)$ is the characteristic polynomial of $A_\Gamma - yJ$. Since J has rank 1, the degree in y of $p(x, y)$ is 1 (this follows from Gaussian elimination in $A_\Gamma - xI - yJ$), so there exist integers a_0, \dots, a_n and b_0, \dots, b_n such that

$$p(x, y) = \sum_{i=0}^n (a_i + b_i y) x^i.$$

Suppose Γ and Γ' are y -cospectral for some $y = y_0$ but not for all y . Then the corresponding polynomials $p(x, y)$ and $p'(x, y)$ are not identical, whilst $p(x, y_0) = p'(x, y_0)$. This implies that $a_i + b_i y_0 = a'_i + b'_i y_0$ with $b_i \neq b'_i$ for some i . So $y_0 = (a'_i - a_i)/(b_i - b'_i)$ is unique and rational. \square

Van Dam, Haemers & Koolen [119] show that there is a pair of nonisomorphic just y -cospectral graphs if and only if y is rational.

Values of y other than $0, \frac{1}{2}, 1$ occur naturally when studying subgraphs of strongly regular graphs.

Proposition 13.1.2 *Let Γ be strongly regular with vertex set X of size n , and let θ be an eigenvalue other than the valency k . Let $y = (k - \theta)/n$. Then for each subset S of X , the spectrum of Γ and the y -spectrum of the graph induced on S determines the y -spectrum of the graph induced on $X \setminus S$.*

Proof. Since $A - yJ$ has only two eigenvalues, this follows immediately from Lemma 2.11.1. \square

This can be used to produce cospectral pairs. For example, let Γ be the Petersen graph, and let S induce a 3-coclique. Then the y -spectrum of the graph induced on $X \setminus S$ is determined by that on S , and does not depend on the coclique chosen. Since θ can take two values, the graphs induced on the complement of a 3-coclique (\hat{E}_6 and $K_1 + C_6$) are y -cospectral for all y .

13.2 Constructing cospectral graphs

Many constructions of cospectral graphs are known. Most constructions from before 1988 can be found in [106, §6.1] and [105, §1.3]; see also [154, §4.6]. More recent constructions of cospectral graphs are presented by Seress [280], who gives an infinite family of cospectral 8-regular graphs. Graphs cospectral to distance-regular graphs can be found in [48], [117], [183], and in §13.2.2. Notice that the mentioned graphs are regular, so they are cospectral with respect to any generalized adjacency matrix, which in this case includes the Laplacian matrix.

There exist many more papers on cospectral graphs. On regular, as well as non-regular graphs, and with respect to the Laplacian matrix as well as the adjacency matrix. We mention [41], [148], [187], [233], [245] and [264], but don't claim to be complete.

Here we discuss four construction methods for cospectral graphs. One used by Schwenk to construct cospectral trees, one from incidence geometry to construct graphs cospectral with distance-regular graphs, one presented by Godsil and McKay, which seems to be the most productive one, and finally one due to Sunada.


13.2.1 Trees

Let Γ and Δ be two graphs, with vertices x and y , respectively. Schwenk [267] examined the spectrum of what he called the *coalescence* of these graphs at x and y , namely, the result $\Gamma +_{x,y} \Delta$ of identifying x and y in the disjoint union $\Gamma + \Delta$. He proved the following (see also [106, p.159] and [154, p.65]).

Lemma 13.2.1 *Let Γ and Γ' be cospectral graphs and let x and x' be vertices of Γ and Γ' respectively. Suppose that $\Gamma - x$ (that is the subgraph of Γ obtained by deleting x) and $\Gamma' - x'$ are cospectral too. Let Δ be an arbitrary graph with a fixed vertex y . Then $\Gamma +_{x,y} \Delta$ is cospectral with $\Gamma' +_{x',y} \Delta$.*

Proof. Let z be the vertex of $Z := \Gamma +_{x,y} \Delta$ that is the result of identifying x and y . A directed cycle in Z cannot meet both $\Gamma - x$ and $\Delta - y$. By §1.2.1 the characteristic polynomial $p(t)$ of Z can be expressed in the numbers of unions of directed cycles with given number of vertices and of components. We find $p(t) = p_{\Gamma-x}(t)p_{\Delta}(t) + p_{\Gamma}(t)p_{\Delta-y}(t) - tp_{\Gamma-x}(t)p_{\Delta-y}(t)$. \square

For example, let $\Gamma = \Gamma'$ be as given below, then $\Gamma - x$ and $\Gamma - x'$ are cospectral, because they are isomorphic.



Suppose $\Delta = P_3$ and let y be the vertex of degree 2. Then Lemma 13.2.1 shows that the graphs in Figure 13.1 are cospectral.




Figure 13.1: Cospectral trees

It is clear that Schwenk's method is very suitable for constructing cospectral trees. In fact, the lemma above enabled him to prove his famous theorem:

Theorem 13.2.2 *With respect to the adjacency matrix, almost all trees are non-DS.*

After Schwenk's result, trees were proved to be almost always non-DS with respect to all kinds of matrices. Godsil and McKay [155] proved that almost all trees are non-DS with respect to the adjacency matrix of the complement \bar{A} , while McKay [243] proved it for the Laplacian matrix L and for the distance matrix D .

13.2.2 Partial linear spaces

A *partial linear space* consists of a (finite) set of points \mathcal{P} , and a collection \mathcal{L} of subsets of \mathcal{P} called lines, such that two lines intersect in at most one point (and consequently, two points are on at most one line). Let $(\mathcal{P}, \mathcal{L})$ be such a partial linear space and assume that each line has exactly q points, and each point is on q lines. Then clearly $|\mathcal{P}| = |\mathcal{L}|$. Let N be the point-line incidence matrix of $(\mathcal{P}, \mathcal{L})$. Then $NN^T - qI$ and $N^TN - qI$ both are the adjacency matrix

of a graph, called the *point graph* (also known as *collinearity graph*) and *line graph* of $(\mathcal{P}, \mathcal{L})$, respectively. These graphs are cospectral, since NN^\top and $N^\top N$ are. But in many examples they are non-isomorphic. An example was given in §11.3.1.

13.2.3 GM switching

Seidel switching was discussed above in §1.8.2. No graph with more than one vertex is DS for the Seidel adjacency matrix. In some cases Seidel switching also leads to cospectral graphs for the adjacency spectrum, for example when graph and switched graph are regular of the same degree.

Godsil and McKay [156] consider a different kind of switching and give conditions under which the adjacency spectrum is unchanged by this operation. We will refer to their method as GM switching. (See also §1.8.3.) Though GM switching has been invented to make cospectral graphs with respect to the adjacency matrix, the idea also works for the Laplacian and the signless Laplacian matrix, as will be clear from the following formulation.

Theorem 13.2.3 *Let N be a $(0,1)$ -matrix of size $b \times c$ (say) whose column sums are 0, b or $b/2$. Define \tilde{N} to be the matrix obtained from N by replacing each column v with $b/2$ ones by its complement $\mathbf{1} - v$. Let B be a symmetric $b \times b$ matrix with constant row (and column) sums, and let C be a symmetric $c \times c$ matrix. Put*

$$M = \begin{bmatrix} B & N \\ N^\top & C \end{bmatrix} \quad \text{and} \quad \tilde{M} = \begin{bmatrix} B & \tilde{N} \\ \tilde{N}^\top & C \end{bmatrix}.$$

Then M and \tilde{M} are cospectral.

Proof. Define $Q = \begin{bmatrix} \frac{2}{b}J - I_b & 0 \\ 0 & I_c \end{bmatrix}$. Then $Q^{-1} = Q$ and $QMQ^{-1} = \tilde{M}$. \square

The matrix partition used in [156] (and in §1.8.3) is more general than the one presented here. But this simplified version suffices for our purposes: to show that GM switching produces many cospectral graphs.

If M and \tilde{M} are adjacency matrices of graphs then GM switching gives cospectral graphs with cospectral complements and hence, by the result of Johnson & Newman quoted in §13.1, it produces cospectral graphs with respect to any generalized adjacency matrix.

If one wants to apply GM switching to the Laplacian matrix L of a graph Γ , take $M = -L$ and let B and C (also) denote the sets of vertices indexing the rows and columns of the matrices B and C , respectively. The requirement that the matrix B has constant row sums means that N has constant row sums, that is, the vertices of B all have the same number of neighbors in C .

For the signless Laplacian matrix, take $M = Q$. Now all vertices in B must have the same number of neighbors in C , and, in addition, the subgraph of Γ induced by B must be regular.

When Seidel switching preserves the valency of a graph, it is a special case of GM switching, where all columns of N have $b/2$ ones. So the above theorem also gives sufficient conditions for Seidel switching to produce cospectral graphs with respect to the adjacency matrix A and the Laplacian matrix L .

If $b = 2$, GM switching just interchanges the two vertices of B , and we call it trivial. But if $b \geq 4$, GM switching almost always produces non-isomorphic graphs. In Figures 13.2 and 13.3 we have two examples of pairs of cospectral



Figure 13.2: Two graphs cospectral w.r.t. any generalized adjacency matrix



Figure 13.3: Two graphs cospectral w.r.t. the Laplacian matrix

graphs produced by GM switching. In both cases $b = c = 4$ and the upper vertices correspond to B and the lower vertices to C . In the example of Figure 13.2, B induces a regular subgraph and so the graphs are cospectral with respect to every generalized adjacency matrix. In the example of Figure 13.3 all vertices of B have the same number of neighbors in C , so the graphs are cospectral with respect to the Laplacian matrix L .

13.2.4 Sunada's method

As a corollary of the discussion in §6.3 we have:

Proposition 13.2.4 *Let Γ be a finite graph, and G a group of automorphisms. If H_1 and H_2 are subgroups of G such that Γ is a cover of Γ/H_i ($i = 1, 2$) and such that each conjugacy class of G meets H_1 and H_2 in the same number of elements, then the quotients Γ/H_i ($i = 1, 2$) have the same spectrum and the same Laplacian spectrum.*

Sunada [289] did this for manifolds, and the special case of graphs was discussed in [187]. See also [41].

Proof. The condition given just means that the induced characters $1_{H_i}^G$ ($i = 1, 2$) are the same. Now apply Lemma 6.3.1 with $M = A$ and $M = L$. \square

Brooks [41] shows a converse: any pair of regular connected cospectral graphs arises from this construction.

13.3 Enumeration

13.3.1 Lower bounds

GM switching gives lower bounds for the number of pairs of cospectral graphs with respect to several types of matrices.

Let Γ be a graph on $n-1$ vertices and fix a set X of three vertices. There is a unique way to extend Γ by one vertex x to a graph Γ' , such that $X \cup \{x\}$ induces

a regular graph in Γ' and that every other vertex in Γ' has an even number of neighbors in $X \cup \{x\}$. Thus the adjacency matrix of Γ' admits the structure of Theorem 13.2.3, where B corresponds to $X \cup \{x\}$. This implies that from a graph Γ on $n - 1$ vertices one can make $\binom{n-1}{3}$ graphs with a cospectral mate on n vertices (with respect to any generalized adjacency matrix) and every such n -vertex graph can be obtained in four ways from a graph on $n - 1$ vertices. Of course some of these graphs may be isomorphic, but the probability of such a coincidence tends to zero as $n \rightarrow \infty$ (see [185] for details). So, if g_n denotes the number of non-isomorphic graphs on n vertices, then:

Theorem 13.3.1 *The number of graphs on n vertices which are non-DS with respect to any generalized adjacency matrix is at least*

$$\left(\frac{1}{24} - o(1)\right) n^3 g_{n-1}.$$

The fraction of graphs with the required condition with $b = 4$ for the Laplacian matrix is roughly $2^{-n} n \sqrt{n}$. This leads to the following lower bound (again see [185] for details):

Theorem 13.3.2 *The number of non-DS graphs on n vertices with respect to the Laplacian matrix is at least*

$$rn\sqrt{n}g_{n-1},$$

for some constant $r > 0$.

In fact, a lower bound like the one in Theorem 13.3.2 can be obtained for any matrix of the form $A + \alpha D$, including the signless Laplacian matrix Q .

13.3.2 Computer results

The mentioned papers [155] and [156] by Godsil and McKay also give interesting computer results for cospectral graphs. In [156] all graphs up to 9 vertices are generated and checked on cospectrality. This enumeration has been extended to 11 vertices by Haemers & Spence [185], and cospectrality was tested with respect to the adjacency matrix A , the set of generalized adjacency matrices $(A \& \bar{A})$, the Laplacian matrix L , and the signless Laplacian matrix Q . The results are in Table 13.1, where we give the fractions of non-DS graphs for each of the four cases. The last three columns give the fractions of graphs for which GM switching gives cospectral non-isomorphic graphs with respect to A , L and Q , respectively. So column GM- A gives a lower bound for column $A \& \bar{A}$ (and, of course, for column A), column GM- L is a lower bound for column L , and column GM- Q is a lower bound for column Q .

Notice that for $n \leq 4$ there are no cospectral graphs with respect to A or to L , but there is one such pair with respect to Q , namely $K_{1,3}$ and $K_1 + K_3$. For $n = 5$ there is just one pair with respect to A . This is of course the Saltire pair $(K_{1,4}$ and $K_1 + C_4)$.

An interesting result from the table is that the fraction of non-DS graphs is nondecreasing for small n , but starts to decrease at $n = 10$ for A , at $n = 9$ for L , and at $n = 6$ for Q . Especially for the Laplacian and the signless Laplacian matrix, these data suggest that the fraction of DS graphs might tend to 1 as

n	# graphs	A	$A \& \bar{A}$	L	Q	GM- A	GM- L	GM- Q
2	2	0	0	0	0	0	0	0
3	4	0	0	0	0	0	0	0
4	11	0	0	0	0.182	0	0	0
5	34	0.059	0	0	0.118	0	0	0
6	156	0.064	0	0.026	0.103	0	0	0
7	1044	0.105	0.038	0.125	0.098	0.038	0.069	0
8	12346	0.139	0.094	0.143	0.097	0.085	0.088	0
9	274668	0.186	0.160	0.155	0.069	0.139	0.110	0
10	12005168	0.213	0.201	0.118	0.053	0.171	0.080	0.001
11	1018997864	0.211	0.208	0.090	0.038	0.174	0.060	0.001
12	165091172592	0.188		0.060	0.027			

Table 13.1: Fractions of non-DS graphs

$n \rightarrow \infty$. In addition, the table shows that the majority of non-DS graphs with respect to $A \& \bar{A}$ and L comes from GM switching (at least for $n \geq 7$). If this tendency continues, the lower bounds given in Theorems 13.3.1 and 13.3.2 will be asymptotically tight (with maybe another constant) and almost all graphs will be DS for all three cases. Indeed, the fraction of graphs that admit a non-trivial GM switching tends to zero as n tends to infinity, and the partitions with $b = 4$ account for most of these switchings (see also [156]). For data for $n = 12$, see [60] and [287].

13.4 DS graphs

In Section 13.2 we saw that many constructions for non-DS graphs are known, and in the previous section we remarked that it seems more likely that almost all graphs are DS, than that almost all graphs are non-DS. Yet much less is known about DS graphs than about non-DS graphs. For example, we do not know of a satisfying counterpart to the lower bounds for non-DS graphs given in §13.3.1. The reason is that it is not easy to prove that a given graph is DS. Below we discuss the graphs known to be DS. The approach is via structural properties of a graph that follow from the spectrum. So let us start with a short survey of such properties.

13.4.1 Spectrum and structure

Let us first investigate for which matrices one can see from the spectrum whether the graph is regular.

Proposition 13.4.1 *Let D denote the diagonal matrix of degrees. If a regular graph is cospectral with a non-regular one with respect to the matrix $R = A + \beta J + \gamma D + \delta I$, then $\gamma = 0$ and $-1 < \beta < 0$.*

Proof. W.l.o.g. $\delta = 0$. Let Γ be a graph with the given spectrum, and suppose that Γ has n vertices and vertex degrees d_i ($1 \leq i \leq n$).

First suppose that $\gamma \neq 0$. Then $\sum_i d_i$ is determined by $\text{tr}(R)$ and hence by the spectrum of R . Since $\text{tr}(R^2) = \beta^2 n^2 + (1 + 2\beta + 2\beta\gamma) \sum_i d_i + \gamma^2 \sum_i d_i^2$, it

follows that also $\sum_i d_i^2$ is determined by the spectrum. Now Cauchy's inequality states that $(\sum_i d_i)^2 \leq n \sum_i d_i^2$ with equality if and only if $d_1 = \dots = d_n$. This shows that regularity of the graph can be seen from the spectrum of R .

Now suppose $\gamma = 0$ and $\beta \neq -1/2$. By considering $\text{tr}(R^2)$ we see that $\sum_i d_i$ is determined by the spectrum of R . The matrix $R = A + \beta J$ has average row sum $r = \beta n + \sum_i d_i/n$ determined by its spectrum. Let R have eigenvalues $\theta_1 \geq \dots \geq \theta_n$. By interlacing, $\theta_1 \geq r \geq \theta_n$, and equality on either side implies that R has constant row sums, and Γ is regular. On the other hand, if $\beta \geq 0$ (resp. $\beta \leq -1$), then R (resp. $-R$) is a nonnegative matrix, hence if Γ is regular, then $\mathbf{1}$ is an eigenvector for eigenvalue $r = \theta_1$ (resp. $r = -\theta_n$). Thus also here regularity of the graph can be seen from the spectrum. \square

It remains to see whether one can see from the spectrum of $A - yJ$ (with $0 < y < 1$) whether the graph is regular. For $y = \frac{1}{2}$ the answer is clearly no: The Seidel adjacency matrix is $S = J - I - 2A$, and for S a regular graph can be cospectral with a non-regular one (e.g. K_3 and $K_1 + K_2$), or with another regular one with different valency (e.g. $4K_1$ and C_4). Chesnokov & Haemers [82] constructed pairs of y -cospectral graphs where one is regular and the other not for all rational y , $0 < y < 1$. Finally, if y is irrational, then one can deduce regularity from the spectrum of $A - yJ$ by Proposition 13.1.1(ii).

Corollary 13.4.2 *For regular graphs, being DS (or not DS) is equivalent for the adjacency matrix, the adjacency matrix of the complement, the Laplacian, and the signless Laplacian matrix.*

Proof. For each of these matrices the above proposition says that regularity can be recognized. It remains to find the valency k . For A , \bar{A} , Q , the largest eigenvalue is k , $n - 1 - k$, $2k$, respectively. For L , the trace is nk . \square

Lemma 13.4.3 *For the adjacency matrix, the Laplacian matrix and the signless Laplacian matrix of a graph Γ , the following can be deduced from the spectrum.*

- (i) *The number of vertices.*
- (ii) *The number of edges.*
- (iii) *Whether Γ is regular.*
- (iv) *Whether Γ is regular with any fixed girth.*

For the adjacency matrix the following follows from the spectrum.

- (v) *The number of closed walks of any fixed length.*
- (vi) *Whether Γ is bipartite.*

For the Laplacian matrix the following follows from the spectrum.

- (vii) *The number of components.*
- (viii) *The number of spanning trees.*

Proof. Part (i) is clear. For L and Q the number of edges is twice the trace of the matrix, while parts (ii) and (v) for A were shown in Proposition 1.3.1. Part (vi) follows from (v), since Γ is bipartite if and only if Γ has no closed walks of odd length. Part (iii) follows from Proposition 13.4.1, and (iv) follows from (iii) and the fact that in a regular graph the number of closed walks of length less than the girth depends on the degree only. Parts (vii) and (viii) follow from Propositions 1.3.7 and 1.3.4. \square



Figure 13.4: Two graphs cospectral w.r.t. the Laplacian matrix
(Laplacian spectrum: $0, 3 - \sqrt{5}, 2, 3, 3, 3 + \sqrt{5}$)

The Saltire pair shows that (vii) and (viii) do not hold for the adjacency matrix. The two graphs of Figure 13.4 have cospectral Laplacian matrices. They illustrate that (v) and (vi) do not follow from the Laplacian spectrum. The graphs $K_1 + K_3$ and $K_{1,3}$ show that (v)–(viii) are false for the signless Laplacian matrix.

13.4.2 Some DS graphs

Lemma 13.4.3 immediately leads to some DS graphs.

Proposition 13.4.4 *The graphs K_n and $K_{m,m}$ and C_n and their complements are DS for any matrix $R = A + \beta J + \gamma D + \delta I$ for which regularity follows from the spectrum of R . In particular this holds for the matrices A, \bar{A}, L and R .*

Proof. Since these graphs are regular, we only need to show that they are DS with respect to the adjacency matrix. A graph cospectral with K_n has n vertices and $n(n-1)/2$ edges and therefore equals K_n . A graph cospectral with $K_{m,m}$ is regular and bipartite with $2m$ vertices and m^2 edges, so it is isomorphic to $K_{m,m}$. A graph cospectral with C_n is 2-regular with girth n , so it equals C_n . \square

Proposition 13.4.5 *The disjoint union of k complete graphs, $K_{m_1} + \dots + K_{m_k}$, is DS with respect to the adjacency matrix.*

Proof. The spectrum of the adjacency matrix A of any graph cospectral with $K_{m_1} + \dots + K_{m_k}$ equals $\{[m_1 - 1]^1, \dots, [m_k - 1]^1, [-1]^{n-k}\}$, where $n = m_1 + \dots + m_k$. This implies that $A + I$ is positive semi-definite of rank k , and hence $A + I$ is the matrix of inner products of n vectors in \mathbb{R}^k . All these vectors are unit vectors, and the inner products are 1 or 0. So two such vectors coincide or are orthogonal. This clearly implies that the vertices can be ordered in such a way that $A + I$ is a block diagonal matrix with all-ones diagonal blocks. The sizes of these blocks are non-zero eigenvalues of $A + I$. \square

In general, the disjoint union of complete graphs is not DS with respect to \bar{A} and L . The Saltire pair shows that $K_1 + K_4$ is not DS for \bar{A} , and $K_5 + 5K_2$ is not DS for L , because it is cospectral with the Petersen graph extended by five isolated vertices (both graphs have Laplacian spectrum $[0]^6 [2]^5 [5]^4$). Note that the above proposition also shows that a complete multipartite graph is DS with respect to \bar{A} .

Proposition 13.4.6 *The path with n vertices is determined by the spectrum of its adjacency matrix. More generally, each connected graph with largest eigenvalue less than 2 is determined by its spectrum.*

Proof. Let Γ be connected with n vertices and have largest eigenvalue less than 2, and let the graph Δ be cospectral. Then Δ does not contain a cycle, and has $n - 1$ edges, so is a tree. By Theorem 3.1.3 (and following remarks) we find that Δ is one of $A_n = P_n, D_n, E_6, E_7, E_8$, and has largest eigenvalue $2 \cos \frac{\pi}{h}$, where h is the Coxeter number. Now Δ is determined by n and h , that is, by its number of vertices and its largest eigenvalue. \square

In fact, P_n is also DS with respect to \bar{A} , L , and Q . The result for \bar{A} , however, is nontrivial and the subject of [134]. The hypothesis ‘connected’ here is needed, but we can describe precisely which pairs of graphs with largest eigenvalue less than 2 are cospectral.

Proposition 13.4.7 (i) $D_{n+2} + P_n$ is cospectral with $P_{2n+1} + P_1$ for $n \geq 2$.

(ii) $D_7 + P_2$ is cospectral with $E_6 + P_3$.

(iii) $D_{10} + P_2$ is cospectral with $E_7 + P_5$.

(iv) $D_{16} + P_4 + P_2$ is cospectral with $E_8 + P_9 + P_5$.

(v) If two graphs Γ and Δ with largest eigenvalue less than 2 are cospectral, then there exist integers a, b, c such that $\Delta + aP_4 + bP_2 + cP_1$ arises from $\Gamma + aP_4 + bP_2 + cP_1$ by (possibly repeatedly) replacing some connected components by some others cospectral with the replaced ones according to (i)–(iv).

For example, $P_{11} + P_2 + P_1$ is cospectral with $E_6 + P_5 + P_3$, and $P_{17} + P_2 + P_1$ is cospectral with $E_7 + P_8 + P_5$, and $P_{29} + P_4 + P_2 + P_1$ is cospectral with $E_8 + P_{14} + P_9 + P_5$, and $E_6 + D_{10} + P_7$ is cospectral with $E_7 + D_5 + P_{11}$, and $E_7 + D_4$ is cospectral with $D_{10} + P_1$, and $E_8 + D_6 + D_4$ is cospectral with $D_{16} + 2P_1$.

It follows that $P_{n_1} + \dots + P_{n_k}$ (with $n_i > 1$ for all i) and $D_{n_1} + \dots + D_{n_k}$ (with $n_i > 3$ for all i) are DS.

We do not know whether $P_{n_1} + \dots + P_{n_k}$ is DS with respect to \bar{A} . But it is easy to show that this graph is DS for L and for Q .

Proposition 13.4.8 The union of k disjoint paths, $P_{n_1} + \dots + P_{n_k}$ each having at least one edge, is DS with respect to the Laplacian matrix L and the signless Laplacian matrix Q .

Proof. The Laplacian and the signless Laplacian eigenvalues of P_n are $2 + 2 \cos \frac{\pi i}{n}$, $i = 1, \dots, n$ (see §1.4.4). Since P_n is bipartite, the signless Laplacian eigenvalues are the same (see Proposition 1.3.10).

Suppose Γ is a graph cospectral with $P_{n_1} + \dots + P_{n_k}$ with respect to L . Then all eigenvalues of L are less than 4. Lemma 13.4.3 implies that Γ has k components and $n_1 + \dots + n_k - k$ edges, so Γ is a forest. Let L' be the Laplacian matrix of $K_{1,3}$. The spectrum of L' equals $[0]^1 [1]^2 [4]^1$. If degree 3 (or more) occurs in Γ then $L' + D$ is a principal submatrix of L for some diagonal matrix D with nonnegative entries. But then $L' + D$ has largest eigenvalue at least 4, a contradiction. So the degrees in Γ are at most two and hence Γ is the disjoint union of paths. The length m (say) of the longest path follows from the largest eigenvalue. Then the other lengths follow recursively by deleting P_m from the graph and the eigenvalues of P_m from the spectrum.

For a graph Γ' cospectral with $P_{n_1} + \dots + P_{n_k}$ with respect to Q , the first step is to see that Γ' is a forest. But a circuit in Γ' gives a submatrix L' in Q with all row sums at least 4. So L' has an eigenvalue at least 4, a contradiction

(by Corollary 2.5.2), and it follows that Γ' is a forest and hence bipartite. Since for bipartite graphs L and Q have the same spectrum, Γ' is also cospectral with $P_{n_1} + \dots + P_{n_k}$ with respect to L , and we are done. \square

The above two propositions show that for A , \bar{A} , L , and Q the number of DS graphs on n vertices is bounded below by the number of partitions of n , which is asymptotically equal to $2^{\alpha\sqrt{n}}$ for some constant α . This is clearly a very poor lower bound, but we know of no better one.

In the above we saw that the disjoint union of some DS graphs is not necessarily DS. One might wonder whether the disjoint union of regular DS graphs with the same degree is always DS. The disjoint union of cycles is DS, as can be shown by an argument similar to that in the proof of Proposition 13.4.8. Also the disjoint union of some copies of a strongly regular DS graph is DS. In general we expect a negative answer, however.

13.4.3 Line graphs

The smallest adjacency eigenvalue of a line graph is at least -2 (see §1.4.5). Other graphs with least adjacency eigenvalue -2 are the cocktailparty graphs ($\overline{mK_2}$, the complement of m disjoint edges) and the so-called generalized line graphs, which are common generalizations of line graphs and cocktailparty graphs (see [105, Ch.1]). We will not need the definition of a generalized line graph, but only use the fact that if a generalized line graph is regular, it is a line graph or a cocktailparty graph. Graphs with least eigenvalue -2 have been characterised by Cameron, Goethals, Seidel and Shult [77] (cf. §7.4). They prove that such a graph is a generalized line graph or is in a finite list of exceptions that comes from root systems. Graphs in this list are called *exceptional graphs*. A consequence of the above characterisation is the following result of Cvetković & Doob [104, Thm.5.1] (see also [105, Thm.1.8]).

Theorem 13.4.9 *Suppose a regular graph Δ has the adjacency spectrum of the line graph $L(\Gamma)$ of a connected graph Γ . Suppose Γ is not one of the fifteen regular 3-connected graphs on 8 vertices, or $K_{3,6}$, or the semiregular bipartite graph with 9 vertices and 12 edges. Then Δ is the line graph $L(\Gamma')$ of a graph Γ' .*

It does not follow that the line graph of a connected regular DS graph, which is not one of the mentioned exceptions, is DS itself. The reason is that it can happen that two non-cospectral graphs Γ and Γ' have cospectral line graphs. For example, both $L(K_6)$ and $K_{6,10}$ have a line graph with spectrum $14^1 8^5 4^9 -2^{45}$, and both $L(\text{Petersen})$ and the incidence graph of the $2-(6, 3, 2)$ design have a line graph with spectrum $6^1 4^5 1^4 0^5 -2^{15}$. The following lemma gives necessary conditions for this phenomenon (cf. [69, Thm.1.7]).

Lemma 13.4.10 *Let Γ be a k -regular connected graph on n vertices and let Γ' be a connected graph such that $L(\Gamma)$ is cospectral with $L(\Gamma')$. Then either Γ' is cospectral with Γ , or Γ' is a semiregular bipartite graph with $n+1$ vertices and $nk/2$ edges, where $(n, k) = (b^2 - 1, ab)$ for integers a and b with $a \leq \frac{1}{2}b$.*

Proof. Suppose that Γ has m edges. Then $L(\Gamma)$ has m vertices.

If N is the point-edge incidence matrix of Γ , then NN^\top is the signless Laplacian of Γ , and $NN^\top - kI$ is the adjacency matrix of Γ , and $N^\top N - 2I$ is

the adjacency matrix of $L(\Gamma)$. Since Γ is connected, the matrix N has eigenvalue 0 with multiplicity 1 if Γ is bipartite, and does not have eigenvalue 0 otherwise. Consequently, $L(\Gamma)$ has eigenvalue -2 with multiplicity $m - n + 1$ if Γ is bipartite, and with multiplicity $m - n$ otherwise. If $\eta \neq 0$, then the multiplicity of $\eta - 2$ as eigenvalue of $L(\Gamma)$ equals the multiplicity of $\eta - k$ as eigenvalue of Γ .

We see that for a regular connected graph Γ , the spectrum of $L(\Gamma)$ determines that of Γ (since $L(\Gamma)$ is regular of valency $2k - 2$ and n is determined by $m = \frac{1}{2}nk$).

Since $L(\Gamma')$ is cospectral with $L(\Gamma)$, also Γ' has m edges. $L(\Gamma')$ is regular and hence Γ' is regular or semiregular bipartite. Suppose that Γ' is not cospectral with Γ . Then Γ' is semiregular bipartite with parameters (n_1, n_2, k_1, k_2) (say), and

$$m = \frac{1}{2}nk = n_1k_1 = n_2k_2.$$

Since the signless Laplacian matrices Q and Q' of Γ and Γ' have the same non-zero eigenvalues, their largest eigenvalues are equal:

$$2k = k_1 + k_2.$$

If $n = n_1 + n_2$ then $k_1 = k_2$, contradiction. So

$$n = n_1 + n_2 - 1.$$

Write $k_1 = k - a$ and $k_2 = k + a$, then $nk = n_1k_1 + n_2k_2$ yields

$$k = (n_1 - n_2)a.$$

Now $n_1k_1 = n_2k_2$ gives

$$(n_1 - n_2)^2 = n_1 + n_2.$$

Put $b = n_1 - n_2$, then $(n, k) = (b^2 - 1, ab)$. Since $2ab = k_1 + k_2 \leq n_2 + n_1 = b^2$, it follows that $a \leq \frac{1}{2}b$. \square

Now the following can be concluded from Theorem 13.4.9 and Lemma 13.4.10.

Theorem 13.4.11 *Suppose Γ is a connected regular DS graph, which is not a 3-connected graph with 8 vertices or a regular graph with $b^2 - 1$ vertices and degree ab for some integers a and b , with $a \leq \frac{1}{2}b$. Then also the line graph $L(\Gamma)$ of Γ is DS.*

Bussemaker, Cvetković, and Seidel [69] determined all connected regular exceptional graphs (see also [110]). There are exactly 187 such graphs, of which 32 are DS. This leads to the following characterisation.

Theorem 13.4.12 *Suppose Γ is a connected regular DS graph with all its adjacency eigenvalues at least -2 , then one of the following occurs.*

- (i) Γ is the line graph of a connected regular DS graph.
- (ii) Γ is the line graph of a connected semiregular bipartite graph, which is DS with respect to the signless Laplacian matrix.
- (iii) Γ is a cocktailparty graph.
- (iv) Γ is one of the 32 connected regular exceptional DS graphs.

Proof. Suppose Γ is not an exceptional graph or a cocktailparty graph. Then Γ is the line graph of a connected graph Δ , say. Whitney [305] has proved that Δ is uniquely determined from Γ , unless $\Gamma = K_3$. If this is the case then $\Gamma = L(K_3) = L(K_{1,3})$, so (i) holds. Suppose Δ' is cospectral with Δ with respect to the signless Laplacian Q . Then Γ and $L(\Delta')$ are cospectral with respect to the adjacency matrix, so $\Gamma = L(\Delta')$ (since Γ is DS). Hence $\Delta = \Delta'$. Because Γ is regular, Δ must be regular, or semiregular bipartite. If Δ is regular, DS with respect to Q is the same as DS. \square

All four cases from Theorem 13.4.12 do occur. For (i) and (iv) this is obvious, and (iii) occurs because the cocktailparty graphs $\overline{mK_2}$ are DS (since they are regular and \bar{A} -cospectral by Proposition 13.4.5). Examples for Case (ii) are the complete graphs $K_n = L(K_{1,n})$ with $n \neq 3$. Thus the fact that K_n is DS implies that $K_{1,n}$ is DS with respect to Q if $n \neq 3$.

13.5 Distance-regular graphs

All regular DS graphs constructed so far have the property that either the adjacency matrix A or the adjacency matrix \bar{A} of the complement has smallest eigenvalue at least -2 . In this section we present other examples.

Recall that a distance-regular graph with diameter d has $d+1$ distinct eigenvalues and that its (adjacency) spectrum can be obtained from the intersection array. Conversely, the spectrum of a distance-regular graph determines the intersection array (see e.g. [117]). However, in general the spectrum of a graph doesn't tell you whether it is distance-regular or not.

For $d \geq 3$ we have constructed graphs cospectral with, but non-isomorphic to $H(d, d)$ in §13.2.2. Many more examples are given in [183] and [120].

In the theory of distance-regular graphs an important question is: 'Which graphs are determined by their intersection array?' For many distance-regular graphs this is known to be the case. Here we investigate in the cases where the graph is known to be determined by its intersection array, whether in fact it is already determined by its spectrum.

13.5.1 Strongly regular DS graphs

The spectrum of a graph Γ determines whether Γ is strongly regular. Indeed, by Proposition 3.3.1 we can see whether Γ is regular. And a regular graph with spectrum $\theta_1 \geq \dots \geq \theta_n$ is strongly regular if and only if $|\{\theta_i \mid 2 \leq i \leq n\}| = 2$.

(That is, a regular graph is strongly regular if and only if either it is connected, and then has precisely three distinct eigenvalues: its valency and two others, or it is the disjoint union aK_ℓ of a ($a \geq 2$) complete graphs of size ℓ .)

Indeed, if Γ has valency k and all eigenvalues θ_i with $i > 1$ are in $\{r, s\}$, then $(A - rI)(A - sI) = cJ$ so that A^2 is a linear combination of A , I and J , and Γ is strongly regular.

By Propositions 13.4.4 and 13.4.5 and Theorem 13.4.11, we find the following infinite families of strongly regular DS graphs.

Proposition 13.5.1 *If $n \neq 8$ and $m \neq 4$, the graphs aK_ℓ , $L(K_n)$ and $L(K_{m,m})$ and their complements are strongly regular DS graphs.*

Note that $L(K_n)$ is the triangular graph $T(n)$, and $L(K_{m,m})$ is the lattice graph $L_2(n)$. For $n = 8$ and $m = 4$ cospectral graphs exist. There is exactly one graph cospectral with $L(K_{4,4})$, the Shrikhande graph ([283]), and there are three graphs cospectral with $L(K_8)$, the so-called Chang graphs ([80]). See also §8.2.

Besides the graphs of Proposition 13.5.1, only a few strongly regular DS graphs are known; these are surveyed in Table 13.2. (Here a *local* graph of a graph Γ is the subgraph induced by the neighbors of a vertex of Γ .)

v		spectrum	name	reference
5	2	$[(-1 \pm \sqrt{5})/2]^2$	pentagon	
13	6	$[(-1 \pm \sqrt{13})/2]^6$	Paley	[275]
17	8	$[(-1 \pm \sqrt{17})/2]^8$	Paley	[275]
16	5	$1^{10} (-3)^5$	folded 5-cube	[274]
27	10	$1^{20} (-5)^6$	GQ(2,4)	[274]
50	7	$2^{28} (-3)^{21}$	Hoffman-Singleton	[183]
56	10	$2^{35} (-4)^{20}$	Gewirtz	[152], [52]
77	16	$2^{55} (-6)^{21}$	M_{22}	[42]
81	20	$2^{60} (-7)^{20}$	Brouwer-Haemers	[51], §8.7
100	22	$2^{77} (-8)^{22}$	Higman-Sims	[152]
105	32	$2^{84} (-10)^{20}$	flags of PG(2,4)	[125]
112	30	$2^{90} (-10)^{21}$	$GQ(3, 9)$	[76]
120	42	$2^{99} (-12)^{20}$	001... in $S(5, 8, 24)$	[125]
126	50	$2^{105} (-13)^{20}$	Goethals	[97]
162	56	$2^{140} (-16)^{21}$	local McLaughlin	[76]
176	70	$2^{154} (-18)^{21}$	01... in $S(5, 8, 24)$	[125]
275	112	$2^{252} (-28)^{22}$	McLaughlin	[161]

Table 13.2: The known sporadic strongly regular DS graphs (up to complements)

Being DS seems to be a very strong property for strongly regular graphs. Most strongly regular graphs have (many) cospectral mates. For example, there are exactly 32548 non-isomorphic strongly regular graphs with spectrum $15, 3^{15}, (-3)^{20}$ (cf. [244]). Other examples can be found in the survey [45]. One might be tempted to conjecture that there are only finitely many strongly regular DS graphs besides the ones from Proposition 13.5.1.

13.5.2 Distance-regularity from the spectrum

If $d \geq 3$ only in some special cases does it follow from the spectrum of a graph that it is distance-regular. The following result surveys the cases known to us.

Theorem 13.5.2 *If Γ is a distance-regular graph with diameter d and girth g satisfying one of the following properties, then every graph cospectral with Γ is also distance-regular, with the same parameters as Γ .*

- (i) $g \geq 2d - 1$,
- (ii) $g \geq 2d - 2$ and Γ is bipartite,

- (iii) $g \geq 2d - 2$ and $c_{d-1}c_d < -(c_{d-1} + 1)(\theta_1 + \dots + \theta_d)$,
- (iv) Γ is a generalized Odd graph, that is, $a_1 = \dots = a_{d-1} = 0$, $a_d \neq 0$,
- (v) $c_1 = \dots = c_{d-1} = 1$,
- (vi) Γ is the dodecahedron, or the icosahedron,
- (vii) Γ is the coset graph of the extended ternary Golay code,
- (viii) Γ is the Ivanov-Ivanov-Faradjev graph.

For parts (i), (iv) and (vi), see [52] (and also [178]), [201], and [183], respectively. Parts (ii), (iii), (v), (vii) are proved in [117] (in fact, (ii) is a special case of (iii)) and (viii) is proved in [120]. Notice that the polygons C_n and the strongly regular graphs are special cases of (i), while bipartite distance-regular graphs with $d = 3$ (these are the incidence graphs of symmetric block designs, see also [106, Thm.6.9]) are a special case of (ii).

An important result on spectral characterisations of distance-regular graphs is the following theorem of Fiol & Garriga [143], a direct consequence of Theorem 11.10.1.

Theorem 13.5.3 *Let Γ be a distance-regular graph with diameter d and $k_d = |\Gamma_d(u)|$ vertices at distance d from any given vertex u . If Γ' is cospectral with Γ and $|\Gamma'_d(x)| = k_d$ for every vertex x of Γ' , then Γ' is distance-regular.*

Let us illustrate the use of this theorem by proving case (i) of Theorem 13.5.2. Since the girth and the degree follow from the spectrum, any graph Γ' cospectral with Γ also has girth g and degree k_1 . Fix a vertex x in Γ' . Clearly $c_{x,y} = 1$ for every vertex y at distance at most $(g-1)/2$ from x , and $a_{x,y} = 0$ (where $a_{x,y}$ is the number of neighbors of y at distance $d(x,y)$ from x) if the distance between x and y is at most $(g-2)/2$. This implies that the number k'_i of vertices at distance i from x equals $k_1(k_1 - 1)^{i-1}$ for $i = 1, \dots, d-1$. Hence $k'_i = k_i$ for these i . But then also $k'_d = k_d$ and Γ' is distance-regular by Theorem 13.5.3.

13.5.3 Distance-regular DS graphs

Brouwer, Cohen & Neumaier [48] gives many distance-regular graphs determined by their intersection array. We only need to check which ones satisfy one of the properties of Theorem 13.5.2. First we give the known infinite families:

Proposition 13.5.4 *The following distance-regular graphs are DS.*

- (i) *The polygons C_n .*
- (ii) *The complete bipartite graphs minus a perfect matching.*
- (iii) *The Odd graphs O_{d+1} .*
- (iv) *The folded $(2d+1)$ -cubes.*

As mentioned earlier, part (i) follows from property (i) of Theorem 13.5.2 (and from Proposition 13.4.4). Part (ii) follows from property (ii) of Theorem 13.5.2, and the graphs of parts (iii) and (iv) are all generalized Odd graphs, so the result follows from property (iv), due to Huang & Liu [201].

Next, there are the infinite families where the spectrum determines the combinatorial or geometric structure, where the graphs are DS if and only if the corresponding structure is determined by its parameters.

Proposition 13.5.5 (i) *A graph cospectral with the incidence graph of a symmetric block design with parameters $2-(v, k, \lambda)$ is itself the incidence graph of a symmetric block design with these same parameters.*

In case (i) the designs known to be uniquely determined by their parameters are the six projective planes $PG(2, q)$ for $q = 2, 3, 4, 5, 7, 8$, and the biplane $2-(11, 5, 2)$, and their complementary designs with parameters $2-(v, v-k, v-2k+\lambda)$.

The remaining known distance-regular DS graphs are presented in Tables 13.3, 13.4, 13.5. For all but one graph the fact that they are unique (that is, determined by their parameters) can be found in [48]. Uniqueness of the Perkel graph has been proved only recently [96]. The last columns in the tables refer to the relevant theorems by which distance-regularity follows from the spectrum. In these tables we denote by $IG(v, k, \lambda)$ the point-block incidence graph of a $2-(v, k, \lambda)$ design, and by GH , GO , and GD the point graph of a generalized hexagon, generalized octagon, and generalized dodecagon, respectively.

Recall that the point graph of a $GH(1, q)$ ($GO(1, q)$, $GD(1, q)$) is the point-line incidence graph of a projective plane (generalized quadrangle, generalized hexagon) of order q . Recall that the point graph of a $GH(q, 1)$ is the line graph of the dual $GH(1, q)$, that is, the line graph of the point-line incidence graph (also known as the *flag graph*) of a projective plane of order q .

Finally, \mathcal{G}_{23} , \mathcal{G}_{21} and \mathcal{G}_{12} denote the binary Golay code, the doubly truncated binary Golay code and the extended ternary Golay code, and HoSi is the Hoffman-Singleton graph.

n	spectrum				g	name	Thm.
12	5	$\sqrt{5}^3$	$(-1)^5$	$(-\sqrt{5})^3$	3	icosahedron	13.5.2vi
14	3	$\sqrt{2}^6$	$(-\sqrt{2})^6$	$(-3)^1$	6	Heawood; $GH(1, 2)$	13.5.2i
14	4	$\sqrt{2}^6$	$(-\sqrt{2})^6$	$(-4)^1$	4	$IG(7, 4, 2)$	13.5.2ii
15	4	2^5	$(-1)^4$	$(-2)^5$	3	$L(\text{Petersen})$	13.4.11
21	4	$(1+\sqrt{2})^6$	$(1-\sqrt{2})^6$	$(-2)^8$	3	$GH(2, 1)$	13.5.2v
22	5	$\sqrt{3}^{10}$	$(-\sqrt{3})^{10}$	$(-5)^1$	4	$IG(11, 5, 2)$	13.5.2ii
22	6	$\sqrt{3}^{10}$	$(-\sqrt{3})^{10}$	$(-6)^1$	4	$IG(11, 6, 3)$	13.5.2ii
26	4	$\sqrt{3}^{12}$	$(-\sqrt{3})^{12}$	$(-4)^1$	6	$GH(1, 3)$	13.5.2i
26	9	$\sqrt{3}^{12}$	$(-\sqrt{3})^{12}$	$(-9)^1$	4	$IG(13, 9, 6)$	13.5.2ii
36	5	2^{16}	$(-1)^{10}$	$(-3)^9$	5	Sylvester	13.5.2i
42	6	2^{21}	$(-1)^6$	$(-3)^{14}$	5	antipodal 6-cover of K_7	13.5.2i
42	5	2^{20}	$(-2)^{20}$	$(-5)^1$	6	$GH(1, 4)$	13.5.2i
42	16	2^{20}	$(-2)^{20}$	$(-16)^1$	4	$IG(21, 16, 12)$	13.5.2ii
52	6	$(2+\sqrt{3})^{12}$	$(2-\sqrt{3})^{12}$	$(-2)^{27}$	3	$GH(3, 1)$	13.5.2v
57	6	$(\frac{3+\sqrt{5}}{2})^{18}$	$(\frac{3-\sqrt{5}}{2})^{18}$	$(-3)^{20}$	5	Perkel	13.5.2i
62	6	$\sqrt{5}^{30}$	$(-\sqrt{5})^{30}$	$(-6)^1$	6	$GH(1, 5)$	13.5.2i
62	25	$\sqrt{5}^{30}$	$(-\sqrt{5})^{30}$	$(-25)^1$	4	$IG(31, 25, 20)$	13.5.2ii
63	8	$\sqrt{8}^{27}$	$(-1)^8$	$(-\sqrt{8})^{27}$	4	antipodal 7-cover of K_9	13.5.2v
105	8	5^{20}	1^{20}	$(-2)^{64}$	3	$GH(4, 1)$	13.5.2v
114	8	$\sqrt{7}^{56}$	$(-\sqrt{7})^{56}$	$(-8)^1$	6	$GH(1, 7)$	13.5.2i
114	49	$\sqrt{7}^{56}$	$(-\sqrt{7})^{56}$	$(-49)^1$	4	$IG(57, 49, 42)$	13.5.2ii

continued...

n	spectrum			g	name	Thm.	
146	9	$\sqrt{8}^{72}$	$(-\sqrt{8})^{72}$	$(-9)^1$	6	$GH(1, 8)$	13.5.2 <i>i</i>
146	64	$\sqrt{8}^{72}$	$(-\sqrt{8})^{72}$	$(-64)^1$	4	$IG(73, 64, 56)$	13.5.2 <i>ii</i>
175	21	7^{28}	2^{21}	$(-2)^{125}$	3	$L(\text{HoSi})$	13.4.11
186	10	$(4+\sqrt{5})^{30}$	$(4-\sqrt{5})^{30}$	$(-2)^{125}$	3	$GH(5, 1)$	13.5.2 <i>v</i>
456	14	$(6+\sqrt{7})^{56}$	$(6-\sqrt{7})^{56}$	$(-2)^{343}$	3	$GH(7, 1)$	13.5.2 <i>v</i>
506	15	4^{230}	$(-3)^{253}$	$(-8)^{22}$	5	M_{23} graph	13.5.2 <i>i</i>
512	21	5^{210}	$(-3)^{280}$	$(-11)^{21}$	4	Coset graph of \mathcal{G}_{21}	13.5.2 <i>iii</i>
657	16	$(7+\sqrt{8})^{72}$	$(7-\sqrt{8})^{72}$	$(-2)^{512}$	3	$GH(8, 1)$	13.5.2 <i>v</i>
729	24	6^{264}	$(-3)^{440}$	$(-12)^{24}$	3	Coset graph of \mathcal{G}_{12}	13.5.2 <i>vii</i>
819	18	5^{324}	$(-3)^{468}$	$(-9)^{26}$	3	$GH(2, 8)$	13.5.2 <i>v</i>
2048	23	7^{506}	$(-1)^{1288}$	$(-9)^{253}$	4	Coset graph of \mathcal{G}_{23}	13.5.2 <i>iii, iv</i>
2457	24	11^{324}	3^{468}	$(-3)^{1664}$	3	$GH(8, 2)$	13.5.2 <i>v</i>

Table 13.3: Sporadic distance-regular DS graphs with diameter 3

By $\text{Biaff}(q)$ we denote the point-line incidence graph of an affine plane of order q minus a parallel class of lines (sometimes called a *biaffine plane*). Any graph cospectral with a graph $\text{Biaff}(q)$ is also such a graph. For prime powers $q < 9$ there is a unique affine plane of order q . ($\text{Biaff}(2)$ is the 8-gon.)

n	nonnegative spectrum	d	g	name	Thm.
18	$3^1 \sqrt{3}^6 0^4$	4	6	Pappus; $\text{Biaff}(3)$	13.5.2ii
30	$3^1 2^9 0^{10}$	4	8	Tutte's 8-cage; $GO(1, 2)$	13.5.2i
32	$4^1 2^{12} 0^6$	4	6	$\text{Biaff}(4)$	13.5.2ii
50	$5^1 \sqrt{5}^{20} 0^8$	4	6	$\text{Biaff}(5)$	13.5.2ii
80	$4^1 \sqrt{6}^{24} 0^{30}$	4	8	$GO(1, 3)$	13.5.2i
98	$7^1 \sqrt{7}^{42} 0^{12}$	4	6	$\text{Biaff}(7)$	13.5.2ii
126	$3^1 \sqrt{6}^{21} \sqrt{2}^{27} 0^{28}$	6	12	$GD(1, 2)$	13.5.2i
128	$8^1 \sqrt{8}^{56} 0^{14}$	4	6	$\text{Biaff}(8)$	13.5.2ii
170	$5^1 \sqrt{8}^{50} 0^{68}$	4	8	$GO(1, 4)$	13.5.2i

Table 13.4: Sporadic bipartite distance-regular DS graphs with $d \geq 4$

n	spectrum	d	g	name	Thm.
20	$3^1 \sqrt{5}^3 1^5 0^4 (-2)^4 (-\sqrt{5})^3$	5	5	dodecahedron	13.5.2vi
28	$3^1 2^8 (-1+\sqrt{2})^6 (-1)^7 (-1-\sqrt{2})^6$	4	7	Coxeter	13.5.2i
45	$4^1 3^9 1^{10} (-1)^9 (-2)^{16}$	4	3	$GO(2, 1)$	13.5.2v
102	$3^1 \left(\frac{1+\sqrt{17}}{2}\right)^9 2^{18} \theta_1^{16} 0^{17}$ $\theta_2^{16} \left(\frac{1-\sqrt{17}}{2}\right)^9 \theta_3^{16}$ $(\theta_1, \theta_2, \theta_3 \text{ roots of } \theta^3 + 3\theta^2 - 3 = 0)$	7	9	Biggs-Smith	13.5.2v
160	$6^1 (2+\sqrt{6})^{24} 2^{30} (2-\sqrt{6})^{24} (-2)^{81}$	4	3	$GO(3, 1)$	13.5.2v
189	$4^1 (1+\sqrt{6})^{21} (1+\sqrt{2})^{27} 1^{28}$ $(1-\sqrt{2})^{27} (1-\sqrt{6})^{21} (-2)^{64}$	6	3	$GD(2, 1)$	13.5.2v
330	$7^1 4^{55} 1^{154} (-3)^{99} (-4)^{21}$	4	5	M_{22} graph	13.5.2v
425	$8^1 (3+\sqrt{8})^{50} 3^{68} (3-\sqrt{8})^{50} (-2)^{256}$	4	3	$GO(4, 1)$	13.5.2v
990	$7^1 5^{42} 4^{55} \left(\frac{-1+\sqrt{33}}{2}\right)^{154} 1^{154} 0^{198}$ $(-3)^{99} \left(\frac{-1-\sqrt{33}}{2}\right)^{154} (-4)^{21}$	8	5	Ivanov-Ivanov- Faradjev	13.5.2viii

Table 13.5: Sporadic non-bipartite distance-regular DS graphs with $d \geq 4$

We finally remark that also the complements of distance-regular DS graphs are DS (but not distance-regular, unless $d = 2$).

13.6 The method of Wang & Xu

Wang & Xu [301] invented a method to show that relatively many graphs are determined by their spectrum and the spectrum of their complement. A sketch.

Let Γ be a graph on n vertices with adjacency matrix A . The *walk matrix* W of Γ is the square matrix of order n with i -th column $A^{i-1}\mathbf{1}$ ($1 \leq i \leq n$). It is nonsingular if and only if A does not have an eigenvector orthogonal to $\mathbf{1}$.

(Indeed, let $u^\top A = \theta u^\top$. Then $u^\top W = (1, \theta, \dots, \theta^{n-1})u^\top \mathbf{1}$. If $u^\top \mathbf{1} = 0$ then this shows that the rows of W are dependent. If for no eigenvector u^\top we have $u^\top \mathbf{1} = 0$, then all eigenvalues have multiplicity 1, and by Vandermonde W is nonsingular.)

Let $p(t) = \sum c_i t^i = \det(tI - A)$ be the characteristic polynomial of A . Let the *companion matrix* $C = (c_{ij})$ be given by $c_{in} = -c_i$ and $c_{ij} = \delta_{i,j+1}$ for $1 \leq j \leq n-1$. Then $AW = WC$.

(Indeed, this follows from $p(A) = 0$.)

Assume that Γ and Γ' are cospectral with cospectral complements. Call their walk matrices W and W' . Then $W^\top W = W'^\top W'$.

(Indeed, $(W^\top W)_{i,j} = \mathbf{1}^\top A^{i+j-2}\mathbf{1}$, and we saw in the proof of Proposition 13.1.1 that if Γ and Γ' , with adjacency matrices A and A' , are y -cospectral for two distinct y , then $\mathbf{1}^\top A^m \mathbf{1} = \mathbf{1}^\top A'^m \mathbf{1}$ for all m .)

Suppose that W is nonsingular. Then W' is nonsingular, and $Q = W'W^{-1}$ is the unique orthogonal matrix such that $A' = QAQ^\top$ and $Q\mathbf{1} = \mathbf{1}$.

(Indeed, since $W^\top W = W'^\top W'$ also W' is nonsingular, and $Q\mathbf{1} = \mathbf{1}$ since $QW = W'$, and $QQ^\top = W'(W^\top W)^{-1}W'^\top = I$. Since Γ and Γ' are cospectral, their companion matrices are equal and $QAQ^\top = QWCW^{-1}Q^\top = W'CW'^{-1} = A'$. If Q is arbitrary with $QQ^\top = I$, $Q\mathbf{1} = \mathbf{1}$ (hence also $Q^\top \mathbf{1} = \mathbf{1}$) and $QAQ^\top = A'$, then $QA^m \mathbf{1} = QA^m Q^\top \mathbf{1} = A'^m \mathbf{1}$ for all m , and $QW = W'$.)

Forget about Γ' and study rational matrices Q with $QQ^\top = I$, $Q\mathbf{1} = \mathbf{1}$ and QAQ^\top a $(0,1)$ -matrix with zero diagonal. Let the *level* of Q be the smallest integer ℓ such that ℓQ is integral. The matrices Q of level 1 are permutation matrices leading to isomorphic graphs. So the graph Γ (without eigenvector orthogonal to $\mathbf{1}$) is determined by its spectrum and the spectrum of its complement when all such matrices Q have level 1.

If Q has level ℓ , then clearly $\ell | \det W$. A tighter restriction on ℓ is found by looking at the Smith Normal Form S of W . Let $S = UWV$ with unimodular integral U and V , where $S = \text{diag}(s_1, \dots, s_n)$ with $s_1 | s_2 | \dots | s_n$. Then $W^{-1} = VS^{-1}U$ so that $s_n W^{-1}$ is integral, and $\ell | s_n$.

Let p be prime, $p | \ell$. There is an integral row vector z , $z \not\equiv 0 \pmod{p}$ such that $zW \equiv 0 \pmod{p}$ and $zz^\top \equiv 0 \pmod{p}$.

(Indeed, let z be a row of ℓQ , nonzero mod p . Now $QW = W'$ is integral and hence $zW \equiv 0 \pmod{p}$. And $QQ^\top = I$, so $zz^\top = \ell^2 \equiv 0 \pmod{p}$.)

This observation can be used to rule out odd prime divisors of ℓ in some cases. Suppose that all numbers s_i are powers of 2, except possibly the last one s_n . Let p be an odd prime divisor of s_n , and suppose that $uu^\top \not\equiv 0 \pmod{p}$, where u is the last row of U . Then $p \nmid \ell$.

(Indeed, $zW \equiv 0 \pmod{p}$ and $W = U^{-1}SV^{-1}$ with unimodular V implies $zU^{-1}S \equiv 0 \pmod{p}$. Assume $p | \ell$, so that $p | s_n$. Let $y = zU^{-1}$. Then all

coordinates of y except for the last one are 0 (mod p). And $z = yU$ is a nonzero constant times u (mod p). This contradicts $uu^\top \not\equiv 0 \pmod{p}$.)

It remains to worry about $p = 2$. Assume that $s_n \equiv 2 \pmod{4}$, so that (with all of the above assumptions) $\ell = 2$. For z we now have $z \not\equiv 0 \pmod{2}$, $zW \equiv 0 \pmod{2}$, $zz^\top = 4$, $z\mathbf{1} = 2$, so that z has precisely four nonzero entries, three 1 and one -1 .

We proved the following:

Theorem 13.6.1 *Let Γ be a graph on n vertices without eigenvector orthogonal to $\mathbf{1}$, and let $S = \text{diag}(s_1, \dots, s_n) = UWV$ be the Smith Normal Form of its walk matrix W , where U and V are unimodular. Let u be the last row of U . If $s_n = 2 \pmod{4}$, and $\gcd(uu^\top, s_n/2) = 1$, and $zW \not\equiv 0 \pmod{2}$ for every $(0,1)$ -vector z with weight 4, then Γ is determined by its spectrum and the spectrum of its complement.* \square

Wang & Xu generate a number of random graphs where this method applies.

Let us abbreviate the condition ‘determined by its spectrum and the spectrum of its complement’ by DGS (determined by the generalized spectrum). Wang & Xu [302] used their approach to find conditions for which a DGS graph remains DGS if an isolated vertex is added.

Theorem 13.6.2 *Let Γ be a graph without eigenvector orthogonal to $\mathbf{1}$. If we have $\gcd(\det A, \det W) = 1$, then the graph obtained from Γ by adding an isolated vertex is DGS if and only if Γ is.*

There is experimental evidence that in most cases where a cospectral mate exists, the level ℓ is 2.

13.7 Exercises

Exercise 1 Show for the adjacency matrix A

- (i) that there is no pair of cospectral graphs on fewer than 5 vertices,
- (ii) that the Saltire pair is the only cospectral pair on 5 vertices,
- (iii) that there are precisely 5 cospectral pairs on 6 vertices.











				
				
$x^6 - x^4$	$x^6 - 4x^4 + 3x^2$	$x^6 - 5x^4 + 4x^2$	$x^6 - 6x^4 - 4x^3 + 5x^2 + 4x$	$x^6 - 7x^4 - 4x^3 + 7x^2 + 4x - 1$

Table 13.6: The cospectral graphs on 6 vertices (with char. pol.)

Chapter 14

Graphs with few eigenvalues

Graphs with few distinct eigenvalues tend to have some kind of regularity. A graph with only one eigenvalue (for A or L or Q) is edgeless, and a connected graph with two distinct adjacency eigenvalues (for A or L or Q) is complete. A connected regular graph Γ has three eigenvalues if and only if Γ is connected and strongly regular. Two obvious next cases are connected regular graphs with four eigenvalues, and general graphs with three eigenvalues. In the latter case the graphs need not be regular, so it matters which type of matrix we consider. For the Laplacian matrix there is an elegant characterization in terms of the structure, which gives a natural generalization of the spectral characterization of strongly regular graphs.

14.1 Regular graphs with four eigenvalues

Suppose Γ is regular with r distinct (adjacency) eigenvalues $k = \lambda_1 > \dots > \lambda_r$. Then the Laplacian matrix has eigenvalues $0 = k - \lambda_1 < \dots < k - \lambda_r$, and the signless Laplacian has eigenvalues $k + \lambda_1 > \dots > k + \lambda_r$. So for regular graphs these three matrices have the same number of distinct eigenvalues. If, in addition, both Γ and its complement $\bar{\Gamma}$ are connected, then $\bar{\Gamma}$ also has r distinct eigenvalues, being $n - k - 1 > -\lambda_r - 1 > \dots > -\lambda_2 - 1$. However, for the Seidel matrix the eigenvalues become $-2\lambda_r - 1 > \dots > -2\lambda_2 - 1$ and $n - 2k - 1$. But $n - 2k - 1$ may be equal to one of the other eigenvalues in which case S has $r - 1$ distinct eigenvalues. For example, the Petersen has three distinct adjacency eigenvalues, but only two distinct Seidel eigenvalues, being ± 3 .

Connected regular graphs with four distinct (adjacency) eigenvalues have been studied by Doob [132, 133], Van Dam [112], and Van Dam & Spence [122]. Many such graphs are known, for example the line graphs of primitive strongly regular graphs, and distance regular graphs of diameter 3. More generally, most graphs defined by a relation of a three-class association scheme have four eigenvalues. There is no nice characterization as for regular graphs with three eigenvalues, but they do possess an interesting regularity property. A graph is *walk-regular*, whenever for every $\ell \geq 2$ the number of closed walks of length ℓ at a vertex v is independent of the choice of v . Note that walk-regularity implies

regularity (take $\ell = 2$). Examples of walk-regular graphs are distance-regular graphs, and vertex-transitive graphs, but there is more.

Proposition 14.1.1 *Let Γ be a connected graph whose adjacency matrix A has $r \geq 4$ distinct eigenvalues. Then Γ is walk-regular if and only if A^ℓ has constant diagonal for $2 \leq \ell \leq r - 2$.*

Proof. We know that the number of closed walks of length ℓ at vertex v equals $(A^\ell)_{v,v}$. Therefore, Γ is walk-regular if and only if A^ℓ has constant diagonal for all $\ell \geq 2$. Suppose A^ℓ has constant diagonal for $2 \leq \ell \leq r - 2$. Then A^2 has constant diagonal, so Γ is regular. The Hoffman polynomial of Γ has degree $r - 1$, and hence $A^{r-1} \in \langle A^{r-2}, \dots, A^2, A, I, J \rangle$. This implies $A^\ell \in \langle A^{r-2}, \dots, A^2, A, I, J \rangle$ for all $\ell \geq 0$. Therefore A^ℓ has constant diagonal for all $\ell \geq 0$. \square

Corollary 14.1.2 *If Γ is connected and regular with four distinct eigenvalues, then Γ is walk-regular.* \square

For a graph Γ with adjacency matrix A , the average number of triangles through a vertex equals $\frac{1}{2n} \text{tr} A^3$. Suppose Γ is walk-regular. Then this number must be an integer. Similarly, $\frac{1}{2n} \text{tr} A^\ell$ is an integer if ℓ is odd, and $\frac{1}{n} \text{tr} A^\ell$ is an integer if n is even. Van Dam and Spence [122] have used these (and other) conditions in their computer generation of feasible spectra for connected regular graphs with four eigenvalues. For constructions, characterizations, and other results on regular graphs with four eigenvalues we refer to Van Dam [112, 113]. Here we finish with the bipartite case, which can be characterized in terms of block designs (see §4.7).

Proposition 14.1.3 *A connected bipartite regular graph Γ with four eigenvalues is the incidence graph of a symmetric 2-design (and therefore distance-regular).*

Proof. Since Γ is connected, bipartite and regular the spectrum is

$$\{k, \lambda_2^{v-1}, (-\lambda_2)^{v-1}, -k\},$$

where $2v$ is the number of vertices. For the adjacency matrix A of Γ , we have

$$A = \begin{bmatrix} O & N \\ N^\top & O \end{bmatrix}, \quad \text{and} \quad A^2 = \begin{bmatrix} NN^\top & O \\ O & N^\top N \end{bmatrix},$$

for some square $(0, 1)$ -matrix N satisfying $N\mathbf{1} = N^\top\mathbf{1} = k\mathbf{1}$. It follows that NN^\top has spectrum $\{(k^2)^1, (\lambda_2^2)^{v-1}\}$, where k^2 corresponds to the row and column sum of NN^\top . This implies that $NN^\top \in \langle J, I \rangle$, and hence N is the incidence matrix of a symmetric design. \square

14.2 Three Laplacian eigenvalues

If a connected graph Γ has three distinct Laplacian eigenvalues $0 < \nu < \nu'$ (say), the complement $\bar{\Gamma}$ has eigenvalues $0 \leq n - \nu' < n - \nu$, so if $\bar{\Gamma}$ is connected, it also has three distinct eigenvalues. To avoid the disconnected exceptions, it is convenient to use the notion of restricted eigenvalues (recall that an eigenvalue

is *restricted* if it has an eigenvector orthogonal to the all-one vector $\mathbf{1}$), and consider graphs with two distinct restricted Laplacian eigenvalues.

We say that a graph Γ has constant $\mu(\Gamma)$ if Γ is not complete and any two distinct nonadjacent vertices of Γ have the same number of common neighbors (equal to $\mu(\Gamma)$).

Theorem 14.2.1 *A graph Γ has two distinct restricted Laplacian eigenvalues ν and ν' if and only if Γ has constant $\mu(\Gamma)$ and its complement $\bar{\Gamma}$ has constant $\mu(\bar{\Gamma})$. If Γ is such a graph, only two vertex degrees d and d' occur, and*

$$\nu + \nu' = d + d' + 1 = \mu(\Gamma) + n - \mu(\bar{\Gamma}), \quad \nu\nu' = dd' + \mu(\Gamma) = \mu(\Gamma)n.$$

Proof. Suppose Γ has just two restricted Laplacian eigenvalues ν and ν' . Then $(L - \nu I)(L - \nu' I)$ has rank 1 and row sum $\nu\nu'$, so

$$(L - \nu I)(L - \nu' I) = \frac{\nu\nu'}{n}J.$$

If u and v are nonadjacent vertices, then $(L)_{uv} = 0$, so $(L^2)_{uv} = \nu\nu'/n$, and $\mu(\Gamma) = \nu\nu'/n$ is constant. Similarly, $\bar{\Gamma}$ has constant $\mu(\bar{\Gamma}) = (n - \nu)(n - \nu')/n$.

Next suppose $\mu = \mu(\Gamma)$ and $\bar{\mu} = \mu(\bar{\Gamma})$ are constant. If u and v are adjacent vertices, then $((nI - J - L)_{uv} = \bar{\mu}$, so $\bar{\mu} = (L^2)_{uv} + n$, and if u and v are nonadjacent, then $(L^2)_{uv} = \mu$. Furthermore $(L^2)_{uu} = d_u^2 + d_u$, where d_u is the degree of u . Writing $D = \text{diag}(d_1, \dots, d_n)$, we obtain

$$\begin{aligned} L^2 &= (\bar{\mu} - n)(D - L) + \mu(J - I - D + L) + D^2 + D = \\ &= (\mu + n - \bar{\mu})L + D^2 - (\mu + n - \bar{\mu} - 1)D - \mu I + \mu J. \end{aligned}$$

Since L and L^2 have zero row sums, it follows that $d_u^2 - d_u(\mu + n - \bar{\mu} - 1) - \mu + \mu n = 0$ for every vertex u . So $L^2 - (\mu + n - \bar{\mu})L + \mu n I = \mu J$. Now let ν and ν' be such that $\nu + \nu' = \mu + n - \bar{\mu}$ and $\nu\nu' = \mu n$, then $(L - \nu I)(L - \nu' I) = \frac{\nu\nu'}{n}J$, so L has distinct restricted eigenvalues ν and ν' . As a side result we obtained that all vertex degrees d_u satisfy the same quadratic equation, so that d_u can only take two values d and d' , and the formulas readily follow. \square

Regular graphs with constant $\mu(\Gamma)$ and $\mu(\bar{\Gamma})$ are strongly regular. So Theorem 14.2.1 generalizes the spectral characterization of strongly regular graphs. Several nonregular graphs with two restricted Laplacian eigenvalues are known. A geodetic graph of diameter three with connected complement provides an example with $\mu(\Gamma) = 1$ (see [48], Theorem 1.17.1). Here we give two other constructions. Both constructions use symmetric block designs (see §4.7). Correctness easily follows by use of Theorem 14.2.1.

Proposition 14.2.2 *Let N be the incidence matrix of a symmetric 2 -(n, k, λ) design. Suppose that N is symmetric (which means that the design has a polarity). Then $L = kI - N$ is the Laplacian matrix of a graph with two restricted eigenvalues, being $k \pm \sqrt{k - \lambda}$. The possible degrees are k and $k - 1$. \square*

If all diagonal elements of N are 0, then the graph Γ is a (n, k, λ) -graph (a strongly regular graph with $\lambda = \mu$), and if all diagonal elements of N are 1, then $\bar{\Gamma}$ is such a graph. Otherwise both degrees k and $k - 1$ do occur. For example the Fano plane admits a symmetric matrix with three ones on the diagonal. The corresponding graph has restricted Laplacian eigenvalues $3 \pm \sqrt{2}$, and vertex degrees 2 and 3. See also §4.8.

Proposition 14.2.3 *Let N be the incidence matrix of a symmetric block design. Write*

$$N = \begin{bmatrix} \mathbf{1} & N_1 \\ \mathbf{0} & N_2 \end{bmatrix}, \text{ and define } L = \begin{bmatrix} vI - J & O & N_1 - J \\ O & vI - J & -N_2 \\ N_1^\top - J & -N_2^\top & 2(k - \lambda)I \end{bmatrix}.$$

Then L has two restricted eigenvalues. □

Other examples, characterizations and a table of feasible spectra can be found in [116] and [113] (see also Exercise 1). See [303] for some more recent results on graphs with three Laplacian eigenvalues.

14.3 Other matrices with at most three eigenvalues

No characterization is known of nonregular graphs with three M -eigenvalues, for a matrix M other than the Laplacian. However several examples and properties are known. Some of these will be discussed below.

14.3.1 Few Seidel eigenvalues

Seidel switching (see §1.8.2) doesn't change the Seidel spectrum, so having few Seidel eigenvalues is actually a property of the switching class of a graph. For example the switching class of \bar{K}_n , the edgeless graph on n vertices, consists of the complete bipartite graphs $K_{m,n-m}$, and all of them have Seidel spectrum $\{(-1)^{n-1}, n-1\}$. Only the one-vertex graph K_1 has one Seidel eigenvalue. Graphs with two Seidel eigenvalues are strong (see §9.1). To be precise, they are the graphs whose associated two-graph is regular (Theorem 9.3.1). The Seidel matrix is a special case of a generalized adjacency matrix. These are matrices of the form $M(x, y, z) = xI + yA + z(J - I - A)$ with $y \neq z$, where A is the adjacency matrix; see also Chapter 13. If A is the adjacency matrix of a strongly regular graph with eigenvalues $k \geq r > s$, then both $nA - (k-r)J$ and $nA - (k-s)J$ (these are basically the nontrivial idempotents of the association scheme) are generalized adjacency matrices with two eigenvalues. We recall that a strong graph either has two Seidel eigenvalues, or is strongly regular. Thus for every strong graph there exist numbers x , y and z , such that $M(x, y, z)$ has two eigenvalues.

Proposition 14.3.1 *A graph is strong if and only if at least one generalized adjacency matrix has two eigenvalues.*

Proof. Correctness of the 'only if' part of the statement has been established already. Without loss of generality we assume that the eigenvalues of $M = M(x, y, z)$ are 0 and 1. So M satisfies $M^2 = M$. Let d_i be the degree of vertex i , then $x = M_{ii} = (M^2)_{ii} = x^2 + d_i y^2 + (n-1-d_i)z^2$, which gives $d_i(y^2 - z^2) = x - x^2 - (n-1)z^2$. So $y = -z$ or Γ is regular. In the first case $S = \frac{1}{z}(M - xI)$ is the Seidel matrix of Γ with two eigenvalues, so Γ is strong. In case Γ is regular, the adjacency matrix $A = \frac{1}{y-z}(M + (z-x)I - zJ)$ has three eigenvalues, so Γ is strongly regular and therefore strong. □

So if a generalized adjacency matrix $M(x, y, z)$ of a nonregular graph has two eigenvalues, then $y = -z$ (and we basically deal with the Seidel matrix).

A strongly regular graph Γ on n vertices with adjacency eigenvalues k, r, s ($k \geq r > s$) has Seidel eigenvalues $\rho_0 = n - 1 - 2k$, $\rho_1 = -2s - 1$, and $\rho_2 = -2r - 1$. If $\rho_0 = \rho_1$, or $\rho_0 = \rho_2$, then Γ has two eigenvalues, otherwise Γ , and all graphs switching equivalent to Γ , have three eigenvalues. For example, the (switching class of the) Petersen graph has two Seidel eigenvalues 3 and -3 , whilst the pentagon C_5 has three Seidel eigenvalues 0 and $\pm\sqrt{5}$. However, not every graph with three Seidel eigenvalues is switching equivalent to a strongly regular graph. Not even if the graph is regular. Indeed, consider a graph Γ whose Seidel matrix S has two eigenvalues ρ_1 and ρ_2 . Then $(S+I) \otimes (S+I) - I$ represents a graph Γ^2 with eigenvalues $(\rho_1 + 1)^2 - 1$, $(\rho_1 + 1)(\rho_2 + 1) - 1$, and $(\rho_2 + 1)^2 - 1$. Moreover, Γ^2 is regular if Γ is.

14.3.2 Three adjacency eigenvalues

Connected regular graphs with three adjacency eigenvalues are strongly regular. The complete bipartite graphs $K_{\ell, m}$ have spectrum $\{-\sqrt{\ell m}, 0^{n-2}, \sqrt{\ell m}\}$. If $\ell \neq m$ they are nonregular with three adjacency eigenvalues. Other nonregular graphs with three adjacency eigenvalues have been constructed by Bridges and Mena [39], Klin and Muzychuk [217], and Van Dam [113, 114]. Chuang and Omid [84] characterized all such graphs with largest eigenvalue at most eight. Many nonregular graphs with three eigenvalues can be made from a strongly regular graphs by introducing one new vertex adjacent to all other vertices. Such a graph is called a *cone* over a strongly regular graph.

Proposition 14.3.2 *Let Γ be a strongly regular graph on n vertices with eigenvalues $k > r > s$. Then the cone $\hat{\Gamma}$ over Γ has three eigenvalues if and only if $n = s(s - k)$.*

Proof. If \hat{A} is the adjacency matrix of $\hat{\Gamma}$, then \hat{A} admits an equitable partition with quotient matrix

$$\begin{bmatrix} 0 & n \\ 1 & k \end{bmatrix}$$

with eigenvalues $(k \pm \sqrt{k^2 + 4n})/2$, which are also eigenvalues of \hat{A} . The other eigenvalues of \hat{A} have eigenvectors orthogonal to the characteristic vectors of the partition, so they remain eigenvalues if the all-one blocks of the equitable partition are replaced by all-zero blocks. Therefore they are precisely the restricted eigenvalues r and s of Γ . So the eigenvalues of \hat{A} are $(k \pm \sqrt{k^2 + 4n})/2$, r and s . Two of these values coincide if and only if $s = (k - \sqrt{k^2 + 4n})/2$. \square

There exist infinitely many strongly regular graphs for which $n = s(s - k)$, the smallest of which is the Petersen graph. The cone over the Petersen graph has eigenvalues 5, 1 and -2 . If a cone over a strongly regular graph has three eigenvalues, then these eigenvalues are integers (see Exercise 3). The complete bipartite graphs provide many examples with nonintegral eigenvalues. In fact:

Proposition 14.3.3 *If Γ is a connected graph with three distinct adjacency eigenvalues of which the largest is not an integer, then Γ is a complete bipartite graph.*

Proof. Assume Γ has $n \geq 4$ vertices. Since the largest eigenvalue ρ is non-integral with multiplicity 1, one of the other two eigenvalues $\bar{\rho}$ (say) also has this property, and the third eigenvalue has multiplicity $n - 2 \geq 2$, so cannot be irrational. Thus the spectrum of Γ is

$$\{\rho = \frac{1}{2}(a + \sqrt{b}), \bar{\rho} = \frac{1}{2}(a - \sqrt{b}), c^{n-2}\},$$

for integer a, b and c . Now $\text{tr} A = 0$ gives $c = -a/(n-2)$. By Perron-Frobenius' theorem, $\rho \geq |\bar{\rho}|$, therefore $a \geq 0$ and $c \leq 0$. If $c = 0$, the eigenvalues of Γ are $\pm\sqrt{b}/2$ and 0, and Γ is bipartite of diameter at most 2, and hence Γ is complete bipartite. If $c \leq -2$, then $\text{tr} A^2 \geq 4(n-2)^2$ so Γ has at least $2(n-2)^2$ edges which is ridiculous. If $c = -1$, then $\rho = \frac{1}{2}(n-2 + \sqrt{b}) \leq n-1$, hence $\sqrt{b} \leq n$ and $\bar{\rho} > -1$. This implies that $A + I$ is positive semi-definite (of rank 2). So $A + I$ is the Gram matrix of a set of unit vectors (in \mathbb{R}^2) with angles 0 and $\pi/2$. This implies that being adjacent is an equivalence relation, so $\Gamma = K_n$, a contradiction. \square

The conference graphs are examples of regular graphs where only the largest eigenvalue is an integer. Van Dam and Spence [72] found a number of nonregular graphs on 43 vertices with eigenvalues 21, $-\frac{1}{2} \pm \frac{1}{2}\sqrt{41}$. It turns out that all these graphs have three distinct vertex degrees: 19, 26 and 35 (which was impossible in case of the Laplacian spectrum).

14.3.3 Three signless Laplacian eigenvalues

Recently, Ayoobi, Omidi and Tayfeh-Rezaie [11] started to investigate nonregular graphs whose signless Laplacian matrix Q has three distinct eigenvalues. They found three infinite families.

(i) The complete K_n with one edge deleted has Q -spectrum

$$\left\{ \frac{1}{2}(3n-6 + \sqrt{n^2+4n-12}), (n-2)^{n-2}, \frac{1}{2}(3n-6 - \sqrt{n^2+4n-12}) \right\}.$$

(ii) The star $K_{1,n-1}$ has Q -spectrum $0^1, 1^{n-2}, n^1$.

(iii) The complement of $K_{m,m} + mK_1$ has Q -spectrum

$$(5m-2)^1, (3m-2)^m, (2m-2)^{2m-2}.$$

In addition there are some sporadic examples (see also Exercise 4). Like in Proposition 14.3.3 the case in which the spectral radius is nonintegral can be characterized.

Proposition 14.3.4 [11] *Let Γ be a connected graph on at least four vertices of which the signless Laplacian has three distinct eigenvalues. Then the largest of these eigenvalues is nonintegral if and only if Γ is the complete graph minus one edge.*

It is not known if there exist other nonregular examples with a nonintegral eigenvalue. We expect that the above list is far from complete.

14.4 Exercises

Exercise 1 Prove that a graph with two restricted Laplacian eigenvalues whose degrees d and d' differ by 1, comes from a symmetric design with a polarity as described in Proposition 14.2.2.

Exercise 2 Let Γ be a strongly regular graph with a coclique C whose size meets Hoffman's bound (4.1.2). Prove that the subgraph of Γ induced by the vertices outside C is regular with at most four distinct eigenvalues. Can it have fewer than four eigenvalues?

Exercise 3 Suppose $\hat{\Gamma}$ is a cone over a strongly regular graph. Show that, if $\hat{\Gamma}$ has three distinct eigenvalues, then all three are integral.

Exercise 4 Show that the cone over the Petersen graph has three signless Laplacian eigenvalues. Find a necessary and sufficient condition on the parameters (n, k, λ, μ) of a strongly regular graph Γ under which the cone over Γ has three signless Laplacian eigenvalues.

Bibliography

- [1] M. Ajtai, J. Komlós & E. Szemerédi, *Sorting in $c \log n$ parallel steps*, Combinatorica **3** (1983) 1–19. (p. 68)
- [2] N. Alon, *Eigenvalues and Expanders*, Combinatorica **6** (1986) 83–96. (pp. 66, 67)
- [3] N. Alon, *The Shannon Capacity of a union*, Combinatorica **18** (1998) 301–310. (p. 65)
- [4] N. Alon, *Large sets in finite fields are sumsets*, J. Number Th. **126** (2007) 110–118. (p. 90)
- [5] N. Alon & F. R. K. Chung, *Explicit Constructions of linear sized tolerant networks*, Discr. Math. **2** (1988) 15–19. (p. 68)
- [6] N. Alon & V. D. Milman, λ_1 , *isoperimetric inequalities for graphs and superconcentrators*, J. Combin. Th. (B) **38** (1985) 73–88. (p. 71)
- [7] N. Alon & P. Seymour, *A Counterexample to the rank-coloring conjecture*, J. Graph Theory **13** (1989) 523–525. (p. 64)
- [8] W. N. Anderson & T. D. Morley, *Eigenvalues of the Laplacian of a graph*, Lin. Multilin. Alg. **18** (1985) 141–145. (p. 48)
- [9] D. M. Appleby, *Symmetric informationally complete-positive operator valued measures and the extended Clifford group*, J. Math. Phys. **46** 052107 (2005). (p. 143)
- [10] M. Aschbacher, *The non-existence of rank three permutation groups of degree 3250 and subdegree 57*, J. Algebra **19** (1971) 538–540. (p. 153)
- [11] F. Ayoobi, G. R. Omid & B. Tayfeh-Rezaie, *A note on graphs whose signless Laplacian has three distinct eigenvalues*, preprint. (p. 200)
- [12] Bhaskar Bagchi, *On strongly regular graphs with $\mu \leq 2$* , Discr. Math. **306** (2006) 1502–1504. (p. 106)
- [13] Hua Bai, *The Grone-Merris Conjecture*, arXiv:0911.2172v3, 1 Jan 2010. (p. 50)
- [14] G. A. Baker, *Drum shapes and isospectral graphs*, J. Math. Phys. **7** (1966) 2238–2242. (p. 26)

- [15] S. Bang, A. Dubickas, J. H. Koolen & V. Moulton, *There are only finitely many distance-regular graphs of fixed valency greater than two*, [arXiv.org:0909.5253](https://arxiv.org/abs/0909.5253), 29 Sep 2009. (p. 160)
- [16] E. Bannai & T. Ito, *On finite Moore graphs*, J. Fac. Sci. Univ. Tokyo, Sect IA **20** (1973) 191–208. (p. 161)
- [17] E. Bannai & T. Ito, *Algebraic Combinatorics I: Association Schemes*, Benjamin/Cummings, London, 1984. (p. 160)
- [18] D. Bauer, H. J. Broersma & H. J. Veldman, *Not every 2-tough graph is Hamiltonian*, Discr. Appl. Math. **99** (2000) 317–321. (p. 69)
- [19] D. Bauer, H. J. Broersma, J. van den Heuvel & H. J. Veldman, *Long cycles in graphs with prescribed toughness and minimum degree*, Discr. Math. **141** (1995) 1–10. (p. 69)
- [20] V. Belevitch, *Theory of $2n$ -terminal networks with application to conference telephony*, Electron. Comm. **27** (1950) 231–244. (p. 140)
- [21] E. R. Berlekamp, J. H. van Lint & J. J. Seidel, *A strongly regular graph derived from the perfect ternary Golay code*, pp. 25–30 in: A survey of combinatorial theory, Symp. Colorado State Univ., 1971, (J.N. Srivastava et al., eds.), North Holland, 1973. (p. 131)
- [22] A. Berman & R. J. Plemmons, *Nonnegative matrices in the mathematical sciences*, Acad. Press, New York, 1979. (p. 30)
- [23] Jochem Berndsén & Mayank, *The Brouwer conjecture for split graphs*, preprint, 2010. (p. 50)
- [24] Monica Bianchini, Marco Gori & Franco Scarselli, *Inside PageRank*, ACM Trans. on Internet Technology **5** (2005) 92–128. (p. 56)
- [25] A. Bigalke & H. A. Jung, *Über Hamiltonsche Kreise und unabhängige Ecken in Graphen*, Monatsh. Math. **88** (1979) 195–210. (p. 112)
- [26] N. L. Biggs, *Algebraic Graph Theory*, Cambridge Tracts in Mathematics, Vol 67, Cambridge Univ. Press, Cambridge, 1974. (Second ed., Cambridge, 1993.) (Not cited)
- [27] A. Blokhuis, *Few-distance sets*, Ph.D. thesis, Eindhoven, 1983. (p. 145)
- [28] A. Blokhuis & A. E. Brouwer, *Uniqueness of a Zara graph on 126 points and non-existence of a completely regular two-graph on 288 points*, pp 6–19 in: *Papers dedicated to J. J. Seidel*, P. J. de Doelder, J. de Graaf & J. H. van Lint (eds.), Eindhoven Univ. Techn. Report 84-WSK-03, Aug 1984. Zbl 554.05049 (p. 139)
- [29] J. A. Bondy, *A graph reconstruction manual*, pp. 221–252 in: *Surveys in Combinatorics*, LMS Lecture Note Series **166**, Cambridge Univ. Press, Cambridge, 1991. (p. 24)
- [30] J. Bosák, A. Rosa & Š. Znám, *On decompositions of complete graphs into factors with given diameters*, pp. 37–56 in: *Theory of Graphs (Proc. Colloq., Tihany, 1966)*, Academic Press, New York, 1968. (p. 20)

- [31] R. C. Bose, *Strongly regular graphs, partial geometries and partially balanced designs*, Pacific J. Math **13** (1963) 389–419. (p. 109)
- [32] R. C. Bose & T. A. Dowling, *A generalization of Moore graphs of diameter two*, J. Combin. Th. **11** (1971) 213–226. (p. 131)
- [33] R. C. Bose & D. M. Mesner, *On linear associative algebras corresponding to association schemes of partially balanced designs*, Ann. Math. Statist. **30** (1959) 21–38. (p. 148)
- [34] R. C. Bose & T. Shimamoto, *Classification and analysis of partially balanced incomplete block designs with two associate classes*, J. Am. Statist. Assoc. **47** (1952) 151–184. (p. 147)
- [35] N. Bourbaki, *Groupes et algèbres de Lie, Chap. 4, 5 et 6*, Hermann, Paris, 1968. (p. 41)
- [36] V. Brankov, P. Hansen & D. Stevanović, *Automated conjectures on upper bounds for the largest Laplacian eigenvalue of graphs*, Lin. Alg. Appl. **414** (2006) 407–424. (p. 48)
- [37] A. Brauer & I. C. Gentry, *On the characteristic roots of tournament matrices*, Bull. Amer. Math. Soc. **74** (1968) 1133–1135. (p. 25)
- [38] W. G. Bridges & R. A. Mena, *Rational circulants with rational spectra and cyclic strongly regular graphs*, Ars. Combin. **8** (1979) 143–161. (p. 104)
- [39] W. G. Bridges & R. A. Mena, *Multiplicative cones - a family of three eigenvalue graphs*, Aequationes Math. **22** (1981) 208–214. (p. 199)
- [40] Sergey Brin & Lawrence Page, *The anatomy of a large-scale hypertextual web search engine*, Proceedings of the Seventh International World Wide Web Conference, Computer Networks and ISDN Systems **30** (1998) 107–117. (p. 55)
- [41] R. Brooks, *Non-Sunada graphs*, Ann. Inst. Fourier (Grenoble) **49** (1999) 707–725. (pp. 176, 179)
- [42] A. E. Brouwer, *The uniqueness of the strongly regular graph on 77 points*, J. Graph Theory **7** (1983) 455–461. (pp. 128, 188)
- [43] A. E. Brouwer, *Toughness and spectrum of a graph*, Lin. Alg. Appl. **226–228** (1995) 267–271. (p. 69)
- [44] A. E. Brouwer, *A non-degenerate generalized quadrangle with lines of size four is finite*, in: Advances in finite geometries and designs, Proc. 3rd Isle of Thorns Conf., Chelwood Gate/UK 1990, pp. 47–49. (p. 115)
- [45] A. E. Brouwer, *Strongly regular graphs*, in: The CRC handbook of combinatorial designs, (C.J. Colbourn and J.H. Denitz eds.), CRC Press, 1996, pp. 667–685. (p. 188)
- [46] A. E. Brouwer, *Small integral trees*, Electr. J. Combin. **15** (2008) N1. (p. 85)

- [47] A. E. Brouwer & A. M. Cohen, *Some remarks on Tits geometries*, with an appendix by J. Tits, *Indag. Math.* **45** (1983) 393–402. (p. 113)
- [48] A. E. Brouwer, A. M. Cohen & A. Neumaier, *Distance-Regular Graphs*, Springer-Verlag, 1989. (pp. 41, 104, 105, 106, 123, 157, 161, 176, 189, 190, 197)
- [49] A. E. Brouwer & C. A. van Eijl, *On the p -rank of the adjacency matrices of strongly regular graphs*, *J. Alg. Combin.* **1** (1992) 329–346. (pp. 169, 170)
- [50] A. E. Brouwer & M. van Eupen, *The correspondence between projective codes and 2-weight codes*, *Designs, Codes and Cryptography* **11** (1997) 261–266. (p. 123)
- [51] A. E. Brouwer & W. H. Haemers, *Structure and uniqueness of the $(81, 20, 1, 6)$ strongly regular graph*, *Discr. Math.* **106/107** (1992) 77–82. (pp. 117, 128, 188)
- [52] A. E. Brouwer & W. H. Haemers, *The Gewirtz graph: An exercise in the theory of graph spectra*, *Europ. J. Combin.* **14** (1993) 397–407. (pp. 127, 188, 189)
- [53] A. E. Brouwer & W. H. Haemers, *Eigenvalues and perfect matchings*, *Lin. Alg. Appl.* **395** (2005) 155–162. (pp. 73, 74)
- [54] A. E. Brouwer & W. H. Haemers, *A lower bound for the Laplacian eigenvalues of a graph—proof of a conjecture by Guo*, *Lin. Alg. Appl.* **429** (2008) 2131–2135. (p. 49)
- [55] A. E. Brouwer & J. H. Koolen, *The vertex connectivity of a distance-regular graph*, *Europ. J. Combin.* **30** (2009) 668–673. (p. 160)
- [56] A. E. Brouwer & J. H. van Lint, *Strongly regular graphs and partial geometries*, in: *Enumeration and designs*, Proc. Silver Jubilee Conf. on Combinatorics, Waterloo 1982, eds. D. M. Jackson & S. A. Vanstone, Academic Press, Toronto. pp. 85–122. (pp. 106, 112, 130, 142, 170)
- [57] A. E. Brouwer & D. M. Mesner, *The connectivity of strongly regular graphs*, *Europ. J. Combin.* **6** (1985) 215–216. (p. 111)
- [58] A. E. Brouwer & A. Neumaier, *A remark on partial linear spaces of girth 5 with an application to strongly regular graphs*, *Combinatorica* **8** (1988) 57–61. (p. 106)
- [59] A. E. Brouwer & A. Neumaier, *The graphs with spectral radius between 2 and $\sqrt{2 + \sqrt{5}}$* , *Lin. Alg. Appl.* **114/115** (1989) 273–276. (p. 41)
- [60] A. E. Brouwer & E. Spence, *Cospectral graphs on 12 vertices*, *Electr. J. Combin.* **16** (2009) N20. (p. 181)
- [61] A. E. Brouwer & H. A. Wilbrink, *Block designs*, in: *Handbook of Incidence Geometry* (F. Buekenhout ed.), Elsevier, 1995, pp. 349–382. (p. 76)

- [62] A. E. Brouwer, R. M. Wilson & Qing Xiang, *Cyclotomy and Strongly Regular Graphs*, J. Alg. Combin. **10** (1999) 25–28. (p. 126)
- [63] R. A. Brualdi & A. J. Hoffman, *On the spectral radius of $(0,1)$ matrices*, Lin. Alg. Appl. **65** (1985) 133–146. (p. 42)
- [64] N. G. de Bruijn, *A combinatorial problem*, Nederl. Akad. Wetensch. **49** (1946) 758–764 = Indag. Math. **8** (1946) 461–467. (p. 25)
- [65] N. G. de Bruijn, *Acknowledgement of priority to C. Flye sainte-Marie on the counting of circular arrangements of 2^n zeros and ones that show each n -letter word exactly once*, Report 75-WSK-06, Techn. Hogeschool Eindhoven. (p. 25)
- [66] Kurt Bryan & Tanya Leise, *The \$25,000,000,000 eigenvector—The linear algebra behind Google*, SIAM Review **48** (2006) 569–581. (p. 56)
- [67] M. Burrow, *Representation theory of finite groups*, Academic Press, New York, 1965. (p. 148)
- [68] F. C. Bussemaker & D. M. Cvetković, *There are exactly 13 connected, cubic, integral graphs*, Univ. Beograd Publ. Elektrotehn. Fak., Ser. Mat. Fiz., No 544–576, (1976) 43–48. (p. 44)
- [69] F. C. Bussemaker, D.M. Cvetković & J. J. Seidel, *Graphs related to exceptional root systems*, T.H.-Report 76-WSK-05, Eindhoven University of Technology, 1976. (pp. 185, 186)
- [70] F. C. Bussemaker, W. H. Haemers, R. Mathon & H. A. Wilbrink, *A $(49,16,3,6)$ strongly regular graph does not exist*, Europ. J. Combin. **10** (1989) 413–418. (p. 127)
- [71] F. C. Bussemaker, R. A. Mathon & J. J. Seidel, *Tables of two-graphs*, pp. 70–112 in: Combinatorics and Graph Theory (S. B. Rao, ed.), Springer LNM 885, 1981. Also Report 79-WSK-05, Techn. Univ. Eindhoven, 1979. (p. 135)
- [72] D. de Caen, E. R. van Dam & E. Spence, *A nonregular analogue of conference graphs*, J. Combin. Th. (A) **88** (1999) 194–204. (p. 200)
- [73] A. R. Calderbank & W. M. Kantor, *The geometry of two-weight codes*, Bull. London Math. Soc. **18** (1986) 97–122. (p. 120)
- [74] P. J. Cameron, *Orbits of permutation groups on unordered sets. II*, J. Lond. Math. Soc., II. Ser. **23** (1981) 249–264. (p. 115)
- [75] P. J. Cameron, *Automorphism groups of graphs*, pp. 89–127 in: Selected Topics in Graph Theory 2, L. W. Beineke & R. J. Wilson, eds., Acad. Press, 1983. (p. 153)
- [76] P. J. Cameron, J.-M. Goethals & J. J. Seidel, *Strongly regular graphs having strongly regular subconstituents*, J. Algebra **55** (1978) 257–280. (pp. 117, 188)

- [77] P. J. Cameron, J.-M. Goethals, J. J. Seidel & E. E. Shult, *Line graphs, root systems, and elliptic geometry*, J. Algebra **43** (1976) 305–327. (pp. 93, 185)
- [78] A. Cayley, *A theorem on trees*, Quart. J. Pure Appl. Math. **23** (1889) 376–378. (p. 15)
- [79] L. C. Chang, *The uniqueness and nonuniqueness of the triangular association scheme*, Sci. Record **3** (1959) 604–613. (p. 110)
- [80] L. C. Chang, *Association schemes of partially balanced block designs with parameters $v = 28$, $n_1 = 12$, $n_2 = 15$ and $p_{11}^2 = 4$* , Sci. Record **4** (1960) 12–18. (pp. 110, 127, 188)
- [81] Gregory Cherlin, *Locally finite generalized quadrangles with at most five points per line*, Discr. Math. **291** (2005) 73–79. (p. 115)
- [82] A. A. Chesnokov & W. H. Haemers, *Regularity and the generalized adjacency spectra of graphs*, Lin. Alg. Appl. **416** (2006) 1033–1037. (p. 182)
- [83] B. Cheyne, V. Gupta & C. Wheeler, *Hamilton cycles in addition graphs*, Rose-Hulman Undergraduate Math. Journal **1** (2003) (electronic). (p. 90)
- [84] H. Chuang, & G. R. Omid, *Graphs with three distinct eigenvalues and largest eigenvalue less than 8*, Lin. Alg. Appl. **430** (2009) 2053–2062. (p. 199)
- [85] F. R. K. Chung, *Diameters and eigenvalues*, J. Amer. Math. Soc. **2** (1989) 187–196. (pp. 70, 90)
- [86] Fan R. K. Chung, *Spectral Graph Theory*, CBMS Lecture Notes 92, Amer. Math. Soc., Providence, 1997. (Not cited)
- [87] V. Chvátal, *Tough graphs and hamiltonian circuits*, Discr. Math. **5** (1973) 215–228. (p. 69)
- [88] V. Chvátal & P. Erdős, *A note on Hamiltonian circuits*, Discr. Math. **2** (1972) 111–113. (p. 112)
- [89] S. M. Cioabă, *Perfect matchings, eigenvalues and expansion*, C. R. Math. Acad. Sci. Soc. R. Can. **27** (4) (2005) 101–104. (p. 74)
- [90] S. M. Cioabă, D. A. Gregory & W. H. Haemers, *Matchings in regular graphs from eigenvalues*, J. Combin. Th. (B) **99** (2009) 287–297. (p. 74)
- [91] Sebastian M. Cioabă, Edwin R. van Dam, Jack H. Koolen & Jae-Ho Lee, *Asymptotic results on the spectral radius and the diameter of graphs*, preprint, 2008. (p. 41)
- [92] W. H. Clatworthy, *Partially balanced incomplete block designs with two associate classes and two treatments per block*, J. Res. Nat. Bur. Standards **54** (1955) 177–190. (p. 126)
- [93] A. Clebsch, *Ueber die Flächen vierter Ordnung, welche eine Doppelcurve zweiten Grades besitzen*, J. für Math. **69** (1868) 142–184. (p. 126)

- [94] L. Collatz & U. Sinogowitz, *Spektren endlicher Grafen*, Abh. Math. Sem. Univ. Hamburg **21** (1957) 63–77. (p. 26)
- [95] K. Coolsaet, J. Degraer & E. Spence, *The Strongly Regular $(45, 12, 3, 3)$ graphs*, Electr. J. Combin. **13** (2006) R32. (p. 127)
- [96] K. Coolsaet & J. Degraer, *A computer assisted proof of the uniqueness of the Perkel graph*, Designs, Codes and Cryptography **34** (2005) 155–171. (p. 190)
- [97] K. Coolsaet & J. Degraer, *Using algebraic properties of minimal idempotents for exhaustive computer generation of association schemes*, Electr. J. Combin. **15** (2008) R30. (p. 188)
- [98] R. Courant & D. Hilbert, *Methoden der Mathematischen Physik*, Springer Verlag, Berlin, 1924. (p. 33)
- [99] H. S. M. Coxeter, *Self-dual configurations and regular graphs*, Bull. Amer. Math. Soc. **56** (1950) 413–455. (p. 126)
- [100] R. Craigen & H. Kharaghani, *Hadamard matrices and Hadamard designs*, in: *Handbook of Combinatorial Designs, 2nd ed.*, (C.J. Colbourn and J.H. Denitz eds.), Chapman & Hall/CRC press, Boca Raton, 2007, Chapter V.1. (p. 141)
- [101] Péter Csikvári, *On a conjecture of V. Nikiforov*, Discr. Math. (2009) **309** 4522–4526. (p. 42)
- [102] Péter Csikvári, *Integral trees of arbitrarily large diameters*, preprint, 2009. (p. 85)
- [103] D. M. Cvetković, *Graphs and their spectra*, Publ. Elektrotehn. Fak. Ser. Mat. Fiz. **354–356** (1971) 1–50. (p. 62)
- [104] D. M. Cvetković & M. Doob, *Root systems, forbidden subgraphs and spectral characterizations of line graphs*, in: *Graph Theory*, Proc. Fourth Yugoslav Sem. *Graph Theory*, Novi Sad, 1983 (Cvetković, Gutman, Pisanski, Tošić eds.), pp. 69–99. (p. 185)
- [105] D. M. Cvetković, M. Doob, I. Gutman & A. Torgašev, *Recent Results in the Theory of Graph Spectra*, Annals of Discr. Math. **36**, North Holland, Amsterdam, 1988. (pp. 176, 185)
- [106] D. M. Cvetković, M. Doob & H. Sachs, *Spectra of Graphs: Theory and Applications*, V.E.B. Deutscher Verlag der Wissenschaften, Berlin, 1979. (Also: Academic Press, New York, 1980. Third edition, Johann Ambrosius Barth Verlag, Heidelberg-Leipzig, 1995. Third rev. enl. ed., Wiley, New York, 1998.) (pp. 21, 176, 177, 189)
- [107] D. Cvetković, M. Lepović & S. K. Simić, *The maximal exceptional graphs*, J. Combin. Th. (B) **86** (2002) 347–363. (p. 66)
- [108] D. Cvetković, P. Rowlinson & S. K. Simić, *Eigenspaces of graphs*, Cambridge University Press, Cambridge, 1997. (p. 66)

- [109] D. Cvetković, P. Rowlinson & S. K. Simić, *Graphs with least eigenvalue -2 : The star complement technique*, J. Alg. Combin. **14** (2001) 5–16. (p. 66)
- [110] Dragoš Cvetković, Peter Rowlinson & Slobodan K. Simić, *Spectral generalizations of line graphs; on graphs with least eigenvalue -2* , LMS Lecture Note Series 314, Cambridge University Press, Cambridge, 2004. (pp. 66, 186)
- [111] Dragoš Cvetković, Peter Rowlinson & Slobodan K. Simić, *Signless Laplacians of finite graphs*, Lin. Alg. Appl. **423** (2007) 155–171. (Not cited)
- [112] E. R. van Dam, *Regular graphs with four eigenvalues*, Lin. Alg. Appl. **226-228** (1995) 139–162. (pp. 195, 196)
- [113] E. R. van Dam, *Graphs with few eigenvalues; an interplay between combinatorics and algebra*, CentER Dissertation **20**, Tilburg University, 1996. (pp. 196, 198, 199)
- [114] E. R. van Dam, *Nonregular graphs with three eigenvalues*, J. Combin. Th. (B) **73** (1998) 101–118. (p. 199)
- [115] E. R. van Dam, *The spectral excess theorem for distance-regular graphs: a global (over)view*, Electr. J. Combin. **15** (2008) R129. (p. 163)
- [116] E. R. van Dam and W. H. Haemers, *Graphs with constant μ and $\bar{\mu}$* , Discr. Math. **182** (1998) 293–307. (p. 198)
- [117] E. R. van Dam & W. H. Haemers, *Spectral characterizations of some distance-regular graphs*, J. Alg. Combin. **15** (2002) 189–202. (pp. 175, 176, 187, 189)
- [118] E. R. van Dam & W. H. Haemers, *Which graphs are determined by their spectrum?*, Lin. Alg. Appl. **373** (2003) 241–272. (p. 175)
- [119] E. R. van Dam, W. H. Haemers & J. H. Koolen, *Cospectral graphs and the generalized adjacency matrix*, Lin. Alg. Appl. **423** (2007) 33–41. (pp. 175, 176)
- [120] E. R. van Dam, W. H. Haemers, J. H. Koolen & E. Spence, *Characterizing distance-regularity of graphs by the spectrum*, J. Combin. Th. (A) **113** (2006) 1805–1820. (pp. 187, 189)
- [121] E. R. van Dam & J. H. Koolen, *A new family of distance-regular graphs with unbounded diameter*, Invent. Math. **162** (2005) 189–193. (pp. 157, 159)
- [122] E. R. van Dam & E. Spence, *Small regular graphs with four eigenvalues*, Discr. Math. **189** (1998) 233–257. (pp. 195, 196)
- [123] R. M. Damerell, *On Moore graphs*, Proc. Cambridge Philos. Soc. **74** (1973) 227–236. (p. 161)
- [124] K. C. Das, *A characterization of graphs which achieve the upper bound for the largest Laplacian eigenvalue of graphs*, Lin. Alg. Appl. **376** (2004) 173–186. (p. 48)

- [125] J. Degraer & K. Coolsaet, *Classification of some strongly regular subgraphs of the McLaughlin graph*, Discr. Math. **308** (2008) 395–400. (p. 188)
- [126] Ph. Delsarte, *Weights of linear codes and strongly regular normed spaces*, Discr. Math. **3** (1972) 47–64. (p. 120)
- [127] Ph. Delsarte, *An algebraic approach to the association schemes of coding theory*, Philips Res. Rep. Suppl. **10** (1973). (pp. 147, 151, 154)
- [128] Ph. Delsarte, J.-M. Goethals & J. J. Seidel, *Orthogonal matrices with zero diagonal, Part II*, Canad. J. Math. **23** (1971) 816–832. (p. 139)
- [129] Ph. Delsarte, J.-M. Goethals & J. J. Seidel, *Bounds for systems of lines and Jacobi polynomials*, Philips Res. Rep. **30** (1975) 91*–105*. (pp. 142, 144, 145)
- [130] Ph. Delsarte, J.-M. Goethals & J. J. Seidel, *Spherical codes and designs*, Geom. Dedicata **6** (1977) 363–388. (p. 145)
- [131] M. DeVos, L. Goddyn, B. Mohar & R. Šámal, *Cayley sum graphs and eigenvalues of (3,6)-fullerenes*, J. Combin. Th. (B) **99** (2009) 358–369. (p. 90)
- [132] M. Doob, *Graphs with a small number of distinct eigenvalues*, Ann. N.Y. Acad. Sci. **175** (1970) 104–110. (p. 195)
- [133] M. Doob, *On characterizing certain graphs with four eigenvalues by their spectrum*, Linear Algebra Appl. **3** (1970) 461–482. (p. 195)
- [134] M. Doob & W. H. Haemers, *The complement of the path is determined by its spectrum*, Lin. Alg. Appl. **356** (2002) 57–65. (p. 184)
- [135] Joshua Ducey & Peter Sin, *The elementary divisors of the incidence matrices of skew lines in $PG(3, p)$* , preprint, 2010. [arXiv:1001.2551v1](https://arxiv.org/abs/1001.2551v1) (p. 174)
- [136] Art M. Duval & Victor Reiner, *Shifted simplicial complexes are Laplacian integral*, Trans. Amer. Math. Soc. **354** (2002) 4313–4344. (p. 54)
- [137] M. N. Ellingham, *Basic subgraphs and graph spectra*, Australasian J. Combin. **8** (1993) 247–265. (p. 66)
- [138] Ky Fan, *On a theorem of Weyl concerning eigenvalues of linear transformations I*, Proc. Nat. Acad. Sci. USA **35** (1949) 652–655. (p. 34)
- [139] W. Feit & G. Higman, *The non-existence of certain generalized polygons*, J. Algebra **1** (1964) 114–131. (p. 161)
- [140] N. C. Fiala & W. H. Haemers, *5-Chromatic strongly regular graphs*, Discr. Math. **306** (2006) 3083–3096. (p. 113)
- [141] M. Fiedler, *Algebraic connectivity of graphs*, Czech. Math. J. **23** (98) (1973) 298–305. (p. 21)
- [142] M. Fiedler, *Eigenvectors of acyclic matrices*, Czech. Math. J. **25** (1975) 607–618. (p. 81)

- [143] M. A. Fiol & E. Garriga, *From local adjacency polynomials to locally pseudo-distance-regular graphs*, J. Combin. Th. (B) **71** (1997) 162–183. (pp. 163, 189)
- [144] M. A. Fiol, S. Gago & E. Garriga, *A simple proof of the spectral excess theorem for distance-regular graphs*, Lin. Alg. Appl. **432** (2010) 2418–2422. (p. 163)
- [145] C. Flye Sainte-Marie, *Solution to question nr. 48*, l'Intermédiaire des Mathématiciens **1** (1894) 107–110. (p. 25)
- [146] P. W. Fowler, P. E. John & H. Sachs, *(3,6)-cages, hexagonal toroidal cages, and their spectra*, Discrete Mathematical Chemistry (New Brunswick, NJ, 1998), DIMACS Ser. Discr. Math. Theoret. Comp. Sci. vol. 51, AMS, Providence, RI, 2000, pp. 139–174. (p. 90)
- [147] J. Friedman, *A proof of Alon's second eigenvalue conjecture and related problems*, Memoirs of AMS 195, 2008, 100pp, and [arXiv:cs/0405020v1](#) (118 pages). See also: *A proof of Alon's second eigenvalue conjecture*, pp. 720–724 in: STOC '03: Proc. 35th annual ACM Symp. on Theory of Computing, San Diego, 2003, ACM, New York, 2003. (p. 67)
- [148] H. Fujii & A. Katsuda, *Isospectral graphs and isoperimetric constants*, Discr. Math. **207** (1999) 33–52. (p. 176)
- [149] F. R. Gantmacher, *Applications of the Theory of Matrices*, Interscience, New York, 1959. (pp. 30, 35)
- [150] A. L. Gavriluyk & A. A. Makhnev, *On Krein Graphs without Triangles*, Dokl. Akad. Nauk 403 (2005) 727–730 (Russian) / Doklady Mathematics 72 (2005) 591–594 (English). (p. 105)
- [151] S. Geršgorin, *Über die Abgrenzung der Eigenwerte einer Matrix*, Izv. Akad. Nauk. SSSR Otd. Fiz.-Mat. Nauk **7** (1931) 749–754. (p. 36)
- [152] A. Gewirtz, *Graphs with maximal even girth*, Canad. J. Math. **21** (1969) 915–934. (pp. 127, 129, 188)
- [153] A. Gewirtz, *The uniqueness of $g(2, 2, 10, 56)$* , Trans. New York Acad. Sci. **31** (1969) 656–675. (p. 127)
- [154] C. D. Godsil, *Algebraic Combinatorics*, Chapman and Hall, New York, 1993. (pp. 31, 176, 177)
- [155] C. D. Godsil & B. D. McKay, *Some computational results on the spectra of graphs*, in: *Combinatorial Mathematics IV* (L.R.A. Casse and W.D. Wallis eds.; Proc. fourth Australian conf. on Combin. Math., Adelaide), Lecture Notes in Mathematics 560, Springer-Verlag, Berlin 1976, pp. 73–92. (pp. 177, 180)
- [156] C. D. Godsil & B. D. McKay, *Constructing cospectral graphs*, Aequat. Math. **25** (1982) 257–268. (pp. 23, 178, 180, 181)
- [157] C. D. Godsil, *Spectra of trees*, Annals of Discr. Math. **20** (1984) 151–159. (p. 80)

- [158] C. D. Godsil & G. Royle, *Algebraic Graph Theory*, Springer-Verlag, New York, 2001. (p. 113)
- [159] J.-M. Goethals & J. J. Seidel, *Orthogonal matrices with zero diagonal*, *Canad. J. Math.* **19** (1967) 1001–1010. (p. 140)
- [160] J.-M. Goethals & J. J. Seidel, *Strongly regular graphs derived from combinatorial designs*, *Canad. J. Math.* **22** (1970) 597–614. (pp. 130, 131, 142)
- [161] J.-M. Goethals & J. J. Seidel, *The regular two-graph on 276 vertices*, *Discr. Math.* **12** (1975) 143–158. (pp. 138, 139, 188)
- [162] R. L. Graham & H. O. Pollak, *On the addressing problem for loop switching*, *Bell System Tech. J.* **50** (1971) 2495–2519. (p. 20)
- [163] M. Grassl, *On SIC-POVMs and MUBs in dimension 6*, [quant-ph/0406175](#). (p. 143)
- [164] B. J. Green, *Counting sets with small sumset, and the clique number of random Cayley graphs*, *Combinatorica* **25** (2005) 307–326. (p. 90)
- [165] R. Grone, *Eigenvalues and degree sequences of graphs*, *Lin. Multilin. Alg.* **39** (1995) 133–136. (p. 48)
- [166] R. Grone & R. Merris, *The Laplacian spectrum of a graph II*, *SIAM J. Discr. Math.* **7** (1994) 221–229. (pp. 48, 50)
- [167] Robert Grone, Russell Merris & William Watkins, *Laplacian unimodular equivalence of graphs*, pp. 175–180 in: *Combinatorial and Graph-Theoretic Problems in Linear Algebra* (R. Brualdi, S. Friedland & V. Klee, eds.), IMA Volumes in Mathematics and Its Applications 50, Springer, NY, 1993. (p. 172)
- [168] D. Gryniewicz, V. F. Lem & O. Serra, *The connectivity of addition Cayley graphs*, *Electr. Notes in Discr. Math.* **29** (2007) 135–139. (p. 90)
- [169] Stephen Guattery & Gary L. Miller, *On the performance of the spectral graph partitioning methods*, pp. 233–242 in: *Proc. 2nd ACM-SIAM Symposium on Discrete Algorithms*, 1995. (p. 58)
- [170] Ji-Ming Guo, *A new upper bound for the Laplacian spectral radius of graphs*, *Lin. Alg. Appl.* **400** (2005) 61–66. (p. 48)
- [171] Ji-Ming Guo, *On the third largest Laplacian eigenvalue of a graph*, *Lin. Multilin. Alg.* **55** (2007) 93–102. (p. 49)
- [172] W. H. Haemers, *An upper bound for the Shannon capacity of a graph*, *Colloq. Math. Soc. János Bolyai Vol 25, Algebraic Methods in Graph Theory* (1978) 267–272. (p. 65)
- [173] W. H. Haemers, *On some problems of Lovász concerning the Shannon capacity of graphs*, *IEEE Trans. Inf. Th.* **25** (1979) 231–232. (p. 65)

- [174] W. H. Haemers, *Eigenvalue methods*, in: Packing and covering in combinatorics (A. Schrijver ed.), Math. Centre Tract 106, Mathematical Centre, Amsterdam, 1979, pp. 15–38. (p. 63)
- [175] W. H. Haemers, *Eigenvalue techniques in design and graph theory*, Reidel, Dordrecht, 1980. Thesis (T.H. Eindhoven, 1979) = Math. Centr. Tract 121 (Amsterdam, 1980). (pp. 33, 63, 112)
- [176] W. H. Haemers, *There exists no $(76, 21, 2, 7)$ strongly regular graph*, pp. 175–176 in: Finite Geometry and Combinatorics, F. De Clerck et al. (eds.), LMS Lecture Notes Series 191, Cambridge University Press, 1993. (p. 128)
- [177] W. H. Haemers, *Interlacing eigenvalues and graphs*, Lin. Alg. Appl. **226–228** (1995) 593–616. (p. 33)
- [178] W. H. Haemers, *Distance-regularity and the spectrum of graphs*, Lin. Alg. Appl. **236** (1996) 265–278. (p. 189)
- [179] W. H. Haemers, *Strongly regular graphs with maximal energy*, Lin. Alg. Appl. **429** (2008) 2719–2723. (p. 142)
- [180] W. H. Haemers, A. Mohammadian & B. Tayfeh-Rezaie, *On the sum of laplacian eigenvalues of graphs*, preprint, 2008. (p. 50)
- [181] W. H. Haemers & C. Roos, *An inequality for generalized hexagons*, Geom. Dedicata **10** (1981) 219–222. (p. 161)
- [182] W. H. Haemers & M. S. Shrikhande, *Some remarks on subdesigns of symmetric designs*, J. Stat. Plann. Infer. **3** (1979) 361–366. (p. 76)
- [183] W. H. Haemers & E. Spence, *Graphs cospectral with distance-regular graphs*, Lin. Multilin. Alg. **39** (1995) 91–107. (pp. 158, 168, 176, 187, 188, 189)
- [184] W. H. Haemers & E. Spence, *The Pseudo-Geometric Graphs for Generalised Quadrangles of Order $(3, t)$* , Europ. J. Combin. **22** (2001) 839–845. (p. 128)
- [185] W. H. Haemers & E. Spence, *Enumeration of cospectral graphs*, Europ. J. Combin. **25** (2004) 199–211. (p. 180)
- [186] W. H. Haemers & Q. Xiang, *Strongly regular graphs with parameters $(4m^4, 2m^4 + m^2, m^4 + m^2, m^4 + m^2)$ exist for all $m > 1$* , Europ. J. Combin. **31** (2010) 1553–1559. (p. 142)
- [187] L. Halbeisen & N. Hungerbühler, *Generation of isospectral graphs*, J. Graph Theory **31** (1999) 255–265. (pp. 176, 179)
- [188] N. Hamada & T. Helleseeth, *A characterization of some $\{3v_2 + v_3, 3v_1 + v_2; 3, 3\}$ -minihypers and some $[15, 4, 9; 3]$ -codes with $B_2 = 0$* , J. Stat. Plann. Infer. **56** (1996) 129–146. (p. 129)
- [189] C. Helmberg, B. Mohar, S. Poljak & F. Rendl, *A spectral approach to bandwidth and separator problems in graphs*, Proceedings third IPCO conf. (G. Rinaldi and L. Wolsey, eds.) (1993) 183–194. (p. 72)

- [190] J. van den Heuvel, *Hamilton cycles and eigenvalues of graphs*, Lin. Alg. Appl. **226–228** (1995) 723–730. (p. 70)
- [191] D. G. Higman, *Partial geometries, generalized quadrangles and strongly regular graphs*, in: Atti Convegno di Geometria, Combinatoria e sue Applicazioni, Perugia 1970, (A. Berlotti, ed.) (1971) pp. 263–293. (p. 161)
- [192] D. G. Higman & C. Sims, *A simple group of order 44,352,000*, Math. Z. **105** (1968) 110–113. (p. 129)
- [193] R. Hill, *Caps and Groups*, pp. 389–394 in: Coll. Intern. Teorie Combin. Acc. Naz. Lincei, Roma 1973, Atti dei convegni Lincei 17, Rome, 1976. (p. 124)
- [194] A. J. Hoffman, *On the polynomial of a graph*, Amer. Math. Monthly **70** (1963) 30–36. (pp. 22, 43)
- [195] A. J. Hoffman, *On eigenvalues and colourings of graphs*, in: Graph theory and its applications, (B. Harris, ed.), Acad. Press, New York, 1970, pp. 79–91. (p. 62)
- [196] A. J. Hoffman & R. R. Singleton, *On Moore graphs with diameter 2 and 3*, IBM J. Res. Develop. **4** (1960) 497–504. (pp. 104, 127)
- [197] A. J. Hoffman & J. H. Smith, *On the spectral radii of topologically equivalent graphs*, pp. 273–281 in: Recent Advances in Graph Theory, Proc. Symp. Prague June 1974, Academia Praha, Prague, 1975. (p. 41)
- [198] S. G. Hoggar, *Two quaternionic 4-polytopes*, pp. 219–230 in: The Geometric Vein, Coxeter Festschrift, C. Davis (ed.), Springer, 1981. (p. 143)
- [199] S. G. Hoggar, *64 Lines from a Quaternionic Polytope*, Geom. Dedicata **69** (1998) 287–289. (p. 143)
- [200] R. A. Horn & C. R. Johnson, *Matrix Analysis*, Cambridge University Press, 1985. (p. 30)
- [201] T. Huang & C. Liu, *Spectral characterization of some generalized odd graphs*, Graphs Comb. **15** (1999) 195–209. (p. 189)
- [202] X. Hubaut, *Strongly regular graphs*, Discr. Math. **13** (1975) 357–381. (p. 106)
- [203] James E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge Univ. Press, Cambridge, 1990. (p. 41)
- [204] Y. J. Ionin & M. S. Shrikhande, *Combinatorics of symmetric designs*, Cambridge University Press, 2006. (p. 142)
- [205] I. Martin Isaacs, *Character Theory of Finite Groups*, Academic Press, 1976. (p. 165)
- [206] A. A. Ivanov & S. V. Shpectorov, *The association schemes of dual polar spaces of type ${}^2A_{2d-1}(p^f)$ are characterized by their parameters if $d > 3$* , Lin. Alg. Appl. **114** (1989) 113–139. (p. 117)

- [207] P. S. Jackson & P. Rowlinson, *On graphs with complete bipartite star complements*, Lin. Alg. Appl. **298** (1999) 9–20. (p. 66)
- [208] C. R. Johnson & M. Newman, *A note on cospectral graphs*, J. Combin. Th. (B) **28** (1980) 96–103. (p. 175)
- [209] L. K. Jørgensen & M. Klin, *Switching of edges in strongly regular graphs. I. A family of partial difference sets on 100 vertices*, Electr. J. Combin. **10** (2003) R17. (pp. 129, 132, 142)
- [210] D. Jungnickel, *On subdesigns of symmetric designs*, Math. Z. **181** (1982) 383–393. (p. 76)
- [211] Taher H. Haveliwala and Sepandar D. Kamvar, *The Second Eigenvalue of the Google Matrix*, Technical Report 2003-20, Stanford University, 2003. (p. 56)
- [212] W. M. Kantor & R. A. Liebler, *The rank three permutation representations of the finite classical groups*, Trans. Amer. Math. Soc. **271** (1982) 1–71. (p. 106)
- [213] P. Kaski & P. R. J. Östergård, *There are exactly five biplanes with $k = 11$* , J. Combin. Designs **16** (2008) 117–127. (p. 105)
- [214] J. B. Kelly, *A characteristic property of quadratic residues*, Proc. Amer. Math. Soc. **5** (1954) 38–46. (p. 104)
- [215] A. K. Kelmans, *On graphs with randomly deleted edges*, Acta Math. Acad. Sci. Hung. **37** (1981) 77–88. (p. 42)
- [216] G. Kirchhoff, *Über die Auflösung der Gleichungen, auf welche man bei der untersuchung der linearen verteilung galvanischer Ströme geführt wird*, Ann. Phys. Chem. **72** (1847) 497–508. (p. 15)
- [217] M. Klin & M. Muzychuk, *On graphs with three eigenvalues*, Discr. Math. **189** (1998) 191–207. (p. 199)
- [218] Yehuda Koren, *On spectral graph drawing*, pp. 496–508 in: COCOON 2003: Computing and Combinatorics, Springer LN Comp. Sci. 2697, 2003. (p. 56)
- [219] M. Krivelevich & B. Sudakov, *Sparse pseudo-random graphs are Hamiltonian*, J. Graph Th. **42** (2003) 17–33. (p. 70)
- [220] C. L. M. de Lange, *Some new cyclotomic strongly regular graphs*, J. Alg. Combin. **4** (1995) 329–330. (p. 126)
- [221] P. W. H. Lemmens & J. J. Seidel, *Equiangular lines*, J. Algebra **24** (1973) 494–512. (pp. 40, 142, 143, 144)
- [222] Jiong-Sheng Li & Yong-Liang Pan, *A note on the second largest eigenvalue of the Laplacian matrix of a graph*, Lin. Multilin. Alg. **48** (2000) 117–121. (p. 49)

- [223] Jiong-Sheng Li & Yong-Liang Pan, *De Caen's inequality and bounds on the largest Laplacian eigenvalue of a graph*, Lin. Alg. Appl. **328** (2001) 153–160. (p. 48)
- [224] Jiong-Sheng Li & Xiao-Dong Zhang, *On the Laplacian eigenvalues of a graph*, Lin. Alg. Appl. **285** (1998) 305–307. (p. 48)
- [225] M. W. Liebeck, *The affine permutation groups of rank three*, Proc. London Math. Soc. (3) **54** (1987) 477–516. (p. 106)
- [226] M. W. Liebeck & J. Saxl, *The finite primitive permutation groups of rank three*, Bull. London Math. Soc. **18** (1986) 165–172. (p. 106)
- [227] J. H. van Lint, *Introduction to coding theory*, Graduate Texts in Math. 86, Springer, New York, 1982. (p. 120)
- [228] J. H. van Lint & A. Schrijver, *Constructions of strongly regular graphs, two-weight codes and partial geometries by finite fields*, Combinatorica **1** (1981) 63–73. (pp. 118, 130)
- [229] J. H. van Lint & J. J. Seidel, *Equilateral point sets in elliptic geometry*, Indag. Math. **28** (1966) 335–348. (pp. 142, 144)
- [230] P. Lisoněk, *New maximal two-distance sets*, J. Combin. Th. (A) **77** (1997) 318–338. (p. 145)
- [231] Huiqing Liu, Mei Lu & Feng Tian, *On the Laplacian spectral radius of a graph*, Lin. Alg. Appl. **376** (2004) 135–141. (p. 48)
- [232] L. Lovász, *On the Shannon capacity of a graph*, IEEE Trans. Inf. Th. **25** (1979) 1–7. (p. 65)
- [233] A. Lubotzky, *Cayley graphs: eigenvalues, expanders and random walks*, pp. 155–189 in: *Surveys in combinatorics, 1995* (Stirling), London Math. Soc. Lecture Note Ser., 218, Cambridge Univ. Press, Cambridge, 1995. (p. 176)
- [234] A. Lubotzky, R. Phillips & P. Sarnak, *Ramanujan Graphs*, Combinatorica **8** (1988) 261–277. (p. 67)
- [235] Ulrike von Luxburg, *A Tutorial on Spectral Clustering*, Statistics and Computing **17** (2007) 395–416. (p. 58)
- [236] F. J. MacWilliams & N. J. A. Sloane, *The Theory of Error-Correcting Codes*, North Holland Publ. Co., Amsterdam, 1977. (pp. 120, 169)
- [237] C. L. Mallows & N. J. A. Sloane, *Two-graphs, switching classes and Euler graphs are equal in number*, SIAM J. Appl. Math. **28** (1975) 876–880. (p. 135)
- [238] M. Marcus & H. Minc, *A survey of matrix theory and matrix inequalities*, Allyn and Bacon, Boston, 1964. (p. 30)
- [239] M. Marcus & H. Minc, *Introduction to linear algebra*, Macmillan, New York, 1965. (p. 148)

- [240] G. A. Margulis, *Explicit group-theoretical constructions of combinatorial schemes and their application to the design of expanders and concentrators*, Probl. Peredachi Inf. **24** (1988) 51–60. English: Problems of Information Transmission **24** (1988) 39–46. (p. 67)
- [241] R. A. Mathon, *Symmetric conference matrices of order $pq^2 + 1$* , Canad. J. Math. **30** (1978) 321–331. (p. 127)
- [242] R. J. McEliece & H. Rumsey, jr., *Euler products, cyclotomy and coding*, J. Number Th. **4** (1972) 302–311. (p. 125)
- [243] B. D. McKay, *On the spectral characterisation of trees*, Ars Combinatoria **3** (1979) 219–232. (p. 177)
- [244] B. D. McKay & E. Spence, *Classification of regular two-graphs on 36 and 38 vertices*, Australas. J. Comb. **24** (2001) 293–300. (pp. 127, 174, 188)
- [245] R. Merris, *Large families of Laplacian isospectral graphs*, Lin. Multilin. Alg. **43** (1997) 201–205. (p. 176)
- [246] R. Merris, *A note on Laplacian graph eigenvalues*, Lin. Alg. Appl. **285** (1998) 33–35. (p. 48)
- [247] B. Mohar, *A domain monotonicity theorem for graphs and Hamiltonicity*, Discr. Appl. Math. **36** (1992) 169–177. (p. 70)
- [248] Oleg R. Musin, *Spherical two-distance sets*, arXiv:0801.3706, 2008. (p. 145)
- [249] A. Neumaier, *Strongly regular graphs with smallest eigenvalue $-m$* , Archiv Math. **33** (1979) 392–400. (pp. 106, 109)
- [250] A. Neumaier, *New inequalities for the parameters of an association scheme*, pp. 365–367 in: Combinatorics and Graph Theory, Proc. Calcutta 1980, Lecture Notes in Math. Vol 885 (S.B. Rao, ed.), Springer, Berlin. (p. 152)
- [251] A. Neumaier, *The second largest eigenvalue of a tree*, Lin. Alg. Appl. **46** (1982) 9–25. (p. 84)
- [252] A. Neumaier, *Completely regular two-graphs*, Arch. Math. **38** (1982) 378–384. (pp. 138, 139)
- [253] R. E. A. C. Paley, *On orthogonal matrices*, J. Math. Phys. Mass. Inst. Technol. **12** (1933) 311–320. (p. 140)
- [254] D. V. Pasechnik, *The triangular extensions of a generalized quadrangle of order $(3, 3)$* , Simon Stevin **2** (1995) 509–518. (p. 139)
- [255] A. J. L. Paulus, *Conference matrices and graphs of order 26*, Technische Hogeschool Eindhoven, report WSK 73/06, Eindhoven, 1983, 89 pp. (p. 126)
- [256] S. E. Payne, *An inequality for generalized quadrangles*, Proc. Amer. Math. Soc. **71** (1978) 147–152. (p. 33)

- [257] M. J. P. Peeters, *Ranks and Structure of Graphs*, PhD thesis, Tilburg University, 1995. (p. 170)
- [258] René Peeters, *On the p -ranks of the adjacency matrices of distance-regular graphs*, J. Alg. Combin. **15** (2002) 127–149. (p. 168)
- [259] J. Petersen, *Sur le théorème de Tait*, L'Intermédiaire des Mathématiciens **5** (1898) 225–227. (p. 126)
- [260] Tomaž Pisanski & John Shawe-Taylor, *Characterizing graph drawing with eigenvectors*, J. Chem. Inf. Comp. Sci. **40** (2000) 567–571. (p. 57)
- [261] L. Pyber, *Large connected strongly regular graphs are Hamiltonian*, preprint, 2009. (p. 70)
- [262] Gregory T. Quenell, *Eigenvalue comparisons in graph theory*, Pacif. J. Math. **176** (1996) 443–461. (p. 67)
- [263] P. Rowlinson, *On the maximal index of graphs with a prescribed number of edges*, Lin. Alg. Appl. **110** (1988) 43–53. (p. 42)
- [264] P. Rowlinson, *The characteristic polynomials of modified graphs*, Discr. Appl. Math. **67** (1996) 209–219. (p. 176)
- [265] B. L. Schader, *On tournament matrices*, Lin. Alg. Appl. **162–164** (1992) 335–368. (p. 25)
- [266] I. Schur, *Über eine Klasse von Mittelbildungen mit Anwendungen auf die Determinantentheorie*, Sitzungsber. Berliner Math. Ges. **22** (1923) 9–20. (p. 33)
- [267] A. J. Schwenk, *Almost all trees are cospectral*, in: *New directions in the theory of graphs* (F. Harary ed.), Academic Press, New York 1973, pp. 275–307. (p. 177)
- [268] A. J. Schwenk, *Exactly thirteen connected cubic graphs have integral spectra*, pp. 516–533 in: *Theory and Applications of graphs* (Proc. Kalamazoo 1976), ed. Y. Alavi & D. Lick, Springer, 1978. (p. 44)
- [269] A. J. Schwenk & O. P. Lossers, *Solution to advanced problem #6434*, Amer. Math. Monthly **94** (1987) 885–886. (p. 20)
- [270] L. L. Scott Jr, *A condition on Higman's parameters*, Notices Amer. Math. Soc. **20**, A-97 (1973) 701-20-45. (pp. 105, 151)
- [271] J. Seberry Wallis, *Hadamard matrices*, Part 4 in: W. D. Wallis, A. P. Street & J. Seberry Wallis, *Combinatorics: Room squares, Sum-free sets, Hadamard matrices*, Springer LNM 292, Berlin etc., 1972. (pp. 141, 142)
- [272] J. Seberry & M. Yamada, *Hadamard matrices, sequences, and block designs*, pp. 431–560 in: *Contemporary design theory*, J. H. Dinitz & D. R. Stinson (eds.), Wiley, New York, 1992. (p. 141)
- [273] J. J. Seidel, *Strongly regular graphs of L_2 -type and of triangular type*, Kon. Nederl. Akad. Wetensch. (A) **70** (1967) 188–196 = Indag. Math. 29 (1967) 188–196. (p. 139)

- [274] J. J. Seidel, *Strongly regular graphs with $(-1, 1, 0)$ adjacent matrix having eigenvalue 3*, Lin. Alg. Appl. **1** (1968) 281–298. (pp. 109, 126, 134, 188)
- [275] J. J. Seidel, *Graphs and two-graphs*, in: *Proc. 5th Southeast. Conf. Comb., Graph Th., Comp. 1974* (F. Hoffman et al., eds.), Utilitas Mathematica Pub., Winnipeg, 1974, pp. 125–143. (p. 188)
- [276] J. J. Seidel, *Strongly regular graphs*, in: *Surveys in Combinatorics*, Proc. 7th British Combinatorial Conf., LMSLN ser. Vol 38, ed. B. Bollobás, Cambridge Univ. Press, Cambridge, 1979, pp.157–180. (p. 151)
- [277] J. J. Seidel, *Discrete Non-Euclidean Geometry*, pp. 843–920 in: *Handbook of Incidence Geometry*, F. Buekenhout, ed., NHPC, 1995. (p. 144)
- [278] J. J. Seidel & S. V. Tsaranov, *Two-graphs, related groups, and root systems*, Bull. Soc. Math. Belgique **42** (1990) 695–711. (p. 135)
- [279] E. Seneta, *Non-negative matrices and Markov chains*, Springer Series in Statistics, Springer, New York, 1981. (p. 30)
- [280] Á. Seress, *Large families of cospectral graphs*, Designs, Codes and Cryptography **21** (2000) 205–208. (p. 176)
- [281] J.-P. Serre, *Répartition asymptotique des valeurs propres de l'opérateur de Hecke T_p* , J. Amer. Math. Soc. **10** (1997) 75–102. (p. 67)
- [282] C. E. Shannon, *The zero-error capacity of a noisy channel*, IRE Trans. Inform. Th. vol. IT-2 (1956) 8–19. (p. 64)
- [283] S. S. Shrikhande, *The uniqueness of the L_2 association scheme*, Ann. Math. Statist. **30** (1959) 781–798. (pp. 110, 126, 188)
- [284] Peter Sin, *The p -rank of the incidence matrix of intersecting linear subspaces*, Des. Codes Cryptogr. **31** (2004) 213–220. (p. 174)
- [285] J. H. Smith, *Some properties of the spectrum of a graph*, pp. 403–406 in: *Combinatorial Structures and their applications* (Proc. Conf. Calgary, 1969), R. Guy et al. (eds.), Gordon and Breach, New York, 1970. (p. 40)
- [286] E. Spence, *The Strongly Regular $(40, 12, 2, 4)$ Graphs*, Electr. J. Combin. **7** 2000, R22. (p. 127)
- [287] E. Spence, <http://www.maths.gla.ac.uk/~es/cospecL/cospecL.html> and [.../cospecSignlessL/cospecSignlessL.html](http://www.maths.gla.ac.uk/~es/cospecSignlessL/cospecSignlessL.html) (p. 181)
- [288] Daniel A. Spielman & Shang-Hua Teng, *Spectral partitioning works: planar graphs and finite element meshes*, pp. 96–105 in: *Proc. 37th Symposium on Foundations of Computer Science* (Burlington, 1996), IEEE 1996. (p. 58)
- [289] Toshikazu Sunada, *Riemannian coverings and isospectral manifolds*, Ann. Math. **121** (1985) 169–186. (p. 179)

- [290] J. J. Sylvester, *A demonstration of the theorem that every homogeneous quadratic polynomial is reducible by real orthogonal substitutions to the form of a sum of positive and negative squares*, Philos. Mag. **4** (1852) 138–142. (Reprinted in: Coll. Math. Papers 1, Cambridge Univ. Press, 1904, pp. 378–381.) (p. 35)
- [291] G. Szegő, *Orthogonal Polynomials*, AMS Colloq. Publ. **23**, 1939 (1st ed.), 1975 (4th ed.). (p. 154)
- [292] R. M. Tanner, *Explicit concentrators from generalized n -gons*, SIAM J. Algebr. Discr. Meth. **5** (1984) 287–293. (p. 68)
- [293] D. E. Taylor, *Some topics in the theory of finite groups*, Ph.D. Thesis, Univ. of Oxford, 1971. (p. 137)
- [294] D. E. Taylor, *Graphs and block designs associated with the three-dimensional unitary groups*, pp. 128–131 in: Combinatorial Math., Proc. 2nd Austr. Conf., D. A. Holton, ed., Springer LNM 403, Berlin 1974. (p. 130)
- [295] D. E. Taylor, *Regular 2-graphs*, Proc. London Math. Soc. **35** (1977) 257–274. (pp. 130, 138)
- [296] W. T. Tutte, *The factorizations of linear graphs*, J. London Math. Soc. **22** (1947) 107–111. (p. 73)
- [297] W. T. Tutte, *All the king's horses. A guide to reconstruction*, pp. 15–33 in: Graph Theory and Related Topics (Proc. Conf. Waterloo, 1977), J. A. Bondy & U. S. R. Murty (eds.), Academic Press, New York, 1979. (p. 24)
- [298] H. Van Maldeghem, *Generalized Polygons*, Birkhäuser, Basel, 1998. (p. 161)
- [299] R. S. Varga, *Matrix iterative analysis*, Prentice-Hall, Englewood Cliffs, New York, 1962. (p. 30)
- [300] R. S. Varga, *Geršgorin and his circles*, Springer, 2004. (p. 36)
- [301] Wei Wang & Cheng-xian Xu, *A sufficient condition for a family of graphs being determined by their generalized spectra*, Europ. J. Combin. **27** (2006) 826–840. (p. 192)
- [302] Wei Wang & Cheng-xian Xu, *Note: On the generalized spectral characterization of graphs having an isolated vertex*, Lin. Alg. Appl. **425** (2007) 210–215. (p. 193)
- [303] Y. Wang, Y. Fan & Y. Tan, *On graphs with three distinct Laplacian eigenvalues*, Appl. Math. J. Chinese Univ. Ser. B **22** (2007) 478–484. (p. 198)
- [304] Mamoru Watanabe, *Note on integral trees*, Math. Rep. Toyama Univ. **2** (1979) 95–100. (p. 85)
- [305] H. Whitney, *Congruent graphs and the connectivity of graphs*, Amer. J. Math. **54** (1932) 150–167. (p. 187)
- [306] H. A. Wilbrink & A. E. Brouwer, *A $(57, 14, 1)$ strongly regular graph does not exist*, Indag. Math. **45** (1983) 117–121. (p. 127)

- [307] H. S. Wilf, *The eigenvalues of a graph and its chromatic number*, J. London Math. Soc. **42** (1967) 330–332. (p. 62)
- [308] R. Woo & A. Neumaier, *On graphs whose spectral radius is bounded by $\frac{3}{2}\sqrt{2}$* , Graphs Comb. **23** (2007) 713–726. (p. 41)
- [309] Chen Yan, *Properties of spectra of graphs and line graphs*, Appl. Math. J. Chinese Univ. Ser. B **17** (2002) 371–376. (p. 48)
- [310] Gerhard Zauner, *Quantendesigns, Grundzüge einer nichtkommutativen Designtheorie*, Ph.D. thesis, Vienna, 1999. (p. 143)
- [311] F. Zhang, *Matrix Theory*, Springer-Verlag, New York, 1999. (p. 34)
- [312] Xiao-Dong Zhang, *Two sharp upper bounds for the Laplacian eigenvalues*, Lin. Alg. Appl. **376** (2004) 207–213. (p. 48)
- [313] Xiao-Dong Zhang & Rong Luo, *The spectral radius of triangle-free graphs*, Australas. J. Comb. **26** (2002) 33–39. (p. 47)

Author Index

- Ajtai, 68
 Alon, 64–67, 71
 Aschbacher, 153
 Ayooobi, 200

 Bagchi, 106
 Bai, 50
 Bang, 160
 Bannai, 160, 161
 Belevitch, 140
 Berman, 30
 Bigalke, 112
 Blokhuis, 139, 145
 Boppana, 66
 Bose, 109, 147, 148
 Brauer, 165
 Bridges, 199
 Brouwer, 41, 49, 85, 106, 115, 123, 161, 189
 Brualdi, 42
 Burrow, 148
 Bussemaker, 44, 186

 Calderbank, 120
 Cameron, 93, 115, 117, 153, 185
 Cauchy, 33
 Cayley, 15
 Chang, 110
 Cherlin, 115
 Chesnokov, 182
 Chuang, 199
 Chung, 70, 90
 Chvátal, 69, 112
 Cioabă, 41, 74
 Cohen, 161, 189
 Courant, 33
 Csikvári, 42, 85
 Cvetković, 21, 44, 62, 185, 186

 van Dam, 41, 120, 157, 159, 175, 176, 195, 196, 199, 200
 Damerell, 161
 Davenport, 125
 Delsarte, 120, 144, 147, 151, 154
 Doob, 21, 185, 195
 Dubickas, 160
 Duval, 54

 Erdős, 112
 van Eupen, 123

 Fan, 34
 Feit, 76, 161
 Fiedler, 21, 81
 Fiol, 162, 163, 189
 Foulser, 106
 Friedman, 67

 Gago, 163
 Gantmacher, 30, 35
 Garriga, 162, 163, 189
 Gavriluk, 105
 Geršgorin, 36
 Gerzon, 143
 Godsil, 31, 176–178, 180
 Goethals, 93, 117, 138, 144, 185
 Graham, 20
 Gregory, 74
 Grone, 48, 50, 172
 Guo, 49

 Haemers, 33, 49, 65, 74, 76, 111, 139, 161, 175, 176, 180, 182
 Hasse, 125
 Helmberg, 72
 Higman, 33, 137, 153, 161
 Hilbert, 33
 Hill, 124
 Hoffman, 22, 41–43, 62, 104
 Horn, 30
 Huang, 189
 Hubaut, 106

 Isaacs, 165
 Ito, 160, 161
 Ivanov, 117

 Johnson, 30, 175
 Jung, 112
 Jungnickel, 76

 Kantor, 106, 115, 120
 Kaski, 105
 Kelly, 24
 Kelmans, 42
 Kirchhoff, 15

- Klin, 199
 Komlós, 68
 Koolen, 41, 157, 159, 160, 175, 176
 Krivelevich, 70

 de Lange, 126
 Lee, 41
 Lemmens, 40, 144
 Liebeck, 106
 Liebler, 106
 van Lint, 106, 118, 120, 144
 Lisoněk, 145
 Liu, 189
 Lovász, 65
 Lubotzky, 67
 Luo, 47

 MacWilliams, 120, 169
 Makhnev, 105
 Marcus, 30, 148
 Margulis, 67
 McEliece, 125
 McKay, 176–178, 180
 Mena, 199
 Merris, 50, 172
 Mesner, 148
 Milman, 71
 Minc, 30, 148
 Mohar, 72
 Moulton, 160
 Musin, 145
 Muzychuk, 199

 Neumaier, 41, 84, 106, 139, 152, 161, 189
 Newman, 175

 Omid, 199, 200
 Östergård, 105

 Page, 55
 Paley, 140
 Pasechnik, 139
 Payne, 33
 Peeters, 168
 Phillips, 67
 Pisanski, 57
 Plemmons, 30
 Poljak, 72
 Pollak, 20
 Pyber, 70

 Quenell, 67

 Reiner, 54
 Rendl, 72
 Roos, 161
 Rowlinson, 42
 Rumsey, 125

 Sachs, 21
 Sarnak, 67
 Saxl, 106
 Schrijver, 118
 Schur, 33
 Schwenk, 44, 176, 177
 Scott, 105, 151
 Seidel, 40, 93, 106, 109, 117, 134, 135, 138, 142, 144, 151, 152, 185, 186
 Seneta, 30
 Seress, 176
 Serre, 66, 67
 Seymour, 64
 Shannon, 64
 Shawe-Taylor, 57
 Shimamoto, 147
 Shpectorov, 117
 Shrikhande, 76, 110
 Shult, 93, 185
 Sims, 33
 Sin, 174
 Singleton, 104
 Sloane, 120, 169
 Smith, 40, 41
 Spence, 180, 195, 196, 200
 Stickelberger, 125
 Sudakov, 70
 Sunada, 176
 Sylvester, 35
 Szegő, 154
 Szemerédi, 68

 Tanner, 68
 Tayfeh-Rezaie, 200
 Taylor, 137
 Tsaranov, 135
 Tutte, 73

 Ulam, 24

 Van Maldeghem, 161
 Varga, 30, 36

Wang, 192, 193

Watanabe, 85

Watkins, 172

Whitney, 187

Wilbrink, 76

Wilf, 62

Witsenhausen, 20

Woo, 41

Xu, 192, 193

Yan, 48

Zhang, 47

Subject Index

- absolute bound, 106, 143, 152
- absolute point, 77
- addition Cayley graphs, 90
- adjacency matrix, 11
- algebraic connectivity, 21
- antipodal, 161
- associated, 134
- association scheme, 147
- automorphism, 20

- bandwidth, 72
- Bannai-Ito conjecture, 160
- Berlekamp-van Lint-Seidel graph, 123, 131
- biaffine plane, 191
- biclique, 71
- Biggs-Smith graph, 160
- bipartite double, 19
- bipartite graph, 15, 43
- biplane, 107
- block graph, 108
- Bose-Mesner algebra, 148
- Brouwer-Haemers graph, 117

- cap, 132
- Cauchy-Binet formula, 15
- Cayley graph, 19, 87
- Cayley sum graph, 90
- Chang graphs, 109, 110, 172
- characteristic matrix, 30
- characteristic polynomial, 12
- chromatic number, 62
- 3-claw, 93
- Clebsch graph, 105, 110
- clique, 61, 138
- clique number, 61
- coalescence, 177
- cocktailparty graph, 93
- coclique, 61
- cofactor, 15
- coherent, 138
- coherent triple, 134
- collinearity graph, 108, 178
- cometric association scheme, 154
- companion matrix, 192
- complement, 14, 102, 135
- completely regular two-graph, 138

- cone, 26, 199
- conference graphs, 104, 140
- conference matrix, 139
- conjunction, 19
- core, 140
- cospectral, 22, 23, 26, 45, 158, 175–183
- Courant-Weyl inequalities, 34
- cover, 88
- covering, 88
- Cvetković bound, 62

- de Bruijn cycle, 25
- de Bruijn graph, 25
- Desargues graph, 45
- descendant, 135
- determinant, 95
- diagonally dominant, 36
- diameter, 14, 70
- dimension, 53
- directed incidence matrix, 11
- discriminant, 95
- distance-regular, 157
- dominance order, 54
- downshifting, 54
- DS graph, 175
- dual, 120, 123
- dual lattice, 94
- Dynkin diagram, 41, 96

- edge expansion, 69
- effective length, 120
- eigenvalue components, 81
- elementary divisors, 171
- endpath, 41
- energy, 57
- equiangular tight frame, 143
- equitable, 30
- equivalent code, 120
- even lattice, 94
- exceptional graph, 99
- expander, 68, 90
- expander mixing lemma, 68
- extended fundamental system, 96

- feasible, 126
- few eigenvalues, 195

- firm generalized n -gon, 114
- flag, 75
- flag graph, 190
- folded 5-cube, 105
- Fowler conjecture, 90
- Frame quotient, 106
- (3, 6)-fullerene, 90
- fundamental system, 96

- Gauss sum, 125
- generalized m -gon, 161
- generalized n -gon, 114
- generalized adjacency matrix, 175
- generalized line graph, 99
- generalized quadrangle, 114
- generalized quadrangles, 109
- geodetic, 162
- geometric srg, 109
- Geršgorin circles, 36
- Gewirtz graph, 105, 107, 112
- Godsil's lemma, 80
- Google, 55
- Gram matrix, 35
- graph, 88
- graphical Hadamard matrix, 141
- Grassmann graph, 159
- Grone-Merris conjecture, 50
- groups (of a design), 107
- Guo conjecture, 49

- Hadamard matrix, 140
- half case, 104, 106
- Hall-Janko graph, 108, 132
- halved graph, 114
- Hamiltonian, 69, 70, 112
- Hamming graph, 158
- Hamming scheme, 155
- Hermitean linear transformation, 27
- Higman-Sims graph, 105
- Higman-Sims technique, 33
- Hill code, 124
- Hoffman bound, 59, 62
- Hoffman coloring, 63
- Hoffman polynomial, 43, 45
- Hoffman-Singleton graph, 72, 104, 105, 112
- homomorphism, 88
- hypercube, 19
- hyperoval, 123

- imprimitive drg, 161
- imprimitive srg, 103
- improper strongly regular graph, 133
- incidence graph, 75
- incidence matrix, 11
- independence number, 61
- index, 39, 107
- inner distribution, 150
- integral lattice, 94
- integral tree, 85
- interlacing, 31
- intersection array, 157
- intersection matrices, 148
- intersection numbers, 147
- invariant factors, 171
- irreducible lattice, 95
- irreducible nonnegative matrix, 27
- isospectral, 22

- Johnson graph, 159
- join, 31
- just y -cospectral, 175

- k -means, 57
- Kelmans operation, 42
- Klein graph, 168
- Kneser graph, 159
- Krein conditions, 105, 151
- Krein parameters, 151
- Kronecker product, 19

- Laplace matrix, 11
- Laplace spectrum, 12
- lattice, 94
- lattice graph, 18, 109
- level of a rational matrix, 192
- line graph, 18, 93, 114, 178
- linear code, 120
- local graph, 188
- loop, 88

- M_{22} graph, 105, 112
- matrix-tree theorem, 15
- Menon design, 141
- metric association scheme, 153
- minimal polynomial, 166
- Moore graph, 113, 153, 161
- μ -bound, 106, 109
- multiplicity free, 88

- net, 108

- Odd graph, 159
- opposite, 114
- orbitals, 106
- orthogonal, 115
- orthogonal array, 108
- orthogonal direct sum, 95
- orthogonality relations, 149
- Paley graph, 67, 101, 140, 168
- partial geometry, 108
- partial linear space, 177
- pentagon, 105
- perfect matching, 72
- period of an irreducible matrix, 28
- Perron-Frobenius, 27
- Perron-Frobenius eigenvector, 55
- Petersen graph, 18, 20, 45, 62, 69, 70, 72, 83, 104, 105, 110, 112
- point graph, 108, 114, 178
- polarity, 77
- positive definite, 35
- positive semidefinite, 35
- primitive nonnegative matrix, 27
- primitive strongly regular graph, 103
- projective code, 120, 122
- projective plane, 77
- proper vertex coloring, 62
- pseudo Latin square graph, 108
- pseudo-geometric, 109
- quasisymmetric design, 107
- quotient graph, 89
- quotient matrix, 30
- Ramanujan graph, 67
- rank of a permutation group, 106
- rationality conditions, 104, 150
- Rayleigh quotient, 31
- reconstruction conjecture, 24
- reduced fundamental system, 96
- regular generalized m -gon, 161
- regular graph, 13
- regular Hadamard matrix, 141
- regular tournament, 26
- restricted eigenvalue, 103, 133, 197
- root lattice, 93, 96
- root system, 96
- roots, 96
- Saltire pair, 180, 183, 193
- Schläfli graph, 65, 110
- Schur complements, 33
- Schur's inequality, 33
- Seidel adjacency matrix, 23, 142
- Seidel spectrum, 23
- Seidel switching, 23, 142
- self-dual lattice, 95
- semi-bipartite graph, 51
- Shannon capacity, 64
- Shrikhande graph, 109, 110, 172
- SICPOVM, 143
- sign changes, 154
- signless Laplace matrix, 11
- simplicial complex, 53
- Singleton bound, 123
- Smith Normal Form, 171
- spectral center, 84
- spectral radius, 39
- spectrum, 11
- split graph, 51
- star complement, 65
- star partition, 66
- star subset, 65
- stepwise matrix, 42
- strength, 107
- strictly diagonally dominant, 36
- strong graph, 133
- strongly connected directed graph, 28
- strongly regular, 101
- subdividing, 41
- sum graphs, 90
- support, 82, 120
- switching class, 23
- symmetric design, 76
- t -design, 75
- ternary Golay code, 118, 123
- tesseract, 22
- thick generalized n -gon, 114
- threshold graph, 50
- tight interlacing, 31
- toughness, 69
- tournament, 25
- transitive tournament, 25
- transversal design, 107
- triangular graph, 18, 102, 109, 159
- twisted Grassmann graphs, 159
- two-graph, 134
- two-weight code, 120

Type I lattice, 95
Type II lattice, 95

unimodular lattice, 95
unital, 76
universal cover, 88

van Dam-Koolen graphs, 159
vertex connectivity, 110

walk matrix, 192
walk-regular, 195
walks, 14
weak generalized quadrangle, 116
weight, 120

 y -cospectral, 175

zero-error channel, 64