

# Human-machine interaction in virtual reality

Paul MacNeilage, Psychology  
Eelke Folmer, Computer Science

# VR Research Topics?

- Interfaces – locomotion, manipulation, eye movements
- Other modalities – haptics, audio, etc.
- Adaptation – advantages, disadvantages
- Sickness – conflict, methods to reduce it
- Presence – methods to increase it

# Experiments on Human Subjects

- Central to VR research and development
- Necessary to prove that a novel method, technology, practice is really beneficial
- All claims should be scientifically proven
- Don't trust introspection!


Experimental psychology

Human factors research

Human-computer interaction


# Scientific Method

## The Scientific Method as an Ongoing Process



From Wikipedia

# Experimental Process




# Psychophysics

- Measuring a psychometric function
- Multisensory interactions
  - Cue integration
  - Conflict detection
- Related to textbook sections:
  - Chapter 12.4: Experiments on human subjects
  - Chapter 6.4: Combining sources of information
  - Chapter 8.4: Mismatched motion and vection




# Signal Detection Theory

- Mathematical model for decision-making in the face of uncertainty
- Ability to tell signal from noise
- Originally developed during WWII to interpret radar readings



- Later applied to psychological research




# Self-motion Lab






# Vestibular Perceptual Sensitivity

One-interval forced choice procedure




# Motion platform and chair



Upright

Side-down

# Vestibular Perceptual Sensitivity




# Relevance for VR

- Rigorous methods for quantifying perception
- May be applied in almost any context
  - Vision: depth, motion, color, etc.
  - Audition: loudness, pitch, location, etc.
  - Haptic: size, shape, etc.
  - Multisensory:
    - Cue integration
    - Conflict detection

# Today's topics


- Measuring perception: psychophysics
- Multisensory interactions
  - Cue integration
  - Conflict detection
- Related to textbook sections:
  - Chapter 12.4: Experiments on human subjects
  - Chapter 6.4: Combining sources of information
  - Chapter 8.4: Mismatched motion and vection

# Human-VR Loop



# Perception as Bayesian Inference

- Brain must infer the state of the outside world from incomplete information



Ernst & DeLuca 2011

# Probabilistic Framework

- Sensory information = evidence ( $I$ )
- Physical states = possible interpretations ( $S$ )
- What is the most likely interpretation given the available evidence?  $P(S|I)$



# Bayes' rule / law / theorem

- Mathematical description of the problem

$$P(S | I) \propto P(I | S)P(S)$$

Posterior

used to make  
perceptual  
judgment

Likelihood

sensory  
information


Prior

knowledge,  
experience

# Bayes' rule / law / theorem

- Mathematical description of the problem

$$P(S | I) \propto P(I | S)P(S)$$




- Decision rule: maximum a posteriori (MAP)

# Priors Responsible for Many Illusions

- Light-from-above prior

a



# Cue Integration

- How does the brain combine different sources of information?
- According to the rules of probability!
- Bayes rule can be applied

# Perception Depends on Many Cues

- Example: multiple visual cues to distance



# Perception Depends on Many Cues

- Example: visual and haptic cues to size

Visual



Haptic



- Different senses referred to as modalities


# Maximum-likelihood Estimation (MLE)

- Bayesian Model of Cue Combination
- Assumptions:
  - Gaussian Noise (i.e. normally distributed)
  - Independence
  - Uniform prior

$$P(S \mid I_1, I_2) \propto P(I_1 \mid S)P(I_2 \mid S)$$


# Maximum-likelihood Estimation (MLE)

- “Optimal” estimate is weighted linear combination of contributing cues
- Weights determined by relative cue reliabilities
- Combined estimate is less variable than any cue in isolation





# Maximum-likelihood Estimation (MLE)



## Predictions

- 1) Weight depends on relative variability

$$\hat{S}_{comb} = \hat{S}_1 w_1 + \hat{S}_2 w_2$$

$$w_1 + w_2 = 1$$

$$w_i = \frac{1}{\sigma_i^2} / \sum_{j=1}^n \frac{1}{\sigma_j^2}$$

- 2) Combined estimate is least variable

$$\sigma_{comb}^2 = \frac{\sigma_1^2 \sigma_2^2}{\sigma_1^2 + \sigma_2^2}$$

---

# Humans integrate visual and haptic information in a statistically optimal fashion

Marc O. Ernst\* & Martin S. Banks


*Vision Science Program/School of Optometry, University of California, Berkeley  
94720-2020, USA*

---


When a person looks at an object while exploring it with their hand, vision and touch both provide information for estimating the properties of the object. Vision frequently dominates the integrated visual–haptic percept, for example when judging size, shape or position<sup>1–3</sup>, but in some circumstances the percept is clearly affected by haptics<sup>4–7</sup>. Here we propose that a general principle, which minimizes variance in the final estimate, determines the degree to which vision or haptics dominates. This principle is realized by using maximum-likelihood estimation<sup>8–15</sup> to combine the inputs. To investigate cue combination quantitatively, we first measured the variances associated with visual and haptic estimation of height. We then used these measurements to construct a maximum-likelihood integrator. This model behaved very similarly to humans in a visual–haptic task. Thus, the nervous system seems to combine visual and haptic information in a fashion that is similar to a maximum-likelihood integrator. Visual dominance occurs when the variance associated with visual estimation is lower than that associated with haptic estimation.

## Conditions

- Visual-only
- Haptic-only
- Combined




# Psychophysical Task




- 2-Interval-Forced-Choice  
“Which bar was bigger?”

# SDT: Two-interval forced choice

- “Which bar was bigger?”




# Haptic VR Equipment



- “Phantom” force-feedback device

# Maximum-likelihood Estimation (MLE)



## Predictions

- 1) Weight depends on relative variability

$$\hat{S}_{comb} = \hat{S}_1 w_1 + \hat{S}_2 w_2$$

$$w_1 + w_2 = 1$$

$$w_i = \frac{1}{\sigma_i^2} / \sum_{j=1}^n \frac{1}{\sigma_j^2}$$

- 2) Combined estimate is least variable

$$\sigma_{comb}^2 = \frac{\sigma_1^2 \sigma_2^2}{\sigma_1^2 + \sigma_2^2}$$

# Additional Examples of MLE

- Disparity & texture cues to surface slant (Knill 1998)
- Visual & proprioceptive cues to location (van Beers et al 1999)
- Visual & auditory cues to location (Gharamani et al. 1998)
- Visual & vestibular cue to heading(Butler et al 2010)
- Et cetera...

# Today's topics

- Measuring perception: psychophysics
- Multisensory interactions
  - Cue integration
  - Conflict detection
- Related to textbook sections:
  - Chapter 12.4: Experiments on human subjects
  - Chapter 6.4: Combining sources of information
  - Chapter 8.4: Mismatched motion and vection





**Self-motion Lab**


# Sensory and Motor Signals Mediating Stationarity Perception

Dr. Paul R. MacNeilage


Department of Psychology and Institute for Neuroscience

University of Nevada, Reno

# Vestibular




# Visual




# Stationarity Perception

## Vestibular



=

## Visual



Perception of a stable world

# Stationarity Perception

- Allows vision to be used for self-motion estimation
- Deficits of stationarity perception are debilitating
  - Disorientation, imbalance, vertigo, sickness
- Underlies effective virtual/augmented reality technology

# Outline

- Signal detection model of stationarity perception
- Psychophysical measurement of stationarity perception
- Factors impacting stationarity perception
  - Motor signals
    - Oculomotor signals
    - Cephalomotor signals
  - Visual stimulus parameters
    - Spatial frequency
    - Retinal stimulus location
- Relation between stationarity perception and sickness


# Outline

- Signal detection model of stationarity perception
- Psychophysical measurement of stationarity perception
- Factors impacting stationarity perception
  - Motor signals
    - Oculomotor signals
    - Cephalomotor signals
  - Visual stimulus parameters
    - Spatial frequency
    - Retinal stimulus location
- Relation between stationarity perception and sickness

# Congruence Evaluation


- Are visual and non-visual self-motion signals congruent?

## Vestibular





≠

## Visual



# Crossmodal Discrimination Model

- How does the nervous system evaluate congruence in support of stationarity perception?





- What factors impact self-motion and crossmodal discrimination?



# Outline

- Signal detection model of stationarity perception
- **Psychophysical measurement of stationarity perception**
- Factors impacting stationarity perception
  - Motor signals
    - Oculomotor signals
    - Cephalomotor signals
  - Visual stimulus parameters
    - Spatial frequency
    - Retinal stimulus location
- Relation between stationarity perception and sickness


# Stationarity Perception Paradigm - Analog



“Range of Immobility”

Wallach & Kravitz (1965)


# Stationarity Perception Paradigm - Hybrid



**Fig. 1** Experimental setup. A Puppetworks mechanical head tracker was used to track head position while subjects viewed a virtual sphere in a Virtual V8 head mounted display. The simulation was run by an SGI 02 computer

- Rotation:
  - Roll, pitch, yaw
- Translation:
  - Naso-occipital, interaural, dorsoventral

# Stationarity Perception Paradigm - Digital



# 2AFC Procedure

Align scene- and head-fixed targets


Head movement target appears

Head movement while fixating

Respond "With" or "Against"


WITH ? AGAINST

Button press



# Accuracy and Precision of Stationarity Judgments

- Accuracy
  - PSE – point of subjective equality
  - Visual gain perceived stationary
- Precision
  - JND – just-noticeable difference
  - Range of immobility/stationarity



# Outline

- Signal detection model of stationarity perception
- Psychophysical measurement of stationarity perception
- **Factors impacting stationarity perception**
  - Motor signals
    - Oculomotor signals
    - Cephalomotor signals
  - Visual stimulus parameters
    - Spatial frequency
    - Retinal stimulus location
- Relation between stationarity perception and sickness


# Role of oculomotor signals

Oculomotor




+

Retinal Flow




=

Vestibular



Head-fixed fixation




Scene-fixed fixation






# Role of cephalomotor signals

Passive



Active




# Movement Conditions


- Adaptive staircase
  - 150 trials per condition
- 20 subjects
- Visual scene – dot cloud

Head-Fixed



Active



Passive




Scene-Fixed




# Passive playback of active trajectories


Active





Passive



# Passive playback of active trajectories



# PSEs and JNDs



Motor signals associated with naturalistic eye and head movement facilitate conflict detection.

# Active vs passive head movement

- Reduced gain during passive head movement
  - Underestimation of head movement
- Reduced sensitivity during passive movement
  - Increased noise because motor signals are absent
- Consistent with spatial updating
  - Mergner et al (1998)
  - Genzel et al (2016)

# Scene- vs head-fixed fixation

- Reduced gain during head-fixed fixation
  - Underestimation of head movement (Clemens et al. 2017)
  - Overestimation of visual scene motion (Garzorz et al 2018)
- Reduced sensitivity during head-fixed fixation
  - Increased signal-dependent noise on retinal image motion? (Garzorz & MacNeilage 2017)

# Outline

- Signal detection model of stationarity perception
- Psychophysical measurement of stationarity perception
- **Factors impacting stationarity perception**
  - Motor signals
    - Oculomotor signals
    - Cephalomotor signals
  - **Visual stimulus parameters**
    - Spatial frequency
    - Retinal stimulus location
- Relation between stationarity perception and sickness




# Spatial frequency



- Speed perception increases with spatial frequency (Diener et al 1976)
- Oculomotor responses may also depend on spatial frequency
- How does stationarity perception depend on spatial frequency?

# Conditions

- Adaptive staircase
  - 300 trials per condition
- 19 subjects



# PSEs and JNDs



More accurate and precise with lower spatial frequency.

# Spatial frequency effects

- More accurate with low spatial frequency
  - High frequencies may lead to overestimation of speed
  - Self-motion estimation may favor low spatial frequency channels
- More precise with low spatial frequency
  - Higher spatial frequencies lead to noisier speed estimates
- These effects are most pronounced during head-fixed fixation
  - Retinal speed estimation depends more on spatial frequency than oculomotor


# Outline

- Signal detection model of stationarity perception
- Psychophysical measurement of stationarity perception
- **Factors impacting stationarity perception**
  - Motor signals
    - Oculomotor signals
    - Cephalomotor signals
  - **Visual stimulus parameters**
    - Spatial frequency
    - Retinal stimulus location
- Relation between stationarity perception and sickness


# Visual motion processing varies with eccentricity

- Motion processing varies with retinal eccentricity
- Vection is driven strongly by peripheral stimulation (Brandt et al 1973)
- How is motion integrated across the visual field to drive stationarity perception?

# Conditions




# Scene-Fixed Condition







# Head-Fixed Condition



# PSEs and JNDs



Sensitivity depends on retinal stimulus location; these effects are mediated by fixation.

# Outline

- Signal detection model of stationarity perception
- Psychophysical measurement of stationarity perception
- Factors impacting stationarity perception
  - Motor signals
    - Oculomotor signals
    - Cephalomotor signals
  - Visual stimulus parameters
    - Spatial frequency
    - Retinal stimulus location
- **Relation between stationarity perception and sickness**

# Stationarity perception and sickness

- Individual differences in:
  - Precision and accuracy of stationarity judgments
  - Susceptibility to sickness
- Are these related?

# Measuring sickness

- Simulator sickness questionnaire
  - Nausea
  - Oculomotor discomfort
  - Disorientation
  - Total score
- Discomfort score
  - Average
  - Ending

No \_\_\_\_\_

Date \_\_\_\_\_

## **SIMULATOR SICKNESS QUESTIONNAIRE**

Kennedy, Lane, Berbaum, & Lilienthal (1993)\*\*\*

Instructions : Circle how much each symptom below is affecting you right now.

1. General discomfort	<u>None</u>	<u>Slight</u>	<u>Moderate</u>	<u>Severe</u>
2. Fatigue	<u>None</u>	<u>Slight</u>	<u>Moderate</u>	<u>Severe</u>
3. Headache	<u>None</u>	<u>Slight</u>	<u>Moderate</u>	<u>Severe</u>
4. Eye strain	<u>None</u>	<u>Slight</u>	<u>Moderate</u>	<u>Severe</u>
5. Difficulty focusing	<u>None</u>	<u>Slight</u>	<u>Moderate</u>	<u>Severe</u>
6. Salivation increasing	<u>None</u>	<u>Slight</u>	<u>Moderate</u>	<u>Severe</u>
7. Sweating	<u>None</u>	<u>Slight</u>	<u>Moderate</u>	<u>Severe</u>
8. Nausea	<u>None</u>	<u>Slight</u>	<u>Moderate</u>	<u>Severe</u>
9. Difficulty concentrating	<u>None</u>	<u>Slight</u>	<u>Moderate</u>	<u>Severe</u>
10. « Fullness of the Head »	<u>None</u>	<u>Slight</u>	<u>Moderate</u>	<u>Severe</u>
11. Blurred vision	<u>None</u>	<u>Slight</u>	<u>Moderate</u>	<u>Severe</u>
12. Dizziness with eyes open	<u>None</u>	<u>Slight</u>	<u>Moderate</u>	<u>Severe</u>
13. Dizziness with eyes closed	<u>None</u>	<u>Slight</u>	<u>Moderate</u>	<u>Severe</u>
14. *Vertigo	<u>None</u>	<u>Slight</u>	<u>Moderate</u>	<u>Severe</u>
15. **Stomach awareness	<u>None</u>	<u>Slight</u>	<u>Moderate</u>	<u>Severe</u>
16. Burping	<u>None</u>	<u>Slight</u>	<u>Moderate</u>	<u>Severe</u>

# Stationarity perception and sickness

- Is accuracy (PSE) correlated with sickness?
- Is precision (JND) correlated with sickness?
- Linear mixed effects modeling approach
  - R library lme4

$$SSQScore \sim PSE + JND + Fixation * RSL + Fixation * Frequency + (1|Subject)$$

# Worse Accuracy -> Increased Sickness

- Simulator sickness questionnaire

- ✓ Nausea
- Oculomotor discomfort
- Disorientation
- Total score

- Discomfort score

- Average
- Ending

Sickness predicted by PSE	Coefficient of Fixed Effect	P value
Nausea	-12.13	p = 0.014
Oculomotor	-8.05	p = 0.289
Disorientation	-8.15	p = 0.311
Total Score	-10.39	p = 0.138
Average Discomfort	-0.100	p = 0.757
End Discomfort	-0.615	p = 0.167

# Worse Precision -> Increased Sickness

- Simulator sickness questionnaire

- Nausea
- ✓ Oculomotor discomfort
- ✓ Disorientation
- ✓ Total score

- Discomfort score

- Average
- Ending

Sickness predicted by JND	Coefficient of Fixed Effect	P value
Nausea	12.54	p =0.289
Oculomotor	36.17	p = 0.045
Disorientation	68.42	p < 0.001
Total Score	41.08	p = 0.013
Average Discomfort	0.361	p = 0.631
End Discomfort	-1.67	p = 0.113



# Conclusions and future directions

- Stationarity paradigm allows identifying self-motion factors
  - Integration of visual signals driving visual self-motion
  - Interactions with motor signals
- Future directions
  - Other motion axes/direction
  - Investigate dynamics; how information is integrated over time?  
(Garzroz & MacNeilage 2019)
  - May be modeled in a causal inference framework
  - Physiological correlates

# Acknowledgments

- Savannah Halow



- Research was supported by NIGMS of NIH under grant number P20 GM103650 and NSF CHS under grant 1911041