

2202-MTL101B Minor Exam

Tatsam

TOTAL POINTS

23 / 30

QUESTION 1

1 Linear Independence 2 / 6

+ 0 pts Not attempted

+ 0 pts Completely wrong/deserves no partial marks

+ 6 pts Correct

+ 4 pts Showing that $\{Av_1, Av_2, \dots, Av_n\}$ is LI implies $\{v_1, v_2, \dots, v_n\}$ is LI.

✓ + 2 pts Showing that converse is not true with a counter example.

+ 1 pts For providing the definition of LI precisely and applying it correctly.

QUESTION 2

2 Basis of Polynomial Space 6 / 6

✓ - 0 pts Correct [$\{x, x^3, 4 - 5x^2 + x^4\}$ is a basis, and Dim 3]

- 1 pts Conclude the basis correctly but did not mention the dimension/ basis is not written in polynomial form/or some minor mistake

- 2 pts Conclude dimension is 3 with the correct procedure but the basis is not correct.

- 4 pts Only conclude that Dimension is 3 but the procedure is not correct/ or procedure is correct but get dimension NOT 3

- 6 pts Wrong Solution

QUESTION 3

3 Basis of Matrix Space 6 / 6

✓ - 0 pts Correct

- 6 pts Unattempted/completely incorrect

- 2 pts In Part (i) did not find the dimension or completely incorrect

- 2 pts Part (ii) completely incorrect or not attempted

- 1 pts Part (ii) partially correct

- 2 pts Part (i) did not show that it is a subspace

- 1 pts In part (i) some mistakes in finding the dimension/Incorrect argument/argument missing

- 1 pts In part (i) incomplete argument to show the subspace

QUESTION 4

4 Solution Space 3 / 6

- 0 pts Correct

- 6 pts Not correct/RRE corresponding to a matrix is unique/not attempted

- 4 pts This just a particular case of 2×2 Matrix/partial marks awarded/second argument is not correct/

✓ - 3 pts Partially correct/ arguments missing (not correct) if A is not invertible/last part is not clear

- 5 pts Major arguments missing

- 1 pts minor arguments missing

QUESTION 5

5 Row Reduced Echelon Form 6 / 6

✓ + 6 pts *Correct*

+ 5 pts RRE form is correct and Rank is 1 but
there is some minor mistake

+ 2 pts Rank is 1 & proof is correct

+ 1 pts Tried but did not converge anywhere

+ 2 pts RRE is not correct but is very close to the
correct one and contains (0,1,0)

+ 0 pts Either totally wrong or not attempted

Prob 1) $M_{n \times n}(\mathbb{R})$.

If $\{AV_1, AV_2, \dots, AV_n\}$ is linearly ind. in $M_{n \times 1}(\mathbb{R})$

this means $\alpha_1 AV_1 + \alpha_2 AV_2 + \alpha_3 AV_3 + \dots + \alpha_n AV_n = 0$
is true if and only if
 $\alpha_1 = \alpha_2 = \alpha_3 = \dots = \alpha_n = 0$

assume $\alpha_1 A = \beta_1, \alpha_2 A = \beta_2, \dots, \alpha_n A = \beta_n$

$\beta_1 V_1 + \beta_2 V_2 + \dots + \beta_n V_n = 0 \quad \text{--- (1)}$
holds true for $\alpha_i A$

$\left. \begin{array}{l} \alpha_1 A \Rightarrow (0)A = 0 = \beta_1 \\ \alpha_2 A \Rightarrow (0)A = 0 = \beta_2 \\ \alpha_n A \Rightarrow (0)A = 0 = \beta_n \end{array} \right\}$ this is an only if
statement.

$\therefore (1)$ holds true if and only if
 $\beta_1 = \beta_2 = \beta_3 = \dots = \beta_n = 0$

$\therefore \{V_1, V_2, \dots, V_n\}$ are linearly independent.

Converse:

given $\alpha_1 V_1 + \alpha_2 V_2 + \dots + \alpha_n V_n = 0$
holds true if and only if $\alpha_1 = \alpha_2 = \dots = \alpha_n = 0$

in order to show $\{AV_1, AV_2, \dots, AV_n\}$ to
be LI we must show

$\alpha_1 AV_1 + \alpha_2 AV_2 + \dots + \alpha_n AV_n = 0$ holds if and only
if $\alpha_1 = \alpha_2 = \dots = \alpha_n = 0$

In order to disprove, we can take any contradicting example.

Say $n=2$

$$A = \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} \quad v_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad v_2 = \begin{pmatrix} 4 \\ 0 \end{pmatrix}$$

given : v_1, v_2 are L.I which true
 $\therefore \alpha v_1 \neq v_2$ for any α .

$$\begin{pmatrix} \alpha \\ \alpha \end{pmatrix} \neq \begin{pmatrix} 4 \\ 0 \end{pmatrix} \Rightarrow \begin{cases} \alpha = 4 \\ \alpha = 0 \end{cases} \text{ not possible.}$$

Now, $AV_1 = \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$

$$AV_2 = \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 4 \\ 0 \end{pmatrix} = \begin{pmatrix} 4 \\ 4 \end{pmatrix}$$

we can easily see $AV_2 = 4(AV_1)$
 which means $\{AV_1, AV_2\}$ are not linearly independent as $\exists \alpha, \beta$ both of which are not 0, such that

$$\alpha AV_1 + \beta AV_2 = 0$$

$$\alpha = 4, \beta = -1$$

Hence the converse is not true.

1 Linear Independence 2 / 6

- + 0 pts Not attempted
- + 0 pts Completely wrong/deserves no partial marks
- + 6 pts Correct
- + 4 pts Showing that $\{Av_1, Av_2, \dots, Av_n\}$ is LI implies $\{v_1, v_2, \dots, v_n\}$ is LI.
- ✓ + 2 pts *Showing that converse is not true with a counter example.*
- + 1 pts For providing the definition of LI precisely and applying it correctly.



भारतीय प्रौद्योगिकी संस्थान दिल्ली
INDIAN INSTITUTE OF TECHNOLOGY DELHI

अनुक्रमांक
Entry No.

2	0	2	2	M	T	6	1	9	6	9
---	---	---	---	---	---	---	---	---	---	---

पाठ सं.
Course No.

M	T	L	1	0	1
---	---	---	---	---	---

ग्रुप संख्या
Group No.

अनुवर्ती पुस्तिका संख्या
CONTINUATION BOOK NO.

1

दिनांक
Date

18	04	2023
----	----	------

Problem 2)

$$p(x) = a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4$$

this vector space is governed by : 5
variables a_0, a_1, a_2, a_3, a_4 .
basis of V : $\{1, x, x^2, x^3, x^4\}$

$$\text{Now, } W = \{p(x) \in V \mid p(1) = -p(-1) \text{ & } p(2) = -p(-2)\}$$

$$p(x) = -p(-1)$$

$$\Rightarrow a_0 + a_1 + a_2 + a_3 + a_4 = a_0 - a_1 + a_2 - a_3 + a_4$$

$$\begin{cases} 2a_1 = -2a_3 \\ a_1 = -a_3 \end{cases} \quad \text{--- (1)}$$

$$p(2) = -p(-2)$$

$$a_0 + 2a_1 + 4a_2 + 8a_3 + 16a_4 = a_0 - 2a_1 + 4a_2 - 8a_3 + 16a_4$$

$$\begin{cases} 4a_1 = -16a_3 \\ a_1 = -4a_3 \end{cases} \quad \text{--- (2)}$$

2

~~Eqn ① and ② give us that~~

~~$a_1 = a_3 = 0$.~~

~~and there is no restriction on a_0, a_2, a_4~~

~~∴ W contains $p(x)$ of the form~~

$$a_0 + a_2 x^2 + a_4 x^4$$

~~if~~

$$p(1) = -p(-1)$$

$$a_0 + a_1 + a_2 + a_3 + a_4 = -a_0 + a_1 - a_2 + a_3 - a_4$$

$$\boxed{a_0 + a_2 + a_4 = 0} \quad \text{--- (1)}$$

$$p(2) = -p(-2)$$

$$a_0 + 2a_1 + 4a_2 + 8a_3 + 16a_4 = -a_0 + 2a_1 - 4a_2 + 8a_3 - 16a_4$$

$$\boxed{a_0 + 4a_2 + 16a_4 = 0} \quad \text{--- (2)}$$

take $a_0 = \lambda_0$ ~~$a_1 = a_3 = 0$~~

$$\text{Eqn (1)} \Rightarrow a_2 + a_4 = -\lambda_0 \quad \text{--- (3)}$$

$$\text{Eq } \textcircled{2} \quad 4a_2 + 16a_4 = -\lambda_0 \quad \textcircled{4}$$

$$\textcircled{4} - 4 \times \textcircled{3}$$

$$12a_4 = 3\lambda_0 \\ a_4 = \frac{1}{4}\lambda_0$$

$$a_2 = -\lambda_0 - a_4 = -\frac{5}{4}\lambda_0$$

There are no restrictions on a_1 & a_3

Let them be $\lambda_1 = a_1$ and $\lambda_2 = a_3$

$$\left(\begin{array}{l} a_0 = \lambda_0 \\ a_1 = \lambda_1 \\ a_2 = -\frac{5}{4}\lambda_0 \\ a_3 = \lambda_2 \\ a_4 = \frac{1}{4}\lambda_0 \end{array} \right)$$

These are the constraints.

$\therefore W$ has the $p(n)$ of the form

$$p(n) = \lambda_0 + \lambda_1 x + \left(-\frac{5}{2}\lambda_0\right)x^2 + \lambda_2 x^3 + \left(\frac{\lambda_0}{4}\right)x^4$$

with 3 free variables λ_0 , λ_1 & λ_2 .

$$\therefore \text{basis of } W = \left\{ x, x^3, \left(1 - \frac{5x^2}{2} + \frac{x^4}{4}\right) \right\}$$

$$\boxed{\text{dimension} = 3}$$

2 Basis of Polynomial Space 6 / 6

✓ - 0 pts Correct [$\{x, x^3, 4 - 5x^2 + x^4\}$ is a basis, and Dim 3]

- 1 pts Conclude the basis correctly but did not mention the dimension/ basis is not written in polynomial form/or some minor mistake

- 2 pts Conclude dimension is 3 with the correct procedure but the basis is not correct.

- 4 pts Only conclude that Dimension is 3 but the procedure is not correct/ or procedure is correct but get dimension NOT 3

- 6 pts Wrong Solution

Problem 3:

$$A = \begin{pmatrix} C_1 & C_2 \\ C_3 & C_4 \end{pmatrix}$$

$$\bar{A}^T = \begin{pmatrix} \bar{C}_1 & \bar{C}_3 \\ \bar{C}_2 & \bar{C}_4 \end{pmatrix}$$

$$A = \bar{A}^T \Rightarrow A : C_1 = \bar{C}_1$$

$$C_2 = \bar{C}_3$$

$$\bar{C}_2 = C_3$$

$$C_4 = \bar{C}_4$$

∴ A must be of the form $\begin{pmatrix} \alpha & \gamma + i\delta \\ \gamma - i\delta & \beta \end{pmatrix}$

where $\alpha, \beta, \gamma, \delta \in \mathbb{R}$

(i) if we take $\alpha = \beta = \gamma = \delta = 0$ $A = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$

∴ Additive identity $0 \in S$

- Now, if we prove
- a. commutative $(A_1 + A_2 = A_2 + A_1)$
 - b. associative $(A_1 + A_2) + A_3 = A_1 + (A_2 + A_3)$
 $(\alpha\beta)A_1 = \alpha(\beta A_1)$
 - c. distributive $\alpha(A_1 + A_2) = \alpha A_1 + \alpha A_2$
 $(\alpha + \beta)A_1 = \alpha A_1 + \beta A_1$
 - d. closure over + and *
 - e. additive inverse
 - f. mult. identity

then S becomes a vector space.

(a), (b), (c), (d) hold true as they are properties of $F = \mathbb{R}$ and $a_{ij} \in \mathbb{C}$.

e. take $\alpha' = -\alpha$, $\beta' = -\beta$, $\gamma' = -\gamma$, $\delta' = \delta'$
we get A'
s.t. $A' + A = 0$

f. take $\gamma = \delta = 0$ and $\alpha = \beta = 1$

$A_1 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ represents mult. identity.

dimension of S is 4 as we need 4 different variables to span S .

basis: $\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}$

(ii) No. let's check closure under

$A = \begin{pmatrix} 1 & i \\ -i & 1 \end{pmatrix}$ (taking any random example to disprove)

$$\bar{A}^T = \begin{pmatrix} 1 & i \\ -i & 1 \end{pmatrix}$$

$\therefore \bar{A} \in S$

take $c_1 = i \in \mathbb{C}$

then $c_1 A$ must also belong to S .

$$B = C_1 A = \begin{pmatrix} i & -1 \\ 1 & i \end{pmatrix}$$

$$\bar{B}^T = \begin{pmatrix} i & 1 \\ -1 & i \end{pmatrix}$$

$$B \neq \bar{B}^T \quad \therefore B \notin S$$

Hence S is not a vector space over field C .

Problem 4)

$$AX = 0$$

in order to solve this, we try to reduce A in the row reduced echelon form by a sequence of elementary operations. $(A : 0) \rightarrow (A' : 0)$ \hookrightarrow RRE form.

$$\text{We know } f(A) = f(I)A$$

where f is any elementary row op.

$$\begin{aligned} \text{Also } f(AB) &= f(I)(AB) \\ &= ((f(I)(A))(B)) \\ &= f(A)B \end{aligned}$$

$$\text{Now, } f(A^2) = f(A)A$$

3 Basis of Matrix Space 6 / 6

✓ - 0 pts Correct

- 6 pts Unattempted/completely incorrect
- 2 pts In Part (i) did not find the dimension or completely incorrect
- 2 pts Part (ii) completely incorrect or not attempted
- 1 pts Part (ii) partially correct
- 2 pts Part (i) did not show that it is a subspace
- 1 pts In part (i) some mistakes in finding the dimension/Incorrect argument/argument missing
- 1 pts In part (i) incomplete argument to show the subspace

$$B = C_1 A = \begin{pmatrix} i & -1 \\ 1 & i \end{pmatrix}$$

$$\bar{B}^T = \begin{pmatrix} i & 1 \\ -1 & i \end{pmatrix}$$

$$B \neq \bar{B}^T \quad \therefore B \notin S$$

Hence S is not a vector space over field C .

Problem 4)

$$AX = 0$$

in order to solve this, we try to reduce A in the row reduced echelon form by a sequence of elementary operations. $(A : 0) \rightarrow (A' : 0)$ \hookrightarrow RRE form.

$$\text{We know } f(A) = f(I)A$$

where f is any elementary row op.

$$\begin{aligned} \text{Also } f(AB) &= f(I)(AB) \\ &= ((f(I)(A))(B)) \\ &= f(A)B \end{aligned}$$

$$\text{Now, } f(A^2) = f(A)A$$

To show both of them have the same solution space, we need to show all the solutions of $AX = 0$ also satisfy $A^2X = 0$ and vice versa.

$$A^2X = 0 \Rightarrow A(AX) = 0 \Rightarrow A(0) = 0$$

∴ All solⁿ of $AX = 0$ satisfy $A^2X = 0$.

We need to prove the converse now.

case (i) $\text{rank}(A) = n \Rightarrow |A| \neq 0 \Rightarrow A^{-1}$ exists.

$$A^2X = 0$$

$$A^{-1}A \cdot AX = A^{-1} \cdot 0$$

$AX = 0$ satisfies..

case (ii) $\text{rank}(A) < n = m$

given $\text{rank}(A^2) = m$.

This means A and A^2 have the same no. of non-zero rows in the RRE form.

While reducing A^2 into its RRE form, let us first use the same sequence of row operations that converted A into its RRE form.

$$P_1 P_2 P_3 \dots P_k (A) \Rightarrow \text{RRE} = \text{say } (A_R)$$

$$\text{Now } P(A^2) = P(A) A \quad \{ \text{proved above} \}$$

$$P_1 P_2 P_3 \dots P_k (A^2) = A_R A$$

Now, since we are given $\text{rank}(A) = \text{rank}(A^2)$

After calculating $A_R A$ and again reducing to RRE form, we are sure to arrive at the same A_R !

The zero rows in A_R would reduce the rows of A to 0 which became 0 while calculating A_R , and the other rows would return the non zero rows as it is.

$$\therefore \text{RRE of } A = \text{RRE of } A^2$$

Hence the solution space of $AX = 0$ and $A^2 X = 0$ is exactly the same.

4 Solution Space 3 / 6

- **0 pts** Correct
- **6 pts** Not correct/RRE corresponding to a matrix is unique/not attempted
- **4 pts** This just a particular case of $\text{Matrix} \backslash \text{partial marks awarded}$ /second argument is not correct/
- ✓ **- 3 pts** *Partially correct/ arguments missing (not correct) if A is not invertible/last part is not clear*
- **5 pts** Major arguments missing
- **1 pts** minor arguments missing

Problem 5) $A X = b$

$$X = \begin{pmatrix} -2 + \alpha \\ 2 \\ \alpha + \beta \end{pmatrix}$$

As we can see the no. of unknowns were 3
and there exist 00 soln $\therefore \text{rank}(A) =$
 $\text{rank}(A|b) < 3$

Let's take variables as x_1, x_2, x_3

$$x_1 = \alpha - 2$$

\therefore this true for $\forall \alpha \in \mathbb{R}$

$$\alpha - 2 = t \quad t \in \mathbb{R}$$

$$x_2 = t$$

$$x_2 = 2$$

$x_3 = \alpha + \beta = t + \beta + 2$, again this takes all \mathbb{R} values for diff β . $\therefore t + \beta + 2 = w \in \mathbb{R}$

$$x_3 = w$$

So, basically the solution set is $(t, 2, w)$
 $t, w \in \mathbb{R}$.

which means x_1 , and x_3 were free variables and ~~part of basis~~ ~~both~~ a leading also $x_2 = 2$ fixed.

Hence, we are sure that the RRE form of $(A|b)$ must look something like :

$$\left(\begin{array}{ccc|c} 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right)$$

given $b = \begin{pmatrix} 4 \\ 2 \\ 6 \end{pmatrix}$, we can estimate A
 could be $\begin{pmatrix} 0 & 2 & 0 & 4 \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 0 \end{pmatrix}$

{ applying $R_2 \rightarrow R_2 + R_1$, $R_1 \rightarrow 2R_1$ } \rightarrow

RRE form of $A = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$

rank(A) = no. of non zero rows = 1

5 Row Reduced Echelon Form 6 / 6

✓ + 6 pts Correct

+ 5 pts RRE form is correct and Rank is 1 but there is some minor mistake

+ 2 pts Rank is 1 & proof is correct

+ 1 pts Tried but did not converge anywhere

+ 2 pts RRE is not correct but is very close to the correct one and contains (0,1,0)

+ 0 pts Either totally wrong or not attempted



भारतीय प्रौद्योगिकी संस्थान दिल्ली
INDIAN INSTITUTE OF TECHNOLOGY DELHI

लघु परीक्षा उत्तर पुस्तिका
MINOR TEST ANSWER BOOK

नाम

Name TATSAM RANJAN SHARMA

अनुक्रमांक

Entry No.

2	0	2	2	M	T	6	1	9	6	9
---	---	---	---	---	---	---	---	---	---	---

पाठ्यक्रम सं.

Course No.

M	T	L	1	0	1	ग्रुप संख्या	1	1
---	---	---	---	---	---	--------------	---	---

ग्रुप संख्या

Group No.

पाठ्यक्रम शीर्षक

Course Title LINEAR ALGEBRA & DIFFERENTIAL EQUATIONS

लघु परीक्षा सं.

Minor Test No.

1

दिनांक
Date

1	8	0	4	2	0	2	3
---	---	---	---	---	---	---	---

प्रयोग किए गए अनुवर्ती पृष्ठों की संख्या

No. of continuation sheets used

1

प्रश्न सं. Q.No.	प्राप्त अंक Marks
1.	
2.	
3.	
4.	
5.	
6.	
7.	
8.	
9.	
10.	
कुल TOTAL	

पाठ्यक्रम निर्धारक के हस्ताक्षर और दिनांक

Signature of Course Co-ordinator and date

अनुचित साधनों का प्रयोग करने वाले छात्रों को निलम्बित/निष्कासित किया जा सकता है।

Students using unfair means are liable to be punished by Suspension/Expulsion.

परीक्षा केन्द्र में सेलफोन, काम्युनिकेटर्स व पीडीए साधनों का प्रयोग करना सख्त मना है।

Use of cell-phones, Communications & PDAs in the Examination Hall is strictly prohibited.

सभी पृष्ठों पर लिखें | Write on all pages.