

Question-1

- Using RRE determine values of a and b for which A is invertible for $b \neq -1$ find the inverse.

$$\left(\begin{array}{cccc|cccc} 1 & 0 & 1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & a & 1 & 1 & 0 & 0 & 1 & 0 \\ 2 & 0 & 0 & b & 0 & 0 & 0 & 1 \end{array} \right)$$

$$R_4 \rightarrow R_4 - 2R_1, R_3 \rightarrow R_3 - aR_2$$

$$\left(\begin{array}{cccc|cccc} 1 & 0 & 1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & -a & 1 & 0 \\ 0 & 0 & -2 & b & -2 & 0 & 0 & 1 \end{array} \right)$$

$$R_1 \rightarrow R_1 - R_3$$

$$R_4 \rightarrow R_4 + 2R_3$$

$$\left(\begin{array}{cccc|cccc} 1 & 0 & 0 & -1 & 1 & a & -1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & -a & 1 & 0 \\ 0 & 0 & 0 & b+2 & -2 & -2a & 2 & 1 \end{array} \right)$$

From above we can conclude that A is invertible for $a \in \mathbb{R}$ & $b \in \mathbb{R} - \{-2\}$.

To compute inverse for $b \neq -1$.

Consider, $b \neq -1$

$$\left(\begin{array}{cccc|cccc} 1 & 0 & 0 & -1 & 1 & a & -1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & -a & 1 & 0 \\ 0 & 0 & 0 & 1 & -2 & -2a & 2 & 1 \end{array} \right)$$

$$R_1 \rightarrow R_1 + R_4$$

$$R_3 \rightarrow R_3 - R_4$$

$$\left(\begin{array}{cccc|cccc} 1 & 0 & 0 & 0 & -1 & -a & 1 & 1 \\ 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 2 & a & -1 & -1 \\ 0 & 0 & 0 & 1 & -2 & -2a & 2 & 1 \end{array} \right)$$

$$\text{Hence } A^{-1} = \begin{pmatrix} -1 & -a & 1 & 1 \\ 0 & 1 & 0 & 0 \\ 2 & a & -1 & -1 \\ -2 & -2a & 2 & 1 \end{pmatrix}$$

Q 2.

$$x + y + z = 3$$

$$ax + 5y + 4z = a$$

$$3x + (a^2 - 8)z = 12$$

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 2 & 5 & 4 & a \\ 3 & 0 & a^2 - 8 & 12 \end{array} \right]$$

$$R_2 \rightarrow R_2 - 2R_1 \quad \text{and} \quad R_3 \rightarrow R_3 - 3R_1$$

$$\sim \left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 0 & 3 & 2 & a-6 \\ 0 & -3 & a^2-11 & 3 \end{array} \right]$$

$$R_3 \rightarrow R_3 + R_2$$

$$\sim \left[\begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 0 & 3 & 2 & a-6 \\ 0 & 0 & a^2-9 & a-3 \end{array} \right]$$

(i) for NO solution

$$\text{Rank}(A) < \text{Rank}(A|b)$$

$$\therefore a^2 - 9 = 0 \quad \text{and} \quad a - 3 \neq 0$$

$$\Rightarrow a = \pm 3 \quad \text{and} \quad a \neq 3$$

$$\therefore a = -3$$

(ii) for Unique solution

$$\text{Rank}(A) = 3$$

$$\therefore a^2 - 9 \neq 0$$

$$\Rightarrow a \neq 3, -3$$

(iii) for Infinitely many solutions

$$\text{Rank}(A) = \text{Rank}(A|b) < 3$$

$$\therefore a^2 - 9 = 0 \quad \text{and} \quad a - 3 = 0$$

$$\Rightarrow a = \pm 3 \quad \text{and} \quad a = 3$$

$$\therefore a = 3$$

Solution 3 :-

(a) let $\alpha, \beta, \gamma \in \mathbb{F}$ then we need to prove that

$$\alpha(v_1 + v_3) + \beta(3v_1 + 2v_2 + v_3) + \gamma(2v_1 + 3v_2 + v_3) = 0 \quad \text{--- (I)}$$
$$\Rightarrow \alpha, \beta, \gamma = 0.$$

On solving (I)

$$\alpha v_1 + \alpha v_3 + 3\beta v_1 + 2\beta v_2 + \beta v_3 + 2\gamma v_1 + 3\gamma v_2 + \gamma v_3 = 0$$
$$\Rightarrow (\alpha + 3\beta + 2\gamma)v_1 + (2\beta + 3\gamma)v_2 + (\alpha + \beta + \gamma)v_3 = 0$$

Since $\{v_1, v_2, v_3\}$ is L.I., the coefficients of v_1, v_2, v_3 in above equations has to be zero.

Thus, we get the system of linear equations on equating the coefficients to zero as :

$$\alpha + 3\beta + 2\gamma = 0 \quad \text{--- (i)}$$

$$2\beta + 3\gamma = 0 \quad \text{--- (ii)}$$

$$\alpha + \beta + \gamma = 0 \quad \text{--- (iii)}$$

$$(i) - (iii), \quad 2\beta + \gamma = 0 \quad \text{--- (iv)}$$

$$(ii) - (iv), \quad 2\gamma = 0 \Rightarrow \gamma = 0$$

$$\text{From (iv),} \quad 2\beta + \gamma = 0 \Rightarrow 2\beta + 0 = 0 \Rightarrow \beta = 0$$

$$\text{From (i),} \quad \alpha + 3\beta + 2\gamma = 0 \Rightarrow \alpha + 0 + 0 = 0 \Rightarrow \alpha = 0$$

$$\therefore \alpha, \beta, \gamma = 0$$

$$\Rightarrow \{v_1 + v_3, 3v_1 + 2v_2 + v_3, 2v_1 + 3v_2 + v_3\} \text{ is L.I.}$$

(b) Let $v = (\frac{1}{2}, \frac{1}{3}, \frac{1}{4}, 1)$.

$v \in W$ since $v_4 = 1 \in \mathbb{Z}$

but for $\alpha = 1/2$, $\alpha v \notin W$ since

$$\alpha v = (\frac{1}{4}, \frac{1}{6}, \frac{1}{8}, \frac{1}{2})$$

$$\alpha v_4 = \frac{1}{2} \notin \mathbb{Z}$$

$\therefore W$ is not a subspace since scalar multiplication doesn't hold.