

(1)  
White

$$\begin{pmatrix} 1 & 0 & \alpha \\ 0 & 1 & \beta \\ 0 & 0 & 0 \end{pmatrix}$$

$$\alpha, \beta \in \mathbb{R}$$

$$\begin{pmatrix} 1 & \mu & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix},$$

$$\mu \in \mathbb{R}$$

$$\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

---

(5)  
Green

$$\begin{pmatrix} 1 & 0 & \alpha \\ 0 & 1 & \beta \\ 0 & 0 & 0 \end{pmatrix}$$

$$\alpha, \beta \in \mathbb{R}$$

$$\begin{pmatrix} 1 & \mu & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix},$$

$$\mu \in \mathbb{R}$$

$$\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

White

2. (a) Since  $AOB = O = BOA$ ,  $O \in W$   
( $O$  denotes the zero matrix of size  $n \times n$ )

$$\therefore W \neq \emptyset$$

If  $X, Y \in W$  and  $\lambda \in \mathbb{R}$ , then  
 $AXB = BXA$  and  $AYB = BYA$

$$\therefore A(X + \lambda Y)B = AXB + \lambda AYB \\ = BXA + \lambda BYA = B(X + \lambda Y)A$$

$$\Rightarrow X + \lambda Y \in W$$

Hence,  $W$  is a subspace.

(b) FALSE: Take  $V = \mathbb{R}$ ,  $A = \{1\}$ ,  $B = \{2\}$   
Then  $\text{span}(A) = \mathbb{R}$ ,  $\text{span}(B) = \mathbb{R}$   
But  $\text{span}(A \cap B) = \{0\}$ .

(c) FALSE:  $(u - 2v) + (2v - w) + (w - u) = 0$   
 $\Rightarrow \{u - 2v, 2v - w, w - u\}$  is linearly dependent.

White

$$3. (a) \quad (x, y, z, w) \in W_1 \cap W_2 \Leftrightarrow \begin{cases} x+y+z+w=0 \\ 2x+3y+4z+5w=0 \\ 4x+3y+2z+w=0 \\ x+2y+2z+w=0 \end{cases}$$

$$\begin{pmatrix} 1 & 1 & 1 & 1 \\ 2 & 3 & 4 & 5 \\ 4 & 3 & 2 & 1 \\ 1 & 2 & 2 & 1 \end{pmatrix} \xrightarrow{\substack{R_2 \rightarrow R_2 - 2R_1 \\ R_3 \rightarrow R_3 - 4R_1 \\ R_4 \rightarrow R_4 - R_1}} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 2 & 3 \\ 0 & -1 & -2 & -3 \\ 0 & 1 & 1 & 0 \end{pmatrix}$$

$$\xrightarrow{\substack{R_1 \rightarrow R_1 - R_4 \\ R_3 \rightarrow R_3 + R_2}} \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 2 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 2 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\xrightarrow{R_3 \rightarrow R_3 - R_2} \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix} \xrightarrow{R_2 \rightarrow R_2 - R_3} \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -3 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\text{rank}(A) = 3 \Rightarrow \dim(\text{soln. space}) = 4 - 3 = 1$$

$$W_1 \cap W_2 = \{ (-\lambda, 3\lambda, -3\lambda, \lambda) : \lambda \in \mathbb{R} \}$$

$$= \text{span} \{ (-1, 3, -3, 1) \}$$

$$\therefore \text{A Basis for } W_1 \cap W_2 \text{ is } \{ (-1, 3, -3, 1) \}$$

$$3. (b) \quad S = \{ (x, y, z) \in \mathbb{R}^3 : x+y+z=1, x+2y+3z=2 \}$$

is a straight line in  $\mathbb{R}^3$  not passing through (0,0,0).

So, any two points on  $S$  spans  $\text{span}(S)$ .

$$\left( \begin{array}{ccc|c} 1 & 1 & 1 & 1 \\ 1 & 2 & 3 & 2 \end{array} \right) \sim \left( \begin{array}{ccc|c} 1 & 1 & 1 & 1 \\ 0 & 1 & 2 & 1 \end{array} \right) \sim \left( \begin{array}{ccc|c} 1 & 0 & -1 & 0 \\ 0 & 1 & 2 & 1 \end{array} \right)$$

$$S = \{ (\lambda, 1-2\lambda, \lambda) : \lambda \in \mathbb{R} \}$$

Putting  $\lambda=0$  &  $\lambda=1$ , we get  $(0, 1, 0), (1, -1, 1) \in S$ .

$\therefore \{ (0, 1, 0), (1, -1, 1) \}$  is a basis for  $\text{span}(S)$ .



White

$$(4) \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ a & b & c & d \end{pmatrix} \sim \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & -1 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ a & b & c & d \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ a & b & c & d \end{pmatrix}$$

$$\sim \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ a & b & c & d \end{pmatrix} \xrightarrow{R_4 \rightarrow R_4 - aR_1 - bR_2 - cR_3} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & d-b \end{pmatrix}$$

Rank of the matrix is 4 if  $d \neq b$  and 3 if  $d = b$ .

For linearly dependent the rank must be less than 4.

$\therefore$  The given set is L.D. iff  $\boxed{b-d=0}$

White

5. (a)  $S_a = \{ \sin 3x, \sin ax \}$

If  $a=0$ ,  $S_a$  is L.D. because  $0 \in S_a$ .

Assume  $a \neq 0$ .

$S_a$  is L.D.  $\Leftrightarrow \sin ax = \lambda \sin 3x \quad \forall x \in \mathbb{R}$   
for some  $\lambda \in \mathbb{R}$ .

Now  $\sin ax = \lambda \sin 3x \Rightarrow a \cos ax = 3\lambda \cos 3x$   
 $\Rightarrow -a^2 \sin ax = -9\lambda \sin 3x$   
 $\Rightarrow a^2 \sin ax = 9\lambda \sin 3x$   
 $= 9 \sin ax$   
 $\Rightarrow (a^2 - 9) \sin ax = 0 \quad \forall x \in \mathbb{R}.$

Clearly if  $a = \pm 3$ ,  $S_a$  is L.D.  $\Rightarrow a^2 = 9 \quad (\because a \neq 0)$

$\therefore S_a$  is L.I. iff  $a \in \mathbb{R} \setminus \{0, 3, -3\}$ .

(b)  $\mathcal{B} = \{ e^{2x}, \sin 3x, \cos 3x \}$ ,  $W = \text{span}(\mathcal{B})$

$f(x) = e^{2x} + \sin 3x - \cos 3x$

(i)  $f'(x) = 2e^{2x} + 3\cos 3x + 3\sin 3x \in \text{span}(\mathcal{B}) = W$

(ii)  $[f'(x)]_{\mathcal{B}} = \begin{bmatrix} 2 \\ 3 \\ 3 \end{bmatrix}$